

Raina Sabharwal NetID: rs1136

Yunqi Shen NetID: ys478

Question 1

Question 2

a) The decision tree in Figure 4 correctly categorizes the provided examples.

b)  $Gain(A) = I(\frac{p}{p+n}, \frac{n}{p+n}) - \sum_{i=1}^n \frac{p_i+n_i}{p+n} I(\frac{p_i}{p_i+n_i}, \frac{n_i}{p_i+n_i})$

$$Gain(GPA) = 1 - [\frac{3}{12}I(1, 0) + \frac{5}{12}I(\frac{3}{5}, \frac{2}{5}) + \frac{4}{12}I(0, 1)] \approx 0.595$$

$$Gain(University) = 1 - [\frac{5}{12}I(\frac{3}{5}, \frac{2}{5}) + \frac{3}{12}I(\frac{2}{3}, \frac{1}{3}) + \frac{4}{12}I(\frac{1}{4}, \frac{3}{4})] \approx 0.095$$

$$Gain(Published) = 1 - [\frac{5}{12}I(\frac{3}{5}, \frac{2}{5}) + \frac{7}{12}I(\frac{3}{7}, \frac{4}{7})] \approx 0.021$$

$$Gain(Recommendation) = 1 - [\frac{8}{12}I(\frac{5}{8}, \frac{3}{8}) + \frac{4}{12}I(\frac{1}{4}, \frac{3}{4})] \approx 0.093$$

1st level: GPA because the information gain is highest

$$Gain(University) = I(\frac{3}{5}, \frac{2}{5}) - [\frac{2}{5}I(\frac{1}{2}, \frac{1}{2}) + \frac{1}{5}I(1, 0) + \frac{2}{5}I(\frac{1}{2}, \frac{1}{2})] \approx 0.171$$

$$Gain(Published) = I(\frac{3}{5}, \frac{2}{5}) - [\frac{2}{5}I(1, 0) + \frac{3}{5}I(\frac{1}{3}, \frac{2}{3})] \approx 0.420$$


$$Gain(Recommendation) = I(\frac{3}{5}, \frac{2}{5}) - [I(\frac{3}{5}, \frac{2}{5})] = 0$$

2nd level: Published because the information gain is highest

$$Gain(University) = I(\frac{1}{3}, \frac{2}{3}) - [\frac{1}{3}I(0, 1) + \frac{1}{3}I(1, 0) + \frac{1}{3}I(0, 1)] \approx 0.918$$

$$Gain(Recommendation) = I(\frac{1}{3}, \frac{2}{3}) - [I(\frac{1}{3}, \frac{2}{3})] = 0$$


3rd level: University because the information gain is highest



The tree is :

c) The trees are equivalent. This is a coincidence because we were provided with the best decision tree. This data set, to me, did not have an obvious decision tree simply by looking at the table so information gains must be calculated.

### Question 3



a)

b) The decision boundary lies on  $y = -x + 4$ . We have a positive support vector at  $(0, 2)$  with line equation  $y = -x + 2$  and a negative support vector at  $(1, 5)$  with line equation  $y = -x + 6$

So  $w = (-\frac{1}{2}, -\frac{1}{2})$  and  $b = 2$

c) The linear SVM is the same as in part b.  $w = (-\frac{1}{2}, -\frac{1}{2})$  and  $b = 2$


### Question 4

a)  $w_0(0) = 0.2, w_1(0) = 1, w_2(0) = -1$

$$w_1 i_1 + w_2 i_2 + w_0 = 0$$

$$i_1 - i_2 + 0.2 = 0$$

Giving us the following initial decision boundary:



4 samples are misclassified

$$i(n) = i(0) = [1, 0.93, 0.45]^T \text{ and } d(0) = -1$$


$$w(n) = w(0) = [0.2, 1, -1]^T$$

$$y(n) = y(0) = \text{sgn}(w^T(0)i(0)) = \text{sgn}(0.2 + 0.93 - 0.45) = \text{sgn}(0.68) = +1 \neq d(0)$$

$$w(n+1) = w(n) + \eta[d(n) - y(n)]i(n)$$

$$y(1) = 1$$

$$w(1) = [0.2, 1, -1]^T + [-1 - 1][1, 0.93, 0.45]^T = [0.2, 1, -1]^T + [-2, -1.86, -0.9]^T = [-1.8, -0.86, -1.9]^T$$




$$i(n) = i(1) = [1, 0.08, 0.73]^T \text{ and } d(0) = 1$$

$$w(n) = w(1) = [-1.8, -0.86, -1.9]^T$$

$$y(n) = y(1) = \text{sgn}(-1.8 - 0.0688 - 1.378) = -1 \neq d(0)$$

$$w(2) = [-1.8, -0.86, -1.9]^T + [1+1][1, 0.08, 0.73]^T = [-1.8, -0.86, -1.9]^T + [2, 0.16, 1.46]^T = [0.2, -0.7, -0.44]^T$$




$$i(n) = i(2) = [1, 0.28, 0.58]^T \text{ and } d(0) = 1$$

$$w(n) = w(2) = [0.2, -0.7, -0.44]^T$$

$$y(n) = y(2) = \text{sgn}(0.2 - 0.196 - 0.2552) = -1 \neq d(0)$$

$$w(3) = [0.2, -0.7, -0.44]^T + [1+1][1, 0.28, 0.58]^T = [0.2, -0.7, -0.44]^T + [2, 0.56, 1.16]^T = [2.2, -0.14, 0.72]^T$$




$$i(n) = i(3) = [1, 0.1, 1]^T \text{ and } d(0) = -1$$

$$w(n) = w(3) = [2.2, -0.14, 0.72]^T$$

$$y(n) = y(3) = \text{sgn}(2.2 - 0.014 + 0.72) = 1 \neq d(0)$$

$$w(4) = [2.2, -0.14, 0.72]^T + [-1 - 1][1, 0.1, 1]^T = [2.2, -0.14, 0.72]^T + [-2, -0.2, -2]^T = [-4.4, -0.028, -1.28]^T$$




$$i(n) = i(4) = [1, 0.6, 0.3]^T \text{ and } d(0) = 1$$


$$w(n) = w(4) = [-4.4, -0.028, -1.28]^T$$

$$y(n) = y(4) = \text{sgn}(-4.4 - 0.0168 - 0.384) = -1 \neq d(0)$$


$$w(5) = [-4.4, -0.028, -1.28]^T + [1+1][1, 0.6, 0.3]^T = [-4.4, -0.028, -1.28]^T + [2, 1.2, 0.6]^T = [-2.4, 1.172, -0.68]^T$$



b)  $w = (-0.875, -1)$  and  $b = 0.95625$



c) In a one dimensional perceptron the equation of the line would be  $x = \frac{x_2 + x_1}{2}$  where  $x_1$  and  $x_2$  are two closest points that are from different classes. So  $w = -2$  and  $b = x_2 + x_1$ .



Question 5

a)

b)

