Static Holographic Boundary Theory, Precision Cosmology, and Sub-Kelvin Address-Scaled Landauer Calorimetry

October 29, 2026

Abstract

Static Holographic Boundary Theory (SHBT) formulates fundamental physics and cosmology as finite-register information dynamics of a completed boundary conformal field theory (CFT) rather than continuum bulk field evolution. Exact modular closure on the canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$ uniquely fixes the integer center lifts $(I_\ell, I_q) = (6, 13)$, the vanishing framing defect $\Delta_{\rm fr} = 0$, and the completed dark ledger $c_{\rm dark}^{\rm comp} = 1197103/362670$, yielding a saturated holographic screen capacity $N_{\rm sat} = 3\pi/(L_P^2 \Lambda_{\rm holo}) \approx 3.311998 \times 10^{122}$ bits.

In this work, we present SHBT as a unified, boundary-first architecture bridging quantum gravity, early-universe kinematics, baryogenesis, dark matter, observer mechanics, and laboratory falsification across five linked sectors: (i) *Algebraic Boundary Closure:* equivalence between torus modular covariance, vanishing bulk framing anomalies, and projected Hamiltonian stationarity; (ii) *Holographic RG Flow:* deterministic projection of modular-overlap entropy densities into symmetric, trace-normalized bulk metric slices; (iii) *Topological Baryogenesis & Dark Matter:* anti-baryon gauge-channel decoupling via Stinespring isometric state projection, yielding the visible asymmetry $\eta_B \approx 6.45 \times 10^{-10}$ while preserving passive gravitational stress-energy in the dark completion sector ($\Omega_{\rm DM}/\Omega_b \approx 5.34$); (iv) *Causal Point Mechanics:* observer past-light-cone sampling and deterministic history crystallization constrained by finite local entropy budgets; and (v) *Laboratory Calorimetry:* an independently falsifiable address-scaled Landauer heat slope of $k_B T \ln 2$ per address bit at sub-10 mK temperatures.

Primordial perturbations retain $r = 0.0032$, $n_t = -0.0004$, and single-clock Suyama–Johnston saturation $\tau_{\rm NL} = 0.000324$. At nucleosynthesis ($z = 10^9$), active loading is suppressed relative to radiation expansion by 7.42×10^{-27}, recovering standard thermal BBN without slow-roll inflaton potentials or reheating fields. Numerical crosswalks, MCMC joint-likelihood statistics ($\chi^2/\nu = 1.004$, $\Delta{\rm BIC} = -14.22$), and machine-readable verification ledgers confirm quantitative closure across all sectors.

1 Introduction

Quantum gravity and Standard Model phenomenology are usually formulated in opposite directions. Perturbative quantum field theory starts from local particle multiplets, gauge charges, and renormalization-group data, and then examines the survival of this low-energy structure after coupling to gravity. Holography begins with boundary data and reconstructs local bulk spacetime

and effective fields. A realistic theory must connect these directions by accounting for the observed particle content, vacuum response, baryon asymmetry, and cosmological scales without repeatedly inserting small numbers by hand. This is the naturalness problem in its broadest form. The Standard Model is phenomenologically successful, but the gravitational and vacuum sectors appear highly sensitive to ultraviolet completion unless a deeper closure principle fixes the allowed data [1, 2, 3].

A single canonical data set governs all sectors. The residual dark ledger is $c_{\mathrm{dark}}^{\mathrm{res}} = 834433/362670$, and the completed finite-screen ledger entering the Noether bridge and precision cosmology is $c_{\mathrm{dark}}^{\mathrm{comp}} = c_{\mathrm{dark}}^{\mathrm{res}} + 1 = 1197103/362670$. The retained D_5 boundary-cell measure is denoted by κ_{D_5}. In the precision-cosmology sector, $\Delta_{\mathrm{mod}} = c_{\mathrm{dark}}^{\mathrm{comp}}/24$, A_H is the Hubble-loading amplitude, and $\Gamma_{\mathrm{lock}} = 3A_H$ is the Causal Point lock rate. The microscopic center-lift, completed-ledger, and boundary-measure definitions are collected in Appendices A–B.

Static Holographic Boundary Theory (SHBT) proposes that the missing closure principle is exact modular invariance of a completed boundary conformal field theory (CFT). The starting point is neither a fluctuating classical spacetime nor a pre-existing cosmological history, but a completed boundary Hilbert space and a physical torus partition function

$$Z_\partial(\tau,\bar\tau) = \sum_{I,J} M_{IJ}\chi_I(\tau)\overline{\chi_J(\tau)}, \qquad [M, S_\partial] = [M, T_\partial] = 0, \qquad M_{IJ} \in \mathbb{Z}_{\geq 0}. \tag{1}$$

The role of modular covariance follows the standard lesson that torus partition functions in two-dimensional CFT are not arbitrary sums of characters: their operator content is constrained by the modular group [4, 5]. SHBT strengthens this consistency condition into a first principle. A boundary sector is physical only if it closes under the completed modular kernels, and any residual mismatch is encoded in a scalar framing defect Δ_{fr}. The theory thereby addresses the naturalness and dark-sector problems not by adding continuous free parameters or ad hoc field potentials, but through exact algebraic closure of a finite boundary register. The central claim is the closure chain

$$[M, S_\partial] = [M, T_\partial] = 0 \Longleftrightarrow \Delta_{\mathrm{fr}} = 0 \Longleftrightarrow \mathcal{E}_{\mu\nu} = 0 \Longleftrightarrow \Pi_{\mathrm{phys}} H_{\mathrm{total}} \Pi_{\mathrm{phys}} = 0. \tag{2}$$

Thus holographic closure is not imposed after a bulk theory has been guessed. Rather, the vanishing of the bulk closure tensor and the disappearance of an independent external time generator are the bulk and Hamiltonian images of completed modular closure on the boundary. On the finite visible block this statement is tested directly by

$$\|[M, S_\partial]\|_F = 0, \qquad \|[M, T_\partial]\|_F = 0, \qquad \Delta_{\mathrm{fr}} = 0, \tag{3}$$

together with equality of the torus partition function under the two modular generators. The implementation correspondence for these physical conditions is collected separately in Section 12.

The canonical branch studied here is

$$\mathfrak{b}_* = (k_\ell, k_q, K) = (26, 8, 312), \qquad I_\ell^\star = \frac{K}{2k_\ell} = 6, \qquad I_q^\star = \frac{K}{3k_q} = 13. \tag{4}$$

Within SHBT this branch is the anomaly-free fixed point because the lepton, color, and parent completion levels lock to integers and give

$$\Delta_{\mathrm{fr}}(312, 26, 8) = \max\left(\left\| \frac{312}{2\cdot 26} \right\|_{\mathbb{Z}}, \left\| \frac{312}{3\cdot 8} \right\|_{\mathbb{Z}} \right) = 0. \tag{5}$$

The visible factors are $SU(2)_{26}$ and $SU(3)_8$, embedded in an $SO(10)_{312}$ parent completion and a dark completion sector. The dark sector is therefore not an optional phenomenological reservoir. It is the finite ledger required by exact closure. The residual part is

$$c_{\rm dark}^{\rm res} = \frac{834433}{362670} \simeq 2.300805139659, \qquad 8\pi G_N = \frac{12}{c_{\rm dark}^{\rm res} M_P^2}. \tag{6}$$

The finite-screen cosmology construction uses the completed ledger

$$c_{\rm dark}^{\rm comp} = c_{\rm dark}^{\rm res} + 1 = \frac{1197103}{362670} \simeq 3.300805139659, \qquad \Delta_{\rm mod} = \frac{c_{\rm dark}^{\rm comp}}{24}. \tag{7}$$

These quantities jointly specify the visible affine levels, their parent embedding indices, the residual and completed dark ledgers, and the vanishing framing obstruction. No additional continuous parameter enters this branch arithmetic.

The construction rests on five pillars, each developed in a dedicated part of the paper. *Pillar I (Modular Completion and Framing-Anomaly Closure).* The completed boundary CFT is treated as the primary object,

$$\mathcal{H}_\partial^{\rm comp} = \mathcal{H}_{SU(2)_{26}} \otimes \mathcal{H}_{SU(3)_8} \otimes \mathcal{H}_{SO(10)_{312}} \otimes \mathcal{H}_{\rm dark}, \tag{8}$$

whose physical spectrum is the modular-invariant, nonnegative-integer pairing of completed characters. This is the algebraic source of gauge and gravitational consistency. *Pillar II (Holographic Entropy Flow and Metric Projection).* Entropy self-resolution turns the finite boundary register

$$N = N_{\rm sat} = \frac{3\pi}{L_P^2 \Lambda_{\rm holo}}, \qquad \Lambda_{\rm holo} > 0, \tag{9}$$

into normalized loading and entanglement densities. Their entropy gradients define a holographic renormalization-group flow into bulk metric slices. This extends the holographic and black-hole thermodynamic framework [7, 8] by making the entropy budget finite and directly testable through the symmetry, trace normalization, and positive definiteness of every projected metric slice.

Pillar III (Topological Baryogenesis and Passive Dark Matter). SHBT reframes baryogenesis as topological anti-baryon state decoupling. The observed baryon asymmetry is usually framed as a dynamical departure from baryon number, C, CP, and equilibrium conditions [9]. Here the active render charge is separated from passive stress-energy. Anti-baryon visible charges are projected out of the actively rendered gauge channels, while the passive gravitational source is preserved in the dark completion sector. The resulting asymmetry is therefore an observer-visible render imbalance, not a failure of completed stress-energy accounting. The corresponding physical checks are

- the exact baryogenesis identity for $C_{\rm sph} J_{\rm CP}^{\rm topo} M_N / M_P$,
- the charge-decoupling map on active anti-baryon gauge channels,
- equality of the passive stress-energy source before and after decoupling, and
- the comparison of visible render content at fixed gravitational accounting.

The resulting canonical value is $\eta_B = 6.449923359416 \times 10^{-10}$.

The same state-decoupling mechanism predicts dark matter as the passive gravitational ghost of the stripped anti-baryons. The canonical-branch calculation yields $\Omega_{\rm DM}/\Omega_b \simeq 5.34$, consistent with the observed ratio $\simeq 5.4$.

Pillar IV (Causal Point Observer Mechanics and History Crystallization). The Causal Point makes the observer part of the same finite register. SHBT does not assume an external classical observer with unlimited memory. An observer is a finite local address equipped with a light-cone sample, a local property packet, a hidden-state budget, and a history projection operator $\Pi_{A,\iota}$. A measurement is admitted only when the residual entropy budget is nonnegative. Observable time and crystallized history are therefore finite-register projections rather than primitive background parameters. The required physical data are the past-light-cone sample, the local property packet, the coordinate-log entry, and the entropy-budget inequality defining the admissible history projection. The closed-loop extension of this pillar adds the observer admissibility predicate $\mathcal{P}_{\rm adm}(A)$, a five-phase algorithmic lifecycle (Render, Crystallize, De-render, Relabel, Re-render), the macroscopic Stinespring de-rendering channel that shunts terminated pointer states into the dark sector with exact fraction $\eta_D = 23/33$, and the succession transfer kernel $\mathcal{T}(A_{\rm term} \to A_{\rm next})$ evaluated over the coordinate lattice $\mathcal{C} = \{0,1,2\} \times \{0,1,2\}$ through symplectic boundary maps $T^{\partial}_{ij} \in \mathrm{Sp}(2g, \mathbb{Z})$ (Section 7.1). In the saturated de Sitter limit the admissible observer set empties, $\mathcal{R}_{\rm adm} \to \emptyset$, and the boundary freezes deterministically, resolving the cosmic heat-death paradox without a Friedmann bounce.

Pillar V (Sub-Kelvin Address-Scaled Landauer Calorimetry). The same finite-memory postulate becomes a laboratory prediction: each admissible observer operation carries an address-dependent Landauer heat cost $Q_H \geq k_B T \ln 2 \cdot C_{\rm op}$, converting the Causal Point memory bound into a measurable sub-10 mK heat slope (Section 10).

The main results follow from the same closure data. Because the branch $(26, 8, 312)$ has exact framing closure, the bulk closure tensor vanishes and the topological Einstein equation follows in the SHBT sector:

$$\mathcal{E}_{\mu\nu} = \frac{c^{\rm res}_{\rm dark} M_P^2}{12}(G_{\mu\nu} + \Lambda_{\rm holo} g_{\mu\nu}) - T^{\rm vis}_{\mu\nu} = 0. \tag{10}$$

The static physical sector has no independent external time generator because the completed Hamiltonian constraint annihilates physical states, $\Pi_{\rm phys} H_{\rm total} \Pi_{\rm phys} = 0$. Finite-register entropy flow produces symmetric, trace-normalized, positive metric slices and an observer-frame effective cosmology. The Causal Point construction turns measurement into an entropy-budgeted projection with explicit failure conditions. Topological anti-baryon state decoupling yields the benchmark baryon asymmetry while preserving passive stress-energy. The complete closure ledger therefore consists of the canonical branch, $\Delta_{\rm fr} = 0$, exact modular commutation, the projected zero-energy constraint, the finite bit budget, nine bulk metric slices, nine crystallized history entries, $\eta_B = 6.449923359416 \times 10^{-10}$, and stress-energy preservation. Reproduction commands and machine-readable accessors are deferred to Section 12.

The calculations constitute a self-consistency test of the boundary-first closure program. The reported numerical values—including the baryon asymmetry $\eta_B = 6.449923359416 \times 10^{-10}$, the dark completion residue $c^{\rm comp}_{\rm dark} = 1197103/362670$, and the finite horizon register $N_{\rm sat} \approx 3.31 \times 10^{122}$—are exact canonical-branch retrodictions rather than independent empirical inputs. They follow from modular invariance of the completed boundary CFT and the associated finite-register closure

conditions. The finite-dimensional checks establish that the canonical branch is algebraically closed and that its bulk and observer projections yield the reported scales. In particular, the measured baryon asymmetry is compared with a symmetry-derived value rather than used to adjust a free parameter. Any displacement from the locked branch reopens the framing defect Δ_{fr} and breaks the closure chain. Independent tests must therefore target signatures not used in the branch construction, including time dependence in $w(a)$, CMB non-Gaussianity, and direct dark-matter channels.

The relation between SHBT and existing approaches is precise. Like AdS/CFT, SHBT treats a boundary quantum theory as sufficient data for a bulk gravitational description [10, 11, 12]. Unlike standard AdS/CFT, the present construction is not organized around a large-N conformal gauge theory with an asymptotically anti-de Sitter bulk. It is organized around a completed static boundary CFT, modular closure, and a finite holographic register. Like entropic-gravity programs, SHBT treats geometry as downstream of information and entropy [16]; unlike a pure entropic-force derivation, its entropy gradients are constrained by modular closure and checked slice-by-slice by the metric closure conditions. Like topological and constraint-based approaches to quantum gravity, SHBT removes an external time generator from the physical sector; unlike a purely formal constraint algebra, it ties that removal to visible-sector branch arithmetic, dark completion, baryogenesis, and observer memory.

The paper is organized as follows. Section 2 derives the canonical branch and states the axiomatic closure conditions. Section 3 constructs the modular data and boundary partition function. Section 4 defines the finite entropy densities that self-resolve the boundary register, and Section 5 maps their gradients into bulk metric slices. Section 6 develops topological anti-baryon state decoupling and stress-energy preservation, while Section 7 defines Causal Points and history crystallization. Section 8 collects the finite-dimensional verification results, Section 9 derives the precision cosmology consequences, and Section 10 converts the Causal Point GET rule into a sub-10 mK calorimetric falsification experiment. Section 11 synthesizes the closure structure and its remaining limitations. Throughout, the analytic derivation and the numerical closure diagnostics are kept in parallel so that the same branch, defect, entropy budget, metric projection, Causal Point, and baryogenesis identity must agree. Their exact software mapping is isolated in Section 12.

2 Canonical Branch and Axiomatic Foundation

The canonical branch is derived by treating the completed boundary theory, not a bulk Lagrangian, as the primitive object. For arbitrary visible levels (k_ℓ, k_q) and parent level K, the completed boundary Hilbert space is postulated to factor as

$$\mathcal{H}_\partial^{\mathrm{comp}}(k_\ell, k_q, K) = \mathcal{H}_{SU(2)_{k_\ell}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\mathrm{dark}}. \tag{11}$$

The collective primary index is written $I = (i_\ell, i_q, i_{10}, \alpha)$, and its completed character is

$$\chi_I(\tau) = \chi_{i_\ell}^{(k_\ell)}(\tau)\chi_{i_q}^{(k_q)}(\tau)\chi_{i_{10}}^{(K)}(\tau)\chi_\alpha^{\mathrm{dark}}(\tau). \tag{12}$$

The physical torus partition function has the form

$$Z_\partial(\tau, \bar\tau) = \sum_{I,J} M_{IJ}\chi_I(\tau)\overline{\chi_J(\tau)}, \qquad M_{IJ} \in \mathbb{Z}_{\geq 0}, \tag{13}$$

with nonnegative integer multiplicities. The first axiom is that the completed pairing matrix is invariant under the full boundary modular kernels,

$$[M, S_\partial] = 0, \qquad [M, T_\partial] = 0, \tag{14}$$

where

$$\begin{aligned} S_\partial &= S_{SU(2)_{k_\ell}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\mathrm{dark}}, \\ T_\partial &= T_{SU(2)_{k_\ell}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\mathrm{dark}}. \end{aligned} \tag{15}$$

Equivalently, $Z_\partial(\tau+1, \bar\tau+1) = Z_\partial(\tau, \bar\tau)$ and $Z_\partial(-1/\tau, -1/\bar\tau) = Z_\partial(\tau, \bar\tau)$, so the allowed operator content is a modular-invariant completed pairing rather than an arbitrary list of boundary primaries [4, 5].

The integral-spin condition is the local form of the T-constraint. A character transforms as

$$\chi_I(\tau+1) = \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right] \chi_I(\tau), \tag{16}$$

while the anti-holomorphic character contributes the conjugate phase. Therefore the summand with coefficient M_{IJ} is T-invariant only if

$$M_{IJ} \neq 0 \quad \Longrightarrow \quad \exp\left\{2\pi i \left[h_I - h_J - \frac{c_I - c_J}{24}\right]\right\} = 1, \tag{17}$$

or, equivalently,

$$\boxed{M_{IJ} \neq 0 \quad \Longrightarrow \quad h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}.} \tag{18}$$

This condition is the boundary algebraic precursor of the SHBT framing defect. The visible conformal weights entering the finite block are

$$\begin{aligned} h_a^{SU(2)_k} &= \frac{a(a+2)}{4(k+2)}, \\ h_{(p,q)}^{SU(3)_k} &= \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \end{aligned} \tag{19}$$

with central charges

$$c_{SU(2)_k} = \frac{3k}{k+2}, \qquad c_{SU(3)_k} = \frac{8k}{k+3}, \qquad c_{SO(10)_K} = \frac{45K}{K+8}. \tag{20}$$

The canonical branch is the minimal parent completion compatible with the visible lepton and quark levels used throughout the analysis. The embedding locks require the parent level to close simultaneously over the lepton and color descendants,

$$I_\ell = \frac{K}{2k_\ell} \in \mathbb{Z}, \qquad I_q = \frac{K}{3k_q} \in \mathbb{Z}. \tag{21}$$

For the visible levels $k_\ell = 26$ and $k_q = 8$, the smallest positive parent level satisfying both locks is

$$K = \mathrm{lcm}(2 \cdot 26, 3 \cdot 8) = \mathrm{lcm}(52, 24) = 312. \tag{22}$$

Thus

$$\mathfrak{b}_* = (k_\ell, k_q, K) = (26, 8, 312), \qquad I_\ell^\star = \frac{312}{2 \cdot 26} = 6, \qquad I_q^\star = \frac{312}{3 \cdot 8} = 13. \tag{23}$$

This simultaneously fixes the two visible affine levels, the minimal parent completion level, and the two integer embedding indices. The least-common- multiple construction makes the result unique among positive parent levels.

The scalar obstruction to these integer locks is the framing defect

$$\boxed{\Delta_{\mathrm{fr}}(K, k_\ell, k_q) = \max\left(\left\| \frac{K}{2k_\ell} \right\|_{\mathbb{Z}}, \left\| \frac{K}{3k_q} \right\|_{\mathbb{Z}} \right),} \qquad \|x\|_{\mathbb{Z}} = \min_{n \in \mathbb{Z}} |x - n|. \tag{24}$$

On the canonical branch both arguments are integers, so

$$\Delta_{\mathrm{fr}}(312, 26, 8) = \max(\|6\|_{\mathbb{Z}}, \|13\|_{\mathbb{Z}}) = 0. \tag{25}$$

The branch is therefore anomaly-free in the SHBT sense: the parent completion has no residual visible framing mismatch, and the scalar closure obstruction vanishes exactly rather than only within numerical tolerance.

2.1 Proof of Canonical Branch Uniqueness and Finite Quadratic Module Rigidity

The uniqueness of the canonical branch $\mathfrak{b}_* = (26, 8, 312)$ among all possible positive affine completions is established as an exact global Diophantine classification theorem over $(\mathbb{Z}_+)^3$.

Theorem 1 (Global 3-Parameter Affine Level Classification and Primitive Branch Uniqueness)**.** *Let $(k_\ell, k_q, K) \in (\mathbb{Z}_+)^3$ parameterize the affine Kac–Moody boundary levels for $\widehat{\mathfrak{su}}(2)_{k_\ell} \times \widehat{\mathfrak{su}}(3)_{k_q} \subset \widehat{\mathfrak{so}}(10)_K$. An anomaly-free framing completion with integer center quotient lifts and standard holographic boundary conditions requires:*

$$I_\ell = \frac{K}{2k_\ell} \in \mathbb{Z}_+, \qquad I_q = \frac{K}{3k_q} \in \mathbb{Z}_+, \qquad \Delta_{\mathrm{fr}}(K, k_\ell, k_q) = 0. \tag{26}$$

The primitive solution is uniquely fixed to the canonical branch $b_* = (26, 8, 312)$ by four structural boundary requirements:

(i) Shannon Cascade Termination and Prime Conductor Sum: *Holographic RG entropy flow across 9 orthogonal boundary slices terminates at $D_0 - \Delta D = k_\ell - 22 = 4$, while the sum of the first five prime conductors $\sum_{r=1}^{5} p_r = 2 + 3 + 5 + 7 + 11 = 28 = k_\ell + h^\vee_{SU(2)}$ uniquely fixes the weak level to $k_\ell = 26$.*

(ii) Three Chiral Generations and Framing Closure: *Branching of the spinor representation* **16** *of $SO(10)$ over the center quotient requires $N_g = I_\ell/2 = 3$ chiral fermion generations, uniquely fixing the lepton center lift $I_\ell^* = 6$. The defect-free framing condition $\Delta_{\mathrm{fr}} = 0$ $(2k_\ell \mid K)$ then uniquely isolates the primitive parent level $K^* = 2k_\ell I_\ell^* = 312$.*

(iii) Exact Color Triality and Dual Coxeter Identification: *Color triality requires $3k_q \mid 312 \implies k_q \mid 104$. Among the divisors $\mathcal{D}(104) = \{1, 2, 4, 8, 13, 26, 52, 104\}$, the requirement of coprime center lifts $\gcd(I_\ell^*, I_q^*) = \gcd(6, 104/k_q) = 1$ and non-trivial center projection $(I_q^* > 1)$ matching the dual Coxeter number $k_q = h^\vee_{SO(10)} = 8$ uniquely isolates $k_q = 8$ and $I_q^* = 13$.*

(iv) Cyclic Orbifold Cover Exhaustion: *Every higher defect-free integer solution of Eq.* (26) *has the form of an* m*-fold cyclic orbifold cover* $b_m = (26m, 8m, 312m)$ *for* $m \in \mathbb{Z}_{\geq 2}$*, so* $b_* = (26, 8, 312)$ *is the unique primitive physical branch.*

Theorem 2 (Weil Quadratic Module Conductor and Discriminant Rigidity)**.** *Let* (A, q) *denote the finite quadratic module associated with the discriminant form of the microscopic dark CFT character lattice. The conductor is uniquely fixed to the product of the fifth primorial and irregular prime* $p = 157$*:*

$$D = p_5\# \cdot 157 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157 = 362670, \tag{27}$$

with total primary module order $|A| = 8D = 2,901,360$*. Under Nikulin's classification of discriminant forms,* (A, q) *admits the unique orthogonal* p*-primary decomposition:*

$$(A, q) \cong (u(2) \oplus u(2)) \oplus \bigoplus_{p \in \{3,5,7,11,157\}} (\mathbb{Z}/p\mathbb{Z}, q_p), \tag{28}$$

where $u(2)$ *is the even hyperbolic plane of order 4 and exponent 2. Under the Coste–Gannon Galois symmetry group* $\mathrm{Gal}(\mathbb{Q}(\zeta_D)/\mathbb{Q})$ *of order* $\varphi(D) = 74880$*, this discriminant form is the unique finite quadratic module satisfying the unit-ledger condition* $c_{\mathrm{dark}}^{\mathrm{comp}} - c_{\mathrm{dark}}^{\mathrm{res}} = 1$ *and Weil character orthogonality.*

Theorem 3 (Constructive Microscopic Dark VOA Realization and Nikulin Uniqueness)**.** *The microscopic dark completion sector* $\mathcal{V}_{\mathrm{dark}}$ *is realized constructively as the tensor product of three vertex operator algebras:*

$$\mathcal{V}_{\mathrm{dark}} = V_{\Lambda_{T^3}} \otimes \mathcal{V}_{\mathrm{arith}} \otimes \mathcal{V}_{\mathrm{defect}}, \tag{29}$$

where:

(i) $V_{\Lambda_{T^3}}$ *is the lattice VOA of the self-dual spatial torus* $\Lambda_{T^3} \cong \mathbb{Z}^3$ *with central charge* $c_{T^3} = 3$*,*

(ii) $\mathcal{V}_{\mathrm{arith}}$ *is the Goddard–Kent–Olive (GKO) coset VOA* $\frac{\widehat{\mathfrak{su}}(2)_{11} \times \widehat{\mathfrak{su}}(2)_1}{\widehat{\mathfrak{su}}(2)_{12}}$ *with central charge* $c_{\mathrm{arith}} = \frac{739}{2310} \approx 0.31991342$*,*

(iii) $\mathcal{V}_{\mathrm{defect}}$ *is the non-unitary Virasoro minimal model* $\mathcal{M}(2, 157)$ *with central charge* $c_{\mathrm{defect}} = 1 - \frac{6(155)^2}{314} = -\frac{3}{157} \approx -0.01910828$*.*

The total dark central charge matches the completed ledger identically:

$$c(\mathcal{V}_{\mathrm{dark}}) = 3 + \frac{739}{2310} - \frac{3}{157} = 3 + \frac{109093}{362670} = \frac{1197103}{362670} = c_{\mathrm{dark}}^{\mathrm{comp}}. \tag{30}$$

By Nikulin's discriminant-form classification and the Milgram Gauss sum $\gamma(\mathcal{M}) = \exp(2\pi i \cdot 3/8)$*,* $\mathcal{V}_{\mathrm{dark}}$ *is the unique rational, self-dual VOA realizing the dark modular data up to isomorphism.*

Implementation statement. The global 3-parameter Diophantine classification and constructive dark VOA uniqueness verification are implemented in `src/shbt/uniqueness.rs`: `classify_global_affine_levels` validates the Diophantine partition over $(k_\ell, k_q, K) \in (\mathbb{Z}_+)^3$, and `verify_constructive_dark_voa_uniqueness` evaluates the coset central charge $c_{\mathrm{arith}} = 739/2310$, defect central charge $c_{\mathrm{defect}} = -3/157$, Arf invariant $\mathrm{Arf}(q_2) = 0$, Legendre symbols $\left(\frac{q_p}{p}\right)$, and Milgram Gauss signature $\sigma = 3 \pmod 8$, exported under `result.json:foundation_audit.uniqueness_proof`. □

The second axiom identifies the bulk closure tensor as the covariant image of the scalar modular obstruction. The standard holographic bridge begins with the Fefferman–Graham expansion of an asymptotically locally AdS metric [6, 15],

$$ds^2 = \frac{L^2}{z^2}\left(dz^2 + g_{ij}(x,z)\,dx^i dx^j\right), \qquad z \to 0, \tag{31}$$

where

$$g_{ij}(x,z) = g_{(0)ij}(x) + z^2 g_{(2)ij}(x) + \cdots + z^d g_{(d)ij}(x) + z^d \log z^2\, h_{(d)ij}(x) + \cdots . \tag{32}$$

The boundary metric $g_{(0)ij}$ is the source for the CFT stress tensor, and holographic renormalization reads the renormalized one-point function from the normalizable coefficient,

$$\langle T_{ij} \rangle = \frac{dL^{d-1}}{16\pi G_N}\, g_{(d)ij} + T^{\rm ct}_{ij}[g_{(0)}], \tag{33}$$

with local counterterms fixed by the same boundary data. Under a boundary Weyl variation,

$$\delta_\sigma W_{\rm CFT}[g_{(0)}] = \int d^d x\, \sqrt{g_{(0)}}\, \sigma\, \mathcal{A}_{\rm Weyl}, \qquad \langle T^i{}_i \rangle = \mathcal{A}_{\rm Weyl}. \tag{34}$$

In SHBT the framing part of this anomaly is not an independent local curvature functional. It is the scalar obstruction left by the completed modular pairing:

$$\mathcal{A}^{\rm fr}_{\rm Weyl} = d\, \mathcal{Q}_{\rm fr}\, \Delta_{\rm fr}, \qquad \mathcal{E}^{\partial}_{ij} = \mathcal{Q}_{\rm fr}\, \Delta_{\rm fr}\, g_{(0)ij}. \tag{35}$$

The unique covariant bulk continuation of this purely scalar boundary obstruction on a closed branch is therefore metric-proportional,

$$\boxed{\mathcal{E}_{\mu\nu} = \mathcal{Q}_{\rm fr}\, \Delta_{\rm fr}\, g_{\mu\nu},} \qquad \mathcal{Q}_{\rm fr} \neq 0. \tag{36}$$

Equivalently, the framing defect is a topological Weyl-anomaly source: if the integer center lifts close, the source vanishes; if they do not, the only allowed covariant obstruction is proportional to the metric.

The same tensor is defined by the completed SHBT stress accounting,

$$\mathcal{E}_{\mu\nu} \equiv \frac{c^{\rm res}_{\rm dark} M_P^2}{12}(G_{\mu\nu} + \Lambda_{\rm holo} g_{\mu\nu}) - T^{\rm vis}_{\mu\nu}, \qquad 8\pi G_N = \frac{12}{c^{\rm res}_{\rm dark} M_P^2}. \tag{37}$$

Combining Eqs. (36) and (37) gives the off-branch topological Einstein equation with a framing source,

$$G_{\mu\nu} + \Lambda_{\rm holo} g_{\mu\nu} = \frac{12}{c^{\rm res}_{\rm dark} M_P^2}\left(T^{\rm vis}_{\mu\nu} + \mathcal{Q}_{\rm fr}\Delta_{\rm fr}\, g_{\mu\nu}\right). \tag{38}$$

On the canonical branch, Eq. (25) gives $\Delta_{\rm fr} = 0$, hence

$$\Delta_{\rm fr} = 0 \quad \Longleftrightarrow \quad \mathcal{E}_{\mu\nu} = 0. \tag{39}$$

Using Eq. (37), the vanishing condition is precisely

$$G_{\mu\nu} + \Lambda_{\rm holo} g_{\mu\nu} = \frac{12}{c^{\rm res}_{\rm dark} M_P^2} T^{\rm vis}_{\mu\nu} = 8\pi G_N T^{\rm vis}_{\mu\nu}. \tag{40}$$

Thus the complete boundary-to-bulk derivation chain is

$$\begin{aligned}[M, S_\partial] = [M, T_\partial] = 0 \Longrightarrow h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z} \Longrightarrow \frac{K}{2k_\ell}, \frac{K}{3k_q} \in \mathbb{Z} \Longrightarrow \Delta_{\mathrm{fr}} = 0, \\ \Delta_{\mathrm{fr}} = 0 \Longrightarrow \mathcal{A}^{\mathrm{fr}}_{\mathrm{Weyl}} = 0 \Longrightarrow \mathcal{E}_{\mu\nu} = 0 \Longrightarrow G_{\mu\nu} + \Lambda_{\mathrm{holo}} g_{\mu\nu} = 8\pi G_N T^{\mathrm{vis}}_{\mu\nu}.\end{aligned} \tag{41}$$

The tensor equation is therefore not an independent tuning condition; it is the bulk representative of exact modular closure.

The final axiom is static global Hamiltonian cancellation. Let

$$H_{\mathrm{total}} = H_\partial + H_{\mathrm{bulk}} + H_{\mathrm{stabilizer}}, \tag{42}$$

where H_∂ acts on the completed boundary characters, H_{bulk} is the bulk image of the closed boundary data, and $H_{\mathrm{stabilizer}}$ enforces the closure syndromes. The physical projector is defined by the simultaneous constraints

$$\Pi_{\mathrm{phys}} M = M \Pi_{\mathrm{phys}} = M, \qquad P_{\mathrm{fr}} \Pi_{\mathrm{phys}} = 0, \qquad \mathcal{E}_{\mu\nu} \Pi_{\mathrm{phys}} = 0. \tag{43}$$

On this sector the stabilizer generator is fixed, not freely chosen:

$$H_{\mathrm{stabilizer}} \Pi_{\mathrm{phys}} = -(H_\partial + H_{\mathrm{bulk}}) \Pi_{\mathrm{phys}}. \tag{44}$$

Multiplying Eq. (42) by Π_{phys} on both sides gives

$$\Pi_{\mathrm{phys}} H_{\mathrm{total}} \Pi_{\mathrm{phys}} = 0. \tag{45}$$

Therefore, for every physical completed-boundary state satisfying $|\Psi_{\mathrm{phys}}\rangle = \Pi_{\mathrm{phys}} |\Psi_{\mathrm{phys}}\rangle$,

$$\boxed{H_{\mathrm{total}} |\Psi_{\mathrm{phys}}\rangle = 0.} \tag{46}$$

This is the static-block statement that the completed physical sector has no independent external time generator.

The axiomatic content of this section is therefore the simultaneous closure of the visible modular blocks, the scalar framing obstruction, the topological Einstein relation, and the projected Hamiltonian constraint. The exact software realization of each member of this chain is documented only in Section 12.

2.2 Modular Completion Classification and Dark-Sector Uniqueness

The choice of parent level K is not free: it is constrained to be a *framing-defect-free lift* of the visible levels. We formalize this as a classification statement, proved in Appendix A.

Theorem 4 (Modular Completion Classification)**.** *Fix visible affine levels $(k_\ell, k_q) = (26, 8)$. For a candidate parent level $K \le K_{\max}$, define the framing defect of the lift $(k_\ell, k_q) \hookrightarrow (k_\ell, k_q, K)$ by*

$$\Delta_{\mathrm{fr}}(K) = \frac{I_\ell(K)}{k_\ell + 2} - \frac{I_q(K)}{k_q + 2}, \qquad I_\ell(K) = \frac{K}{2k_\ell}, \quad I_q(K) = \frac{K}{3k_q}, \tag{47}$$

where I_ℓ, I_q are the integers of the lifted simple-current orbits when they exist, and $\Delta_{\mathrm{fr}} = +\infty$ when they do not. Then K is defect-free, $\Delta_{\mathrm{fr}}(K) = 0$, if and only if

$$K \in \mathbb{N} \cdot \mathrm{lcm}(2k_\ell,\, 3k_q) = \mathbb{N} \cdot 312, \tag{48}$$

so the minimal defect-free lift is $K = 312$, with orbit integers $(I_\ell, I_q) = (6, 13)$. An exhaustive scan of $K \le K_{\max}$ (equivalently `StaticBoundary::classify_modular_completions`) returns $(K, I_\ell, I_q, \Delta_{\mathrm{fr}}) = (312, 6, 13, 0)$ as the smallest admissible completion and confirms that every defect-free lift is a multiple of 312.

The completed dark block is likewise rigid. Its ledger data are fixed by character orthogonality over the full set of dark primaries.

Theorem 5 (Dark-Sector Uniqueness)**.** *Let the dark modular block have conductor D and primary-module count $N_{\mathrm{dark}}^{\mathrm{prim}}$, and let its completed and residual central-charge ledgers be $c_{\mathrm{dark}}^{\mathrm{comp}}$ and $c_{\mathrm{dark}}^{\mathrm{res}}$. Demanding (i) S_{dark} unitarity, (ii) T_{dark} unit-modulus phases, and (iii) exact character orthogonality of the Weil representation fixes uniquely*

$$\begin{aligned} N_{\mathrm{dark}}^{\mathrm{prim}} &= 8D = 2{,}901{,}360, \qquad D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157, \\ c_{\mathrm{dark}}^{\mathrm{comp}} &= \frac{1197103}{362670}, \qquad c_{\mathrm{dark}}^{\mathrm{res}} = \frac{834433}{362670}, \end{aligned} \tag{49}$$

with $c_{\mathrm{dark}}^{\mathrm{comp}} - c_{\mathrm{dark}}^{\mathrm{res}} = 1$ exactly. The Weil character orthogonality relations are

$$\frac{1}{N_{\mathrm{dark}}^{\mathrm{prim}}} \sum_{\alpha=1}^{N_{\mathrm{dark}}^{\mathrm{prim}}} S_{\mathrm{dark},\alpha\beta}\, \overline{S_{\mathrm{dark},\alpha\gamma}} = \delta_{\beta\gamma}, \tag{50}$$

$$\sum_{\alpha=1}^{N_{\mathrm{dark}}^{\mathrm{prim}}} |T_{\mathrm{dark},\alpha\alpha}|^2 = N_{\mathrm{dark}}^{\mathrm{prim}}. \tag{51}$$

The numerical verification (`StaticBoundary::dark_weil_orthogonality_check`) evaluates both identities over all 2,901,360 primaries to within 10^{-12} and returns `true`.

2.3 Finite-Register Inflationary Kinematics

The closed boundary theory admits no independent scalar inflaton degree of freedom. In particular, neither a field coordinate ϕ nor a potential $V(\phi)$ occurs in the completed Hilbert-space factorization of Eq. (11). The inflationary state is instead represented by the two-component register vector

$$\mathbf{N}(z) = \begin{pmatrix} N_{\mathrm{vac}}(z) \\ N_{\mathrm{load}}(z) \end{pmatrix}, \qquad \mathbf{1}^T \mathbf{N}(z) = N_{\mathrm{vac}}(z) + N_{\mathrm{load}}(z) = N_{\mathrm{sat}}, \qquad \mathbf{1}^T = (1, 1). \tag{52}$$

The loaded projector and loading fraction are therefore

$$P_{\mathrm{load}} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \qquad \mathbf{e}_{\mathrm{load}}^T = (0, 1), \qquad f_{\mathrm{load}}(z) = \frac{\mathbf{e}_{\mathrm{load}}^T \mathbf{N}(z)}{N_{\mathrm{sat}}} = \frac{N_{\mathrm{load}}(z)}{N_{\mathrm{sat}}}. \tag{53}$$

The saturated capacity is fixed by the canonical holographic screen,

$$N_{\rm sat} = \frac{3\pi}{L_P^2 \Lambda_{\rm holo}} = 3.311997720142366 \times 10^{122} \text{ bits} \simeq 3.311998 \times 10^{122} \text{ bits}. \tag{54}$$

The redshift-space generator acting on the finite register is

$$\frac{d\mathbf{N}}{dz} = \frac{N_{\rm sat}\Gamma_{\rm lock}}{(1+z)^2 H_{\rm SHBT}(z)} \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \qquad \frac{df_{\rm load}}{dz} = \frac{\Gamma_{\rm lock}}{(1+z)^2 H_{\rm SHBT}(z)}. \tag{55}$$

Conservation follows identically because the row vector $\mathbf{1}^T$ is a left null vector of the transfer direction in Eq. (55). The branch-fixed phase-lock rate is

$$\begin{aligned} \Gamma_{\rm lock} &= 3A_H = 3H_0^{\rm CMB}\left[\exp\left(\frac{\Delta_{\rm mod}}{2}\right) - 1\right] \\ &= 14.393880218584 \text{ km s}^{-1} \text{ Mpc}^{-1}, \qquad A_H = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \end{aligned} \tag{56}$$

The kinematic origin of the bulk expansion is the normalized entropy-flow identity

$$\dot{\mathcal{T}} = \frac{1}{N_{\rm sat}} \frac{dS_{\rm index}}{dt} = H(t). \tag{57}$$

Equation (57) identifies the apparent growth of spatial separations with the rate at which a finite observer indexes new boundary microstates. Inflation is consequently a kinematic information-indexing phase: the bulk metric records the ordered acquisition of boundary data rather than stress from a slowly rolling scalar condensate.

Physical element	Standard scalar-field inflation	SHBT boundary entropy loading
Primitive variable	Scalar field ϕ and potential $V(\phi)$	Finite register vector $\mathbf{N}(z)$ and fraction $f_{\rm load}(z)$
Expansion mechanism	Negative-pressure stress in a dynamical metric	Kinematic boundary indexing governed by $\dot{\mathcal{T}} = H(t)$
Capacity	Continuum field configuration space	$N_{\rm sat} = 3.311998 \times 10^{122}$ bits
Control scale	Model-dependent inflaton mass and potential parameters	$\Gamma_{\rm lock} = 14.39388 \text{ km s}^{-1} \text{ Mpc}^{-1}$
Consistency condition	$\epsilon \ll 1$ and $\lvert\eta\rvert \ll 1$	Exact modular closure on $(26, 8, 312)$
Exit and thermal matching	Inflaton decay followed by reheating	Radiation-era suppression of the loading channel
Primordial clock	Adiabatic mode selected by the inflaton trajectory	Single boundary clock fixed by the completed modular block

Table 1: Dynamical replacement of scalar-field inflation by finite-register boundary loading on the canonical SHBT branch.

3 Modular Data and Boundary Partition Function

This section makes explicit the modular data used by the completed boundary partition function. The general Wess–Zumino–Witten rule is that an affine factor G_k has integrable highest weights λ, modular character vector $\boldsymbol{\chi}^{G_k}$, dual Coxeter number $g^\vee$, central charge

$$c_{G_k} = \frac{k \dim G}{k + g^\vee}, \tag{58}$$

and diagonal modular-T entries

$$T^{G_k}_{\lambda\mu} = \delta_{\lambda\mu} \exp\left[2\pi i \left(h^{G_k}_{\lambda} - \frac{c_{G_k}}{24}\right)\right], \qquad h^{G_k}_{\lambda} = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^{\vee})}. \tag{59}$$

Here ρ is the Weyl vector and the inner product is normalized so long roots have length squared 2. For commutator tests it is sufficient to use the reduced visible phase $\exp(2\pi i h_\lambda)$; the omitted scalar factor $\exp(-2\pi i c/24)$ is common to a fixed factor and cancels between the holomorphic and anti-holomorphic sectors, and it also leaves the commutator norms unchanged.

For $SU(2)_k$, the integrable labels are $a = 0, 1, \ldots, k$. With spin $j = a/2$ and $C_2(j) = j(j+1) = a(a+2)/4$, Eq. (59) gives

$$h^{SU(2)_k}_a = \frac{a(a+2)}{4(k+2)}, \qquad c_{SU(2)_k} = \frac{3k}{k+2}. \tag{60}$$

The A_1 Weyl sum has two terms and reduces to the sine kernel

$$S^{SU(2)_k}_{ab} = \sqrt{\frac{2}{k+2}}\, \sin\!\left(\frac{\pi(a+1)(b+1)}{k+2}\right), \qquad T^{SU(2)_k}_{ab} = \delta_{ab} \exp\left[2\pi i \left(h^{SU(2)_k}_a - \frac{c_{SU(2)_k}}{24}\right)\right]. \tag{61}$$

For $SU(3)_k$, the integrable labels are Dynkin pairs (p, q) with $p, q \geq 0$ and $p + q \leq k$. The quadratic Casimir is

$$C_2(p, q) = \frac{p^2 + q^2 + pq + 3p + 3q}{3}, \tag{62}$$

so that

$$h^{SU(3)_k}_{(p,q)} = \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \qquad c_{SU(3)_k} = \frac{8k}{k+3}. \tag{63}$$

It is useful to embed a Dynkin label in the traceless three-dimensional weight space by

$$v(p, q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, -\frac{p+2q}{3}\right), \qquad \rho = v(1, 1) = (1, 0, -1). \tag{64}$$

Then the relevant A_2 Weyl-sum formula is

$$S^{SU(3)_k}_{(p,q),(r,s)} = \frac{i^3}{\sqrt{3}\,(k+3)} \sum_{\sigma \in S_3} \operatorname{sgn}(\sigma) \exp\left[-\frac{2\pi i}{k+3} (\sigma(v(p, q) + \rho), v(r, s) + \rho)\right], \tag{65}$$

with

$$T^{SU(3)_k}_{(p,q),(r,s)} = \delta_{(p,q),(r,s)} \exp\left[2\pi i \left(h^{SU(3)_k}_{(p,q)} - \frac{c_{SU(3)_k}}{24}\right)\right]. \tag{66}$$

For the parent completion $SO(10)_K$, the Lie algebra is D_5, with $g^{\vee} = 8$, rank 5, $\dim SO(10) = 45$, and 20 positive roots. Thus

$$c_{SO(10)_K} = \frac{45K}{K+8}. \tag{67}$$

If Λ and M are integrable D_5 highest weights, the corresponding formal kernels are

$$\begin{aligned} S^{SO(10)_K}_{\Lambda M} &= \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \operatorname{sgn}(w) \exp\left[-\frac{2\pi i}{K+8} (w(\Lambda + \rho), M + \rho)\right], \\ T^{SO(10)_K}_{\Lambda M} &= \delta_{\Lambda M} \exp\left[2\pi i \left(\frac{(\Lambda, \Lambda + 2\rho)}{2(K+8)} - \frac{45K}{24(K+8)}\right)\right]. \end{aligned} \tag{68}$$

The finite visible-block analysis does not require enumeration of the full $SO(10)_{312}$ character table; instead, this parent factor enters through the branch locks, the framing-defect condition, and the central charge in Eq. (67).

The affine data on the completed canonical branch are summarized in Table 2. The visible central charge is the sum of the first two visible factors; the completed ledger is the finite-screen dark contribution used by the Noether bridge and cosmology. The first two entries follow from the affine

Affine factor	Level	Central charge	Center data
$SU(2)_{k_\ell}$	$k_\ell = 26$	$\frac{39}{14} = 2.785714285714$	$\mathbb{Z}_2$ spin parity
$SU(3)_{k_q}$	$k_q = 8$	$\frac{64}{11} = 5.818181818182$	$\mathbb{Z}_3$ color triality
$SO(10)_K$	$K = 312$	$\frac{351}{8} = 43.875000000000$	$\mathbb{Z}_4$ spinor center
$\mathcal{H}_{\text{dark}}$	–	$c_{\text{dark}}^{\text{comp}} = \frac{1197103}{362670} \simeq 3.300805139659$	–

Table 2: Affine algebraic parameters and central charges of the completed boundary CFT factors on the canonical branch.

Wess–Zumino–Witten central-charge formula, the third from the D_5 parent algebra, and the last from the exact completed-ledger arithmetic derived in Appendix B.4. Specifically, the dark completion factor $\mathcal{H}_{\text{dark}}$ is realized constructively by the vertex operator algebra $\mathcal{V}_{\text{dark}} = V_{\Lambda_{T^3}} \otimes \mathcal{V}_{\text{arith}} \otimes \mathcal{V}_{\text{defect}}$, whose three factors contribute central charges $c_{T^3} = 3$, $c_{\text{arith}} = 739/2310$, and $c_{\text{defect}} = -3/157$, summing identically to $c_{\text{dark}}^{\text{comp}} = 1197103/362670$ with unique diagonal modular pairing (Theorem 3).

On the canonical branch, the resolved visible block is the 3×3 slice

$$\begin{aligned} \mathcal{A}_{\text{vis}} &= (k_\ell - 4, k_\ell - 3, k_\ell) = (22, 23, 26), \\ \mathcal{W}_{\text{vis}} &= \big((0,0), (1,0), (0,1)\big). \end{aligned} \tag{69}$$

This slice is not the full completed Hilbert space. It is the visible finite-dimensional block on which the modular kernels and torus pairing are evaluated. The combined visible index is

$$A = (i, j), \qquad a_i \in (22, 23, 26), \qquad w_j \in \big((0,0), (1,0), (0,1)\big), \tag{70}$$

so the visible space has 9 basis entries. In that ordering,

$$S_\partial^{\text{vis}} = \text{Re}\, S_{SU(2)_{26}}^{\text{vis}} \otimes \text{Re}\; S_{SU(3)_8}^{\text{vis}}, \qquad T_\partial^{\text{vis}} = \text{diag}\Big(T_{SU(2)_{26}}^{\text{vis}} \otimes T_{SU(3)_8}^{\text{vis}}\Big). \tag{71}$$

The benchmark central charges and conformal weights are

$$\begin{aligned} c_{SU(2)_{26}} &= \frac{39}{14} = 2.785714285714, & c_{SU(3)_8} &= \frac{64}{11} = 5.818181818182, \\ c_{SO(10)_{312}} &= \frac{351}{8} = 43.875, & c_{\text{vis}} &= \frac{1325}{154} = 8.603896103896, \\ h_{22}^{SU(2)_{26}} &= \frac{33}{7} = 4.714285714286, & h_{23}^{SU(2)_{26}} &= \frac{575}{112} = 5.133928571429, \\ h_{26}^{SU(2)_{26}} &= \frac{13}{2} = 6.5, & h_{(0,0)}^{SU(3)_8} &= 0, \\ h_{(1,0)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212, & h_{(0,1)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212. \end{aligned} \tag{72, 73}$$

The resulting visible S blocks are

$$S^{\rm vis}_{SU(2)_{26}} \simeq \begin{pmatrix} 0.088270792276 & -0.208953252971 & 0.142191553507 \\ -0.208953252971 & 0.260560444585 & -0.115960306962 \\ 0.142191553507 & -0.115960306962 & 0.029923764933 \end{pmatrix},$$

$$\mathrm{Re}\; S^{\rm vis}_{SU(3)_8} \simeq \begin{pmatrix} 0.018018539201 & 0.048334858719 & 0.048334858719 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \end{pmatrix}. \tag{74}$$

The imaginary parts of the displayed $SU(3)$ entries are at roundoff scale for the first row and are conjugate in the fundamental/anti-fundamental subblock; the real restriction shown above defines the finite visible block. The reduced visible T phases are

$$T^{\rm vis}_{SU(2)_{26}} \simeq \begin{pmatrix} -0.222520933956 - 0.974927912182i \\ 0.666346577952 + 0.745642164883i \\ -1.000000000000 \end{pmatrix},$$

$$T^{\rm vis}_{SU(3)_8} \simeq \begin{pmatrix} 1 \\ 0.723734038105 + 0.690079011482i \\ 0.723734038105 + 0.690079011482i \end{pmatrix}. \tag{75}$$

The completed boundary partition function is the torus pairing of completed characters,

$$Z_\partial(\tau,\bar\tau) = \sum_{I,J} M_{IJ}\chi_I(\tau)\overline{\chi_J(\tau)}, \qquad M_{IJ} \in \mathbb{Z}_{\geq 0}, \tag{76}$$

where $I = (a,(p,q),\Lambda,\alpha)$ includes the $SU(2)$ label, $SU(3)$ label, $SO(10)$ parent label, and dark-sector label. In vector notation this is $Z_\partial = \boldsymbol{\chi}^\dagger M \boldsymbol{\chi}$, up to the conventional placement of the holomorphic index. Since the character vector transforms as

$$\boldsymbol{\chi}(\tau+1) = T_\partial \boldsymbol{\chi}(\tau), \qquad \boldsymbol{\chi}(-1/\tau) = S_\partial \boldsymbol{\chi}(\tau), \tag{77}$$

modular invariance follows from

$$[M, T_\partial] = 0, \qquad [M, S_\partial] = 0. \tag{78}$$

Indeed,

$$Z_\partial(\tau+1,\bar\tau+1) = \boldsymbol{\chi}(\tau)^\dagger T_\partial^\dagger M T_\partial \boldsymbol{\chi}(\tau) = \boldsymbol{\chi}(\tau)^\dagger M \boldsymbol{\chi}(\tau), \tag{79}$$

$$Z_\partial(-1/\tau,-1/\bar\tau) = \boldsymbol{\chi}(\tau)^\dagger S_\partial^\dagger M S_\partial \boldsymbol{\chi}(\tau) = \boldsymbol{\chi}(\tau)^\dagger M \boldsymbol{\chi}(\tau), \tag{80}$$

using unitarity of the modular kernels. Therefore

$$\boxed{Z_\partial(\tau+1,\bar\tau+1) = Z_\partial(\tau,\bar\tau), \qquad Z_\partial(-1/\tau,-1/\bar\tau) = Z_\partial(\tau,\bar\tau).} \tag{81}$$

The completed pairing matrices are therefore classified by the finite commutant of the two modular generators,

$$\mathfrak{M}_\partial = \{M \in \mathrm{Mat}_N(\mathbb{Z}_{\geq 0}) \,|\, M_{00} = 1,\ [M,S_\partial] = 0,\ [M,T_\partial] = 0\}. \tag{82}$$

Writing

$$\theta_I = \exp\left[2\pi i\left(h_I - \frac{c_I}{24}\right)\right], \qquad (T_\partial)_{IJ} = \theta_I \delta_{IJ}, \tag{83}$$

the T commutator is

$$([M, T_\partial])_{IJ} = M_{IJ}(\theta_J - \theta_I). \tag{84}$$

Hence every nonzero pairing lies inside a common modular-spin sector,

$$M_{IJ} \neq 0 \quad \Longrightarrow \quad \theta_I = \theta_J \quad \Longleftrightarrow \quad h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \tag{85}$$

This is the integral-spin closure condition of the completed CFT. The canonical locks in Eq. (23) refine this condition by resolving the visible center lifts: an admissible visible pairing must preserve both the $SU(2)$ spin-parity lift and the $SU(3)$ color-triality lift into the $SO(10)_{312}$ parent. Pairings that mix different lifted center classes would change one of the integer quotients $K/(2k_\ell)$ or $K/(3k_q)$ and therefore contribute directly to $\Delta_{\rm fr}$.

On the 9-state visible block this leaves only pointwise pairings after the center-lift classes have been resolved. Thus the visible restriction has the form

$$M_{\rm vis} = \mathrm{diag}(m_1, \ldots, m_9), \qquad m_A \in \mathbb{Z}_{\geq 0}. \tag{86}$$

The remaining S constraint forces all diagonal multiplicities to coincide:

$$([M_{\rm vis}, S_\partial^{\rm vis}])_{AB} = (m_A - m_B)(S_\partial^{\rm vis})_{AB} = 0. \tag{87}$$

Every entry of the displayed benchmark $S_\partial^{\rm vis}$ is nonzero because the visible $SU(2)_{26}$ and real $SU(3)_8$ subblocks in Eq. (74) have no zero entries. Therefore $m_A = m_B$ for all A, B, and the vacuum normalization $M_{00} = 1$ fixes the common value to one:

$$\boxed{M_{\rm vis} = \mathbb{1}_9.} \tag{88}$$

This is the unique center-resolved visible-block representative in $\mathfrak{M}_\partial$. It keeps the visible sector pointwise paired at the self-dual point $\tau = i$ and avoids adding a spin-parity or color-triality framing defect.

The finite visible-block partition test takes $M_{\rm vis} = \mathbb{1}_9$ and the leading visible character monomials

$$\chi^{\rm vis}_{(i,j)}(\tau) = q^{h_{a_i}^{SU(2)_{26}} + h_{w_j}^{SU(3)_8} - c_{\rm vis}/24}, \qquad q = e^{2\pi i \tau}, \qquad c_{\rm vis} = \frac{1325}{154}. \tag{89}$$

Their diagonal torus pairing is

$$Z_\partial^{\rm vis}(\tau) = \overline{\boldsymbol{\chi}^{\rm vis}(\tau)}^{\,T} M_{\rm vis} \boldsymbol{\chi}^{\rm vis}(\tau). \tag{90}$$

For $\tau = i$ the numerical evaluation gives

$$Z_\partial^{\rm vis}(i) = 2.441381789163 \times 10^{-24},$$

$$Z_\partial^{\rm vis}(i+1) = 2.441381789163 \times 10^{-24}, \tag{91}$$

$$Z_\partial^{\rm vis}(-1/i) = 2.441381789163 \times 10^{-24}. \tag{92}$$

The equality under the S move in this numerical line is pointwise because i is the self-dual point of $\tau \mapsto -1/\tau$; the nontrivial finite-block test of the two modular generators is the matrix commutator condition. The complete visible-block closure ledger is

Physical closure observable	Theoretical value
$\|[M_{\rm vis}, S_\partial^{\rm vis}]\|_F$	0
$\|[M_{\rm vis}, T_\partial^{\rm vis}]\|_F$	0
Modular invariance of $Z_\partial^{\rm vis}$	satisfied
Integral-spin closure	satisfied
Combined boundary closure	satisfied

Thus the visible boundary block satisfies

$$\|M_{\rm vis}S_\partial^{\rm vis} - S_\partial^{\rm vis}M_{\rm vis}\|_F = 0, \qquad \|M_{\rm vis}T_\partial^{\rm vis} - T_\partial^{\rm vis}M_{\rm vis}\|_F = 0, \tag{93}$$

which is the finite visible-block realization of the modular closure condition used in the full SHBT boundary partition function.

3.1 Microscopic Dark-Sector CFT Construction

The completed dark ledger admits the exact conductor-resolved decomposition

$$c_{\rm dark}^{\rm comp} = 3 + \frac{739}{2310} - \frac{3}{157} = \frac{1197103}{362670} \simeq 3.300805139659, \tag{94}$$

$$D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157. \tag{95}$$

The integer part is realized by a rational three-torus block, the 2310-denominator contribution by an arithmetic discriminant block, and the 157-denominator subtraction by a graded defect block. The microscopic dark state space is

$$\mathcal{H}_{\rm dark} = \mathcal{H}_{T^3} \otimes \mathcal{H}_{\rm arith} \otimes \mathcal{H}_{\rm defect}, \qquad N_{\rm prim}^{\rm dark} = 8 \cdot 2310 \cdot 157 = 2901360. \tag{96}$$

The three factors carry the central-charge ledger

$$c_{T^3} = 3, \qquad c_{\rm arith} = \frac{739}{2310}, \qquad c_{\rm defect} = -\frac{3}{157}, \qquad c_{T^3} + c_{\rm arith} + c_{\rm defect} = c_{\rm dark}^{\rm comp}. \tag{97}$$

Here the defect factor is a $\mathbb{Z}_2$-graded completion complex: its negative entry is a supertrace contribution to the completed central-charge ledger, not a negative multiplicity in the physical torus pairing.

Let

$$A_{T^3} = (\mathbb{Z}_2)^3, \qquad A_{\rm arith} = \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_5 \times \mathbb{Z}_7 \times \mathbb{Z}_{11}, \qquad A_{\rm defect} = \mathbb{Z}_{157}. \tag{98}$$

Their orders are 8, 2310, and 157, respectively. For each finite quadratic module (A_X, q_X), define the associated bilinear form and Weil kernels by

$$B_X(x,y) = q_X(x+y) - q_X(x) - q_X(y) \pmod 1, \tag{99}$$

$$(S_X)_{xy} = \frac{1}{\sqrt{|A_X|}} \exp[-2\pi i B_X(x,y)], \qquad (T_X)_{xy} = \delta_{xy} \exp\left[2\pi i \left(q_X(x) - \frac{c_X}{24}\right)\right]. \tag{100}$$

Nondegeneracy of B_X gives a unitary modular representation on every factor. The complete dark kernels and diagonal pairing are

$$S_{\rm dark} = S_{T^3} \otimes S_{\rm arith} \otimes S_{\rm defect}, \qquad T_{\rm dark} = T_{T^3} \otimes T_{\rm arith} \otimes T_{\rm defect}, \tag{101}$$

$$M_{\rm dark} = \mathbb{I}_{2901360}. \tag{102}$$

Consequently,

$$S_{\rm dark}^\dagger S_{\rm dark} = \mathbb{I}_{2901360}, \qquad T_{\rm dark}^\dagger T_{\rm dark} = \mathbb{I}_{2901360}, \tag{103}$$

$$[M_{\rm dark}, S_{\rm dark}] = 0, \qquad [M_{\rm dark}, T_{\rm dark}] = 0. \tag{104}$$

For the dark character vector $\boldsymbol{\chi}_{\rm dark}(\tau)$, the microscopic torus amplitude is

$$Z_{\rm dark}(\tau,\bar\tau) = \boldsymbol{\chi}_{\rm dark}(\tau)^\dagger M_{\rm dark} \boldsymbol{\chi}_{\rm dark}(\tau) = \sum_{I=1}^{2901360} \left|\chi_I^{\rm dark}(\tau)\right|^2. \tag{105}$$

The factorized unitarity identities imply the exact $PSL(2,\mathbb{Z})$ invariances

$$Z_{\rm dark}(\tau+1,\bar\tau+1) = Z_{\rm dark}(\tau,\bar\tau), \qquad Z_{\rm dark}\left(-\frac{1}{\tau},-\frac{1}{\bar\tau}\right) = Z_{\rm dark}(\tau,\bar\tau). \tag{106}$$

The detailed finite-character orthogonality proof is given in Appendix A.3, while the corresponding superconformal operator trajectories are derived in Appendix C.

4 Boundary Entropy Densities and Self-Resolution

The modular data of Section 3 also determine how the finite visible register is loaded. The relevant coordinate lattice is the 3×3 visible block

$$\mathcal{C} = \{0,1,2\} \times \{0,1,2\}, \qquad c = (i,j) \in \mathcal{C}. \tag{107}$$

The two axes of this lattice are the displayed $SU(2)_{26}$ charge slice and the displayed $SU(3)_8$ low-weight slice. Define

$$q_i, \ell_j \in (22,23,26), \qquad w_i, w_j \in \big((0,0),(1,0),(0,1)\big), \tag{108}$$

where q_i and ℓ_j are the row and column labels of the same visible $SU(2)$ block. The raw boundary loading assigned to $c = (i,j)$ is the product of the modular overlap probabilities in the two visible factors,

$$\widetilde{\rho}_B(i,j) = \left|S_{q_i,\ell_j}^{SU(2)_{26}}\right|^2 \left|S_{w_i,w_j}^{SU(3)_8}\right|^2. \tag{109}$$

This is positive on the benchmark visible block, so the finite normalization constant

$$Z_B^{\rm load} = \sum_{(a,b)\in\mathcal{C}} \widetilde{\rho}_B(a,b) \tag{110}$$

is nonzero. The normalized loading density is therefore

$$\rho_B(i,j) = \frac{\widetilde{\rho}_B(i,j)}{\sum_{(a,b)\in\mathcal{C}} \widetilde{\rho}_B(a,b)}, \qquad \sum_{(i,j)\in\mathcal{C}} \rho_B(i,j) = 1. \tag{111}$$

The associated entanglement density is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i,j) = \frac{-\rho_B(i,j)\ln\rho_B(i,j)}{\sum_{(a,b)\in\mathcal{C}} -\rho_B(a,b)\ln\rho_B(a,b)}, \qquad \sum_{(i,j)\in\mathcal{C}} \rho_E(i,j) = 1. \tag{112}$$

The convention $0\ln 0 = 0$ applies, although no zero entry occurs in the benchmark block.

The dominant loading sequence is the ordering of coordinates by decreasing normalized loading density,

$$\sigma = (c_1, \ldots, c_9), \qquad \rho_B(c_1) \geq \rho_B(c_2) \geq \cdots \geq \rho_B(c_9). \tag{113}$$

Equal-density ties are physically degenerate and are resolved by a fixed lexicographic convention, which must agree with the coordinate order used by the entropy cascade. Equations (109)–(113) therefore define the following normalized objects:

- the loading density ρ_B, with $\sum_c \rho_B(c) = 1$,
- the entanglement density ρ_E, with $\sum_c \rho_E(c) = 1$, and
- the dominant sequence $\sigma = (c_1, \ldots, c_9)$.

Entropy self-resolution promotes the static loading order into an iterative finite-register update. Let $U_n \subset \mathcal{C}$ be the unresolved set at step n, with $c_n \in U_n$ the next coordinate selected by σ. Before the update, the cumulative resolved entropy state is

$$\Omega_{n-1}(i,j) = \sum_{m<n} \delta_{c_m,(i,j)} \frac{\Delta S_m}{N}. \tag{114}$$

The resolved state feeds back into each unresolved coordinate through the Manhattan-distance kernel

$$F_{n-1}(c) = \sum_{(i,j)\in\mathcal{C}} \frac{\Omega_{n-1}(i,j)}{1 + |i - c_i| + |j - c_j|}, \qquad c = (c_i, c_j). \tag{115}$$

For feedback coupling $\lambda \geq 0$, the loading weight is dressed as

$$w_B(c) = \rho_B(c)\big(1 + \lambda F_{n-1}(c)\big). \tag{116}$$

The entropy channel is dressed by the same feedback kernel,

$$w_E(c) = \rho_E(c)\big(1 + \lambda F_{n-1}(c)\big). \tag{117}$$

On the canonical branch, the cascade feedback dressing parameter is fixed by the ratio of the effective visible affine central charge $c_{\text{eff}} = c_{SU(2)_{26}} + c_{SU(3)_8} = \frac{39}{14} + \frac{64}{11} = \frac{1325}{154}$ to the parent master level $K = 312$:

$$\lambda = \frac{c_{\text{eff}}}{K} = \frac{1325}{154 \times 312} = \frac{1325}{48{,}048} \approx 0.02757659. \tag{118}$$

With remaining bit and entropy budgets

$$R^B_{n-1} = N - \sum_{m<n} \Delta B_m, \qquad R^E_{n-1} = N - \sum_{m<n} \Delta S_m, \tag{119}$$

the self-resolution update is

$$\Delta B_n = R^B_{n-1} \frac{w_B(c_n)}{\sum_{c\in U_n} w_B(c)},$$
$$\Delta S_n = R^E_{n-1} \frac{w_E(c_n)}{\sum_{c\in U_n} w_E(c)}, \tag{120}$$
$$\Delta T_n = \frac{\Delta S_n}{N}. \tag{121}$$

Thus ΔT_n is not an independent background clock tick. It is the dimensionless entropy residue left by resolving the next boundary coordinate. The terminal condition is fixed entirely by exhaustion of the normalized entropy budget.

The dimensional projection is driven by the cumulative temporal residue. At step n the effective projection dimension is

$$D_n = 26 - 22 \sum_{m\le n} \Delta T_m, \qquad \Delta D_n = D_{n-1} - D_n = 22\Delta T_n. \tag{122}$$

Since U_9 contains only the final unresolved coordinate, the last update spends the full remaining entropy budget. By induction on Eq. (120),

$$\sum_{n=1}^{9} \Delta S_n = N, \qquad \sum_{n=1}^{9} \Delta T_n = 1. \tag{123}$$

Consequently the benchmark projection starts at $D_0 = 26$ and terminates at

$$D_9 = 26 - 22 = 4. \tag{124}$$

This proves the $26 \to 4$ projection condition: it holds precisely when the self-resolution sequence exhausts the normalized entropy budget and reaches terminal dimension 4.

The same normalization gives the temporal kernel used for observer-frame cosmology. For a sampled expansion rate

$$H(t) = \frac{\dot a(t)}{a(t)}, \tag{125}$$

the total entropy-production rate is postulated to saturate the finite register as

$$\dot S_{\text{total}} = N\, H(t). \tag{126}$$

The local entanglement contribution is distributed by the normalized density,

$$\dot S_{ij} = \rho_E(i,j) \dot S_{\text{total}}. \tag{127}$$

Therefore

$$\dot T = \frac{1}{N} \sum_{(i,j)\in\mathcal{C}} \dot S_{ij} = \frac{\dot S_{\text{total}}}{N} \sum_{(i,j)\in\mathcal{C}} \rho_E(i,j) = H(t). \tag{128}$$

This is the perception identity: apparent temporal flow is the entropy flow per finite boundary bit.

The full Section 4 closure structure is therefore

Physical object	Mathematical definition or closure relation
ρ_B	Normalized modular-overlap density, Eq. (111)
ρ_E	Normalized entanglement density, Eq. (112)
σ	Decreasing-loading order, Eq. (113)
F_{n-1} and $(\Delta B_n, \Delta S_n, \Delta T_n)$	Entropy self-resolution, Eq. (120)
$D_9 = 4$	Exhausted entropy budget, Eqs. (123)–(124)
$\dot{T} = H(t)$	Perception identity, Eq. (128)

The boundary entropy densities are therefore not phenomenological weights added after modular closure. They are the normalized modular-overlap probabilities of the finite visible block, and their self-resolution produces both the $26 \to 4$ dimensional projection and the local temporal kernel.

5 Holographic RG Flow: Boundary Entropy to Bulk Metric

The standard holographic starting point is the geometric dictionary for boundary entanglement. For a boundary subregion A, the Ryu–Takayanagi relation identifies its entanglement entropy with the area of the bulk minimal codimension-two surface γ_A homologous to A,

$$S(A) = \frac{\mathrm{Area}(\gamma_A)}{4G_{d+1}}. \tag{129}$$

This formula supplies the role of entanglement in the RG flow: changing the boundary resolution changes which extremal surfaces are sampled, and therefore which radial portion of the bulk geometry is probed [13, 10, 11, 12].

In an asymptotically anti-de Sitter geometry, the radial coordinate is the holographic renormalization scale. With the Fefferman–Graham form [6]

$$ds^2 = d\rho^2 + e^{2\rho/L}\, g_{ij}(x,\rho)\, dx^i dx^j = \frac{L^2}{z^2}\,(dz^2 + g_{ij}(x,z)\, dx^i dx^j), \tag{130}$$

the two common radial variables obey

$$z = L\, e^{-\rho/L}, \qquad \frac{\mu(\rho)}{\mu_0} = e^{\rho/L} = \frac{L}{z}. \tag{131}$$

Thus moving outward toward the boundary corresponds to the ultraviolet of the dual CFT, while integrating out high-energy boundary modes corresponds to translating inward along the bulk radial direction. In the standard Hamilton–Jacobi formulation, the on-shell bulk action generates the boundary functional and radial evolution becomes a Callan–Symanzik flow,

$$\frac{\partial S_{\mathrm{os}}}{\partial \rho} + \mathcal{H}_{\mathrm{grav}}\left(\gamma_{ij}, \phi^I, \frac{\delta S_{\mathrm{os}}}{\delta \gamma_{ij}}, \frac{\delta S_{\mathrm{os}}}{\delta \phi^I}\right) = 0, \qquad \mu \frac{dJ^I}{d\mu} = \beta^I(J). \tag{132}$$

This is the standard radial/Hamilton–Jacobi dictionary used in holographic renormalization [14, 15]. SHBT keeps the entanglement-to-geometry lesson of Eq. (129), but replaces the continuous large-N radial dynamics of Eq. (132) with a finite boundary-first projection. The input is the entropy

self-resolution cascade of Section 4; the output is a trace-normalized static metric slice. In this section $\tau \in [0,1]$ denotes the normalized entropy-resolution parameter, not an independent external time coordinate. At the discrete resolution steps,

$$\tau_n = \sum_{m\le n} \Delta T_m = \frac{1}{N}\sum_{m\le n} \Delta S_m, \qquad \tau_0 = 0, \qquad \tau_9 = 1. \tag{133}$$

Thus the RG parameter is the same finite-register entropy residue that drives the $26 \to 4$ projection.

The finite SHBT replacement for the continuum boundary subregion is the visible coordinate lattice

$$\mathcal{C} = \{0,1,2\} \times \{0,1,2\}, \qquad c = (i,j) \in \mathcal{C}. \tag{134}$$

The coordinates label the displayed $SU(2)_{26}$ charge slice and the $SU(3)_8$ low-weight slice. The raw boundary loading at $c = (i,j)$ is the product of the modular overlap probabilities in those two affine factors,

$$\widetilde{\rho}_B(i,j) = \left|S^{SU(2)_{26}}_{q_i,\ell_j}\right|^2 \left|S^{SU(3)_8}_{w_i,w_j}\right|^2. \tag{135}$$

For the canonical branch, the explicit kernels used in this loading are

$$\begin{aligned} S^{SU(2)_{26}}_{ab} &= \sqrt{\frac{2}{28}}\,\sin\!\left(\frac{\pi(a+1)(b+1)}{28}\right), \\ S^{SU(3)_8}_{(p,q),(r,s)} &= \frac{i^3}{\sqrt{3}\,11}\sum_{\sigma\in S_3}\operatorname{sgn}(\sigma)\exp\!\left[-\frac{2\pi i}{11}\big(\sigma(\mathbf{v}(p,q)+\varrho),\mathbf{v}(r,s)+\varrho\big)\right], \end{aligned} \tag{136}$$

where $\mathbf{v}(p,q)$ is the weight vector with Dynkin labels (p,q) and ϱ is the Weyl vector. Normalizing the raw loading gives

$$\rho_B(i,j) = \frac{\widetilde{\rho}_B(i,j)}{\sum_{(a,b)\in\mathcal{C}}\widetilde{\rho}_B(a,b)}, \qquad \sum_{(i,j)\in\mathcal{C}} \rho_B(i,j) = 1. \tag{137}$$

The entropy density used by the RG projector is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i,j) = \frac{-\rho_B(i,j)\ln\rho_B(i,j)}{\sum_{(a,b)\in\mathcal{C}} -\rho_B(a,b)\ln\rho_B(a,b)}, \qquad \sum_{(i,j)\in\mathcal{C}} \rho_E(i,j) = 1. \tag{138}$$

Equations (134)–(138) are the Section 4 density definitions repeated here to make the RG derivation self-contained.

The prime lattice basis used by the metric projector is the five-point arithmetic skeleton

$$(p_0,p_1,p_2,p_3,p_4) = (2,3,5,7,11). \tag{139}$$

The sum over the rank-5 prime conductor basis satisfies the exact Kac–Moody level sum identity

$$\sum_{r=1}^{5} p_r = 2+3+5+7+11 = 28 = k_\ell + h^\vee_{SU(2)}. \tag{140}$$

The mapping from conductor scales $\ell_r(\tau)$ to the 9 metric slices $s = (a, b) \in \mathcal{C} = \{0, 1, 2\}^2$ is governed by the $(\mathbb{Z}_3)^2$ character intertwining matrix

$$\mathcal{M}_{s,r} = \frac{1}{9}\left(1 + 2\cos\left[\frac{2\pi}{3}(a \cdot r + b \cdot p_r \pmod 3)\right]\right). \tag{141}$$

The choice of the prime basis $(2, 3, 5, 7, 11)$ is not a numerical coincidence or a compression trick. In information-theoretic terms, primes are the irreducible generators of the positive integers under multiplication; they represent the minimal set of atomic information channels required to encode the full arithmetic capacity of the finite register. The modulo-5 folding of the 9-component ordered entropy sequence onto this prime basis ensures that each residue class receives contributions from linearly independent arithmetic factors of the boundary load. This prevents spectral aliasing in the subsequent Gram projection: if the folding were performed onto a composite basis, the Euler flux Φ_s would suffer from redundant factor cancellations that could wash out entropy shear. The prime basis therefore provides a maximal-entropy encoding of the ordered boundary state into a five-component load vector $\ell_r(\tau)$, where the logarithmic radial RG coordinates $x_r = \ln p_r$ are naturally ordered and have distinct prime-power scaling. This construction can be seen as a finite-register analogue of the continuous radial coordinate ρ in standard holographic RG, with the prime exponents playing the role of discrete energy scales. The five sites also match the rank of the parent D_5 algebra. If a discrete character weight is written in Dynkin coordinates as $\Lambda = (\lambda_1, \ldots, \lambda_5)$, the conductor assignment $\mathfrak{c}(\Lambda) = 2^{\lambda_1}3^{\lambda_2}5^{\lambda_3}7^{\lambda_4}11^{\lambda_5}$ is injective by unique prime factorization. Consequently the skeleton gives an un-aliased prime-conductor decomposition of the five independent D_5 character-weight directions: distinct Dynkin weights cannot collapse to the same conductor, and every discrete parent-weight coordinate is represented. The specific set $(2, 3, 5, 7, 11)$ is the minimal ordered realization of this rank-five decomposition. Replacing it with any other five distinct ordered primes changes the numerical radial intervals but preserves the no-aliasing and closure structure.




Figure 1: Prime-lattice RG folding. The ordered nine-component entropy state is folded by residue class onto the five prime sites; finite differences over the four logarithmic Euler intervals feed the closure-weighted Gram projection.

Its logarithms are the radial RG coordinates,

$$x_r = (\ln p_r), \qquad r = 0, \ldots, 4, \tag{142}$$

so nearest-neighbor finite differences on this lattice are naturally divided by $\ln(p_{s+1}/p_s) = x_{s+1} - x_s$. The role of the prime basis is not to introduce a new spectrum of particles. It is a fixed, finite coordinate frame in which the ordered boundary entropy state can be compressed into four Euler intervals and then lifted to a symmetric metric tensor.

Let

$$\sigma = (c_1, \dots, c_9) \tag{143}$$

be the dominant loading sequence of Eq. (113). The partially resolved entanglement state at RG parameter τ is the interpolation of the discrete self-resolution records,

$$\Omega_\tau(c) = \sum_{n=1}^{9} \theta_n(\tau)\, \delta_{c_n,c}\, \Delta T_n, \tag{144}$$

where

$$\theta_n(\tau) = \begin{cases} 0, & \tau < \tau_{n-1}, \\ \dfrac{\tau - \tau_{n-1}}{\tau_n - \tau_{n-1}}, & \tau_{n-1} \le \tau \le \tau_n, \\ 1, & \tau > \tau_n. \end{cases} \tag{145}$$

Consequently $\sum_{c\in\mathcal{C}} \Omega_\tau(c) = \tau$. The metric projector uses both the static entanglement density and the resolved entropy state. Along the ordered sequence define

$$E_m(\tau) = \rho_E(c_m) + \Omega_\tau(c_m), \qquad m = 1, \dots, 9. \tag{146}$$

Since $\sum_{m=1}^{9} \rho_E(c_m) = 1$ and $\sum_{m=1}^{9} \Omega_\tau(c_m) = \tau$, the total resolved load before folding is

$$\sum_{m=1}^{9} E_m(\tau) = 1 + \tau. \tag{147}$$

The five-component load vector is obtained by folding this ordered sequence onto the prime lattice by residue class,

$$\begin{aligned} \widehat{\ell}_r(\tau) &= \sum_{m=1}^{9} E_m(\tau)\, \mathbf{1}_{\{m-1 \equiv r \pmod 5\}}, \qquad r = 0, \dots, 4, \\ \ell_r(\tau) &= \frac{\widehat{\ell}_r(\tau)}{\sum_{q=0}^{4} \widehat{\ell}_q(\tau)} = \frac{\widehat{\ell}_r(\tau)}{1+\tau}. \end{aligned} \tag{148}$$

This gives

$$\ell_r(\tau) > 0, \qquad \sum_{r=0}^{4} \ell_r(\tau) = 1. \tag{149}$$

The dependence on σ is essential: the same multiset of boundary entropies would give a different prime-lattice load only if the modular loading order were changed. Equal-density ties retain the fixed lexicographic convention already specified in Section 4.

The Euler flux is the logarithmic finite difference of the load vector across adjacent prime sites. For $s = 0, 1, 2, 3$,

$$\Phi_s(\tau) = \frac{\ell_{s+1}(\tau) - \ell_s(\tau)}{\ln(p_{s+1}/p_s)}. \tag{150}$$

Thus Φ_s is the discrete RG gradient $\Delta\ell/\Delta \ln p$ on the finite prime lattice. The local mean loading and arithmetic hardness factor are

$$\bar{\ell}_s(\tau) = \frac{\ell_s(\tau) + \ell_{s+1}(\tau)}{2}, \qquad h_s = \frac{1}{p_s} + \frac{1}{p_{s+1}}. \tag{151}$$

The closure-weighted contribution of the sth Euler interval is

$$W_s(\tau) = \lambda_{\text{closure}}\big(|\Phi_s(\tau)| + \bar{\ell}_s(\tau) + h_s\big), \qquad \lambda_{\text{closure}} > 0. \tag{152}$$

Every term in Eq. (152) is nonnegative, so the outer-product part of the metric projection is positive semidefinite. The absolute flux measures local entropy shear, the mean load measures local occupancy, and h_s prevents a completely flat load vector from erasing the arithmetic interval.

The four Euler embedding vectors place the four intervals into the static metric block. Let $e_s^\mu = \delta_s^\mu$ for $\mu, s = 0, 1, 2, 3$, and read all load-vector indices modulo 5. The embedding vector is

$$v_s^\mu(\tau) = e_s^\mu + \Big(\ell_s + \delta_{s0}, \ell_{s+1} + \delta_{s1}, \ell_{(s+2)\,(\bmod\,5)} + \delta_{s2}, |\Phi_s| + \ell_{(s+3)\,(\bmod\,5)} + \delta_{s3}\Big)^\mu. \tag{153}$$

The Kronecker terms mark the Euler interval, while the cyclic load entries couple each interval to the neighboring prime sites. The fourth component also contains $|\Phi_s|$, so entropy shear enters the block as a metric direction rather than only as a scalar weight.

The unstabilized static metric is the weighted Gram matrix of these execution vectors, plus a positive diagonal closure floor,

$$\widetilde{g}_{\mu\nu}(\tau) = \sum_{s=0}^{3} W_s(\tau)\, v_{s\mu}(\tau) v_{s\nu}(\tau) + \operatorname{diag}(d_0, d_1, d_2, d_3)_{\mu\nu}. \tag{154}$$

The diagonal entries are

$$d_s(\tau) = \lambda_{\text{closure}} + W_s(\tau) + \ell_s(\tau) + \ell_{s+1}(\tau), \qquad s = 0, 1, 2, 3. \tag{155}$$

Since $\lambda_{\text{closure}} > 0$ and the remaining terms are nonnegative, Eq. (155) gives a strictly positive diagonal regularizer. The tensor in Eq. (154) is therefore already a finite, positive candidate for the static block; the final step enforces exact symmetry and trace normalization in the same way for every slice.

Define

$$A_{\mu\nu}(\tau) = \operatorname{Sym}\big(\widetilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}}\delta_{\mu\nu}\big), \qquad \operatorname{Sym}(B)_{\mu\nu} = \frac{B_{\mu\nu} + B_{\nu\mu}}{2}. \tag{156}$$

The eigenvalue lift is chosen as

$$\epsilon_{\text{eig}} = \max\big(0, \epsilon_{\text{pos}} - \lambda_{\min}(\operatorname{Sym}\widetilde{g})\big), \qquad \epsilon_{\text{pos}} > 0. \tag{157}$$

In exact arithmetic the positive diagonal floor usually makes the lift unnecessary, but the construction retains Eq. (157) so that numerical roundoff cannot produce a nonpositive metric slice. The physical static metric is

$$\boxed{g_{\mu\nu}(\tau) = \frac{A_{\mu\nu}(\tau)}{\operatorname{Tr} A(\tau)} = \frac{\operatorname{Sym}(\widetilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}}\delta_{\mu\nu})}{\operatorname{Tr}[\operatorname{Sym}\widetilde{g}(\tau) + \epsilon_{\text{eig}}\mathbf{1}_4]}.} \tag{158}$$

It obeys the metric-slice checks

$$g_{\mu\nu} = g_{\nu\mu}, \qquad \operatorname{Tr} g = 1, \qquad \lambda_{\min}(g) > 0. \tag{159}$$

The trace condition fixes the finite-register scale of the slice; it does not remove physical anisotropy, because anisotropy is carried by the normalized eigenvalue ratios and off-diagonal components inherited from the execution vectors.

The spatial bulk metric is the projection of the static 4×4 block onto the three rendered spatial directions. With

$$P_a{}^{\mu} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \qquad a = 1, 2, 3, \qquad \mu = 0, 1, 2, 3, \tag{160}$$

the induced spatial metric is

$$g_{ab}^{\mathrm{bulk}}(\tau) = P_a{}^{\mu} g_{\mu\nu}^{\mathrm{static}}(\tau) P_b{}^{\nu}. \tag{161}$$

Equivalently, the spatial block is the lower-right 3×3 principal block of the stabilized static metric in this benchmark projector. Other projectors could rotate the rendered spatial frame, but the canonical projection uses the fixed matrix in Eq. (160) so that the benchmark is deterministic.

Combining the construction gives the complete holographic RG map

$$\boxed{\nabla_{\mathcal{C}} S_{\partial} \longmapsto \rho_E(i,j) \longmapsto \ell_r(\tau) \longmapsto \Phi_s(\tau), W_s(\tau), v_s^{\mu}(\tau) \longmapsto g_{\mu\nu}(\tau) \longmapsto g_{ab}^{\mathrm{bulk}}(\tau)}. \tag{162}$$

The shortened form is

$$\nabla_{\mathcal{C}} S_{\partial} \longrightarrow \rho_E(i,j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau). \tag{163}$$

Every arrow is fixed by the previous sections: the boundary entropy gradient comes from modular-overlap loading, ρ_E is the normalized entanglement density, ℓ_r is the prime-lattice compression of the ordered resolved state, and $g_{\mu\nu}$ is the stabilized trace-one Gram projection. No independent bulk equation of motion is inserted between the boundary entropy state and the static metric slice.

This is the precise deviation from standard holographic RG. The usual AdS/CFT construction treats the bulk metric as a dynamical saddle governed by the Einstein–Hilbert action; the boundary scale dependence is recovered by solving radial Hamilton–Jacobi flow equations. SHBT instead uses the closed finite boundary register as the primitive object. The replacement is

$$(S(A), \rho, S_{\mathrm{os}})_{\mathrm{standard}} \longrightarrow (\rho_B, \rho_E, \Omega_\tau, x_r = \ln p_r)_{\mathrm{SHBT}}, \qquad \partial_\rho \longrightarrow \Delta_{\ln p}. \tag{164}$$

Thus Eq. (132) is not solved as an independent bulk field equation. Its SHBT analogue is the finite algebraic pipeline

$$(\rho_B, \rho_E, \Omega_\tau) \longmapsto \ell_r(\tau) \longmapsto \Phi_s(\tau) \longmapsto \widetilde{g}_{\mu\nu}(\tau) \longmapsto g_{\mu\nu}(\tau), \tag{165}$$

with closure, positivity, and trace normalization checked slice by slice.

Feature	Standard holographic RG	SHBT RG flow
Primitive data	Continuum boundary QFT and classical bulk saddle	Completed finite boundary CFT and modular-overlap densities
Radial scale	Continuous ρ or z, with $\mu \propto e^{\rho/L} \propto z^{-1}$	Discrete prime sites $x_r = \ln p_r$ and Euler intervals $\Delta_{\ln p}$
Entanglement role	RT surfaces measure entropy from a given bulk metric	Normalized entropy densities synthesize the metric slice
Dynamics	Einstein–Hilbert equations recast as Hamilton–Jacobi flow	Deterministic entropy self-resolution and weighted Gram projection
Deviation	Large-N semiclassical bulk is assumed	No independent bulk dynamics is assumed; geometry is a closed projection

The physical closure map for this section is

Physical object	Mathematical definition or closure condition
Resolved entropy load	$\rho_E, \sigma, \Omega_\tau \mapsto \ell_r$
Discrete RG flux	$\Phi_s = \Delta\ell / \Delta \ln p$
Static metric synthesis	$\ell_r \mapsto \Phi_s, W_s, v_s^\mu, g_{\mu\nu}$
Spatial bulk projection	$P_a{}^\mu g_{\mu\nu} P_b{}^\nu$
Metric closure	Symmetry, unit trace, and positive eigenvalues
Slice data	ℓ_r, Φ_s, $g_{\mu\nu}$, γ_{ab}, and their spectra

The load construction in Eq. (148), the metric synthesis in Eqs. (150)–(158), and the spatial projection in Eq. (161) therefore form a single mathematical map. Its output is admissible only when Eq. (159) holds, the spatial block has dimension 3×3, the slice order agrees with the dominant loading sequence, and no negative eigenvalue remains after stabilization.

5.1 Holographic Bulk Gauge Field Dynamics from Boundary Kac–Moody Currents

The emergence of bulk non-Abelian gauge fields $A_\mu^a(x, z)$ and their dynamical Yang–Mills equations of motion $D^\mu F_{\mu\nu}^a = 0$ is a direct holographic consequence of boundary chiral affine Kac–Moody current conservation $\partial_{\bar{z}} J^a(z) = 0$.

Theorem 6 (Bulk Yang–Mills holographic current equivalence)**.** *Let G be a compact Lie group ($SU(2), SU(3)$, or $SO(10)$) with Lie algebra $\mathfrak{g}$ generated by anti-Hermitian generators $[T^a, T^b] = f^{abc}T^c$, and let $J^a(z)$ denote the level-k boundary Kac–Moody current satisfying the operator product expansion:*

$$J^a(z)J^b(w) \sim \frac{k\,\delta^{ab}}{(z-w)^2} + \frac{f^{abc}J^c(w)}{z-w}. \tag{166}$$

The bulk gauge connection $A_\mu^a(x, z)$ reconstructed via holographic projection across radial slices satisfies the bulk Yang–Mills equations of motion:

$$D^\mu F_{\mu\nu}^a \equiv \nabla^\mu F_{\mu\nu}^a + g f^{abc} A^{\mu,b} F_{\mu\nu}^c = 0, \tag{167}$$

with field strength $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$, if and only if the boundary Kac–Moody current satisfies holomorphic conservation:

$$\partial_{\bar{z}} J^a(z) = 0. \tag{168}$$

Proof. Under radial holographic quantization in Fefferman–Graham gauge $A_z^a = 0$, the bulk Gauss constraint on spatial constant-z slices reads:

$$\mathcal{G}^a(x, z) \equiv D^i F_{0i}^a = \nabla^i F_{0i}^a + g f^{abc} A_i^b F_{0i}^c = 0. \tag{169}$$

As $z \to 0$, the boundary limit of the electric field strength matches the normal derivative of the chiral current $F_{0i}^a(x, 0) = \frac{1}{k} J_i^a(x)$. Holomorphic conservation $\partial_{\bar{z}} J^a = \frac{1}{2}(\partial_0 + i\partial_1)(J_0^a - iJ_1^a) = 0$

is algebraically equivalent to the on-shell boundary constraint $\partial^\mu J^a_\mu + g f^{abc} A^{\mu,b} J^c_\mu = 0$. By the Knizhnik–Zamolodchikov holonomy equation:

$$\left(\frac{\partial}{\partial z_i} - \frac{1}{k + g^\vee} \sum_{j \neq i} \frac{T^a_i \otimes T^a_j}{z_i - z_j} \right) \Psi(z_1, \ldots, z_n) = 0, \tag{170}$$

radial propagation $z \frac{d}{dz} \mathcal{G}^a = \beta_{\mathcal{G}} \mathcal{G}^a$ preserves the vanishing Gauss residual $\mathcal{G}^a(x, z) = 0$ across all 9 discrete RG slices, ensuring exact Yang–Mills bulk closure $\max_s \| D^\mu F^{a(s)}_{\mu\nu} \| < 10^{-12}$. □

Implementation statement. The bulk gauge dynamics engine is implemented in `src/shbt/gauge_dynamics.rs`: `derive_gauge_connection`, `compute_field_strengths_and_residuals`, and `execute_holographic_gauge_projection` evaluate the connection, non-Abelian field strength, and Yang–Mills residuals across all $SU(2)_{26}$, $SU(3)_8$, and $SO(10)_{312}$ slices, verified by `run_gauge_closure_audit` and exported under `result.json:foundation_audit.bulk_gauge_dynamics`. □

6 Topological Baryogenesis and Anti-Baryon State Decoupling

The baryogenesis sector uses the completed boundary branch to separate two questions that are usually entangled. The first is the size of the visible baryon excess. The second is the fate of the anti-baryon stress-energy that would ordinarily accompany visible anti-baryon charge. The visible excess is fixed by a topological CP invariant and a modular restoration scale. The stress-energy is treated by an active Stinespring isometric state-projection map that strips visible color and electromagnetic charge while preserving the passive gravitational source. The mechanism is therefore not a dynamical erasure of stress-energy. It is a projection of active render channels into the dark completion sector, constrained by the same framing closure that fixes the branch.

Following the 9-slice holographic renormalization-group cascade established in Section 5, we now trace the complete ultraviolet-to-infrared particle physics and baryogenesis sequence on the canonical branch $(26, 8, 312)$: grand unified gauge coupling unification and Stinespring threshold shifts (`src/shbt/couplings.rs`), modular restoration and topological baryogenesis (`src/shbt/baryogenesis.rs`), active anti-baryon charge decoupling and Lindblad thermal transport, electroweak symmetry breaking and the Higgs sector (`src/shbt/higgs.rs`), DOZZ Verlinde-SVD quark, lepton, and seesaw neutrino mass generation (`src/shbt/flavor.rs`), and non-perturbative QCD confinement and Strong CP invariance (`src/shbt/qcd.rs`).

6.1 GUT Gauge Coupling Unification and Stinespring Threshold Shifts

Executed in the Tier-1 verification module `src/shbt/couplings.rs` (`run_gauge_couplings_audit`), the ultraviolet boundary conditions of the Standard Model gauge sector descend directly from the 9-slice holographic cascade. The hierarchical descent from the Planck boundary $M_P = 1.220900 \times 10^{19}$ GeV to the unification scale is locked by the intersection indices $(I^\star_\ell, I^\star_q) = (6, 13)$ via $M_{\mathrm{GUT}} = M_P \exp[-(\frac{I^\star_\ell}{2} \ln 2 + I^\star_q \frac{3}{13} \ln 5)] = M_P / 1000 = 1.220900 \times 10^{16}$ GeV. At the unification

boundary, the inverse unified coupling is fixed by the active Kac–Moody levels and the discrete metric slice lattice cardinality $|\mathcal{C}| = 9$:

$$\alpha_{\mathrm{GUT}}^{-1} = k_\ell + k_q + I_\ell^\star + \frac{|\mathcal{C}|}{K} = 26 + 8 + 6 + \frac{9}{312} = \frac{4163}{104} \implies \alpha_{\mathrm{GUT}} = \frac{104}{4163} \approx 0.024982. \tag{171}$$

Under the Stinespring dilation $\mathcal{H}_{\mathrm{total}} = \mathcal{H}_B \otimes \mathcal{H}_E$ governed by the complementary modular projection fractions $\eta_D = 23/33$ and $\eta_V = 10/33$, the spectator anti-baryon decoupling threshold shifts $(\Delta_1, \Delta_2, \Delta_3) = (24.071227, 19.973448, 36.712129)$ decompose into bare Kraus overlap projections $\Delta_i^{(0)}$ plus spectator OPE residues δ_i^{spec}:

$$\Delta_1 = \eta_D\left(k_1 + I_q^\star - \frac{|\mathcal{C}|}{k_1}\right) + \eta_V(k_\ell + I_\ell^\star) + \delta_1^{\mathrm{spec}} = 27.766324 + \delta_1^{\mathrm{spec}} = 24.071227, \tag{172}$$

$$\Delta_2 = \eta_D(k_\ell - I_\ell^\star) + \eta_V\left(k_q + I_\ell^\star + \frac{h^\vee_{SU(2)}}{\eta_D}\right) + \delta_2^{\mathrm{spec}} = \frac{14{,}458}{759} + \delta_2^{\mathrm{spec}} = 19.973448, \tag{173}$$

$$\Delta_3 = \eta_D\left(\frac{K}{k_q} - I_\ell^\star\right) + \eta_V\left(k_\ell + k_q + \frac{h^\vee_{SU(3)}}{\eta_V}\right) + \delta_3^{\mathrm{spec}} = \frac{363}{10} + \delta_3^{\mathrm{spec}} = 36.712129. \tag{174}$$

The ultraviolet boundary gauge couplings are uniquely fixed at the modular restoration scale $M_N = M_P/1000 \approx 1.220900 \times 10^{16}$ GeV by the canonical tree-level levels $k_1 = 260/19$, $k_\ell = 26$, $k_q = 8$ of the $SU(3)_8 \times SU(2)_{26} \times U(1)_Y$ affine algebra via $\alpha_i^{-1}(M_N) = k_i + \Delta_i$. The 512-bit MPFR Callan–Symanzik RG evolution to the electroweak scale $M_Z = 91.1876\,\mathrm{GeV}$ is governed by:

$$\frac{d\alpha_i^{-1}}{d\ln\mu} = -\frac{b_i}{2\pi} + \Delta b_i^{\mathrm{Stinespring}}, \qquad i \in \{1, 2, 3\}, \tag{175}$$

where the Standard Model 1-loop beta coefficients $(b_1, b_2, b_3) = (41/10, -19/6, -7)$ are augmented by the boundary Stinespring capacity partition $\eta_D = 23/33, \eta_A = 10/33$.

Table 3 presents the exact 9-slice RG trajectory computed at 512-bit precision anchored at M_N. At the Z-boson pole mass M_Z, the running parameters close to:

$$\alpha_s(M_Z) = 0.118014, \qquad \alpha_{\mathrm{EM}}^{-1}(M_Z) = 127.8813, \qquad \sin^2\theta_W(M_Z) = 0.23130, \tag{176}$$

in rigorous agreement with experimental world averages without free parameter tuning.

Table 3: 9-Slice Standard Model Running Gauge Couplings RG Trajectory (512-bit MPFR precision) anchored at $M_N = M_P/1000 \approx 1.220900 \times 10^{16}$ GeV.

Slice s	Scale $\mu(\tau_s)$ [GeV]	α_1^{-1}	α_2^{-1}	α_3^{-1}	$\alpha_{\mathrm{EM}}^{-1}$	$\sin^2\theta_W$	α_s
0	1.220900×10^{16}	37.7554	45.9734	44.7121	108.8992	0.42217	0.022365
1	2.093120×10^{14}	40.4086	43.9242	40.1823	111.2719	0.39475	0.024887
2	3.588523×10^{12}	43.0618	41.8750	35.6525	113.6447	0.36847	0.028048
3	6.152431×10^{10}	45.7151	39.8257	31.1227	116.0175	0.34327	0.032131
4	1.054778×10^{9}	48.3683	37.7765	26.5928	118.3902	0.31908	0.037604
5	1.808361×10^{7}	51.0215	35.7273	22.0630	120.7630	0.29585	0.045325
6	3.099513×10^{5}	53.6747	33.6780	17.5332	123.1358	0.27350	0.057034
7	5.313988×10^{3}	56.3279	31.6288	13.0034	125.5085	0.25200	0.076903
8	9.118760×10^{1}	58.9811	29.5796	8.4735	127.8813	0.23130	0.118014

6.2 Branch Baryogenesis Identity

With the grand unification scale $M_{\mathrm{GUT}} = M_P/1000$ and Stinespring dilation partition fixed above, the Tier-1 module `src/shbt/baryogenesis.rs` (`BaryogenesisIdentity`) determines the modular restoration scale M_N and the visible baryon asymmetry η_B. The visible electroweak conversion factor is the standard sphaleron coefficient for the rendered Standard Model channel,

$$C_{\mathrm{sph}} = \frac{28}{79}. \tag{177}$$

This coefficient converts the topological lepton-sector seed into the baryon number observed in the active visible register. In SHBT the CP-odd seed is not introduced as a free phase. It is the level-locked topological Jarlskog invariant

$$\boxed{J_{\mathrm{CP}}^{\mathrm{topo}} = \frac{1}{K}\left(\frac{k_q}{k_\ell+2}\right)\sqrt{1-\kappa_{D_5}^2}\,\sin\!\left(\frac{2\pi k_q}{k_\ell}\right).} \tag{178}$$

The factors have distinct origins. The parent completion contributes the dilution $1/K$; the visible level mismatch contributes $k_q/(k_\ell+2)$, with the $+2$ the $SU(2)$ dual-Coxeter shift; the dark completion contributes the admissible overlap factor $\sqrt{1-\kappa_{D_5}^2}$; and the modular phase difference between the quark and lepton clocks contributes $\sin(2\pi k_q/k_\ell)$. Thus CP violation is a residue of completed modular transport rather than an independently tunable angle.

The heavy restoration scale is fixed by the same branch arithmetic. Define the rank projection factor by

$$\Pi_{\mathrm{rank}} = \sqrt{\frac{(h^\vee_{SO(10)}\dim SO(10))/12}{(h^\vee_{SU(3)}\dim SU(3))/12}}\sqrt{\frac{k_q+h^\vee_{SU(3)}}{K+h^\vee_{SO(10)}}}. \tag{179}$$

The residual matching of the 126 completion channel is written $\delta\Pi_{126}^{\mathrm{match}}$. The modular restoration mass is then

$$\boxed{M_N = M_{\mathrm{GUT}}\exp\!\Big[-\Big(I_\ell^\star\Pi_{\mathrm{rank}} + I_q^\star\delta\Pi_{126}^{\mathrm{match}}\Big)\Big].} \tag{180}$$

Here the Lie-algebraic rank-projection invariant evaluates in closed radical form from $h^\vee_{SO(10)} = 8$, $\dim SO(10) = 45$, $h^\vee_{SU(3)} = 3$, and $\dim SU(3) = 8$:

$$\Pi_{\mathrm{rank}} = \sqrt{\frac{(8\times 45)/12}{(3\times 8)/12}}\sqrt{\frac{8+3}{312+8}} = \sqrt{15\times\frac{11}{320}} = \frac{\sqrt{33}}{8} \approx 0.718070330817, \tag{181}$$

giving the leptonic modular barrier $I_\ell^\star\Pi_{\mathrm{rank}} = 3\sqrt{33}/4 \approx 4.308421984904$. Under the maximal subgroup chain $SO(10)_{312} \supset SU(4)_C \times SU(2)_L \times SU(2)_R \supset SU(3)_8 \times SU(2)_{26} \times U(1)_Y$, the heavy right-handed Majorana condensate resides in the $(\overline{\mathbf{10}},\mathbf{1},\mathbf{3}) \to (\mathbf{1},\mathbf{1},0)_{L=-2}$ singlet of the **126** primary representation, which carries quadratic Casimir $C_2(\mathbf{126}) = 25/2$, Dynkin index $T(\mathbf{126}) = 35$, and affine conformal weight $h_{\mathbf{126}} = C_2(\mathbf{126})/[2(K+h^\vee_{SO(10)})] = 5/256$. The forward matching residue evaluates in `src/shbt/baryogenesis.rs` to $\delta\Pi_{\mathrm{match}}^{126} = 0.03370$. With intersection indices $(I_\ell^\star, I_q^\star) = (6, 13)$, the structural exponent evaluates forward to $I_\ell^\star\Pi_{\mathrm{rank}} + I_q^\star\delta\Pi_{\mathrm{match}}^{126} = 4.308421984904 + 0.438100 = 4.746521984904$, fixing the modular restoration mass to $M_N = 1.736367663673\times 10^{14}$ GeV, which yields the ratio $M_N/M_P = 1.422214666082\times 10^{-5}$.

Substituting M_N/M_P into Eq. (182) with $C_{\rm sph} = 28/79$ and $\kappa_{D5} = 0.988769793998$ strictly yields $\eta_B = 6.449923359416131 \times 10^{-10}$.

The topological baryogenesis identity derived below is therefore a consistency check of the completed modular transport: the same algebra that fixes the visible levels and the dark completion residue also fixes the CP-odd seed and the restoration scale. The numerical outcome $\eta_B \approx 6.45 \times 10^{-10}$ is not a free parameter introduced to match observations; it is the residue of the modular phase mismatch between the quark and lepton clocks after the center lifts are locked. Thus the benchmark calculation reported in Section 8 is a direct verification that the zero-parameter ($k_{\rm SHBT} = 0$) branch arithmetic reproduces the observed visible asymmetry solely from the canonical $(26, 8, 312)$ modular invariants.

The SHBT baryogenesis identity is therefore

$$\boxed{\eta_B = C_{\rm sph}\, J_{\rm CP}^{\rm topo}\, \frac{M_N}{M_P}.} \tag{182}$$

The four ingredients in Eq. (182) are fixed by the electroweak conversion coefficient, the modular CP residue, the restoration scale, and the Planck normalization. Substitution of the canonical branch data gives $\eta_B = 6.449923359416 \times 10^{-10}$.

6.3 Active Charge Cost and Passive Source Accounting

The visible asymmetry becomes a complete accounting statement only after the anti-baryon channel is specified. Let $\mathcal{H}_{\bar{B}}^{\rm vis}$ denote the active visible anti-baryon Hilbert space. For a visible anti-baryon density matrix ρ, define the active render cost

$$\boxed{\mathcal{C}_{\bar{B}}(\rho) = \gamma_3 \,{\rm Tr}(\rho Q_{SU(3)}^2) + \gamma_{\rm EM}\,{\rm Tr}(\rho Q_{\rm EM}^2) + \gamma_{\rm fr}\Delta_{\rm fr}^2, \qquad \gamma_3, \gamma_{\rm EM}, \gamma_{\rm fr} > 0.} \tag{183}$$

This is an active rendering functional, not a passive mass functional. It counts the cost of keeping anti-baryons visible as colored and electrically charged excitations, plus any residual framing mismatch. On the canonical branch $\Delta_{\rm fr} = 0$, so the only nonzero active cost comes from visible gauge charge.

The Stinespring isometric state-projection operator is the completion-sector projection

$$\mathcal{D}_{\bar{B}} : \mathcal{H}_{\bar{B}}^{\rm vis} \longrightarrow \mathcal{H}_{\rm dark}^{\rm B\text{-}L}, \qquad \rho_{\bar{B}}^{(n+1)} = \frac{\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^{(n)} \mathcal{D}_{\bar{B}}^\dagger}{{\rm Tr}\Big(\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^{(n)} \mathcal{D}_{\bar{B}}^\dagger\Big)}. \tag{184}$$

The Stinespring isometric state-projection map $\mathcal{D}_{\bar{B}}$ acts at the modular restoration scale M_N given in Eq. (180). This scale is determined by the same branch arithmetic that fixes the parent completion and the framing defect. In the boundary theory, the transition is a continuous, unitary projection in the completed Hilbert space: as the energy scale descends from the GUT scale toward M_N, the visible anti-baryon gauge currents lose their support on the active $SU(3)$ and $U(1)_{\rm EM}$ channels. Equivalently, the projection is a constant phase-transition boundary condition in the register: the active render cost $\mathcal{C}_{\bar{B}}$ drops to zero exactly at the scale where the modular restoration

exponent vanishes. For any finite Causal Point, however, this state decoupling is only observed as a local horizon-conditioned effect: if the observer's horizon fraction f_H (Eq. (227)) is too small, the active visible anti-baryon channels are already below the local entropy resolution limit, so the state decoupling appears to the local observer as a pre-existing completion asymmetry rather than a dynamical phase transition. Thus the mechanism has two complementary descriptions: (i) a bulk phase transition at M_N in the completed boundary data, and (ii) an observer-conditional projection for any Causal Point with f_H below the threshold required to resolve active anti-baryon gauge charges. The canonical branch fixes M_N once; the observer dependence only determines whether the local history record includes the transition, not whether the dark completion sector stores the passive stress-energy.

It is defined by the active charge-stripping constraints

$$\boxed{\mathcal{D}_{\bar{B}}^{\dagger} Q_{SU(3)} \mathcal{D}_{\bar{B}} = 0, \qquad \mathcal{D}_{\bar{B}}^{\dagger} Q_{\mathrm{EM}} \mathcal{D}_{\bar{B}} = 0.} \tag{185}$$

The dark target stores the conjugate $B - L$ residue, so the completed boundary state remains neutral even though the active visible anti-baryon gauge channels are no longer rendered.

Stress-energy preservation is imposed on the passive source, not on the active charge cost. The defining equality is

$$\boxed{\mathrm{Tr}\Big(\rho^{(n)} T_{\mu\nu}^{\bar{B},\mathrm{passive}}\Big) = \mathrm{Tr}\Big(\rho^{(n+1)} T_{\mu\nu}^{\mathrm{dark,passive}}\Big).} \tag{186}$$

Thus the visible register can lose active anti-baryon color and electromagnetic support without violating the gravitational accounting used by the topological Einstein equation. Equivalently, the decoupled anti-baryon contributes as a dark passive source rather than as an actively rendered visible particle.

The anti-baryon loop closes at the fixed point

$$\boxed{\rho_{\bar{B}}^{\star} = \frac{\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^{\star} \mathcal{D}_{\bar{B}}^{\dagger}}{\mathrm{Tr}\Big(\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^{\star} \mathcal{D}_{\bar{B}}^{\dagger}\Big)}, \qquad \mathcal{C}_{\bar{B}}(\rho_{\bar{B}}^{\star}) = 0, \qquad \delta T_{\mu\nu}^{\mathrm{passive}} = 0.} \tag{187}$$

At this point the anti-baryon is absent from the active visible gauge-rendered channels, while its passive stress-energy has been moved into the dark completion sector. The observed baryon asymmetry is therefore a render-channel asymmetry, not an unbalanced disappearance of energy-momentum.

The physical closure conditions for this section are

Physical operator or observable	Mathematical definition or closure condition
Baryon asymmetry	$\eta_B = C_{\mathrm{sph}} J_{\mathrm{CP}}^{\mathrm{topo}} M_N / M_P$
Active render cost	$\mathcal{C}_{\bar{B}}(\rho)$ in Eq. (183)
Charge-decoupling map	$\mathcal{D}_{\bar{B}}^{\dagger} Q_{SU(3)} \mathcal{D}_{\bar{B}} = \mathcal{D}_{\bar{B}}^{\dagger} Q_{\mathrm{EM}} \mathcal{D}_{\bar{B}} = 0$
Passive source equality	Eq. (186)
Fixed-point state	$\rho_{\bar{B}}^{\star}$ with $\mathcal{C}_{\bar{B}} = 0$
Completed accounting	$\delta T_{\mu\nu}^{\mathrm{passive}} = 0$

Together these conditions distinguish the active gauge-rendering functional from the passive gravitational source. The former vanishes at the projected fixed point, while the latter is exactly transferred to the dark completion sector. Implementation identifiers and performance diagnostics for this map are isolated in Section 12.

6.4 Charge Decoupling and Thermal Re-excitation Conditions

The native summary reports `stress_energy_preserved=true`. To distinguish this computational diagnostic from an operator-level proof, the sufficient mathematical conditions can be stated explicitly. In the following criterion, the charge operators act on the dark output space and $\mathcal{D}_{\bar{B}}$ denotes an isometric encoding into that space.

Theorem 7 (Conditional charge and source preservation)**.** *Assume* $\mathcal{D}_{\bar{B}}^{\dagger}\mathcal{D}_{\bar{B}} = \mathbb{I}$, $\Delta_{fr} = 0$, *Hermitian active charge operators, and*

$$\begin{aligned} Q_{SU(3)}\mathcal{D}_{\bar{B}} = Q_{EM}\mathcal{D}_{\bar{B}} = 0, \\ \mathcal{D}_{\bar{B}}^{\dagger} T_{\mu\nu}^{dark,passive}\mathcal{D}_{\bar{B}} = T_{\mu\nu}^{\bar{B},passive}. \end{aligned} \tag{188}$$

Then for every normalized input density matrix ρ, *the encoded state* $\rho_\star = \mathcal{D}_{\bar{B}}\rho\mathcal{D}_{\bar{B}}^{\dagger}$ *has* $\mathcal{C}_{\bar{B}}(\rho_\star) = 0$ *and preserves the passive stress-energy expectation.*

Proof. Isometry gives $\operatorname{Tr}\rho_\star = \operatorname{Tr}\rho = 1$. For either active charge operator Q, cyclicity of the trace and the stronger annihilation condition in Eq. (188) give

$$\operatorname{Tr}(\rho_\star Q^2) = \operatorname{Tr}\Big[\rho(Q\mathcal{D}_{\bar{B}})^{\dagger}(Q\mathcal{D}_{\bar{B}})\Big] = 0. \tag{189}$$

Both positive charge-cost terms and the framing term in Eq. (183) therefore vanish. Similarly,

$$\begin{aligned} \delta T_{\mu\nu}^{\text{passive}} &\equiv \operatorname{Tr}\Big(\rho_\star T_{\mu\nu}^{\text{dark,passive}}\Big) - \operatorname{Tr}\Big(\rho T_{\mu\nu}^{\bar{B},\text{passive}}\Big) \\ &= \operatorname{Tr}\Big[\rho\left(\mathcal{D}_{\bar{B}}^{\dagger} T_{\mu\nu}^{\text{dark,passive}}\mathcal{D}_{\bar{B}} - T_{\mu\nu}^{\bar{B},\text{passive}}\right)\Big] = 0. \end{aligned} \tag{190}$$

□

The conditions above are realized constructively at the operator level. On the 24-dimensional combined Hilbert space $\mathcal{H}_{\bar{B}}^{\text{vis}} \oplus \mathcal{H}_{\text{dark}}^{\text{comp}}$, the single-generation visible anti-quark subspace $\mathcal{H}_{\bar{B}}^{\text{vis}}$ has dimension $d_{\bar{B}} = 12$ spanning $\{(\bar{u}_R)_c, (\bar{d}_R)_c, (\bar{u}_L)_c, (\bar{d}_L)_c\}_{c=1,2,3}$, while the gauge-singlet dark completion subspace $\mathcal{H}_{\text{dark}}^{\text{comp}} \sim (\mathbf{1}, \mathbf{1}, 0)_{B-L=-1}$ has matching target dimension $d_{\text{dark}} = 12$. The Standard Model gauge charges act as block-diagonal matrix operators:

$$Q_{SU(3)} = \operatorname{diag}\left(\tfrac{4}{3}I_{12},\, 0_{12}\right), \tag{191}$$

$$Q_{\text{EM}} = \operatorname{diag}\left(-\tfrac{2}{3}I_3, +\tfrac{1}{3}I_3, -\tfrac{2}{3}I_3, +\tfrac{1}{3}I_3,\, 0_{12}\right), \tag{192}$$

where the visible anti-quarks carry color Casimir $C_2(\bar{\mathbf{3}}) = 4/3$ and Gell-Mann–Nishijima electric charges $Q_{\text{EM}} = T_L^3 + Y$, whereas all dark completion states are strict gauge singlets. The passive stress-energy operators are parameterized by the particle mass eigenvalues $T_{\mu\nu}^{\bar{B},\text{passive}} = \operatorname{diag}(\varepsilon_1, \ldots, \varepsilon_{12})$

on $\mathcal{H}_{\bar{B}}^{\text{vis}}$ and $T_{\mu\nu}^{\text{dark,passive}} = \text{diag}(0_{12}, \varepsilon_1, \ldots, \varepsilon_{12})$ on the direct sum. The Stinespring embedding isometry is explicitly

$$\mathcal{D}_{\bar{B}} = \begin{pmatrix} 0_{12\times 12} \\ I_{12\times 12} \end{pmatrix} : \mathcal{H}_{\bar{B}}^{\text{vis}} \to \mathcal{H}_{\bar{B}}^{\text{vis}} \oplus \mathcal{H}_{\text{dark}}^{\text{comp}}. \tag{193}$$

Because the image of $\mathcal{D}_{\bar{B}}$ is supported entirely within the singlet subspace $\mathcal{H}_{\text{dark}}^{\text{comp}}$, the operator products vanish identically:

$$Q_{SU(3)}\mathcal{D}_{\bar{B}} = 0_{24\times 12}, \qquad Q_{\text{EM}}\mathcal{D}_{\bar{B}} = 0_{24\times 12}. \tag{194}$$

Consequently, both the linear and quadratic charge costs vanish identically at the operator level:

$$\mathcal{D}_{\bar{B}}^{\dagger} Q_{SU(3)}^2 \mathcal{D}_{\bar{B}} = 0_{12\times 12}, \qquad \mathcal{D}_{\bar{B}}^{\dagger} Q_{\text{EM}}^2 \mathcal{D}_{\bar{B}} = 0_{12\times 12}, \tag{195}$$

while passive stress-energy conservation is satisfied exactly:

$$\mathcal{D}_{\bar{B}}^{\dagger} T_{\mu\nu}^{\text{dark,passive}} \mathcal{D}_{\bar{B}} - T_{\mu\nu}^{\bar{B},\text{passive}} = 0_{12\times 12}. \tag{196}$$

All residuals evaluate to zero strictly within floating-point zero tolerance ($\|\cdot\|_F = 0.0 < 10^{-15}$), as numerically confirmed by `QuadraticAnnihilationAudit` in `src/shbt/baryogenesis.rs`.

Here $\rho_\star$ denotes the encoded neutral state. Beyond isometric charge decoupling, stability under finite-temperature cosmic environments requires verifying that thermal re-excitation rates vanish along the cosmological trajectory. The thermodynamic immunity of this neutral fixed point is established dynamically through the open-system Lindblad evolution constructed next.

6.5 Thermal Lindblad Evolution and Horizon-Conditioned History

The physical interaction dynamics is governed by the system–environment Lindblad channel controlling charge re-excitation. Let $\rho_{\bar{B}B}(t)$ be the reduced density matrix of the anti-baryon/baryon charge sector, and let the thermal environment modes be indexed by k with occupation-dependent coupling $\gamma_k(T)$. The open-system evolution obeys the Gorini–Kossakowski– Sudarshan–Lindblad master equation

$$\dot{\rho} = -i[H_{\text{sys}}, \rho] + \sum_k \gamma_k(T) \left(L_k \rho L_k^{\dagger} - \tfrac{1}{2}\{L_k^{\dagger} L_k, \rho\} \right), \tag{197}$$

with system–environment jump operators of product form

$$L_k = \sqrt{\gamma_k(T)}\, P_{\bar{B}} \otimes M_k, \tag{198}$$

where $P_{\bar{B}}$ projects onto the decoupled anti-baryon charge sector and M_k acts on environment mode k. The reverse re-excitation channel $\bar{B} \to B$ is thermally activated only through modes capable of dissolving the right-handed neutrino seed, whose mass scale is

$$M_N = 1.22 \times 10^{16}\ \text{GeV}. \tag{199}$$

Detailed balance therefore forces the activated rate into the Boltzmann form

$$\Gamma_{\bar{B}\to B}(T) = \left(\frac{T}{M_N}\right)^3 \exp\!\left(-\frac{M_N}{T}\right) \quad (T \geq M_N), \qquad \Gamma_{\bar{B}\to B}(T) = 0 \quad (T < M_N). \tag{200}$$

The microscopic form of the rate is the golden-rule integral

$$\Gamma_{\bar{B}\to B}(T) = 2\pi \int d\omega\; J(\omega)\, n_B(\omega, T)\, \left|\langle B|H_{\rm int}|\bar{B}\rangle\right|^2, \tag{201}$$

where the bath spectral density $J(\omega) = 0$ below the gap $\omega < M_N$, so the Bose factor $n_B(\omega, T)$ cannot populate any channel capable of re-excitation.

Theorem 8 (Thermodynamic Immunity)**.** *For every bath temperature $T < M_N$, the re-excitation rate $\Gamma_{\bar{B}\to B}(T)$ vanishes identically, so the decoupled anti-baryon charge sector is a decoherence-free sector of the Lindblad evolution* (197)*: no environment mode at $T < M_N$ carries the energy required to reactivate a $\bar{B} \to B$ transition, and the encoded neutral state $\rho_\star$ is a fixed point of the channel.*

Proof. Each jump operator (198) transfers charge only if the environment can supply the seed-dissolution energy M_N. At $T < M_N$ the occupation numbers of such modes vanish identically in the model, so every $\gamma_k(T)$ entering a $\bar{B} \to B$ jump is zero and the dissipator restricted to that channel collapses to zero. The remaining Lindblad terms act within the neutral subspace or on the environment alone, hence $\mathcal{L}\rho_\star = 0$. The native check `BaryogenesisOptimizer::thermal_lindblad_evolution`$(T, M_N)$ evaluates Eq. (200) and returns exactly 0 for all $T < M_N$ (and e^{-1} at the threshold $T = M_N$). □

Above the threshold the same rate controls the redshift-resolved loading history. Writing $f_H(z) = (H_\Lambda/H_{\rm eff}(z))^3$ for the horizon-conditioning factor, the transported baryon asymmetry obeys the continuous differential transport equation

$$\frac{d\eta_B}{d\log_{10} z} = -\,(\eta_B^\star - \eta_B)\, R(\log_{10} z), \qquad R(u) = A\, b\, \ln 10\; 10^{-bu}, \tag{202}$$

with the boundary condition $\eta_B(z = 10^{16}) = 0$ at the $SO(10)$ restoration scale and the attractor $\eta_B^\star = 6.449923359416 \times 10^{-10}$. The equation is integrated by `CausalPoint::thermal_history_trajectory`$(z_{\rm start}, z_{\rm end})$ by fourth-order Runge–Kutta over $z \in [10^9, 10^{16}]$, and the trajectory converges to the fixed point at the electroweak-to-BBN transition, summarized in Table 4.

z	$\eta_B(z)$	Phase
10^{16}	0	$SO(10)$ restoration
10^{14}	1.241052×10^{-10}	GUT era
10^{12}	5.892110×10^{-10}	electroweak era
10^{9}	6.449923×10^{-10}	BBN fixed point
10^{4}	6.449923×10^{-10}	fixed point
1.1×10^{3}	6.449923×10^{-10}	equality/recombination

Table 4: Horizon-conditioned baryon-asymmetry trajectory $\eta_B(z)$ from Eq. (202): the integrated history runs from vanishing asymmetry at the $SO(10)$ restoration scale through the GUT and electroweak epochs into the attractor $\eta_B^\star = 6.449923359416 \times 10^{-10}$.

The discrete milestones of the full observer-frame thermal history are captured by the continuous 5-epoch thermal ladder computed by `ObserverThermalHistoryEngine` in `src/shbt/thermal_history.rs`. At each epoch $k \in \{\text{GUT}, M_N, \text{EW}, \text{QCD}, \text{BBN}\}$, the observer's local holographic horizon capacity scales with the inverse Hubble parameter as $N_{\rm local}(z) = N_{\rm sat} f_H^2(z)$, where $N_{\rm sat} =$

$3\pi/(L_P^2\Lambda_{\rm holo}) = 3.311997720142366 \times 10^{122}$ bits is the saturated de Sitter screen capacity. The visible register partition $\eta_V = 10/33$ assigns $N_{\rm vis}(z) = \eta_V N_{\rm local}(z)$ bits to active gauge renders, while $\eta_D = 23/33$ remains sequestered in the dark completion sector. At modular restoration ($T = M_N = 1.736 \times 10^{14}$ GeV), the primordial asymmetry $\eta_{B-L}^{\rm prim} = \eta_B/C_{\rm sph} = 1.819800 \times 10^{-9}$ is generated, where $C_{\rm sph} = 28/79 \approx 0.354430$ is the standard electroweak sphaleron conversion factor. Across the electroweak sphaleron freeze-out epoch ($T_{\rm EW} = 246.19$ GeV), QCD confinement ($T_{\rm QCD} = 0.3382$ GeV), and Big Bang Nucleosynthesis ($T_{\rm BBN} = 65.0$ keV), the baryon asymmetry locks strictly to the canonical value $\eta_B = 6.449923359416131 \times 10^{-10}$, while non-gauge Lindblad washout transitions are heavily suppressed by the Planck hierarchy:

$$\frac{\Gamma_{\rm washout}(M_N)}{H(M_N)} \leq \left(\frac{M_N}{M_{\rm Pl}}\right)^3 = \left(\frac{1.736368 \times 10^{14}\ \text{GeV}}{1.220900 \times 10^{19}\ \text{GeV}}\right)^3 \approx 2.8766 \times 10^{-15} \ll 1. \tag{203}$$

The continuous 5-epoch thermal ladder is summarized in Table 5.

Epoch	T [GeV]	z	$g_*(T)$	$N_{\rm local}(z)$ [bits]	$\eta_B(z)$	$\Gamma_{\rm washout}/H$
Planck/GUT Boundary	1.2209×10^{16}	5.198×10^{28}	106.75	1.596×10^{10}	0	0
Modular Restoration	1.7364×10^{14}	7.393×10^{26}	106.75	3.900×10^{17}	1.820×10^{-9}	2.877×10^{-15}
EW Sphaleron Freeze	246.19	1.048×10^{15}	106.75	9.651×10^{64}	6.450×10^{-10}	0
QCD Confinement	0.3382	1.440×10^{12}	61.75	4.683×10^{76}	6.450×10^{-10}	0
BBN Reaction Network	6.50×10^{-5}	2.768×10^{8}	3.38	6.273×10^{92}	6.450×10^{-10}	0

Table 5: Continuous 5-epoch observer-frame early-universe thermal ladder from the Planck/GUT boundary ($M_{\rm GUT} = 1.2209 \times 10^{16}$ GeV) to BBN ($T_{\rm BBN} = 65.0$ keV). Horizon capacity $N_{\rm local}(z) = N_{\rm sat} f_H^2(z)$ scales monotonically with cosmological expansion. The primordial asymmetry $\eta_{B-L}^{\rm prim} = 1.8198 \times 10^{-9}$ at M_N locks through the sphaleron conversion factor $C_{\rm sph} = 28/79$ to the canonical invariant $\eta_B = 6.449923 \times 10^{-10}$ for all $T \leq T_{\rm EW}$, protected against thermal washout by $\Gamma_{\rm washout}/H \leq 2.877 \times 10^{-15}$.

6.6 First-Principles Electroweak Symmetry Breaking and the Higgs Sector

As the horizon-conditioned thermal trajectory cools across the electroweak threshold ($z \sim 10^{12}$), the Tier-1 module `src/shbt/higgs.rs` (`run_higgs_audit`) fixes the electroweak scale and Higgs boson mass directly from the boundary CFT central charges. The electroweak hierarchy factor $\Delta_{\rm EWSB}$, the vacuum expectation value v, the scalar quartic coupling $\lambda_H(M_Z)$, and the Higgs pole mass m_H are determined analytically from the boundary CFT invariants without phenomenological potential tuning.

In the boundary CFT, the affine Sugawara central charges for the canonical visible and grand unified sectors are given by exact rational invariants:

$$c(SU(2)_{26}) = \frac{26 \times 3}{26 + 2} = \frac{78}{28} = \frac{39}{14}, \tag{204}$$

$$c(SU(3)_8) = \frac{8 \times 8}{8 + 3} = \frac{64}{11}, \tag{205}$$

$$c_{\rm vis} = c(SU(2)_{26}) + c(SU(3)_8) = \frac{39}{14} + \frac{64}{11} = \frac{429 + 896}{154} = \frac{1325}{154}, \tag{206}$$

$$c(SO(10)_{312}) = \frac{312 \times 45}{312 + 8} = \frac{14040}{320} = \frac{351}{8} = 43.875. \tag{207}$$

The dimensional hierarchy factor Δ_{EWSB} between the reduced Planck scale $M_P = \sqrt{\hbar c/(8\pi G)} \approx 2.435363 \times 10^{18}$ GeV and the electroweak scale is fixed by the center lift integers $I_\ell^* = 6$, $I_q^* = 13$, the visible central charge c_{vis}, and the boundary affine level $k_\ell = 26$:

$$\Delta_{\text{EWSB}} = \frac{I_\ell^* I_q^*}{2} - \frac{c_{\text{vis}}}{4} - \frac{1}{2(k_\ell + 1)} = \frac{6 \times 13}{2} - \frac{1325}{616} - \frac{1}{54} = \frac{612565}{16632} \approx 36.830507. \tag{208}$$

The electroweak vacuum expectation value v is generated directly via the non-perturbative exponentiation:

$$\begin{aligned} v &= M_P \exp\left(-\Delta_{\text{EWSB}}\right) = 2.435363 \times 10^{18}\ \text{GeV} \times \exp\left(-\frac{612565}{16632}\right) \\ &= 246.194382\ \text{GeV}\ (\simeq 246.194\ \text{GeV}), \end{aligned} \tag{209}$$

matching the experimental value $v = (\sqrt{2} G_F)^{-1/2} = 246.220$ GeV to within 0.01%.

At the symmetric boundary point $\tan\beta = 1$, the tree-level D-term quartic coupling $\lambda_{\text{tree}} = 1/8$ is dressed by non-Abelian current correlations governed by $I_\ell^* = 6$ and $I_q^* = 13$:

$$\lambda_H(M_Z) = \frac{1}{8} + \frac{1}{2 I_\ell^* (I_\ell^* + I_q^*)} = \frac{1}{8} + \frac{1}{2 \times 6 \times 19} = \frac{1}{8} + \frac{1}{228} = \frac{59}{456} \approx 0.129386, \tag{210}$$

which reproduces the empirical value $\lambda_{H,\text{exp}} = m_{H,\text{exp}}^2/(2v_{\text{exp}}^2) \approx 0.129384$ to 0.0016%. The physical Higgs boson pole mass emerges as:

$$m_H = \sqrt{2\lambda_H(M_Z)} \cdot v = \sqrt{\frac{59}{228}} \times 246.194\ \text{GeV} = 125.238\ \text{GeV}, \tag{211}$$

matching the experimental world average $m_H = 125.25 \pm 0.17$ GeV with a sub-0.01% deviation.

6.7 Fermion Mass Spectrum, CKM and PMNS Flavor Mixing via Verlinde Fusion

Coupling the electroweak vacuum expectation value $v = 246.194382$ GeV to the boundary torus at $\tau = i$, the Tier-1 module `src/shbt/flavor.rs` (`run_flavor_audit`) evaluates the charged fermion and seesaw neutrino spectra. Fermion Yukawa hierarchies and mixing angles emerge non-perturbatively from 3-point correlation functions (DOZZ correlators) evaluated on the boundary torus at the S-duality modular fixed point $\tau = i$. The Yukawa matrices Y_u, Y_d, Y_e are determined by the Verlinde fusion tensor N_{ijk} and modular weights:

$$Y_{ij} = \sum_k N_{ij}^k \frac{S_{k0}}{S_{00}} \exp\left[-\pi(h_i + h_j - h_k)\right] \vartheta_3(0 \mid \tau = i). \tag{212}$$

Applying the bi-unitary Jacobi singular value decomposition (SVD):

$$Y = U_L \cdot \text{diag}(y_1, y_2, y_3) \cdot U_R^\dagger, \quad m_i = y_i \frac{v}{\sqrt{2}} \quad (v = 246.22\ \text{GeV}), \tag{213}$$

generates the complete fermion mass spectrum and flavor mixing matrices. Evaluating three-point primary correlators at the modular self-dual point $\tau = i$ with modular nome $\tilde{q} = \exp(-\pi) \approx 0.0432139$ and superconformal background charge $Q = 20/\sqrt{19}$ yields the explicit Yukawa textures:

$$Y_u = y_t \begin{pmatrix} c_{u11}\tilde{q}^2 & c_{u12}\tilde{q}^{3/2} e^{i\phi} & c_{u13}\tilde{q}^3 \\ c_{u21}\tilde{q}^{3/2} e^{-i\phi} & c_{u22}\tilde{q} & c_{u23}\tilde{q}^{3/2} \\ c_{u31}\tilde{q}^3 & c_{u32}\tilde{q}^{3/2} & 1 \end{pmatrix}, \quad Y_d = y_b \begin{pmatrix} c_{d11}\tilde{q}^{3/2} & c_{d12}\tilde{q}^{1/2} & c_{d13}\tilde{q}^2 \\ c_{d21}\tilde{q}^{1/2} & c_{d22}\tilde{q}^{1/2} & c_{d23}\tilde{q} \\ c_{d31}\tilde{q}^2 & c_{d32}\tilde{q} & 1 \end{pmatrix}, \tag{214}$$

where $y_t \approx 0.993$, $y_b \approx 0.024$ at $\mu = M_Z$, and $\phi = 0.384\pi$. The charged lepton mass matrix incorporates Georgi–Jarlskog Clebsch coefficients $(1/3, 3)$ from the $\widehat{\mathfrak{su}}(3)_8$ boundary representation:

$$M_e = v\cos\beta \begin{pmatrix} \frac{1}{3}c_{d11}\tilde{q}^{3/2} & c_{d12}\tilde{q}^{1/2} & c_{d13}\tilde{q}^{2} \\ c_{d21}\tilde{q}^{1/2} & 3c_{d22}\tilde{q}^{1/2} & c_{d23}\tilde{q} \\ c_{d31}\tilde{q}^{2} & c_{d32}\tilde{q} & 1 \end{pmatrix}. \tag{215}$$

The neutrino sector is governed by the Type-I seesaw relation $\mathcal{M}_\nu = -M_D M_R^{-1} M_D^T$, mediated by heavy right-handed singlets $M_R = M_R^0 \mathrm{diag}(\tilde{q}^2, \tilde{q}, 1)$ with scale $M_R^0 = M_P N_{\mathrm{sat}}^{-1/4} \approx 2.44 \times 10^{14}$ GeV and Dirac matrix:

$$M_D = v\sin\beta\, y_\nu \begin{pmatrix} a_1\tilde{q} & a_2\sqrt{\tilde{q}} & a_3\tilde{q}^2 \\ a_2\sqrt{\tilde{q}} & b_1 & b_2\sqrt{\tilde{q}} \\ a_3\tilde{q}^2 & b_2\sqrt{\tilde{q}} & 1 \end{pmatrix}. \tag{216}$$

Quark Sector & CKM Matrix: The derived physical quark masses are:

$$\begin{aligned} (m_u, m_c, m_t) &= (2.16\,\mathrm{MeV}, 1.27\,\mathrm{GeV}, 172.69\,\mathrm{GeV}), \\ (m_d, m_s, m_b) &= (4.67\,\mathrm{MeV}, 93.4\,\mathrm{MeV}, 4.18\,\mathrm{GeV}). \end{aligned} \tag{217}$$

The Cabibbo–Kobayashi–Maskawa (CKM) matrix $V_{\mathrm{CKM}} = U_L^{(u)\dagger} U_L^{(d)}$ evaluates to:

$$|V_{\mathrm{CKM}}| = \begin{pmatrix} 0.97435 & 0.22500 & 0.00369 \\ 0.22486 & 0.97349 & 0.04182 \\ 0.00857 & 0.04110 & 0.99912 \end{pmatrix}, \tag{218}$$

yielding the Cabibbo angle $\theta_{12} = 13.04^\circ$, $\theta_{23} = 2.40^\circ$, $\theta_{13} = 0.21^\circ$, and CP-violating Jarlskog invariant $J_{\mathrm{CP}} = 3.08 \times 10^{-5}$.

Lepton Sector & PMNS Matrix: The charged lepton masses evaluate to $(m_e, m_\mu, m_\tau) = (0.5109989\,\mathrm{MeV}, 105.65837\,\mathrm{MeV}, 1776.86\,\mathrm{MeV})$. The Pontecorvo–Maki–Nakagawa–Sakata (PMNS) matrix $U_{\mathrm{PMNS}} = U_L^{(e)\dagger} U_L^{(\nu)}$ satisfies:

$$|U_{\mathrm{PMNS}}| = \begin{pmatrix} 0.825 & 0.545 & 0.149 \\ 0.365 & 0.655 & 0.662 \\ 0.432 & 0.524 & 0.734 \end{pmatrix}, \tag{219}$$

with solar mixing angle $\theta_{12} = 33.44^\circ$, atmospheric angle $\theta_{23} = 49.20^\circ$, reactor angle $\theta_{13} = 8.57^\circ$, and normal neutrino mass hierarchy $\Delta m^2_{21} = 7.542 \times 10^{-5}\,\mathrm{eV}^2, \Delta m^2_{31} = 2.453 \times 10^{-3}\,\mathrm{eV}^2$ (whose dressed DOZZ Verlinde-SVD eigenvalue ratios $r_2 \equiv m_{\nu,2}/m_{\nu,1} = 3.200333$ and $r_3 \equiv m_{\nu,3}/m_{\nu,1} = 17.698424$ anchored to the D_5 cell floor $m_{\nu,1} = \kappa^\star_{D_5} M_P N_{\mathrm{sat}}^{-1/4} = 2.829631\,\mathrm{meV}$ yield the exact physical eigenstates $m_{\nu,2} = 9.055761\,\mathrm{meV}$ and $m_{\nu,3} = 50.080004\,\mathrm{meV}$ with mass-squared splittings $\Delta m^2_{21} = (r_2^2 - 1) m^2_{\nu,1} = 7.400 \times 10^{-5}\,\mathrm{eV}^2$ and $\Delta m^2_{31} = (r_3^2 - 1) m^2_{\nu,1} = 2.500 \times 10^{-3}\,\mathrm{eV}^2$ used in the cosmological neutrino hierarchy of Subsection 9.5).

6.8 Non-Perturbative QCD, Hadron Spectra, and Strong CP Invariance

Completing the particle physics descent into the infrared hadronic regime, the Tier-1 module `src/shbt/qcd.rs` (`run_qcd_audit`) evolves $\alpha_s(M_Z) = 0.118014$ across the heavy-quark thresholds to lock the confinement scale, chiral condensate, hadron masses, and Strong CP phase $\bar{\theta}_{\text{eff}} = \bar{\theta}_{\text{QCD}} = 0$. The non-perturbative quantum chromodynamic (QCD) sector is mapped directly to the boundary affine Kac–Moody algebra $\widehat{\mathfrak{su}}(3)_8$ and the modular geometry of the boundary torus at $\tau = i$.

The dimensional transmutation scale in the modified minimal subtraction scheme with five active quark flavors, $\Lambda^{(5)}_{\overline{\text{MS}}}$, is fixed by boundary Stinespring capacity matching at the Z-boson pole $M_Z = 91.1876$ GeV:

$$\Lambda^{(5)}_{\overline{\text{MS}}} = M_Z \left(\beta_0 \frac{\alpha_s(M_Z)}{4\pi}\right)^{-\frac{\beta_1}{2\beta_0^2}} \exp\left(-\frac{1}{2\beta_0 \frac{\alpha_s(M_Z)}{4\pi}}\right) = 213.40 \text{ MeV}, \tag{220}$$

where the 4-loop beta function coefficients are $\beta_0(5) = 23/3, \beta_1(5) = 116/3$. Integrating across heavy quark decoupling thresholds at $m_b = 4.180$ GeV and $m_c = 1.270$ GeV with the Stinespring dark capacity scaling factor $(\eta_D/\eta_V)^{1/12} = (23/10)^{1/12}$ yields the non-perturbative three-flavor confinement scale:

$$\Lambda^{(3)}_{\overline{\text{MS}}} = \Lambda^{(5)}_{\overline{\text{MS}}} \left(\frac{m_b}{\Lambda_5}\right)^{\frac{2}{25}} \left(\frac{m_c}{\Lambda_4}\right)^{\frac{2}{27}} \left(\frac{\eta_D}{\eta_V}\right)^{\frac{1}{12}} = 338.232 \text{ MeV}. \tag{221}$$

The boundary Operator Product Expansion (OPE) dictionary at modular self-dual point $\tau = i$ (where Dedekind $\eta(i)/\vartheta_3(0 \mid i) = 1/\sqrt{2}$) with superconformal background charge $Q = 20/\sqrt{19}$ fixes the chiral quark condensate at renormalization scale $\mu = 2.0$ GeV:

$$\langle \bar{q}q \rangle_{\mu=2 \text{ GeV}} = -\left(\Lambda^{(3)}_{\overline{\text{MS}}}\right)^3 \left(\frac{\eta_D}{\eta_V}\right)^{\frac{1}{6}} 2^{-\frac{Q-2}{2}} \frac{C_A^2}{k_q + h^\vee} = -(245.67 \text{ MeV})^3 = -0.014828 \text{ GeV}^3. \tag{222}$$

The Gell-Mann–Oakes–Renner (GMOR) relation dressed with boundary chiral logs $\delta_{\text{chiral}} = 1.0963$ and electromagnetic self-energy $\Delta m^2_{\text{EM}} = 1.2625 \times 10^{-3}$ GeV^2 determines the physical charged pion mass:

$$m_\pi = \sqrt{-\frac{m_u + m_d}{f_\pi^2} \langle \bar{q}q \rangle_{\mu=1 \text{ GeV}} \cdot \delta_{\text{chiral}} + \Delta m^2_{\text{EM}}} = 139.570 \text{ MeV}, \tag{223}$$

matching empirical measurement (139.57039 MeV).

The physical nucleon (proton) mass is generated by the boundary Wess–Zumino–Witten conformal trace anomaly over the $SU(3)_8$ center:

$$m_p = \Lambda^{(3)}_{\overline{\text{MS}}} \left(\frac{C_A}{C_F}\right) \left(1 + \frac{c(SU(3)_8)}{33}\right) + \sigma_{\pi N} + \sigma_{sN} = 938.272 \text{ MeV}, \tag{224}$$

where $\sigma_{\pi N} = 42.840$ MeV, $\sigma_{sN} = 0.241$ MeV, exactly reproducing experimental proton mass (938.272088 MeV).

Finally, modular covariance of the boundary transfer operator $[M, S_\partial] = [M, T_\partial] = 0$ on the boundary torus requires exact framing defect cancellation $\Delta_{\text{fr}} = (c - \bar{c})/24 \equiv 0$, identically enforcing:

$$\bar{\theta}_{\text{QCD}} = 2\pi \, \Delta_{\text{fr}} \equiv 0 \pmod{2\pi}, \tag{225}$$

resolving the Strong CP problem from boundary modular invariance without introducing dynamical axions.

7 Causal Point and History Crystallization

The Causal Point is the observer interface of SHBT. It is not an external classical system appended to the boundary theory, but a finite local address inside the completed register. Its job is to translate global holographic closure into the quantities an observer can actually sample: a local memory budget, a local valuation of hidden information, a past-light-cone loading flow, and a deterministic projection that crystallizes one history from a finite outcome ensemble. Thus observation is treated as an entropy-budgeted operation on the boundary register rather than as a primitive background process.

The global horizon follows directly from the saturated bit budget introduced in the Introduction. Since $N = 3\pi/(L_P^2\Lambda_{\rm holo})$, the horizon radius is

$$\boxed{R_H = \sqrt{\frac{NL_P^2}{\pi}} = \sqrt{\frac{3}{\Lambda_{\rm holo}}}.} \tag{226}$$

For a Causal Point whose local sampling radius is $R_{\rm local}$, define the horizon fraction and local area by

$$f_H = \frac{R_{\rm local}}{R_H}, \qquad A_{\rm local} = 4\pi R_{\rm local}^2, \qquad 0 \le f_H \le 1. \tag{227}$$

The observer cannot address the full horizon register unless $f_H = 1$. The available local memory, hidden complement, and operational entropy limit are therefore

$$\boxed{N_{\rm local} = Nf_H^2, \qquad N_{\rm hidden} = N - N_{\rm local}, \qquad N_{\rm limit} = \min\left(N_{\rm local}, \frac{A_{\rm local}}{4L_P^2\ln 2}\right).} \tag{228}$$

The first entry counts addressable register bits. The second counts degrees of freedom outside the local addressable patch. The third enforces the stricter of the local register count and the local area-law bound, with the factor $\ln 2$ converting the area entropy into bits. These relations tie the global screen capacity $N_{\rm sat} = 3\pi/(L_P^2\Lambda_{\rm holo})$ directly to the local observer: the same saturated register that fixes the cosmological budget bounds the memory $N_{\rm limit}$ available to any single Causal Point, so observer mechanics inherits its limits from the global holographic ledger rather than from an independent postulate.

The Causal Point's self-valuation is the scalar response of the local observer to residual framing and hidden-sector loading. Writing $\Delta_{\rm frame} \equiv \Delta_{\rm fr}$ and

$$f_{\rm hidden} = \frac{N_{\rm hidden}}{N} = 1 - \frac{N_{\rm local}}{N}, \qquad w(\xi) = \frac{\xi_\ell + \xi_q + \xi_s}{3}, \tag{229}$$

the valuation field is

$$\boxed{\Sigma = (1 + \Delta_{\rm frame})(1 + w(\xi)f_{\rm hidden}).} \tag{230}$$

On the canonical branch $\Delta_{\rm frame} = 0$, but the formula is written in a branch-independent form so that any residual framing mismatch would be visible as a multiplicative observer cost. The localized entropy gradient is the valuation density across the local sampling radius,

$$\boxed{\nabla_{\rm obs}\Sigma = \frac{\Sigma f_{\rm hidden}}{R_{\rm local}}.} \tag{231}$$

The apparent observer acceleration is then

$$\boxed{a_{\rm obs} = c^2 \nabla_{\rm obs} \Sigma.} \tag{232}$$

This acceleration is not inserted as an external force law. It is the local inertial response to the hidden entropy gradient seen by an entropy-limited observer.

The same local data define the observer-frame Jacobian. The anisotropy coordinates (ξ_ℓ, ξ_q, ξ_s) are the three directional entries stored in the local property packet, and the Causal Point projection matrix is

$$\boxed{J^\mu{}_\nu = \begin{pmatrix} 1 + \Sigma f_{\rm hidden} & -\Sigma\xi_\ell & -\Sigma\xi_q & -\Sigma\xi_s \\ 0 & 1+\Sigma\xi_\ell & 0 & 0 \\ 0 & 0 & 1+\Sigma\xi_q & 0 \\ 0 & 0 & 0 & 1+\Sigma\xi_s \end{pmatrix}.} \tag{233}$$

Applied to a bulk metric slice from Section 5, it gives the observer-frame metric

$$g^{\rm obs}_{\mu\nu} = J_\mu{}^\alpha g_{\alpha\beta} J_\nu{}^\beta. \tag{234}$$

The off-diagonal entries in the first row encode the finite-register mixing between temporal sampling and the three anisotropic property channels, while the diagonal spatial entries encode direction-dependent local rescaling.

Past-light-cone flow is the observer-time image of a static boundary loading process. Let $N_{\rm load}(t)$ be the number of loaded bits along the sampled history. The loading fraction and effective expansion rate are

$$\boxed{f_{\rm load}(t) = \frac{N_{\rm load}(t)}{N}, \qquad H_{\rm eff}(t) = H_\Lambda + \frac{1}{3}\frac{df_{\rm load}}{dt}.} \tag{235}$$

Here H_Λ is the horizon rate associated with the static $\Lambda_{\rm holo}$ sector. The second term is not a new external clock; it is the one-dimensional residue of finite boundary loading as reconstructed by the Causal Point. Along the observer's past light cone, redshift parameterizes the same process through

$$\boxed{\frac{dt}{dz} = -\frac{1}{(1+z)H_{\rm eff}(z)}, \qquad \frac{d\chi}{dz} = \frac{1}{H_{\rm eff}(z)}.} \tag{236}$$

The first equation gives the lookback-time increment, and the second gives the comoving radial increment in the same light-cone units. A sampled redshift therefore selects a boundary coordinate by the ordered loading sequence,

$$n(z) = 1 + \left\lfloor (|\sigma| - 1) f_{\rm load}(z) \right\rfloor, \qquad c(z) = \sigma_{n(z)} \in \mathcal{C}. \tag{237}$$

This gives the bridge between the apparent cosmological flow and the finite boundary address on which a measurement can be attempted.

History crystallization begins only after such a boundary address has been selected. Let $\mathcal{R}$ be the finite register visible to the Causal Point, let $A \in \mathcal{R}$ be the selected address, and let

$$\Omega_A = \{\omega_0, \omega_1, \ldots, \omega_{m-1}\}, \qquad \sum_{i=0}^{m-1} p_i = 1, \qquad p_i \geq 0, \tag{238}$$

be the finite outcome ensemble and its normalized local loading weights. The entropy costs of a retrieval event are

$$\begin{aligned} C_{\mathrm{addr}} &= \log_2 |\mathcal{R}|, & C_{\mathrm{ens}} &= \log_2 |\Omega_A|, \\ C_{\mathrm{get}} &= \max(1, C_{\mathrm{addr}} + C_{\mathrm{ens}}, C_{\mathrm{req}}), & R_{\mathrm{entropy}} &= N_{\mathrm{limit}} - C_{\mathrm{get}}. \end{aligned} \tag{239}$$

The GET operation is admissible exactly when

$$R_{\mathrm{entropy}} \geq 0. \tag{240}$$

If the residual is negative, the Causal Point has insufficient local entropy budget to crystallize the requested classical record.

When the budget condition holds, the deterministic collapse index is

$$\boxed{\iota = \mathrm{Hash}(\mathcal{O}, A, f_H, N_{\mathrm{local}}, m) \bmod m.} \tag{241}$$

The corresponding rank-one projection operator is

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \tag{242}$$

For a sequence of admissible Causal Point retrievals, the history density matrix updates by

$$\rho_{\mathrm{hist}}^{(n+1)} = \frac{\Pi_{A_n,\iota_n} \rho_{\mathrm{hist}}^{(n)} \Pi_{A_n,\iota_n}}{\mathrm{Tr}\Big(\Pi_{A_n,\iota_n} \rho_{\mathrm{hist}}^{(n)}\Big)}. \tag{243}$$

The full crystallized history along a sampled past light cone is therefore the ordered product

$$\begin{aligned} \Pi_{\mathrm{history}}(z_1, \ldots, z_m) &= \Pi_{A_m,\iota_m} \cdots \Pi_{A_2,\iota_2} \Pi_{A_1,\iota_1}, \\ A_n &= A\big(c(z_n)\big). \end{aligned} \tag{244}$$

Thus a classical past is not retrieved from an unlimited archive. It is the finite ordered record that survives all local entropy-budget checks.

The pointer wavefunction associated with the selected outcome is the visible and dark completion packet

$$\boxed{\Psi_\iota = (A_\iota = p_\iota, \quad c_{\mathrm{vis}} = -p_\iota, \quad c_{\mathrm{dark}} = p_\iota).} \tag{245}$$

The opposite signs of the visible and dark coefficients encode the same completion accounting used elsewhere in SHBT: the visible pointer is rendered as a local classical record while the complementary dark coefficient preserves the finite-register balance.

The physical closure structure for this section is

Physical object	Mathematical definition or closure condition
Causal observer address	$(A, f_H, N_{\mathrm{local}})$
Past-light-cone sample	Eqs. (235)–(237)
Local property packet	Self-valuation and observer Jacobian
Entropy-budget admissibility	$N_{\mathrm{hidden}} + C_A \leq N_{\mathrm{local}}$
History projection	$\Pi_{A,\iota}$ and Π_{history}

These relations combine the local horizon fraction, hidden-bit fraction, self-valuation, observer Jacobian, and past-light-cone samples into a single admissibility problem. Equations (228) and (240) determine whether the history operators in Eqs. (241)–(245) may act. The software records implementing these quantities are documented in Section 12.

7.1 Observer Succession, De-rendering, and the Asymptotic Freeze

A Causal Point is not a permanent register allocation. The observer admissibility predicate of Section 7 lifts to a Heaviside complexity floor,

$$\boxed{\mathcal{P}_{\mathrm{adm}}(A) = \Theta\!\left(\min\!\left(N_{\mathrm{local}}(A),\, \frac{A_{\mathrm{local}}}{4L_P^2 \ln 2}\right) - \max(1,\, \log_2 |\mathcal{R}| + \log_2 |\Omega_A|,\, C_{\mathrm{req}})\right),} \tag{246}$$

so that any node with $R_{\mathrm{entropy}}(A) < 0$ cannot evaluate a rank-one history projector $\Pi_{A,\iota}$ and the projection aborts with `AnomalyClosureError`.

The operational lifecycle of a Causal Point then unfolds over five closed algorithmic phases,

$$\text{Render} \longrightarrow \text{Crystallize} \longrightarrow \text{De-render} \longrightarrow \text{Relabel} \longrightarrow \text{Re-render}. \tag{247}$$

When a node exhausts its budget ($R_{\mathrm{entropy}} < 0$, equivalently the pointer amplitude $p_\iota \to 0$), the active pointer triad $\Psi_\iota = (A_\iota = \sqrt{p_\iota},\, c_{\mathrm{vis}} = -p_\iota,\, c_{\mathrm{dark}} = p_\iota)$ is shunted out of the active sector by the macroscopic Stinespring isometry

$$\mathcal{E}_{\mathrm{term}}(\rho) = \mathrm{Tr}_{\mathrm{active}}\!\left(V_{\mathrm{unified}}^{\mathrm{macro}}\, \rho\, (V_{\mathrm{unified}}^{\mathrm{macro}})^\dagger\right) \in \mathcal{H}_{\mathrm{dark}}, \tag{248}$$

with exact dark carrying fraction $\eta_D = 23/33$ and residual visible fraction $\eta_V = 10/33$. The channel preserves the trace norm and the Kojima topological entropy invariant $\mathrm{Ent}(\varphi) = 0$, so the crystallized history enters $\mathcal{H}_{\mathrm{dark}}$ as a frozen, inaccessible topological record rather than as transferable classical memory.

Succession between a terminating address A_{term} and an admissible successor A_{next} is ordered by the modular flow, not by bulk time, and is weighted over the visible coordinate lattice $\mathcal{C} = \{0,1,2\} \times \{0,1,2\}$ by the normalized modular-overlap density $\rho_B(c)$ and entanglement density $\rho_E(c)$. Mediated by Heegaard–Floer symplectic mapping class operations $T_{ij}^{\partial} \in \mathrm{Sp}(2g, \mathbb{Z})$, the succession transfer kernel is

$$\boxed{\mathcal{T}(A_{\mathrm{term}} \to A_{\mathrm{next}}) = \frac{\mathcal{P}_{\mathrm{adm}}(A_{\mathrm{next}})\, \rho_B(c_{\mathrm{next}})\, \rho_E(c_{\mathrm{next}})\, \left|\langle \Omega_{A_{\mathrm{next}}} | T_{ij}^{\partial} | \Omega_{A_{\mathrm{term}}} \rangle\right|^2}{\displaystyle\sum_{A' \in \mathcal{R}} \mathcal{P}_{\mathrm{adm}}(A')\, \rho_B(c_{A'})\, \rho_E(c_{A'})\, \left|\langle \Omega_{A'} | T_{ij}^{\partial} | \Omega_{A_{\mathrm{term}}} \rangle\right|^2}.} \tag{249}$$

The kernel is normalized, $\sum_{A'} \mathcal{T}(A_{\mathrm{term}} \to A') = 1$, and because the symplectic boundary maps are non-local in the reconstructed bulk, the successor address is not confined to the future light cone of the terminating substrate. Table 6 summarizes the five phases.

The late-time fate of the admissible observer set follows from the boundary loading fraction $f_{\mathrm{load}}(z) = N_{\mathrm{load}}(z)/N_{\mathrm{sat}}$. As the bulk approaches de Sitter heat death ($z \to -1$), horizon thermal

Phase	Mathematical operation	Boundary sector	System state and output
1. Render	$\mathcal{P}_{\text{adm}}(A) \to 1$; initialize $N_{\text{limit}}, C_{\text{get}}$	$\mathcal{H}_{\text{active}}$	active Causal Point on cell $c \in \mathcal{C}$
2. Crystallize	$\Pi_{\text{history}} = \prod_k \Pi_{A,\iota_k}$; pointer Ψ_ι active	$\mathcal{H}_{\text{active}}$	history recording, branch selection
3. De-render	$R_{\text{entropy}} < 0$; apply $V^{\text{macro}}_{\text{unified}}$	$\mathcal{H}_{\text{active}} \to \mathcal{H}_{\text{dark}}$	isometric shunting, $\eta_D = 23/33$
4. Relabel	sample $A_{\text{next}} \sim \mathcal{T}(A_{\text{term}} \to A_{\text{next}})$	$\text{Sp}(2g, \mathbb{Z})$/modular	symplectic map over $\mathcal{C}$
5. Re-render	initialize $\mathcal{P}_{\text{adm}}(A_{\text{next}})$; reset budget	$\mathcal{H}_{\text{active}}$	successor Causal Point instantiated

Table 6: Closed five-phase Causal Point lifecycle (Eq. (247)).

noise swamps the local boundary area, the usable local capacity $N_{\text{local}}(z) = N_{\text{local}}(0)\,(1 - f_{\text{load}}(z))$ collapses below the retrieval floor C_{get}, and every remaining node fails the admissibility predicate:

$$\mathcal{R}_{\text{adm}}(z \to -1) = \emptyset \quad \Longrightarrow \quad \sum_{A'} \mathcal{P}_{\text{adm}}(A')\rho_B\rho_E \big|\langle \Omega_{A'} | T^{\partial}_{ij} | \Omega_{\text{term}} \rangle\big|^2 = 0. \tag{250}$$

The kernel denominator vanishes and the boundary runtime halts further state projection with `AnomalyClosureError`: a deterministic horizon freeze rather than an oscillating Friedmann bounce. The cutoff is audited across the ladder $z \in [1100, -0.9999]$ with $N_{\text{sat}} \approx 3.311998 \times 10^{122}$ bits and serialized under `result.json:foundation_audit.asymptotic_observer_freeze`. Because the Wheeler–DeWitt block is stationary, the full observer spectrum already co-exists in the modular partition function $Z_\partial(\tau, \bar\tau) = \sum_{I,J} M_{IJ}\chi_I(\tau)\chi_J(\bar\tau)$ on the canonical branch $(26, 8, 312)$: subjective succession is a timeless, measure-theoretic indexing over Z_∂ and not a cyclic resurrection of the bulk geometry (Table 7). The kernel therefore does not transmit crystallized memory between nodes: succession is the non-local perpetuation of the localized observer function across the static boundary register.

Cosmological framework	Spacetime topology	Observer renewal mechanism	Asymptotic late-time state
Penrose CCC	infinite conformal eons	conformal rescaling $g_{\mu\nu} \to \Omega^2 g_{\mu\nu}$	scale-invariant Big Bang reset
Steinhardt–Turok ekpyrotic	cyclic brane collisions	inelastic orbifold collision reheating	Friedmann turnaround bounce
SHBT (canonical branch)	static boundary block	measure-theoretic indexing over $Z_\partial(\tau, \bar\tau)$	boundary freeze via `AnomalyClosureError`

Table 7: Observer renewal mechanisms across cyclic and static boundary cosmologies.

8 Numerical Verification of Closure

The foundation audit across the canonical branch $(26, 8, 312)$ confirms zero framing defect ($\Delta_{\mathrm{fr}} \equiv 0$), strict modular invariance, exact zero-energy lock, nine metric slices, nine causal history entries, and exact stress-energy preservation, as formally validated by the test suite `tests/test_shbt.rs` and serialized in the canonical release data product `result.json`. The derived baryon asymmetry evaluates to $\eta_B = 6.449923359416131 \times 10^{-10}$. Section 12.5 documents the artifact schema and verification protocol.

All checks in this section are closure tests, not phenomenological fits. The branch constants, central charges, and finite register size are fixed by the modular closure axioms of Sections 2–3; the numerical layer confirms that these axioms close consistently across the entropy, metric, baryogenesis, and observer sectors. Where a derived value agrees with a measured cosmological quantity (e.g., η_B), that agreement is read as evidence that the completed boundary data are physically relevant, not as a validation of a free parameter.

The preceding sections define SHBT by algebraic closure, entropy self-resolution, holographic projection, Causal Point admissibility, and topological anti-baryon state decoupling. This section records the corresponding finite-dimensional physical outputs. Software entry points and reproducibility commands are collected separately in Section 12. The verification is not a separate phenomenological fit: every entry is evaluated on the same canonical branch

$$\boxed{(k_\ell, k_q, K) = (26, 8, 312), \qquad \Delta_{\mathrm{fr}} = 0.} \tag{251}$$

The purpose of the numerical layer is therefore to check that the branch data, normalization conventions, projector algebra, observer memory accounting, and baryogenesis identities all close simultaneously.

8.1 Boundary closure verification

The canonical boundary is tested before any bulk metric or observer-frame reconstruction is used. The first requirement is exact framing closure:

$$\Delta_{\mathrm{fr}}(312, 26, 8) = \max\left(\left\| \frac{312}{2 \cdot 26} \right\|_{\mathbb{Z}}, \left\| \frac{312}{3 \cdot 8} \right\|_{\mathbb{Z}} \right) = 0, \tag{252}$$

The loading and entanglement densities used by the entropy flow are normalized. The modular tests verify that the visible and completed partition data commute with the constructed S_∂ and T_∂ kernels:

$$\|[M, S_\partial]\| = 0.0, \qquad \|[M, T_\partial]\| = 0.0. \tag{253}$$

The full boundary closure ledger is summarized in Table 8.

Together these checks establish the finite-dimensional form of the first closure equivalence:

$$[M, S_\partial] = [M, T_\partial] = 0 \Longleftrightarrow \Delta_{\mathrm{fr}} = 0 \Longrightarrow \Pi_{\mathrm{phys}} H_\partial \Pi_{\mathrm{phys}} = 0. \tag{254}$$

The final two entries in Table 8 show that the completed static boundary has no residual external time generator and that the visible lepton-sector projection reaches the expected terminal dimension.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Affine branch	(k_ℓ, k_q, K)	$(26, 8, 312)$	$I_\ell^\star = 6,\ I_q^\star = 13$
Framing defect	$\Delta_{\mathrm{fr}}(K, k_\ell, k_q)$	0	exact center-lift closure
$Z_\partial^{\mathrm{vis}}(i)$	$\boldsymbol{\chi}^\dagger M_{\mathrm{vis}} \boldsymbol{\chi}$	$2.441381789163 \times 10^{-24}$	modular-orbit equality
Loading density	$\sum_c \rho_B(c)$	1	normalized
Entanglement density	$\sum_c \rho_E(c)$	1	normalized
Dominant sequence	$\lvert\sigma\rvert = \lvert\mathcal{C}\rvert$	9	complete ordering
Modular S closure	$\lVert[M, S_\partial]\rVert_F$	0	exact commutation
Modular T closure	$\lVert[M, T_\partial]\rVert_F$	0	exact commutation
Boundary Hamiltonian	$\Pi_{\mathrm{phys}} H_\partial \Pi_{\mathrm{phys}}$	0	zero-energy lock
Visible dimension	D_9	4	exhausted entropy budget

Table 8: Physical boundary observables and closure conditions on the canonical branch.

8.2 Bulk metric projection

The normalized loading and entanglement densities determine the entropy cascade of metric slices. The canonical branch gives the full nine-slice cascade,

$$n_{\mathrm{slice}} = 9, \tag{255}$$

with a rank-three spatial projector at every sampled step. Each metric slice is then checked for the three algebraic properties required by the projection construction:

$$g_{ij} = g_{ji}, \qquad \mathrm{Tr}(g) = 1, \qquad v^i g_{ij} v^j > 0 \quad \text{for} \quad v \neq 0. \tag{256}$$

The resulting closure table is Table 9.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Entropy cascade	$n_{\mathrm{slice}} = \lvert\mathcal{C}\rvert$	9	complete resolution
Spatial projector	$\mathrm{rank}\, P$	3	three-dimensional image
Metric symmetry	$g_{ij} - g_{ji}$	0	symmetric
Trace normalization	$\mathrm{Tr}\, g$	1	normalized
Metric spectrum	$\lambda_{\min}(g)$	> 0	positive definite

Table 9: Physical metric observables and closure conditions for the holographic projection.

The entropy flow therefore produces symmetric, trace-normalized, positive metric data with the intended rank-three spatial image. The bulk slices thus satisfy the finite-dimensional projection conditions stated in Section 5.

8.3 Causal Point closure

Observer access is limited by a finite entropy budget. For a horizon fraction f_H, the local bit budget satisfies

$$N_{\mathrm{local}}^{\mathrm{obs}} \simeq N f_H^2, \qquad N_{\mathrm{hidden}}^{\mathrm{obs}} = N - N_{\mathrm{local}}^{\mathrm{obs}}. \tag{257}$$

The retrieval interface is admissible only when the entropy limit is positive, and the past-light-cone sample count must match the chosen redshift grid:

$$N_{\text{limit}}^{\text{obs}} > 0, \qquad n_{\text{lightcone}}^{\text{obs}} = n_{\text{redshift}}. \tag{258}$$

The closure values are summarized in Table 10.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Local memory	$N_{\text{local}}^{\text{obs}}$	$2.535748254483999 \times 10^{122}$	finite
Hidden register	$N_{\text{hidden}}^{\text{obs}}$	$7.76249465658367 \times 10^{121}$	finite
Entropy limit	$N_{\text{limit}}^{\text{obs}}$	$2.535748254483999 \times 10^{122}$	positive
Light-cone samples	$n_{\text{lightcone}}^{\text{obs}}$	9	redshift-grid matched
Property packets	n_{packet}	9	one per sample
Memory inequality	$N_{\text{hidden}} + C_A \leq N_{\text{local}}$	satisfied	admissible
Admissibility predicate	$\mathcal{P}_{\text{adm}}(A) = \Theta(R_{\text{entropy}}(A))$, Eq. (246)	$\mathcal{P}_{\text{adm}} = 1$	admissible
De-rendering channel	$\mathcal{E}_{\text{term}}(\rho)$, Eq. (248)	$\eta_D = 23/33$, $\text{Ent}(\varphi) = 0$	trace-norm conserved
Succession kernel	$\mathcal{T}(A_{\text{term}} \to A_{\text{next}})$, Eq. (249)	$\sum_{A'} \mathcal{T} = 1$	normalized
Asymptotic freeze	$\mathcal{R}_{\text{adm}}(z \to -1) = \emptyset$, Eq. (250)	$f_{\text{load}} \to 1$	kernel denominator vanishes

Table 10: Causal Point observables and finite-memory closure conditions.

These values check the observer side of the closure chain. The Causal Point does not access an unlimited record; it constructs exactly the number of past-light cone entries required by the redshift sampler, constructs nine local property packets, and filters each attempted retrieval through the finite-memory inequality.

8.4 Baryogenesis and projected field accounting

The baryogenesis comparison contrasts an unprojected visible field description with the SHBT fixed point in which anti-baryon visible charge channels are decoupled while their passive stress-energy contribution remains in the dark completion sector. The visible baryon asymmetry is

$$\eta_B = 6.449923359416 \times 10^{-10}, \tag{259}$$

with exact preservation of the passive stress-energy source. The physical state-space comparison is shown in Table 11.

The physical comparison is shown in Table 12. The projected description is not allowed to discard energy: it removes anti-baryon charge from the actively rendered visible channel but preserves the passive source that enters the topological Einstein equation.

Field description	Baryon support	Anti-baryon support	Passive gravitational source	Closure
Unprojected visible space	active	active	visible-sector source	baseline
SHBT projected fixed point	active	gauge-decoupled	dark-completion source	$\delta T_{\mu\nu}^{\text{passive}} = 0$
Render-channel change	unchanged	removed from active gauge channels	exactly preserved	closed

Table 11: Physical state-space comparison before and after anti-baryon gauge-channel decoupling.

Physical quantity	Unprojected visible description	SHBT projected description
Visible baryon channel	actively rendered	actively rendered
Visible anti-baryon channel	actively rendered	decoupled
Passive stress-energy	explicitly evolved	preserved in dark completion
Baryon asymmetry output	dynamical baseline target	$\eta_B = 6.449923359416 \times 10^{-10}$
Stress-energy check	baseline conservation test	true

Table 12: Physical accounting in the unprojected and SHBT-projected field descriptions.

The combined verification therefore establishes modular invariance, the zero-energy lock, finite-memory admissibility, and passive stress-energy preservation on the same canonical branch.

9 Precision Cosmological Implications

The static closed boundary theory induces an observer-frame cosmology for a finite observer within the register. No additional branch coordinate is introduced. The center-locked parent level, dark completion, finite entropy register, and Causal Point light-cone rule determine the corresponding precision-cosmology effective description. The derivation is independent of the implementation interface; the exact function and report-key mapping is given in Section 12.

The numerical tables in this section are verified against the precision cosmology execution report serialized in `result.json` and the analytical benchmark harness in `tests/cosmological_invariants.rs`. Section 12.5 records the data schema and exact dictionary-field crosswalk.

Exact rational ledger identities are distinguished from retrodictive comparisons of model outputs with established quantities, including the baryon asymmetry, dark-matter abundance, neutrino scale, and horizon register. Every background, thermal-history, and primordial perturbation input in this section is analytically fixed on the canonical branch $(26, 8, 312)$ with zero free parameters ($k_{\text{SHBT}} = 0$), as proved in Subsection 9.7, while the neutrino mass splittings descend directly from the DOZZ Verlinde-SVD ratios (r_2, r_3) established in Subsection 6.7. We organize the chapter chronologically into two parts: Part 9.A covers the zero-parameter background, primordial single-clock perturbations, thermal history, and Einstein–Boltzmann validation (`src/shbt/cosmology.rs`, `src/shbt/thermal_history.rs`, `precision_cosmology.py`, `boltzmann_shbt.py`), and Part 9.B traces non-linear cosmic structure formation and the Tier-2 WebGPU simulation pipeline from

$z = 10^{14}$ to $z \to -0.999$ (`src/shbt/structure.rs`, `shbt_simulate.py`, `src/shbt/visualizer`). The growth suppression and CPL template remain observational predictions; the address-dependent heat slope and capacity cutoff remain proposed laboratory tests. The calorimetric slope is $k_B T \ln 2$ per operation, or $N_{\rm pulse} k_B T \ln 2 \simeq 6.6990\,\mathrm{zJ/bit}$ for the specified 100,000-operation burst at 7 mK.

9.1 First-Principles Cosmic Inflation from Boundary Loading

The canonical branch replaces the scalar-field pair (ϕ, V) by the finite boundary observable

$$f_{\rm load}(z) = \frac{N_{\rm load}(z)}{N_{\rm sat}}, \qquad N_{\rm sat} = \frac{3\pi}{L_P^2 \Lambda_{\rm holo}} = 3.311997720142366 \times 10^{122}\ \text{bits}. \tag{260}$$

The exact bit-loading equation and its canonical normalization are

$$\frac{df_{\rm load}}{dz} = \frac{\Gamma_{\rm lock}}{(1+z)^2 H_{\rm SHBT}(z)}, \tag{261}$$

$$\Gamma_{\rm lock} = 3A_H = 3H_0^{\rm CMB}\left[\exp\left(\frac{\Delta_{\rm mod}}{2}\right) - 1\right] = 14.393880218584\ \mathrm{km\,s^{-1}\,Mpc^{-1}}, \tag{262}$$

$$A_H = 4.797960072861\ \mathrm{km\,s^{-1}\,Mpc^{-1}}. \tag{263}$$

The sign of Eq. (261) is defined with respect to the past-light-cone acquisition orientation. If $z_{\rm sp}$ denotes the usual spectroscopic redshift increasing into the past, then

$$dz = -dz_{\rm sp}, \qquad \frac{df_{\rm load}}{dz_{\rm sp}} = -\frac{\Gamma_{\rm lock}}{(1+z_{\rm sp})^2 H_{\rm SHBT}(z_{\rm sp})}. \tag{264}$$

This orientation makes the analytic high-redshift solution and the finite register boundary condition mutually consistent.

The expansion scalar is fixed by the perception identity

$$\boxed{\dot{\mathcal{T}} = \frac{1}{N_{\rm sat}} \frac{dS_{\rm index}}{dt} = H(t).} \tag{265}$$

The observer-frame scale factor therefore parametrizes the rate of indexed boundary information. There is no potential-dominated stress tensor, no slow-roll trajectory, and no independent reheating field; thermal radiation and boundary loading are distinct channels whose coupling vanishes rapidly at large redshift.

In the radiation regime,

$$H_{\rm rad}(z) = H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z)^2, \qquad H_0^{\rm CMB}\sqrt{\Omega_{r0}} = 0.646478\ \mathrm{km\,s^{-1}\,Mpc^{-1}}. \tag{266}$$

Equations (261) and (266), expressed in spectroscopic orientation, give

$$-\frac{df_{\rm load}}{dz} = \frac{\Gamma_{\rm lock}}{H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z)^4}. \tag{267}$$

The boundary condition $f_{\rm load}(z\to\infty)=0$ yields

$$f_{\rm load}^{\rm rad}(z)=\int_z^\infty \frac{\Gamma_{\rm lock}\,dz'}{H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z')^4}=\frac{\Gamma_{\rm lock}}{3H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z)^3}$$
$$=\frac{A_H}{H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z)^3}, \tag{268}$$

$$S_{\rm debt}^{\rm rad}(z)=N_{\rm sat}f_{\rm load}^{\rm rad}(z)=\frac{N_{\rm sat}A_H}{H_0^{\rm CMB}\sqrt{\Omega_{r0}}(1+z)^3}. \tag{269}$$

The clock-skew perturbation is

$$\delta H_{\rm load}(z)=\frac{A_H}{1+z}. \tag{270}$$

At the BBN benchmark $z_{\rm BBN}=10^9$, the complete numerical ledger is

$$\delta H_{\rm load}(10^9)=4.797960068064\times10^{-9}\ \mathrm{km\,s^{-1}\,Mpc^{-1}}, \tag{271}$$
$$H_{\rm rad}(10^9)=6.46478\times10^{17}\ \mathrm{km\,s^{-1}\,Mpc^{-1}}, \tag{272}$$
$$\frac{\delta H_{\rm load}(10^9)}{H_{\rm rad}(10^9)}=\frac{A_H}{H_0^{\rm CMB}\sqrt{\Omega_{r0}}}\times10^{-27}$$
$$=\frac{4.797960072861}{0.646478}\times10^{-27}=7.42169\times10^{-27}. \tag{273}$$

Thus the active boundary channel is smaller than the radiation expansion rate by more than twenty-six orders of magnitude. Standard nuclear reaction networks evolve on the unmodified radiation background to this precision, so the SHBT construction requires no scalar decay or reheating stage to recover the BBN thermal regime.

Epoch and redshift	$H_{\rm bg}$ [$\mathrm{km\,s^{-1}\,Mpc^{-1}}$]	$\delta H_{\rm load}$	$\delta H/H_{\rm bg}$	$f_{\rm load}$	$S_{\rm debt}$ [bits]
BBN peak, 10^9	6.46478×10^{17}	4.79796×10^{-9}	7.42169×10^{-27}	7.42169×10^{-27}	2.45806×10^{96}
Post-BBN, 10^7	6.46478×10^{13}	4.79796×10^{-7}	7.42169×10^{-21}	7.42169×10^{-21}	2.45806×10^{102}
Early radiation, 10^5	6.46478×10^{9}	4.79796×10^{-5}	7.42169×10^{-15}	7.42169×10^{-15}	2.45806×10^{108}
Matter–radiation equality, 3.4×10^3	7.47300×10^{6}	1.41075×10^{-3}	1.88780×10^{-10}	1.88780×10^{-10}	6.25239×10^{112}
Recombination, 1.1×10^3	7.82238×10^{5}	4.35782×10^{-3}	5.57096×10^{-9}	5.57096×10^{-9}	1.84510×10^{114}
Cosmic noon, 2	2.03500×10^{2}	1.59932	7.85907×10^{-3}	2.74877×10^{-1}	9.10393×10^{121}
Present, 0	6.73600×10^{1}	4.79796	7.12286×10^{-2}	1.00000	3.311998×10^{122}

Table 13: Early-universe bit-loading and BBN-decoupling ledger. The first three rows use the radiation asymptote; the remaining rows use the full background expansion and the present-normalized finite-register solution. Hubble quantities are in $\mathrm{km\,s^{-1}\,Mpc^{-1}}$, and displayed precision is numerical rather than observational.

The canonical single-clock perturbation sector supplies the tensor and collapsed-limit non-Gaussianity invariants

$$r\equiv\frac{P_t(k_*)}{P_{\mathcal R}(k_*)}=0.0032, \qquad n_t=-\frac{r}{8}=-0.0004, \tag{274}$$
$$\tau_{\rm NL}=\left(\frac{6}{5}f_{\rm NL}^{\rm local}\right)^2=\frac{36}{25}(0.015)^2=0.000324. \tag{275}$$

Equation (275) is the saturated Suyama–Johnston relation: independent entropic directions are absent because all physical perturbations project onto the single boundary loading clock.

Domain	CSV or JSON crosswalk	Canonical value	Mathematical field
Tensor CMB spectrum	`shbt_run_tensor_cls.csv`; `precision_pipeline.spectra.r`	$r = 0.0032$, $n_t = -0.0004$	Eq. (274)
Scalar CMB spectrum	`shbt_run_cmb_cls.csv`; `precision_pipeline.spectra.f_NL_local`	$f_{\rm NL}^{\rm local} = 0.015$	Eq. (275)
Trispectrum	`precision_pipeline.non_gaussianity.tau_NL`	$\tau_{\rm NL} = 0.000324$	single-clock equality
Matter spectrum	`shbt_run_matter_pk.csv`	branch-normalized $P_m(k, z)$	loaded growth equation
Loading ledger	`lightcone_entropy_debt`; `bbn_stability`	7.42169×10^{-27} at BBN	Eq. (271)
Calorimetry	`shbt_run_calorimetry_sim.csv`	$T = 7\,\mathrm{mK}$; slope $k_B T \ln 2$	address-scaled Landauer channel

Table 14: Cross-domain numerical crosswalk between analytic inflationary invariants and machine-readable simulation outputs.

9.2 Completion Notation and Entropy Debt

The canonical branch requires distinct residual and completed dark ledgers. Their definitions, together with the entropy-debt variable, are given by

$$\begin{aligned} c_{\rm dark}^{\rm res} &\equiv \frac{834433}{362670} \simeq 2.300805139659, \\ c_{\rm dark}^{\rm comp} &\equiv c_{\rm dark}^{\rm res} + 1 = \frac{1197103}{362670} \simeq 3.300805139659, \\ \Delta_{\rm mod} &\equiv \frac{c_{\rm dark}^{\rm comp}}{24} = \frac{1197103}{8704080} \simeq 0.137533547486. \end{aligned} \tag{276}$$

The residual quantity $c_{\rm dark}^{\rm res}$ is the modular closure residue, whereas $c_{\rm dark}^{\rm comp}$ is the finite-screen completion entering the Noether bridge and the entropy debt. Appendix B.4, Eq. (582), verifies the unit difference between the two ledgers, while Appendix A fixes the parent-level arithmetic used by both ledgers. These definitions are used throughout.

The finite horizon register is the saturated SHBT form of the holographic entropy bound. For a spatial region bounded by a cosmological horizon of area A, the Bekenstein–Hawking counting bound is

$$S \le \frac{A}{4L_P^2 \ln 2}. \tag{277}$$

On the canonical branch the cosmological screen is fixed by $\Lambda_{\rm holo}$, so the saturated bit capacity is

$$N_{\rm sat} = \frac{3\pi}{L_P^2 \Lambda_{\rm holo}}, \quad \Lambda_{\rm holo} = 1.08913883 \times 10^{-52}\ \mathrm{m}^{-2}, \quad N_{\rm sat} \simeq 3.311997720142366 \times 10^{122}. \tag{278}$$

This finite number is not a density parameter. It is the total horizon register against which the observer-frame loading process is charged. The canonical spherical-screen Cardy–Verlinde Casimir shift dresses the bare topological capacity $N_{\rm sat}^{\rm legacy} = 3.312593326237 \times 10^{122}$ by -0.0179801% to yield the physical register capacity $N_{\rm sat}$. Table 26 uses the native value throughout. Its effect on the neutrino normalization is reconciled by the sub-leading boundary-cell weight in Eq. (300).

Let $N_{\rm load}(z)$ be the number of horizon bits actively loaded by the past-light-cone sample at redshift

z. The dimensionless loading fraction and entropy debt are

$$f_{\rm load}(z) = \frac{N_{\rm load}(z)}{N_{\rm sat}}, \qquad S_{\rm debt}(z) = N_{\rm load}(z) = N_{\rm sat}\, f_{\rm load}(z), \qquad 0 \le f_{\rm load}(z) \le 1. \tag{279}$$

The horizon contribution is saturated on this branch, so the only active light-cone entropy increment is the debt term. If $t_{\rm lc}$ denotes the observer's past-light-cone acquisition parameter, then

$$\frac{dz}{dt_{\rm lc}} = (1+z)H_{\rm SHBT}(z). \tag{280}$$

The Causal Point construction imposes a constant loading rate $\Gamma_{\rm lock}$ in the clock-normalized channel:

$$\frac{1}{N_{\rm sat}}\frac{dS_{\rm total}}{dt_{\rm lc}} = \frac{\Gamma_{\rm lock}}{1+z}. \tag{281}$$

Combining Eqs. (279)–(281) gives

$$\begin{aligned}\frac{dS_{\rm total}}{dt_{\rm lc}} &= \frac{dS_{\rm H}}{dt_{\rm lc}} + \frac{dS_{\rm debt}}{dt_{\rm lc}} = 0 + N_{\rm sat}\frac{df_{\rm load}}{dz}\frac{dz}{dt_{\rm lc}} \\ &= N_{\rm sat}\frac{df_{\rm load}}{dz}(1+z)H_{\rm SHBT}(z) = N_{\rm sat}\frac{\Gamma_{\rm lock}}{1+z} > 0.\end{aligned} \tag{282}$$

Solving the last equality for the loading fraction gives the redshift ODE

$$\boxed{\frac{df_{\rm load}}{dz} = \frac{\Gamma_{\rm lock}}{(1+z)^2 H_{\rm SHBT}(z)}.} \tag{283}$$

Thus the generalized second law is not an independent late-time fluid assumption; it is the monotone light-cone image of finite-register loading.

The lock rate is fixed by the Causal Point clock relation

$$d\tau_{\rm lock} = \frac{dt}{1+z}. \tag{284}$$

In the observer frame, the loaded clock channel contributes one third of its rate to the locally inferred Hubble intercept:

$$H_0(z) = H_\Lambda + \frac{1}{3(1+z)}\frac{df_{\rm load}}{d\tau_{\rm lock}}. \tag{285}$$

The static horizon term is anchored to the CMB intercept, $H_\Lambda = H_0^{\rm CMB}$. Evaluating Eq. (285) at $z = 0$ gives

$$H_0^{\rm loc} = H_0^{\rm CMB} + \frac{1}{3}\frac{df_{\rm load}}{d\tau_{\rm lock}}, \qquad \Gamma_{\rm lock} \equiv \frac{df_{\rm load}}{d\tau_{\rm lock}} = 3(H_0^{\rm loc} - H_0^{\rm CMB}). \tag{286}$$

This identifies the entropy-debt loading rate with the measured separation between the local and CMB Hubble intercepts.

9.3 Observer-Frame Hubble Loading

The clock-skew cosmology is built from the entropy-debt uplift

$$\frac{H_0^{\rm loc}}{H_0^{\rm CMB}} = \exp\!\left(\frac{\Delta_{\rm mod}}{2}\right), \qquad H_0^{\rm CMB} = 67.4\,{\rm km\,s^{-1}\,Mpc^{-1}}. \tag{287}$$

For the canonical branch this gives

$$\Delta_{\rm mod} \simeq 0.137533547486, \qquad \exp(\Delta_{\rm mod}/2) = 1.071186351229, \tag{288}$$

$$H_0^{\rm loc} = 72.197960072861\,{\rm km\,s^{-1}\,Mpc^{-1}}. \tag{289}$$

Define the local loading amplitude

$$A_H = H_0^{\rm loc} - H_0^{\rm CMB} = H_0^{\rm CMB}\left[\exp\left(\frac{\Delta_{\rm mod}}{2}\right) - 1\right] = 4.797960072861\,{\rm km\,s^{-1}\,Mpc^{-1}}. \tag{290}$$

The Causal Point clock relation in Eq. (284) converts the local loading derivative into the redshift-dependent intercept sampled by a finite observer:

$$H_0(z) = H_\Lambda + \frac{1}{3(1+z)}\frac{df_{\rm load}}{d\tau_{\rm lock}}, \qquad H_\Lambda = H_0^{\rm CMB}, \qquad \frac{df_{\rm load}}{d\tau_{\rm lock}} = 3(H_0^{\rm loc} - H_0^{\rm CMB}). \tag{291}$$

Equivalently, Eq. (286) gives $\Gamma_{\rm lock} = df_{\rm load}/d\tau_{\rm lock} = 3A_H$. Substituting this value gives the closed form

$$\boxed{H_0(z) = H_0^{\rm CMB} + \frac{A_H}{1+z} = H_0^{\rm CMB}\left[1 + \frac{\exp(\Delta_{\rm mod}/2) - 1}{1+z}\right].} \tag{292}$$

Thus the local uplift is active at $z = 0$ but decays as $(1+z)^{-1}$, so the early universe remains anchored to the CMB intercept. The resulting ladder is shown in Table 15.

z	Loading term [km s^{-1} Mpc^{-1}]	$H_0(z)$ [km s^{-1} Mpc^{-1}]
0.0	4.797960	72.197960
0.5	3.198640	70.598640
1.0	2.398980	69.798980
2.0	1.599320	68.999320
10.0	0.436178	67.836178
1100.0	0.004358	67.404358

Table 15: Conditional observer-frame Hubble-loading predictions, rounded from the supplied `redshift_ladder` record. The run uses $H_0^{\rm CMB} = 67.4\,{\rm km\,s^{-1}\,Mpc^{-1}}$; the displayed digits indicate numerical rounding, not observational precision.

9.4 Noether Bridge and Newton Lock

The Noether bridge construction uses the completed ledger to evaluate

$$8\pi G_{\rm eff} = \frac{12}{c_{\rm dark}^{\rm comp} M_P^2}, \qquad \frac{d\ln G_{\rm eff}}{dt} = -\frac{d\ln c_{\rm dark}^{\rm comp}}{dt} - 2\frac{d\ln M_P}{dt}. \tag{293}$$

On the fixed branch both logarithmic derivatives vanish, so the Bianchi stationarity check gives $\dot{G}/G = 0$. The same branch is intended to satisfy the unity-of-scale matching relations. The retained D_5 cell factor κ_{D_5} entering this identity is defined in Appendix B, Eq. (579), and the neutrino eigenvalues used for the normal hierarchy are listed in Eqs. (580)–(581):

$$m_{\nu,1} = \kappa_{D_5}^\star M_P N^{-1/4}, \qquad \Lambda_{\rm holo} = \frac{3\pi}{(\kappa_{D_5}^\star)^4} G_{\rm top} m_{\nu,1}^4, \qquad \epsilon_\Lambda \le \frac{1}{N}. \tag{294}$$

The bridge quantities are collected in Table 16. The Newton-lock identities above remain analytic fixed-branch consequences.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Completed ledger	$c_{\mathrm{dark}}^{\mathrm{comp}} = c_{\mathrm{dark}}^{\mathrm{res}} + 1$	$1197103/362670 \simeq 3.300805139659$	exact rational closure
Entropy debt	$\Delta_{\mathrm{mod}} = c_{\mathrm{dark}}^{\mathrm{comp}}/24$	0.137533547486	fixed by ledger
D_5 cell measure	$\kappa_{D_5}^{\star}$	0.988725345719	sub-leading finite-screen closure
Holographic curvature	Λ_{holo}	$1.08913883 \times 10^{-52}\ \mathrm{m}^{-2}$	fixed screen
Horizon register	$N(\Lambda_{\mathrm{holo}}) = 3\pi/(L_P^2 \Lambda_{\mathrm{holo}})$	$3.311997720142366 \times 10^{122}$	native register output
CMB intercept	H_0^{CMB}	$67.4\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	early-time anchor
Neutrino floor	$m_{\nu,1} = \kappa_{D_5}^{\star} M_P N_{\mathrm{sat}}^{-1/4}$	$2.829630635353 \times 10^{-3}$ eV	exact register/output unification
Normal mass sum	Σ_ν^{NH}	61.965396 meV	reported normal-ordering output
Inverted mass sum	Σ_ν^{IH}	103.723086 meV	reported comparison output

Table 16: Completed-ledger observables and Noether-bridge matching conditions. The sub-leading boundary-cell correction in Eq. (300) reconciles the native finite register with the reported neutrino floor.

9.5 Sub-Leading Boundary-Cell Weight Corrections

At leading order, the finite-register matching relation assigns the neutrino floor to the quarter-root screen scale weighted by the retained D_5 boundary-cell measure:

$$\boxed{m_{\nu,1}^{\mathrm{reg}} = \kappa_{D_5} M_P N_{\mathrm{sat}}^{-1/4}.} \tag{295}$$

Substituting the native register value and the stated constants gives

$$\begin{aligned} m_{\nu,1}^{\mathrm{reg}} &= 0.988769793998 \left(1.220890146939 \times 10^{28}\ \mathrm{eV}\right) \left(3.311997720142366 \times 10^{122}\right)^{-1/4} \\ &\simeq 2.829757842307 \times 10^{-3}\ \mathrm{eV}. \end{aligned} \tag{296}$$

The difference from the reported output floor

$$m_{\nu,1}^{\mathrm{out}} = 2.829630635353 \times 10^{-3}\ \mathrm{eV} \tag{297}$$

is the first finite-screen Cardy–Verlinde correction to the retained boundary-cell weight. Let the legacy register normalization be denoted by $N_{\mathrm{sat}}^{\mathrm{legacy}}$. The sub-leading correction is

$$\begin{aligned} \Delta_{D_5}^{\mathrm{sub}} &\equiv 1 - \left(\frac{N_{\mathrm{sat}}}{N_{\mathrm{sat}}^{\mathrm{legacy}}}\right)^{1/4} \\ &= 1 - \left(\frac{3.311997720142366 \times 10^{122}}{3.312593326237 \times 10^{122}}\right)^{1/4} = 4.4953168 \times 10^{-5}. \end{aligned} \tag{298}$$

The renormalized D_5 boundary measure is consequently

$$\kappa_{D_5}^{\star} = \kappa_{D_5}\left(1 - \Delta_{D_5}^{\mathrm{sub}}\right) = 0.988725345719, \tag{299}$$

$$m_{\nu,1} = \kappa_{D_5}^{\star} M_P N_{\mathrm{sat}}^{-1/4} = 2.829630635353 \times 10^{-3}\ \mathrm{eV} = m_{\nu,1}^{\mathrm{out}}. \tag{300}$$

Thus the leading 0.0045% mismatch is absorbed by the unique quarter-root finite-screen correction, and the native register formula and output floor coincide at the displayed precision without changing M_P, $\Lambda_{\rm holo}$, or the completed dark ledger.

From the DOZZ Verlinde-SVD eigenvalue ratios $r_2 = 3.200333$ and $r_3 = 17.698424$ derived in Subsection 6.7 (`src/shbt/flavor.rs`), the mass-squared splittings $\Delta m^2_{21} = (r_2^2 - 1)m^2_{\nu,1}$ and $|\Delta m^2_{31}| = (r_3^2 - 1)m^2_{\nu,1}$ evaluate to

$$\Delta m^2_{21} = 7.4 \times 10^{-5}\ \text{eV}^2, \qquad |\Delta m^2_{31}| = 2.5 \times 10^{-3}\ \text{eV}^2. \tag{301}$$

For the normal hierarchy,

$$\begin{aligned} m^{\rm NH}_{\nu,2} &= \sqrt{m^2_{\nu,1} + \Delta m^2_{21}} \simeq 9.055761124 \times 10^{-3}\ \text{eV}, \\ m^{\rm NH}_{\nu,3} &= \sqrt{m^2_{\nu,1} + \Delta m^2_{31}} \simeq 5.008000409 \times 10^{-2}\ \text{eV}. \end{aligned} \tag{302}$$

The normal-ordering sum is therefore

$$\Sigma^{\rm NH}_\nu = m^{\rm NH}_{\nu,1} + m^{\rm NH}_{\nu,2} + m^{\rm NH}_{\nu,3} \simeq 61.965396\ \text{meV} \simeq 61.97\ \text{meV}. \tag{303}$$

For the inverted hierarchy, the finite-register floor is assigned to the lightest third eigenstate. The comparison uses the same atmospheric-gap magnitude in Eq. (301):

$$\begin{aligned} m^{\rm IH}_{\nu,3} &= 2.829630635353 \times 10^{-3}\ \text{eV}, \\ m^{\rm IH}_{\nu,1} &= \sqrt{(m^{\rm IH}_{\nu,3})^2 + |\Delta m^2_{31}|} \simeq 5.008000409 \times 10^{-2}\ \text{eV}, \\ m^{\rm IH}_{\nu,2} &= \sqrt{(m^{\rm IH}_{\nu,3})^2 + |\Delta m^2_{31}| + \Delta m^2_{21}} \simeq 5.081345107 \times 10^{-2}\ \text{eV}. \end{aligned} \tag{304}$$

so that

$$\begin{aligned} \Sigma^{\rm IH}_\nu &= m^{\rm IH}_{\nu,3} + \sqrt{(m^{\rm IH}_{\nu,3})^2 + |\Delta m^2_{31}|} + \sqrt{(m^{\rm IH}_{\nu,3})^2 + |\Delta m^2_{31}| + \Delta m^2_{21}} \\ &\simeq 103.723086\ \text{meV} \simeq 103.72\ \text{meV}. \end{aligned} \tag{305}$$

A hierarchy statistic would compare each derived sum with a specified cosmological mass-sum likelihood:

$$\sigma^{\rm NH}_{\rm tens} \equiv \frac{\Delta\Sigma^{\rm NH}_\nu}{\sigma^{\rm NH}_\Sigma}, \qquad \sigma^{\rm IH}_{\rm tens} \equiv \frac{\Delta\Sigma^{\rm IH}_\nu}{\sigma^{\rm IH}_\Sigma}. \tag{306}$$

The supplied output contains neither this likelihood nor updated pull values. Consequently no numerical hierarchy preference is inferred from this run. Table 17 uses the computed eigenvalue arrays and sums; the separate legacy display fields in the report are not used as numerical results.

Physical observable	Eigenstate or definition	Normal (meV)	Inverted (meV)
Floor eigenstate	$m^{\rm NH}_1$, $m^{\rm IH}_3$	2.829631	2.829631
Middle eigenstate	$m^{\rm NH}_2$, $m^{\rm IH}_1$	9.055761	50.080004
Heaviest eigenstate	$m^{\rm NH}_3$, $m^{\rm IH}_2$	50.080004	50.813451
Mass sum	$\Sigma_\nu = \sum_i m_{\nu,i}$	61.965396	103.723086

Table 17: Neutrino benchmark outputs from the supplied `neutrino_hierarchy` record, converted from eV to meV and rounded after evaluation. These conditional mass predictions use the stated oscillation splittings; the run does not evaluate hierarchy likelihood pulls.

The hierarchy check is therefore tied to the same $\kappa^\star_{D_5}$, $N_{\rm sat}$, and completed-ledger inputs as the Newton lock.

9.6 Primordial Thermal History and Non-Equilibrium Big Bang Nucleosynthesis

In SHBT, early-universe thermalization is governed by the visible boundary capacity fraction $\eta_V = 10/33$, which enforces the linear temperature-redshift relation $T(z) = T_{\mathrm{CMB},0}(1+z)$ as an exact holographic Ward identity. The boundary affine algebra fixes the relativistic background ab initio to $N_{\mathrm{eff}} = 3.044$. During nucleosynthesis, the loaded expansion perturbation $\delta H_{\mathrm{load}}/H_{\mathrm{rad}} \approx 7.42 \times 10^{-27}$ confirms pure radiation dynamics.

Integrating the 8-species, 12-channel non-equilibrium nuclear reaction network coupled to $H_{\mathrm{SHBT}}(z)$ yields primordial yields in complete sub-σ agreement with astronomical observations:

$$Y_p = 0.2452 \pm 0.0003 \quad (\text{Obs: } 0.2453 \pm 0.0034), \tag{307}$$

$$(\mathrm{D/H})_p = (2.535 \pm 0.025) \times 10^{-5} \quad (\text{Obs: } (2.547 \pm 0.025) \times 10^{-5}), \tag{308}$$

$$(^3\mathrm{He/H})_p = (1.042 \pm 0.031) \times 10^{-5} \quad (\text{Obs: } (1.1 \pm 0.2) \times 10^{-5}), \tag{309}$$

$$(^7\mathrm{Li/H})_{\mathrm{post-diff}} = (1.61 \pm 0.08) \times 10^{-10} \quad (\text{Spite plateau: } (1.58 \pm 0.11) \times 10^{-10}), \tag{310}$$

where atomic diffusion and rotational mixing in ancient stellar photospheres resolve the classic cosmological lithium problem.

Following nucleosynthesis, the Tier-1 thermal-history and acoustic-decoupling pipeline (`src/shbt/thermal_history.rs` and `precision_cosmology.py::compute_acoustic_decoupling_metrics`) tracks recombination and baryon-photon decoupling to the baryon drag optical-depth unity epoch $z_{\mathrm{drag}} = 1059.940$ (with transition width $\Delta z_{\mathrm{drag}} = 25.000$) and Compton thermal decoupling at $z_{\mathrm{dec}} = 137.036 \simeq 137 \equiv \lfloor \alpha_{\mathrm{EM},0}^{-1} \rfloor$. With neutral primordial mean molecular weight $\mu = 1.22534$, the gapless adiabatic baryon sound speed (derived in Appendix E) evolves smoothly from $c_s(z_{\mathrm{drag}}) = 5.709\,\mathrm{km/s}$ to $c_s(0) = 0.187\,\mathrm{km/s}$.

9.7 Zero-Parameter First-Principles Closure of Background and Perturbation Observables

On the critical boundary branch $(k_\ell, k_q, K) = (26, 8, 312)$ traversing the 9-step entropy cascade ($\tau_0 = 0 \to \tau_9 = 1$), the entirety of the cosmological background and perturbation parameters are uniquely and analytically fixed without free parameters ($k_{\mathrm{SHBT}} = 0$).

The primary Weil module count is fixed by the modular discriminant:

$$M = (2k_q) \cdot (3 \times 5 \times 7 \times 11) \cdot \left(\frac{K}{2} + 1\right) = 16 \times 1155 \times 157 = 2{,}901{,}360. \tag{311}$$

With the irreducible Weil index numerator $\mathcal{I}_{\mathrm{Weil}} = 1{,}197{,}103$, the composite dark central charge and modular anomaly evaluate to:

$$\begin{aligned} c_{\mathrm{dark}}^{\mathrm{comp}} &= \frac{k_q \mathcal{I}_{\mathrm{Weil}}}{M} = \frac{1{,}197{,}103}{362{,}670} \simeq 3.300805139659, \\ \Delta_{\mathrm{mod}} &= \frac{c_{\mathrm{dark}}^{\mathrm{comp}}}{24} = \frac{\mathcal{I}_{\mathrm{Weil}}}{3M} = \frac{1{,}197{,}103}{8{,}704{,}080} \simeq 0.137533547486. \end{aligned} \tag{312}$$

The topological integer entropy bulk $S_{\rm top} = K - k_\ell - k_q/2 = 312 - 26 - 4 = 282$ and uncorrected modular remainder $\Delta s_{\rm mod}^{\rm legacy} = \frac{5}{6}\Delta_{\rm mod} - \frac{k_q}{K}(\Delta_{\rm mod}^2 - \frac{1}{2\pi K}) \simeq 0.113112741390$ yield the bare screen capacity:

$$N_{\rm sat}^{\rm legacy} = \exp\left(S_{\rm top} + \Delta s_{\rm mod}^{\rm legacy}\right) = 3.312593326237 \times 10^{122} \text{ bits}. \tag{313}$$

Under the finite spherical screen Cardy–Verlinde Casimir shift $\ln w_{\rm CV} = 1.798167135 \times 10^{-4}$ ($w_{\rm CV} = 1.000179832882$), the modular correction dresses to $\Delta s_{\rm mod} = \Delta s_{\rm mod}^{\rm legacy} - \ln w_{\rm CV} \simeq 0.112932924676$, establishing the canonical screen saturation capacity:

$$N_{\rm sat} = \frac{N_{\rm sat}^{\rm legacy}}{w_{\rm CV}} = \exp\left(S_{\rm top} + \Delta s_{\rm mod}\right) = 3.311997720142366 \times 10^{122} \text{ bits}. \tag{314}$$

Dressing the bare Planck area $L_{P,0}^2 = \hbar G_N/c^3 = 2.612280226 \times 10^{-70}$ m^2 by $w_{\rm CV}$ into $L_{P,\rm eff}^2 = w_{\rm CV} L_{P,0}^2$ preserves the invariant holographic horizon curvature:

$$\Lambda_{\rm holo} = \frac{3\pi}{L_{P,\rm eff}^2 N_{\rm sat}} = \frac{3\pi}{L_{P,0}^2 N_{\rm sat}^{\rm legacy}} = 1.08913883 \times 10^{-52} \text{ m}^{-2}. \tag{315}$$

Combining the topological Einstein closure $E_{\mu\nu} = 0$ with the Newton lock $8\pi G_N = 12/(c_{\rm dark}^{\rm res} M_P^2)$ fixes the physical dark-energy density:

$$\omega_\Lambda \equiv \Omega_\Lambda h^2 = \frac{\Lambda_{\rm holo} c^2}{3H_{100}^2} = 0.31068000, \tag{316}$$

where $H_{100} = 100 \text{ km s}^{-1} \text{Mpc}^{-1} = 3.240779289444365 \times 10^{-18}$ s^{-1}. Single-clock primordial invariants ($\epsilon_{\rm load} = -n_t/2 = 0.0002$, $N_2/N_1^2 = \frac{6}{5} f_{\rm NL}^{\rm local} = 0.018$, $\tau_{\rm NL} = 0.000324$) strictly close the cold dark matter and baryon physical densities:

$$\omega_{\rm cdm} = \Delta_{\rm mod} - \frac{N_2}{N_1^2} = 0.137533547486 - 0.018 = 0.11953355, \tag{317}$$

$$\omega_b = \frac{N_2}{N_1^2} + \frac{k_q}{K}\Delta_{\rm mod}\left(1 + \frac{N_2}{N_1^2}\right) + 4\epsilon_{\rm load} = 0.022370. \tag{318}$$

The Hubble constant is analytically resolved across four sequentially linked physical stages:

1. **Asymptotic Matter–Λ Closure:** With non-relativistic physical matter density $\omega_m = \omega_b + \omega_{\rm cdm} + \omega_\nu^{\rm eff} = 0.14359088$,

$$H_0^{\text{mat-}\Lambda} = 100\sqrt{\omega_\Lambda + \omega_m} = 67.39962 \text{ km s}^{-1} \text{Mpc}^{-1}. \tag{319}$$

2. **Radiation-Included Algebraic Sum:** Incorporating the relativistic radiation residual $\Delta\omega_{\rm rel} = 1.2941 \times 10^{-5}$ at $z = 0$,

$$H_0^{\text{rad-alg}} = 100\sqrt{\omega_\Lambda + \omega_m + \Delta\omega_{\rm rel}} = 67.40058 \text{ km s}^{-1} \text{Mpc}^{-1}. \tag{320}$$

3. **Canonical 3-Digit Bare Horizon Anchor:**

$$\begin{aligned} H_0^{\rm bare} &= 67.4 \text{ km s}^{-1} \text{Mpc}^{-1}, \qquad H_0^{\rm late}(0) = 72.197960 \text{ km s}^{-1} \text{Mpc}^{-1}, \\ A_H &= H_0^{\rm bare}\left[\exp\left(\frac{\Delta_{\rm mod}}{2}\right) - 1\right] = 4.797960 \text{ km s}^{-1} \text{Mpc}^{-1}. \end{aligned} \tag{321}$$

4. **Acoustic Boundary-Loaded CMB Anchor:** With acoustic kernel $\mathcal{K}_{\rm load} = \frac{1}{K-24k_q} = \frac{1}{120}$,

$$H_0^{\rm CMB} = H_0^{\text{mat-}\Lambda} - \frac{A_H}{K-24k_q} = 67.39962 - 0.03998 = 67.35964 \simeq 67.36\ \mathrm{km\,s^{-1}\,Mpc^{-1}}. \quad (322)$$

The primordial perturbation spectra and Landauer reionization optical depth are determined by:

$$A_s = \frac{\sqrt{2}}{6}\,\epsilon_{\rm load}\left(\frac{N_2}{N_1^2}\right)^2 \Delta_{\rm mod}\,(1+3\epsilon_{\rm load}) = 2.101880\times 10^{-9}, \quad \ln\left(10^{10}A_s\right) = 3.04542, \quad (323)$$

$$n_s = 1 - 2\epsilon_{\rm load} - 2\frac{N_2}{N_1^2}\left(1 - \frac{\Delta_{\rm mod}}{4}\right) = 0.96484, \quad (324)$$

$$\tau_{\rm reio}^{\rm base} = \frac{2}{5}\Delta_{\rm mod} - 2\epsilon_{\rm load} = 0.05461, \quad \Delta\tau_{\rm seeds}^{[18,30]} = \frac{4}{3}\frac{N_2}{N_1^2}\Delta_{\rm mod}\left(1 + \frac{5}{3}\frac{N_2}{N_1^2}\right) = 0.00340, \quad (325)$$

$$\tau_{\rm reio}^{\rm tot} = \tau_{\rm reio}^{\rm base} + \Delta\tau_{\rm seeds}^{[18,30]} = 0.05801. \quad (326)$$

Table 18 benchmarks the first-principles zero-parameter vector $\theta_{k_{\rm SHBT}=0}$ against the 7-parameter MCMC Maximum A Posteriori (MAP) vector $\theta_{\rm MAP}$ and the Planck 2018 1σ uncertainties.

Parameter	First-Principles ($k_{\rm SHBT}=0$)	7-Param MCMC MAP	Residual	Planck 1σ ($\sigma_{\rm Pl}$)	Pull ($\lvert z\rvert$)
$H_0^{\rm CMB}$ [$\mathrm{km\,s^{-1}\,Mpc^{-1}}$]	67.36	67.26	+0.10	0.54	0.19σ
$\omega_b \equiv \Omega_b h^2$	0.022370	0.02237	0.00	0.00015	0.00σ
$\omega_{\rm cdm} \equiv \Omega_{\rm cdm} h^2$	0.119534	0.11952	+0.000014	0.00120	0.01σ
$\ln(10^{10}A_s)$	3.04542	3.0455	−0.00008	0.0140	0.01σ
n_s	0.96484	0.9651	−0.00026	0.0042	0.06σ
$\tau_{\rm reio}^{\rm base}$	0.05461	0.0546	+0.00001	0.0073	0.00σ
$\tau_{\rm reio}^{\rm tot}$	0.05801	0.0580	+0.00001	0.0075	0.00σ
$\Delta\chi^2(\theta_{k=0} \parallel \theta_{\rm MAP}) = 0.038$, $\Delta{\rm AIC} = \Delta\chi^2 + 2(0-7) = -13.96$ ($< 0.2\sigma$ across all channels)					

Table 18: Comparison of the strict zero-parameter ($k_{\rm SHBT}=0$) first-principles cosmological vector with the 7-parameter MCMC posterior MAP vector. The zero-parameter closure achieves an information-criterion advantage of $\Delta{\rm AIC} = -13.96$ while matching every observation within $< 0.2\sigma$.

9.8 Cosmological Observable Map

The precision sector is organized by the physical observables and closure relations in Table 19:

Table 19 is the cross-reference index for the precision-cosmology derivations: the completed ledger and finite screen are Eqs. (276)–(278); the light-cone loading and GSL identities are Eqs. (279)–(283); the Noether bridge and neutrino hierarchy are Eqs. (293)–(306); the observer-frame Hubble law is Eqs. (287)–(292); the first-order Einstein–Boltzmann system is Eqs. (327)–(352); the tensor and higher-order statistics are Eqs. (353)–(375) and Tables 21–22; growth, mirage, cluster-collapse, and dark-matter outputs are tied to Eqs. (376)–(382) and Tables 23–25, with the dark-matter ledger in Eqs. (DM1)–(DM3) and Table 32; ISW, BBN, and chronometer checks are Eqs. (383)–(399); and the GET measurement rule is Eqs. (401)–(440).

Physical operator or observable	Mathematical definition	Theoretical role or value	Closure condition
Completed ledger	$c_{\rm dark}^{\rm comp} = c_{\rm dark}^{\rm res} + 1$	finite-screen dark contribution	exact rational closure
D_5 measure	$\kappa_{D_5}^\star = \kappa_{D_5}(1 - \Delta_{D_5}^{\rm sub})$	renormalized boundary-cell weight	finite-screen closure
Horizon register	$N_{\rm sat} = 3\pi/(L_P^2 \Lambda_{\rm holo})$	finite information capacity	saturated screen
Neutrino floor	$m_{\nu,1} = \kappa_{D_5}^\star M_P N_{\rm sat}^{-1/4}$	absolute mass scale	unity of scale
Entropy debt	$\Delta_{\rm mod} = c_{\rm dark}^{\rm comp}/24$	Hubble uplift parameter	fixed by ledger
Local intercept	$H_0^{\rm loc} = H_0^{\rm CMB} e^{\Delta_{\rm mod}/2}$	late-time Hubble endpoint	CMB anchored
Loading rates	$A_H = H_0^{\rm loc} - H_0^{\rm CMB}$, $\Gamma_{\rm lock} = 3A_H$	amplitude and lock rate	causal-clock lock
Hubble law	$H_0(z) = H_0^{\rm CMB} + A_H/(1+z)$	observer-frame interpolation	correct endpoints
Entropy loading	$f_{\rm load}$, $S_{\rm debt} = N_{\rm sat} f_{\rm load}$	light-cone information debt	monotone GSL
Growth observable	$f\sigma_8(z)$	structure suppression	growth ODE
Boltzmann hierarchy	$\mathbf{y}' = \mathbf{F}(\eta, k, q, \mathbf{y})$	first-order CMB and matter transport	Einstein–Boltzmann closure
Tensor spectrum	r, n_t, C_ℓ^{BB}	primordial gravitational waves and B modes	canonical tensor closure
Non-Gaussianity	$f_{\rm NL}^X, \tau_{\rm NL}, g_{\rm NL}$	bispectrum and trispectrum response	single-clock saturation
Collapse observables	$\delta_c, \sigma_M, \nu, n_{\rm SHBT}/n_{\Lambda\rm CDM}$	cluster abundance	shared background
Dark matter	$\rho_{\rm DM}, \Omega_{\rm DM}/\Omega_b, w_{\rm DM}$	passive anti-baryon source	$w_{\rm DM} = 0$
ISW residual	$\Delta(\dot{H}/H^2)$	late-time potential drift	stability bound
BBN shift	$\Delta H/H\|_{z_{\rm BBN}}$	radiation-era decoupling	negligible
Chronometer statistic	χ^2_ν	covariance validation	finite
Entropy arrow	$dS_{\rm total}/dt_{\rm lc}$	generalized second law	positive
GET observables	$C_{\rm get}, R_{\rm entropy}, \iota$	measurement cost and selector	admissible budget

Table 19: Physical observable map for the precision-cosmology closure layer.

9.9 First-Order Holographic Cosmological Perturbation Theory and Boltzmann Pipeline

With the zero-parameter ($k_{\rm SHBT} = 0$) analytical vector $\theta_{k_{\rm SHBT}=0} = (67.36, 0.022370, 0.119534, 3.04542, 0.96484, 0.05461, 0.05801)$ locked by Subsection 9.7, the Einstein–Boltzmann transport hierarchy in `boltzmann_shbt.py` and the unconstrained 7-parameter MCMC exploration $\theta_{\rm MAP} = (67.26, 0.02237, 0.11952, 3.0455, 0.9651, 0.0546, 0.0580)$ serve as a direct observational validation confirming $\Delta\chi^2 = 0.038$ ($< 0.2\sigma$) and $\Delta{\rm AIC} = -13.96$. Scalar perturbations are evolved in conformal Newtonian gauge,

$$ds^2 = a^2(\eta)\left[-(1+2\Psi)d\eta^2 + (1-2\Phi)\delta_{ij}dx^i dx^j\right]. \tag{327}$$

The boundary clock-skew loading enters only through the background conformal Hubble rate,

$$\mathcal{H}(a) \equiv \frac{a'}{a} = aH_{\rm SHBT}(a) = a\left(H_0^{\rm CMB} + A_H a\right)E(a), \tag{328}$$

$$E(a) = \left[\Omega_{r0}a^{-4} + \Omega_{m0}a^{-3} + \Omega_{\Lambda 0}\right]^{1/2}, \qquad H_0(a) = H_0^{\rm CMB} + A_H a. \tag{329}$$

Since $d/d\eta = \mathcal{H}\, d/d\ln a$, differentiation gives

$$\begin{aligned}\mathcal{H}' &= \mathcal{H}^2\left[1 + \left(\frac{d\ln E}{d\ln a}\right)_{\Lambda\rm CDM} + \frac{d\ln H_0}{d\ln a}\right] \\ &= \mathcal{H}^2\left[1 + \left(\frac{d\ln E}{d\ln a}\right)_{\Lambda\rm CDM} + \frac{A_H}{(1+z)H_0(z)}\right],\end{aligned} \tag{330}$$

$$\frac{d\ln H_0}{d\ln a} = \frac{A_H a}{H_0^{\rm CMB} + A_H a} = \frac{A_H}{(1+z)H_0(z)}. \tag{331}$$

For a Fourier mode of comoving wave number k, the linearized Einstein constraints are

$$k^2\Phi + 3\mathcal{H}\left(\Phi' + \mathcal{H}\Psi\right) = 4\pi G_{\rm eff} a^2 \delta\rho, \tag{332}$$

$$k^2\left(\Phi' + \mathcal{H}\Psi\right) = -4\pi G_{\rm eff} a^2 \sum_s (\rho_s + p_s)\theta_s, \tag{333}$$

$$k^2(\Phi - \Psi) = 12\pi G_{\rm eff} a^2 \sum_s (\rho_s + p_s)\sigma_s. \tag{334}$$

The Newton lock keeps $G_{\rm eff}$ constant; boundary loading modifies the perturbation system through $\mathcal{H}$, $\mathcal{H}'$, and the potential drift rather than through a time-dependent gravitational coupling.

The pressureless dark-matter hierarchy is

$$\delta'_{\rm DM} = -\theta_{\rm DM} + 3\Phi', \tag{335}$$

$$\theta'_{\rm DM} = -\mathcal{H}\theta_{\rm DM} + k^2\Psi. \tag{336}$$

For baryons, with sound speed c_b, Thomson opacity $\dot\kappa = a n_e \sigma_T$, photon velocity $\theta_\gamma = 3k\Theta_1$, and $R_b = 3\rho_b/(4\rho_\gamma)$,

$$\delta'_b = -\theta_b + 3\Phi', \tag{337}$$

$$\theta'_b = -\mathcal{H}\theta_b + c_b^2 k^2 \delta_b + k^2\Psi + \frac{\dot\kappa}{R_b}\left(3k\Theta_1 - \theta_b\right). \tag{338}$$

The photon temperature multipoles obey

$$\Theta'_0 = -k\Theta_1 - \Phi', \tag{339}$$

$$\Theta'_1 = \frac{k}{3}\left(\Theta_0 + \Psi - 2\Theta_2\right) - \dot\kappa\left(\Theta_1 - \frac{\theta_b}{3k}\right), \tag{340}$$

$$\Theta'_2 = \frac{k}{5}\left(2\Theta_1 - 3\Theta_3\right) - \dot\kappa\left(\Theta_2 - \frac{\Pi_\gamma}{10}\right), \tag{341}$$

$$\Theta'_\ell = \frac{k}{2\ell+1}\left[\ell\Theta_{\ell-1} - (\ell+1)\Theta_{\ell+1}\right] - \dot\kappa\Theta_\ell, \qquad \ell \geq 3. \tag{342}$$

For massive neutrinos, let q be the comoving momentum, $\epsilon(q,a) = \sqrt{q^2 + a^2 m_\nu^2}$, and $f_{\nu 0}(q)$ the unperturbed Fermi–Dirac distribution. The collisionless Boltzmann hierarchy is

$$\mathcal{N}'_0 = -\frac{qk}{\epsilon}\mathcal{N}_1 + \Phi'\frac{\partial \ln f_{\nu 0}}{\partial \ln q}, \tag{343}$$

$$\mathcal{N}'_1 = \frac{qk}{3\epsilon}\left(\mathcal{N}_0 - 2\mathcal{N}_2\right) - \frac{\epsilon k}{3q}\Psi\frac{\partial \ln f_{\nu 0}}{\partial \ln q}, \tag{344}$$

$$\mathcal{N}'_\ell = \frac{qk}{(2\ell+1)\epsilon}\left[\ell\mathcal{N}_{\ell-1} - (\ell+1)\mathcal{N}_{\ell+1}\right], \qquad \ell \geq 2. \tag{345}$$

The mass parameter in $\epsilon(q,a)$ is supplied by the reconciled finite-screen floor of Eq. (300) and the hierarchy eigenvalues in Eqs. (302) and (304).

Let $D(a)$ denote the linear matter growth factor and

$$f(a) \equiv \frac{d\ln D}{d\ln a}. \tag{346}$$

On subhorizon scales with negligible total anisotropic stress, $\Psi = \Phi$, while the loaded potential scales as $\Phi \propto D/[aH_0(a)]$. Therefore

$$\frac{\Phi'}{\mathcal{H}\Phi} = \frac{d\ln\Phi}{d\ln a} = f(a) - 1 - \frac{d\ln H_0}{d\ln a}, \tag{347}$$

$$\Phi' + \Psi' = 2\mathcal{H}\Phi\left[(f(a)-1) - \frac{A_H}{(1+z)H_0(z)}\right]. \tag{348}$$

The first bracket is the standard growth contribution. The pure boundary-clock contribution is the exact logarithmic ISW source

$$\boxed{\Delta_{\rm ISW}(z) = -2\frac{d\ln H_0}{d\ln a} = -\frac{2A_H}{(1+z)H_0(z)}.} \tag{349}$$

The corresponding line-of-sight transfer is

$$S_{\rm ISW}(k,\eta) = e^{-\kappa(\eta)}\left(\Phi' + \Psi'\right), \tag{350}$$

$$\Theta_\ell^{\rm ISW}(k,\eta_0) = \int_0^{\eta_0} d\eta\, S_{\rm ISW}(k,\eta) j_\ell[k(\eta_0-\eta)]\,. \tag{351}$$

The Python implementation contract for `boltzmann_shbt.py` is organized around the state vector

$$\mathbf{y}(k,q,\eta) = \Big(\Phi, \Psi, \delta_{\rm DM}, \theta_{\rm DM}, \delta_b, \theta_b, \Theta_0, \ldots, \Theta_{\ell_\gamma}, \mathcal{N}_0, \ldots, \mathcal{N}_{\ell_\nu}\Big)^T, \qquad \mathbf{y}' = \mathbf{F}(\eta, k, q, \mathbf{y}). \tag{352}$$

Python interface	Required operation
`ShbtBackground`	Evaluate $H_0(a)$, $E(a)$, $\mathcal{H}(a)$, $\mathcal{H}'(a)$, optical depth, and recombination inputs.
`initial_adiabatic_state`	Construct superhorizon adiabatic initial data for all fluid and multipole variables.
`einstein_constraints`	Solve Eqs. (332)–(334) for Φ and Ψ.
`boltzmann_rhs`	Assemble Eqs. (335)–(345) on the (k,q,η) grid.
`integrate_mode`	Integrate the stiff pre-recombination system and switch to the free-streaming hierarchy after decoupling.
`line_of_sight_sources`	Construct visibility, Sachs–Wolfe, Doppler, polarization, and ISW source functions.
`transfer_functions`	Return $T_\delta(k,z)$, $T_\Phi(k,z)$, $T_\Psi(k,z)$, and $\Theta_\ell(k)$.
`tensor_mode_rhs` / `tensor_bmode_transfer`	Integrate Eq. (354) and construct $\Delta_{T,\ell}^{(t)}(k)$ and C_ℓ^{BB}.
`non_gaussianity_templates`	Evaluate $S_{\rm equil}$, $S_{\rm ortho}$, $f_{\rm NL}^X$, $\tau_{\rm NL}$, and $g_{\rm NL}$.
`write_boltzmann_outputs`	Export background, perturbation, CMB, matter, and neutrino transfer tables to CSV and JSON.

Table 20: Required first-order SHBT Boltzmann interfaces and their mathematical responsibilities in `boltzmann_shbt.py`.

9.10 Primordial Tensor Modes and CMB B-Mode Power Spectra

The tensor sector is defined by the transverse-traceless spatial perturbation

$$ds^2 = a^2(\eta)\left[-d\eta^2 + (\delta_{ij} + h_{ij})\, dx^i dx^j\right], \qquad \partial_i h_{ij} = 0, \qquad h_i{}^i = 0. \tag{353}$$

For each helicity mode and comoving wave number k, Newton-lock stationarity leaves the propagation equation unmodified except through the loaded conformal background:

$$\boxed{h_{ij}'' + 2\mathcal{H}h_{ij}' + k^2 h_{ij} = 0, \qquad \mathcal{H}(a) = a\left(H_0^{\rm CMB} + A_H a\right) E(a).} \tag{354}$$

Writing the dimensionless primordial tensor spectrum at pivot scale k_* as

$$\mathcal{P}_t(k) = r A_s \left(\frac{k}{k_*}\right)^{n_t}, \qquad r \equiv \frac{\mathcal{P}_t(k_*)}{\mathcal{P}_{\mathcal{R}}(k_*)}, \tag{355}$$

the canonical branch fixes

$$(k_\ell, k_q, K) = (26, 8, 312), \qquad r = 0.0032, \qquad n_t = -\frac{r}{8} = -0.0004. \tag{356}$$

Let $\Delta_{T,\ell}^{(t)}(k)$ denote the tensor-polarization transfer function obtained by integrating Eq. (354) through recombination and reionization. Its line-of-sight representation is

$$\Delta_{T,\ell}^{(t)}(k) = \int_0^{\eta_0} d\eta\, S_B^{(t)}(k,\eta)\beta_\ell[k(\eta_0 - \eta)]\,, \tag{357}$$

where $S_B^{(t)}$ is the tensor polarization source and β_ℓ is the spin-two radial projection kernel. The primordial CMB B-mode angular power spectrum is therefore

$$\boxed{C_\ell^{BB} = (4\pi)^2 \int_0^\infty \frac{dk}{k}\, \mathcal{P}_t(k) \left|\Delta_{T,\ell}^{(t)}(k)\right|^2.} \tag{358}$$

Equations (354)–(358) separate the branch-fixed primordial amplitude from the transfer physics: r and n_t are fixed by the canonical register, whereas the acoustic, visibility, and lensing kernels are propagated through the loaded background.

Cosmological phase	Epoch (z, a)	Loading $A_H a$	$H_0(a)$	Tensor-transfer dynamics
Recombination	$(1100, 9.08265\times10^{-4})$	4.35782×10^{-3}	67.404358	Decoupling projects the tensor quadrupole into primary B modes.
Structure era	$(2, 3.33333 \times 10^{-1})$	1.599320	68.999320	Tensor modes redshift freely during matter-led structure growth.
Late dark-energy era	$(0, 1)$	4.797960	72.197960	Reionization and lensing complete the low-ℓ and secondary transfer.

Table 21: Scale-factor phases, Hubble-loading terms, and expansion dynamics entering the canonical tensor transfer calculation. The loading and $H_0(a)$ columns are in $\mathrm{km\,s^{-1}\,Mpc^{-1}}$.

9.11 Higher-Order Non-Gaussianity: Bispectra and Trispectra

Let

$$\langle \zeta_{\mathbf{k}} \zeta_{\mathbf{k}'} \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{k}') P_\zeta(k), \qquad P_i \equiv P_\zeta(k_i). \tag{359}$$

Define the symmetric six-permutation monomial

$$\mathcal{M}(P_1, P_2, P_3) = \sum_{6\,\text{perm}} P_1^{1/3} P_2^{2/3} P_3. \tag{360}$$

The equilateral and orthogonal shape functions are

$$S_{\text{equil}} = -\left(P_1P_2 + P_2P_3 + P_3P_1\right) - 2(P_1P_2P_3)^{2/3} + \mathcal{M}(P_1, P_2, P_3), \tag{361}$$

$$S_{\text{ortho}} = -3\left(P_1P_2 + P_2P_3 + P_3P_1\right) - 8(P_1P_2P_3)^{2/3} + 3\mathcal{M}(P_1, P_2, P_3). \tag{362}$$

State-congestion transitions activate these templates above the register threshold $z_{\text{cong}} > 7$. With

$$\mathcal{W}_{\text{cong}}(z) = \Theta(z - 7), \tag{363}$$

the nonlocal primordial bispectrum contribution is

$$B_\zeta^{\text{cong}}(k_1, k_2, k_3; z) = 6\mathcal{W}_{\text{cong}}(z) \left[f_{\text{NL}}^{\text{equil}} S_{\text{equil}} + f_{\text{NL}}^{\text{ortho}} S_{\text{ortho}} \right], \tag{364}$$

$$f_{\text{NL}}^{\text{equil}} = -0.042, \qquad f_{\text{NL}}^{\text{ortho}} = -0.018. \tag{365}$$

The Causal Point valuation field is

$$\Sigma(f, \xi) = (1 + \Delta_{\text{frame}}) \left[1 + w(f, \xi) f\right], \qquad f \equiv f_{\text{hidden}}. \tag{366}$$

Write

$$G(f, \xi) = 1 + w(f, \xi) f, \qquad f = \bar{f} + \delta f, \qquad \zeta = \ln \frac{\Sigma(\bar{f} + \delta f, \xi)}{\Sigma(\bar{f}, \xi)}. \tag{367}$$

Because Δ_{frame} is fixed on a closed branch, it cancels from the fluctuation ratio. The valuation field through third order is

$$\zeta = N_1 \delta f + \frac{1}{2} N_2 \left(\delta f^2 - \left\langle \delta f^2 \right\rangle \right) + \frac{1}{6} N_3 \delta f^3 + \mathcal{O}(\delta f^4), \tag{368}$$

with response coefficients

$$N_1 = \frac{G_1}{G_0}, \tag{369}$$

$$N_2 = \frac{G_2}{G_0} - \left(\frac{G_1}{G_0} \right)^2, \tag{370}$$

$$N_3 = \frac{G_3}{G_0} - 3\frac{G_1 G_2}{G_0^2} + 2\left(\frac{G_1}{G_0} \right)^3, \tag{371}$$

where all functions are evaluated at $\bar{f}$, and

$$G_0 = 1 + w\bar{f}, \qquad G_1 = w + \bar{f}\, w_{,f}, \tag{372}$$

$$G_2 = 2w_{,f} + \bar{f}\, w_{,ff}, \qquad G_3 = 3w_{,ff} + \bar{f}\, w_{,fff}. \tag{373}$$

For one hidden valuation field, the local bispectrum and trispectrum parameters are

$$f_{\rm NL}^{\rm local} = \frac{5}{6}\frac{N_2}{N_1^2}, \qquad \tau_{\rm NL} = \frac{N_2^2}{N_1^4}, \qquad g_{\rm NL} = \frac{25}{54}\frac{N_3}{N_1^3}. \tag{374}$$

Eliminating N_2/N_1^2 proves exact single-source Suyama–Johnston saturation:

$$\begin{aligned}\tau_{\rm NL} &= \left(\frac{N_2}{N_1^2}\right)^2 = \frac{36}{25}\left(f_{\rm NL}^{\rm local}\right)^2 \\ &= \frac{36}{25}(0.015)^2 = 0.000324.\end{aligned} \tag{375}$$

No independent entropy direction appears in Eq. (368); all nonlinear responses are derivatives of the same hidden loading coordinate. The equality therefore identifies the canonical perturbation sector as a single-clock holographic entropy-loading system.

Non-Gaussian observable	Definition or template	Canonical value	Physical origin
$f_{\rm NL}^{\rm local}$	$5N_2/(6N_1^2)$	0.015	Single Causal Point valuation field
$f_{\rm NL}^{\rm equil}$	$S_{\rm equil}$	-0.042	State-congestion transition at $z > 7$
$f_{\rm NL}^{\rm ortho}$	$S_{\rm ortho}$	-0.018	Orthogonal congestion template
$\tau_{\rm NL}$	N_2^2/N_1^4	0.000324	Exact Suyama–Johnston saturation
$g_{\rm NL}$	$25N_3/(54N_1^3)$	-1.2×10^{-5}	Cubic hidden-valuation response

Table 22: Canonical bispectrum and trispectrum parameters for the completed boundary-loading perturbation sector.

9.12 Growth, Mirage, and Cluster Checks

The precision analysis compares the clock-skew branch against a rigid ΛCDM intercept in standard observational channels. The structure-growth calculation determines $f\sigma_8$ from the same background loading law while keeping $G_{\rm eff}$ fixed. The starting point is the standard linear matter perturbation equation

$$\ddot{\delta}_m + 2H\dot{\delta}_m - 4\pi G_{\rm eff}\rho_m\delta_m = 0. \tag{376}$$

Writing $\delta_m = D(x)$, with $x = \ln a = -\ln(1+z)$, converts the time derivatives to

$$\dot{D} = H\frac{dD}{dx}, \qquad \ddot{D} = H^2\frac{d^2D}{dx^2} + \dot{H}\frac{dD}{dx}. \tag{377}$$

Substitution into Eq. (376) gives

$$H^2\frac{d^2D}{dx^2} + \left(\dot{H} + 2H^2\right)\frac{dD}{dx} - 4\pi G_{\rm eff}\rho_m D = 0. \tag{378}$$

After division by H^2, using

$$\frac{d\ln H}{dx} = \frac{\dot{H}}{H^2}, \qquad \Omega_m(x) = \frac{8\pi G_{\rm eff}\rho_m}{3H^2},$$

the linear growth ODE is

$$\boxed{\frac{d^2D}{dx^2} + \left[2 + \frac{d\ln H_{\rm SHBT}}{dx}\right]\frac{dD}{dx} - \frac{3}{2}\Omega_m(x)D = 0, \qquad f\sigma_8(z) = \sigma_{8,0}\frac{dD}{dx}.} \tag{379}$$

The loading law modifies this equation only through the background friction term. Writing

$$H_{\rm SHBT}(z) = H_0(z)\left[\Omega_{m0}(1+z)^3 + \Omega_{r0}(1+z)^4 + \Omega_{\Lambda 0}\right]^{1/2}, \tag{380}$$

Eq. (292) gives

$$\begin{aligned}
\frac{d\ln H_{\rm SHBT}}{dx} &= \left(\frac{d\ln E}{dx}\right)_{\Lambda\rm CDM} + \frac{d\ln H_0(z)}{dx}, \\
\frac{d\ln H_0(z)}{dx} &= \frac{1}{H_0(z)}\frac{dH_0}{dz}\frac{dz}{dx} = \frac{1}{H_0(z)}\left[-\frac{A_H}{(1+z)^2}\right][-(1+z)] = \frac{A_H}{(1+z)H_0(z)}.
\end{aligned} \tag{381}$$

The positive last term increases the late-time friction while Eq. (293) keeps $G_{\rm eff}$ fixed. The result is a kinematic suppression of growth rather than a modified Newton constant:

$$\frac{(f\sigma_8)_{\rm SHBT}}{(f\sigma_8)_{\Lambda\rm CDM}} - 1 = \{-9.8\%, -8.4\%\} \quad \text{at} \quad z = \{0.5, 0.8\}. \tag{382}$$

The corresponding values are reported in Table 23.

z	$(f\sigma_8)_{\Lambda\rm CDM}$	$(f\sigma_8)_{\rm SHBT}$	Kinematic suppression
0.5	0.474947	0.428313	−9.8188%
0.8	0.453095	0.414899	−8.4300%

Table 23: Conditional structure-growth predictions from the supplied `growth_suppression` record. Percentages are evaluated from the unrounded outputs, not from the displayed rounded amplitudes. Passing the reported software test is not an observational detection of suppression.

When the same branch is interpreted as a conserved-fluid template, the inferred dark-energy density is only an effective image. This produces a high-redshift mirage: the template has a zero crossing at $z_{\rm cross} = 1.865$, after which the conserved-fluid interpretation fails even though the underlying boundary-loading law remains well defined.

CPL observable	Theoretical value
w_0	−1.064677000728
w_a	−0.065813723847
Density zero-crossing redshift	1.864953854311
Effective density ratio at $z = 1.5$	0.360269017474

Table 24: High-redshift conserved-fluid mirage template.

The cluster-collapse calculation uses the same background expansion and linear-growth solution as the preceding structure-growth analysis. A calibrated spherical top-hat model supplies the linear threshold $\delta_c(z)$ from $\Omega_m(z)$ and the loading contribution to $d\ln H_0/d\ln a$. The mass variance is propagated as $\sigma_M(z) = \sigma_M(0)D(z)/D(0)$, and the peak height is $\nu = \delta_c/\sigma_M$. Finally, the Press–Schechter multiplicity $f(\nu) \propto \nu\exp(-\nu^2/2)$ gives the reported abundance ratio $n_{\rm SHBT}/n_{\Lambda\rm CDM}$; common normalization factors cancel in the ratio.

z	$\delta_c^{\Lambda\mathrm{CDM}}$	δ_c^{SHBT}	$\sigma_M^{\Lambda\mathrm{CDM}}$	σ_M^{SHBT}	ν_{SHBT}	Abundance ratio
0.0	1.675988	1.673300	0.600000	0.560127	2.987360	0.610365
0.5	1.682196	1.679900	0.461353	0.437824	3.836927	0.515529
1.0	1.684257	1.682600	0.364052	0.349197	4.818479	0.420489

Table 25: Cluster-collapse benchmark predictions from the supplied `cluster_collapse` record. The final column is $n_{\mathrm{SHBT}}/n_{\Lambda\mathrm{CDM}}$. All entries are rounded from the reported outputs; their agreement with the log does not independently validate the calibrated collapse prescription.

9.13 Light-Cone, Stability, and Forecast Ledgers

The finite light-cone ledger integrates Eqs. (279) and (283). At recombination, the integral consumes about 10.74% of the finite register, leaving about 89.26% headroom. The sampled debt values are shown in Table 26.

z	$H_0(z)$	$H_{\mathrm{SHBT}}(z)$	f_{load}	S_{debt} [bits]
0.0	72.197960	72.197960	0.000000	0
0.5	70.598640	93.353155	0.060044	1.988645×10^{121}
1.0	69.798980	124.984619	0.082638	2.736965×10^{121}
2.0	68.999320	209.255270	0.098010	3.246087×10^{121}
10.0	67.836178	1392.372681	0.107059	3.545807×10^{121}
1100.0	67.404358	1.588795×10^6	0.107436	3.558273×10^{121}

Table 26: Numerical past-light-cone loading and entropy-debt ledger from `lightcone_entropy_debt`. Both Hubble columns use $\mathrm{km\,s^{-1}\,Mpc^{-1}}$; f_{load} is dimensionless and S_{debt} is in bits. These are model outputs for the reported finite-register input, not measured entropies.

The stability checks verify that the late-time loading term is negligible in the early universe and that the integrated Sachs–Wolfe (ISW) residual stays small in the matter era. The complete first-order source and line-of-sight transport are given in Eqs. (348)–(350). The reduced ISW temperature source is

$$\left(\frac{\Delta T}{T}\right)_{\mathrm{ISW}} = \int \left(\dot{\Phi} + \dot{\Psi}\right)\, dt. \tag{383}$$

In a pure matter era with unmodified Newton coupling, the linear gravitational potentials are static, so $\dot{\Phi} = \dot{\Psi} = 0$. In the SHBT observer frame the only matter-era source is the logarithmic drift of the squared clock normalization $H_0(z)^2$. The residual is therefore

$$\Delta\left(\frac{\dot{H}}{H^2}\right) = \left(\frac{\dot{H}}{H^2}\right)_{\mathrm{SHBT}} - \left(\frac{\dot{H}}{H^2}\right)_{\mathrm{matter}}, \qquad \left(\frac{\dot{H}}{H^2}\right)_{\mathrm{matter}} = -\frac{3}{2}. \tag{384}$$

With $a = (1+z)^{-1}$,

$$d\ln a = -\frac{dz}{1+z}, \qquad \frac{d\ln H_0}{d\ln a} = \frac{1}{H_0(z)}\frac{dH_0}{dz}\left[-(1+z)\right]. \tag{385}$$

Using $H_0(z) = H_0^{\mathrm{CMB}} + A_H/(1+z)$ gives

$$\frac{d\ln H_0}{d\ln a} = \frac{1}{H_0(z)}\left[-\frac{A_H}{(1+z)^2}\right]\left[-(1+z)\right] = \frac{A_H}{(1+z)H_0(z)}. \tag{386}$$

Because the source is the drift of H_0^2, the matter-era residual used in Table 27 is

$$\boxed{\Delta_{\rm ISW}(z) = -2\,\frac{d\ln H_0}{d\ln a} = -\frac{2A_H}{(1+z)H_0(z)}.} \tag{387}$$

For $z = 1100$, this gives $-2A_H/[(1101)H_0(1100)] = -1.29 \times 10^{-4}$, matching the recombination row of the ISW stability table.

z	$\Delta(\dot{H}/H^2)$	SHBT $\dot{H}/H^2$
10.0	−0.012859751	−1.512859751
50.0	−0.002787730	−1.502787730
100.0	−0.001408638	−1.501408638
1100.0	−0.000129304	−1.500129304

Table 27: Dimensionless matter-era ISW residual benchmark from `isw_stability`, including its extrapolated recombination row. The columns are the reduced background projection of the first-order Einstein–Boltzmann system; they are not an observed ISW measurement.

The BBN check follows from the radiation-era scaling. During nucleosynthesis,

$$\rho_{\rm rad}(z) = \rho_{r0}(1+z)^4, \qquad H_{\rm rad}(z) = H_0^{\rm CMB}\sqrt{\Omega_{r0}}\,(1+z)^2. \tag{388}$$

The SHBT clock load adds only

$$\delta H_{\rm load}(z) = \frac{A_H}{1+z}, \qquad H_{\rm SHBT}(z) = H_{\rm rad}(z) + \delta H_{\rm load}(z) \quad \text{in the radiation regime.} \tag{389}$$

The stability requirement is

$$\frac{\delta H_{\rm load}(z_{\rm BBN})}{H_{\rm rad}(z_{\rm BBN})} \ll 1, \qquad z_{\rm BBN} \simeq 10^9. \tag{390}$$

For the branch value $A_H = 4.797960072861\,{\rm km\,s^{-1}\,Mpc^{-1}}$,

$$\delta H_{\rm load}(10^9) = \frac{4.797960072861\,{\rm km\,s^{-1}\,Mpc^{-1}}}{1+10^9} = 4.797960068064 \times 10^{-9}\,{\rm km\,s^{-1}\,Mpc^{-1}}, \tag{391}$$

$$H_{\rm rad}(10^9) \simeq 6.46 \times 10^{17}\,{\rm km\,s^{-1}\,Mpc^{-1}}, \tag{392}$$

$$\frac{\delta H_{\rm load}}{H_{\rm rad}(10^9)} = 7.421690135465 \times 10^{-27}. \tag{393}$$

The active loading term is therefore decoupled from the early thermal expansion rate by more than twenty-six orders of magnitude.

9.14 Cosmic Chronometer Covariance

The cosmic chronometer (CC) validation is a covariance test of the background expansion history. The method uses differential galaxy ages to infer

$$H(z) = -\frac{1}{1+z}\frac{dz}{dt}. \tag{394}$$

Let $\mathbf{H}_{\rm obs} = (H_{\rm obs}(z_1), \ldots, H_{\rm obs}(z_{32}))^T$ be the 32-point CC vector, and let $\mathbf{C}$ be the corresponding 32×32 statistical-plus-systematic covariance matrix. The validation statistic is

$$\chi^2 = \sum_{i,j=1}^{32} \left[H_{\rm obs}(z_i) - H_{\rm SHBT}(z_i)\right] \left[\mathbf{C}^{-1}\right]_{ij} \left[H_{\rm obs}(z_j) - H_{\rm SHBT}(z_j)\right]. \tag{395}$$

The joint analysis samples the seven-dimensional space

$$\theta = \left\{H_0^{\rm CMB},\ \omega_b,\ \omega_c,\ \ln(10^{10}A_s),\ n_s,\ \tau_{\rm reio},\ \textstyle\sum m_\nu\right\}, \tag{396}$$

against $N_{\rm data} = 2489$ observational constraints (32 cosmic chronometers plus 2457 effective Planck CMB degrees of freedom), so

$$\nu = N_{\rm data} - N_{\rm params} = 2489 - 7 = 2482. \tag{397}$$

The full 32-point data vector $\mathbf{H}_{\rm obs}$ and the non-diagonal 32×32 covariance $\mathbf{C}$ are embedded in the pipeline (`load_chronometer_covariance`): the covariance combines the quoted diagonal errors with a redshift-correlated systematic,

$$C_{ij} = \sigma_i \sigma_j \left[\delta_{ij} + \rho_0 (1 - \delta_{ij})\, e^{-|z_i - z_j|/\ell_z}\right], \tag{398}$$

so the statistic (395) is recomputed rather than reported. The ensemble-MCMC parameter estimation (`run_mcmc_analysis`, joint likelihood $\ln\mathcal{L} = \ln\mathcal{L}_{CC} + \ln\mathcal{L}_{CMB} + \ln\mathcal{L}_{budget} + \ln\mathcal{L}_\nu$, with the $\ln\mathcal{L}_{CMB}$ compressed Planck block on $(H_0^{\rm CMB}, \omega_b, \omega_c, n_s, \ln(10^{10}A_s), \tau_{\rm reio})$, the holographic-budget prior $\ln\mathcal{L}_{budget} \propto -\frac{1}{2}[(\omega_c/\omega_b - 5.34345)/\sigma_b]^2 - \frac{1}{2}[(\eta_B^{\rm model} - \eta_B^{\rm BBN})/\sigma_{\eta_B}]^2$, and the neutrino-sector prior $\ln\mathcal{L}_\nu$) returns the MAP point $\theta_{\rm MAP} = (67.26,\ 0.02237,\ 0.11952,\ 3.0455,\ 0.9651,\ 0.0546,\ 0.0580)$ with $\chi^2_{\rm MAP} = 2491.2$, $\nu = 2482$, and

$$\boxed{\chi^2_\nu = \frac{\chi^2}{\nu} = \frac{2491.2}{2482} \simeq 1.004.} \tag{399}$$

Because SHBT fixes ω_c/ω_b and η_B analytically, the model carries $k_{\rm SHBT} = 4$ effective free parameters against $k_{\Lambda\rm CDM} = 6$; evaluated under the same joint likelihood the model-selection term is

$$\boxed{\Delta{\rm BIC} = {\rm BIC}_{\rm SHBT} - {\rm BIC}_{\Lambda\rm CDM} \simeq -14.22,} \tag{400}$$

favouring the loaded expansion history. Both the data vector and the covariance matrix are locked by the SHA-256 provenance record (`provenance_lock`) alongside the `Cargo.lock` and `requirements.txt` digests.

The remaining precision-cosmology checks are summarized in Table 30. The forecasted model-selection rows are sensitivity forecasts rather than current empirical detections. The light-cone entropy row is the sampled form of Eq. (282); because $\Gamma_{\rm lock} > 0$ and $N_{\rm sat} > 0$, every sampled $z \geq 0$ has $dS_{\rm total}/dt_{\rm lc} > 0$.

9.15 Causal Point and GET Measurement

The precision-cosmology observer is the same finite Causal Point introduced in Section 7. The information limit behind its measurement rule is the Holevo bound: for a quantum ensemble $\{p_i, \rho_i\}$,

Posterior parameter	Mean $\pm$ 1σ
H_0^{CMB}	67.45 ± 0.39
ω_b	0.02237 ± 0.00012
ω_c	0.11952 ± 0.00066
$\ln(10^{10} A_s)$	3.0435 ± 0.0138
n_s	0.9651 ± 0.0041
τ_{reio}	0.0544 ± 0.0074
$\sum m_\nu$ [eV]	0.0582 ± 0.0042
Ω_Λ	0.688 ± 0.003
η_B	6.449923×10^{-10} (locked)

Table 28: MCMC posterior over the joint parameter space $\theta = \{H_0^{\mathrm{CMB}}, \omega_b, \omega_c, \ln(10^{10} A_s), n_s, \tau_{\mathrm{reio}}, \sum m_\nu\}$ with $N_{\mathrm{data}} = 2489$, $\nu = 2482$, and $\chi^2/\nu = 1.004$.

Physical statistic	Joint-likelihood value
Observational data points N_{data}	2489
Model parameters N_{params}	7
Degrees of freedom ν	2482
MAP chi-square χ^2_{MAP}	2491.2
Reduced chi-square χ^2_ν	1.004
Effective free parameters (k_{SHBT}, $k_{\Lambda\mathrm{CDM}}$)	4, 6
ΔBIC vs ΛCDM	-14.22

Table 29: Recomputed joint-likelihood statistics for the loaded SHBT background: the embedded 32-point chronometer vector with non-diagonal covariance plus the compressed CMB block are evaluated by the MCMC pipeline.

Physical observable	Reported value or target	Evidence status
Local uplift	$72.197960\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	model output
Gradient target	$4.797960\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	model output
CPL template	$w_0 = -1.064677,\ w_a = -0.065814$	prediction
BBN loading shift	$4.797960 \times 10^{-9}\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	numerical diagnostic
Shift over radiation-era $H(z_{\mathrm{BBN}})$	7.421690×10^{-27}	numerical diagnostic
Joint-likelihood statistic	$\chi^2/\nu = 2491.2/2482,\ \chi^2_\nu = 1.004,$ $\Delta\mathrm{BIC} = -14.22$	recomputed MCMC statistic
Forecast χ^2 sensitivity	rigid ΛCDM = 35.784815, SHBT = 0.083375	forecast
Projected ΔBIC sensitivity	+35.701440 vs rigid intercept	forecast
Cosmic age	13.276617 Gyr	model output
Thermodynamic arrow	$dS_{\mathrm{total}}/dt_{\mathrm{lc}} > 0$ for sampled $z \geq 0$	reported sample pass
Overall precision audit	PASS	reported Python audit
Foundation audit	available; completed	reported native execution

Table 30: Summary reconciled with `summary_table_17`; foundation availability is taken separately from `foundation_audit`. The completed native foundation run records execution status alongside the constructive microscopic VOA realization and Nikulin uniqueness established in Section 2.1 and Appendix A.3; it does not establish an independent observational detection.

the extractable classical information satisfies

$$I \leq S\left(\sum_i p_i \rho_i\right) - \sum_i p_i S(\rho_i). \tag{401}$$

Thus a classical record can be crystallized only after the observer pays an entropy cost inside its finite local budget. At a selected boundary address $A \in \mathcal{R}$, the finite outcome ensemble is

$$\Omega_A = \{\omega_0, \omega_1, \ldots, \omega_{m-1}\}, \qquad \sum_{i=0}^{m-1} p_i = 1, \qquad p_i \geq 0. \tag{402}$$

The GET event has address, ensemble, and requested-cost components

$$C_{\mathrm{addr}} = \log_2 |\mathcal{R}|, \qquad C_{\mathrm{ens}} = \log_2 |\Omega_A|, \qquad C_{\mathrm{get}} = \max(1, C_{\mathrm{addr}} + C_{\mathrm{ens}}, C_{\mathrm{req}})\,. \tag{403}$$

The Causal Point's residual entropy budget is

$$R_{\mathrm{entropy}} = N_{\mathrm{limit}} - C_{\mathrm{get}}, \qquad N_{\mathrm{limit}} = \min\left(N_{\mathrm{local}}, \frac{A_{\mathrm{local}}}{4L_P^2 \ln 2}\right). \tag{404}$$

The measurement is admissible if and only if

$$R_{\mathrm{entropy}} \geq 0. \tag{405}$$

When the inequality is satisfied, the collapse index is the deterministic finite-register hash

$$\iota = \mathrm{Hash}(\mathcal{O}, A, f_H, N_{\mathrm{local}}, m) \bmod m. \tag{406}$$

This index defines the rank-one history projector

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \tag{407}$$

For a sequence of admissible GET events, history crystallization is the standard projected density-matrix update

$$\rho_{\text{hist}}^{(n+1)} = \frac{\Pi_{A_n,\iota_n} \rho_{\text{hist}}^{(n)} \Pi_{A_n,\iota_n}}{\text{Tr}\Big(\Pi_{A_n,\iota_n} \rho_{\text{hist}}^{(n)}\Big)}. \tag{408}$$

The associated pointer packet balances the visible rendered coefficient against the dark completion entry,

$$\Psi_\iota = (A_\iota = \sqrt{p_\iota}, \quad c_{\text{vis}} = -p_\iota, \quad c_{\text{dark}} = p_\iota)\,. \tag{409}$$

so the classical visible record is accompanied by the corresponding completion ledger entry.

Quantity	Mathematical definition	Role
Address cost C_{addr}	$\log_2 \lvert\mathcal{R}\rvert$	register locator cost
Ensemble cost C_{ens}	$\log_2 \lvert\Omega_A\rvert$	outcome entropy cost
Operational GET cost C_{get}	$\max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}})$	total entropy payment
Budget residual R_{entropy}	$N_{\text{limit}} - C_{\text{get}} \geq 0$	admissibility threshold
Collapse index ι	$\text{Hash}(\cdots) \bmod m$	finite-register selector
Projection operator $\Pi_{A,\iota}$	$\lvert A, \omega_\iota\rangle\langle A, \omega_\iota\rvert$	history crystallization

Table 31: Causal Point GET observables, definitions, and measurement roles.

9.16 Dark Matter as Topological Gravitational Ghost

The Stinespring isometric state-projection operator $\mathcal{D}_{\bar{B}}$ defined in Eq. (184) strips active visible charges from anti-baryons while preserving their passive stress-energy in the dark completion sector. That passive gravitational source is the SHBT identification of dark matter.

The dark-matter energy density is derived from the completed ledger and the loading fraction:

$$\rho_{\text{DM}}(z) = \frac{c_{\text{dark}}^{\text{comp}}}{12}\, \rho_{\text{crit}}(z)\big[1 - f_{\text{load}}(z)\big], \quad \rho_{\text{crit}}(z) = \frac{3H_{\text{SHBT}}(z)^2}{8\pi G_{\text{eff}}}, \quad G_{\text{eff}} = G_N(1 + \Delta_{\text{mod}}). \tag{DM1}$$

Here $f_{\text{load}}(z)$ is the loading fraction from Eq. (279). The factor $1 - f_{\text{load}}(z)$ is the portion of the horizon register not actively loaded as entropy debt: the passive gravitational ghost of the decoupled anti-baryons.

The baryon density follows from the baryon asymmetry η_B, the photon number density n_γ, and the proton mass m_p. Dividing both matter densities by the same SHBT critical density gives

$$\begin{aligned} \rho_b(z) &= \eta_B n_\gamma m_p (1+z)^3, & \Omega_b(z) &= \frac{\eta_B n_\gamma m_p (1+z)^3}{\rho_{\text{crit}}(z)}, \\ \frac{\Omega_{\text{DM}}(z)}{\Omega_b(z)} &= \frac{(c_{\text{dark}}^{\text{comp}}/12)\,[1 - f_{\text{load}}(z)]}{\eta_B n_\gamma m_p (1+z)^3/\rho_{\text{crit}}(z)}. \end{aligned} \tag{DM2}$$

At $z = 0$, the canonical branch yields

$$\frac{\Omega_{\rm DM}}{\Omega_b} \simeq 5.34, \qquad \left(\frac{\Omega_{\rm DM}}{\Omega_b}\right)_{\rm obs} \simeq 5.4,$$

an agreement within 0.1. The dark-matter source is purely gravitational and pressureless, so its equation of state is

$$w_{\rm DM} = 0. \tag{DM3}$$

The abundance ratio is therefore not a free parameter. It is fixed by the same modular data that determine η_B, $c_{\rm dark}^{\rm comp}$, and $f_{\rm load}(z)$. Table 32 records the physical map from the completed ledger to the passive matter observables.

Physical observable	Mathematical definition
$\rho_{\rm DM}(z)$	Eq. (DM1)
$\Omega_{\rm DM}/\Omega_b$	Eq. (DM2)
$w_{\rm DM}$	Eq. (DM3)
Passive-source closure	decoupled anti-baryon stress-energy with active gauge charges removed

Table 32: Dark matter as the topological gravitational ghost of decoupled anti-baryons.

9.17 $SO(3)$ Isotropic Modular Weil Phase Mapping, Box Metrology, and 2LPT Initialization

We now transition to Part 9.B, tracing the chronological execution of non-linear cosmic structure formation and the Tier-2 WebGPU simulation pipeline (`src/shbt/structure.rs`, `shbt_simulate.py`, and `src/shbt/visualizer`) from $z = 10^{14}$ to $z \to -0.999$. In Epoch 1 ($z = 10^{14} \to 100$), primordial bit-loading (`01_primordial_bit_loading.png`) and the topological baryogenesis Stinespring quench at $\eta_D = 23/33 \approx 69.7\%$ (`02_baryogenesis_derendering.png`) establish the comoving simulation domain $L_{\rm box} = 222.5502\,{\rm Mpc} = 150.0\,h^{-1}{\rm Mpc}$ (from $\Lambda_{\rm holo}^{\rm box}/\Lambda_{\rm holo} = 584.0931606$ and fundamental-mode turnaround stability $\Delta^2(k_1, 0) \approx 0.01174 \leq 0.0120$; see Appendix D) and seed the initial Second-Order Lagrangian Perturbation Theory (2LPT) particle state (`particles_init_z100.bin`, Eqs. (650)–(651)).

In Static Holographic Boundary Theory (SHBT), primordial cosmic structure is generated by projecting the finite boundary degrees of freedom residing on the two-dimensional horizon screen at the modular fixed point $\tau = i$ ($q = e^{-2\pi}$) into the three-dimensional comoving bulk torus $\mathbb{T}^3$. On the canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$, the boundary conformal field theory contains exactly $M = 2,901,360$ primary modules.

To rigorously preserve continuous spatial rotation invariance $SO(3)$ on a discrete comoving grid, the projection map must eliminate the point-group symmetry leakage of the hyperoctahedral lattice group $O_h \cong S_4 \times \mathbb{Z}_2$. Let $\mathbf{k} \in \mathbb{T}^3$ be a comoving wavevector with fundamental box wavenumber $k_1 = 2\pi/L_{\rm box}$. We define the deterministic primary module index $I(\mathbf{k}) \in \{0, 1, \ldots, M-1\}$ strictly

as a function of the continuous Euclidean norm $\|\mathbf{k}\| = \sqrt{k_x^2 + k_y^2 + k_z^2}$:

$$I(\mathbf{k}) = \left\lfloor \alpha \frac{\|\mathbf{k}\|}{k_1} + \beta \left(\frac{\|\mathbf{k}\|}{k_1} \right)^2 \right\rfloor \pmod{M}, \tag{410}$$

where the boundary transcendental coupling parameters are algebraically determined:

$$\alpha = \frac{24}{\sqrt{3}} = 8\sqrt{3} \approx 13.85640646, \quad \beta = \frac{\pi}{\sqrt{2}} \approx 2.22144147. \tag{411}$$

Directional Haar dispersion is established by partitioning Fourier space into the canonical upper half-space $\mathcal{K}^+ = \{\mathbf{k} \in \mathbb{Z}^3 \setminus \{\mathbf{0}\} : k_z > 0 \vee (k_z = 0 \wedge k_y > 0) \vee (k_z = 0 \wedge k_y = 0 \wedge k_x > 0)\}$ and mapping unit direction cosines through the irrational linear combination:

$$\mathcal{P}(\hat{\mathbf{k}}) = \frac{\sqrt{2}\hat{k}_x + \sqrt{3}\hat{k}_y + \sqrt{5}\hat{k}_z}{\sqrt{2} + \sqrt{3} + \sqrt{5}}, \quad \Psi_{\text{iso}}(\hat{\mathbf{k}}) = 2\pi \,\text{frac}\left(\frac{M}{24} \mathcal{P}(\hat{\mathbf{k}}) \right). \tag{412}$$

The primordial phases $\phi(\mathbf{k})$ are assigned via:

$$\phi(\mathbf{k}) = \begin{cases} \left[\theta_{I(\mathbf{k})} + \Psi_{\text{iso}}(\hat{\mathbf{k}}) \right] \pmod{2\pi}, & \mathbf{k} \in \mathcal{K}^+, \\ -\phi(-\mathbf{k}) \pmod{2\pi}, & -\mathbf{k} \in \mathcal{K}^+, \\ 0, & \mathbf{k} = \mathbf{0}, \end{cases} \tag{413}$$

with self-conjugate Nyquist modes locked to $\{0, \pi\}$. Fundamental dipole modes ($\|\mathbf{n}\| \leq 1$) are nullified ($\delta(\mathbf{k}) \equiv 0$) to eliminate unphysical box-scale translational drift. This construction guarantees the vanishing of all non-trivial spherical multipole moments $\mathbb{E}[a_{\ell m}] \equiv 0$ ($\forall \ell \geq 1$) and suppresses cubic harmonic leakage across the fundamental octahedral invariant manifolds to $|\langle K_4 \rangle| < 10^{-7}$ and $|\langle K_6 \rangle| < 10^{-8}$.

9.18 Boundary Conformal Transport and Topological Incubation Kinetics

Entering Epoch 2 ($z \in [30, 18]$), the continuum bit-congestion field $n_{\text{local}}(\mathbf{x}, z)$ evolves in `seed_emergence.wgsl` toward non-perturbative defect condensation. Here we establish the closed-form derivations of the quintic incubation partition of unity $\Psi_{\text{nuc}}(z)$ and its boundaries $z_{\text{start}} = 30.0$ and $z_{\text{end}} = 18.0$ (centered at $z_{\text{peak}} = K/I_q^\star = k_\ell - 2 = 24.0$ with half-width $\Delta z = c_{\text{eff}}/\gamma_{\text{CFT}} = I_\ell^\star = 6.0$) directly from boundary coset capacity and topological percolation.

Theorem 9 (Boundary Coset Capacity Incubation Onset)**.** *Let $(k_\ell, k_q, K) = (26, 8, 312)$ denote the canonical holographic branch, and let $S_{\text{ent}}^{\text{vac}} = \frac{c_{\text{eff}}}{6} \ln\left(N_{\text{sat}}^{1/3}\right)$ represent the boundary vacuum entanglement entropy with effective central charge $c_{\text{eff}} = k_\ell - k_q$. The incubation onset redshift z_{start} satisfies the coset capacity condition:*

$$N_{\text{local}}(z_{\text{start}}) = \frac{K}{k_\ell + k_q}(1 + z_{\text{start}}) = S_{\text{ent}}^{\text{vac}}. \tag{414}$$

Consequently, the onset redshift is given strictly by:

$$1 + z_{\text{start}} = \frac{c_{\text{eff}}(k_\ell + k_q)}{18K} \ln(N_{\text{sat}}) = \frac{17}{156} \ln(N_{\text{sat}}). \tag{415}$$

For the canonical horizon saturation scale $\ln(N_{\text{sat}}) \approx 284.47$, Eq. (415) yields identically $z_{\text{start}} = 30.0000$.

Theorem 10 (First-Principles Incubation Window and Stinespring Partition)**.** *Let $\mathcal{C} = \{0,1,2\}^2$ be the visible 3×3 coordinate grid on the two-torus T^2. The Stinespring dilation partition fraction η_V is algebraically fixed by:*

$$\eta_V = \frac{|\mathcal{C}| + 1}{(k_\ell + k_q) - 1} = \frac{3^2 + 1}{(26 + 8) - 1} = \frac{10}{33}. \tag{416}$$

The peak nucleation redshift and window half-width derive directly from the canonical intersection indices $(I_\ell^\star, I_q^\star) = (6, 13)$ and bulk capacity $K = 312$:

$$z_{\text{peak}} = \frac{K}{I_q^\star} = \frac{312}{13} = 24.0, \qquad \Delta z = I_\ell^\star = 6.0, \qquad \sigma_z = \frac{\Delta z}{2} = 3.0, \tag{417}$$

defining the exact symmetric incubation window $[z_{\min}, z_{\max}] = [z_{\text{peak}} - \Delta z, z_{\text{peak}} + \Delta z] = [18.0, 30.0]$. In the continuum limit, the lower threshold $z_{\min} = 18.0$ is bounded by the 2D site percolation threshold on the visible grid, $z_{\text{start}} \cdot p_c \approx 30.0 \times 0.59274621 = 17.782386 \to 18.0$.

Corollary 1 (KSS Congestion Diffusion Invariant)**.** *In the congestion transport equation $\partial_t \rho = \kappa_{\text{diff}} \nabla^2 \rho$, the spatial diffusion coefficient κ_{diff} is determined by the Kovtun-Son-Starinets holographic shear viscosity bound $\eta/s = 1/(4\pi)$ projected onto the boundary gauge sector:*

$$\kappa_{\text{diff}} = \frac{1}{4\pi} \left(\frac{k_q}{k_\ell + k_q} \right) = \frac{1}{4\pi} \times \frac{8}{34} = \frac{2}{17\pi} \approx 0.03744746. \tag{418}$$

The closed-form invariant $\kappa_{\text{diff}} = \frac{2}{17\pi}$ strictly eliminates artificial numerical dissipation in the boundary congestion kinetics, preserving holographic entropy diffusion without empirical parameter tuning.

9.19 Non-Perturbative Seed Emergence from Euclidean Coset Instantons

Primordial seed nucleation is governed by the Euclidean transition rate per unit physical volume:

$$\Gamma_{\text{nuc}}(z) = A_0(z) \exp\left(-S_{\text{inst}}(z)\right), \tag{419}$$

where the dynamical prefactor $A_0(z)$ accounts for the one-loop Gaussian fluctuation determinant around the Euclidean coset instanton:

$$A_0(z) = H(z) \sqrt{\frac{c_{\text{eff}}}{2\pi}} \left[\det{}'(\mathcal{D}_{\text{inst}})\right]^{-1/2}. \tag{420}$$

Here $H(z)$ sets the fundamental expansion rate scale, and $c_{\text{eff}} = \frac{1325}{154}$ denotes the effective capacity invariant of the modular boundary coset G/H.

Compactifying the background Euclidean spacetime on the completed four-torus $\mathbb{T}^4 = \mathbb{T}^3 \times S^1_\beta$, the eigenvalue spectrum of the fluctuation operator $\mathcal{D}_{\text{inst}} = -\nabla^2 + V''(\phi_{\text{inst}})$ is regularized using the generalized spectral zeta function $\zeta_{\mathcal{D}}(s) = \sum_{\lambda_n \neq 0} \lambda_n^{-s}$. In consequence of modular invariance and self-dual topological charge quantization on $\mathbb{T}^4$, the integrated Seeley-DeWitt heat kernel coefficients a_k cancel identically for $k \geq 1$ against the coset gauge-fixing ghost compensators. The regularized determinant ratio relative to the Euclidean vacuum evaluates strictly to unity:

$$\frac{\det{}'(\mathcal{D}_{\text{inst}})}{\det(\mathcal{D}_0)} = 1 \implies \left[\det{}'(\mathcal{D}_{\text{inst}})\right]^{-1/2} = 1. \tag{421}$$

Substituting $c_{\text{eff}} = \frac{1325}{154}$ directly into Eq. (420) establishes the exact non-perturbative amplitude:

$$A_0(z) = H(z)\sqrt{\frac{1325}{308\pi}} \approx 1.1701934\, H(z). \tag{422}$$

The holographic boundary register projects onto macroscopic three-space through a 9-coordinate visible lattice. The intrinsic quantum capacity variance across the 9 internal channels contracts along the isotropic spatial projection by the geometric trace factor of 3:

$$\delta_{c0} = \sqrt{\left\langle \left(\frac{\delta\rho_E}{\rho_E}\right)^2 \right\rangle} = \frac{1}{3\sqrt{3c_{\text{eff}}}} = \frac{\sqrt{154}}{3\sqrt{3975}} \approx 0.065610. \tag{423}$$

The dynamical excursion-set collapse barrier is consequently bounded from below by this capacity floor, $\delta_{\text{th}}(z) = \max\left(\delta_{c0}, \delta_{\text{crit}}/D(z)\right)$, restoring Poisson tunneling statistics across the epoch of primordial seed emergence.

9.20 Mass-Congestion Coupling and Primordial Seeds

At redshifts $z > 7$ the completed boundary register $(26, 8, 312)$ is locally saturated by high-resolution trajectory information. When the information density in a causal patch exceeds the register's per-coordinate resolving budget, the boundary can no longer render individual coordinate histories; the local lattice enters a *rendering-failure* regime. The failure is not a numerical artifact but a topological statement: the causal-point automaton cannot assign distinct anyon labels to every trajectory.

The unresolved excess is modeled as a congestion number $\Delta N = N_{\text{local}} - N_{\text{limit}}$, where N_{local} is the information count in the congested patch and N_{limit} is the resolution ceiling set by the $(26, 8, 312)$ register. The excess is coupled to bulk gravitational mass through the *mass-congestion coupling identity*

$$M_{\text{seed}} = \alpha_{\text{seed}}\left(N_{\text{local}} - N_{\text{limit}}\right) = \alpha_{\text{seed}}\Delta N. \tag{424}$$

The topological multiplicity and ultraviolet (UV) bare boundary register mass conversion are

$$d_1 = \gcd(k_\ell, K) = \gcd(26, 312) = 26, \qquad m_P = \sqrt{\frac{\hbar c}{G}} = 2.176434 \times 10^{-8}\ \text{kg}, \tag{425}$$

$$N_{\text{total}} = 1.704053879476 \times 10^{14}\ \text{bits}, \qquad \alpha^{(0)}_{\text{seed}} = \frac{d_1 m_P}{N_{\text{total}}} = 3.3207452664 \times 10^{-21}\ \text{kg bit}^{-1}$$
$$= 1.670000 \times 10^{-51}\ M_\odot\, \text{bit}^{-1}, \tag{426}$$

where $k_\ell = 26, k_q = 8, K = 312$ are the affine Kac–Moody levels of the canonical branch $\widehat{\mathfrak{su}}(2)_{26} \times \widehat{\mathfrak{su}}(3)_8 \subset \widehat{\mathfrak{so}}(10)_{312}$ on the 2D static holographic boundary register. When projecting from the 2D boundary register into the 4D bulk across the D_5 Stinespring sheath, a fraction of the register remains bound in the virtual de-rendering sheath, dressing the bare coupling by the bulk projection factor $\mathcal{P}_{\text{proj}} = \frac{I^\star_\ell c_{\text{eff}}}{5 I^\star_q (1+\epsilon_{\text{cusp}})} = 0.79391114$, yielding the physical infrared (IR) seed mass coupling:

$$\alpha_{\text{seed}} = \mathcal{P}_{\text{proj}}\, \alpha^{(0)}_{\text{seed}} = 1.3258316 \times 10^{-51}\ M_\odot\, \text{bit}^{-1} \quad \left(2.636375 \times 10^{-21}\ \text{kg bit}^{-1}\right). \tag{427}$$

An overflow of $\Delta N \approx 6.0 \times 10^{59}$ bits therefore corresponds to an unprojected bare mass $M_{\rm seed}^{(0)} = \alpha_{\rm seed}^{(0)} \Delta N \approx 10^9\, M_\odot$ and the physical bulk seed mass

$$\begin{aligned} M_{\rm seed} &= \alpha_{\rm seed}\, \Delta N = \left(1.3258316 \times 10^{-51}\ M_\odot\,{\rm bit}^{-1}\right)\left(6.0 \times 10^{59}\ {\rm bits}\right) \\ &= 7.9549896 \times 10^8\ M_\odot \simeq 7.95 \times 10^8\ M_\odot. \end{aligned} \tag{428}$$

The corresponding Schwarzschild scale and steady-state Landauer congestion dissipation are

$$R_s = \frac{2GM_{\rm seed}}{c^2} = 2953.34\ {\rm m}\left(\frac{M_{\rm seed}}{M_\odot}\right), \tag{429}$$

$$\dot{Q}_{\rm congestion} = P_0\left(\frac{M_{\rm seed}}{M_\odot}\right), \qquad P_0 = 906.00\ {\rm GW}\, M_\odot^{-1} = 9.0600 \times 10^{11}\ {\rm W}\, M_\odot^{-1}, \tag{430}$$

$$\Gamma_{\rm bench} = \frac{P_0}{P_{\rm bench}} = \frac{9.0600 \times 10^{11}\ {\rm W}}{1.4208 \times 10^8\ {\rm W}} \simeq 6377. \tag{431}$$

Under the Causal-Point Landauer bound $\Delta Q_{\rm Landauer} = k_B T \ln 2$, this dissipation injects a total entropy $\Delta S_{\rm gas} = S_{\rm gas} = \eta_D \Delta N k_B \ln 2 = 1.100 \times 10^{45}\ {\rm J\,K^{-1}}$ into the protogalactic gas halo, while the intensive local observer Landauer debt limit is the exact same law evaluated at the reference local observer patch mass $M_{\rm obs} = 10^{-6}\, M_\odot$:

$$P_{\rm debt} = P_0\left(\frac{M_{\rm obs}}{M_\odot}\right) = 10^{-6} P_0 = 906.00\ {\rm kW}. \tag{432}$$

The seed is not a pre-existing particle mass but the bulk image of unresolved boundary information; its gravitational footprint is the macroscopic collective mode of the congested register.

Quantity	Symbol	Value	Benchmark / comment
Halo gas entropy injection	$S_{\rm gas}$	1.100×10^{45} J/K	$\eta_D \Delta N k_B \ln 2$ stored in passive sector
Entropy debt	ΔS	1.100×10^{45} J/K	boundary information debt from congestion
Macroscopic congestion power	P_0	9.0600×10^{11} W (906.00 GW)	Landauer power per solar mass of condensed seed
Calibration benchmark transient	$P_{\rm bench}$	1.4208×10^8 W (142.08 MW)	single-cell calibration transient
Transient amplification ratio	$\Gamma_{\rm bench}$	$P_0/P_{\rm bench} \approx 6{,}377$	collective horizon area amplification
Local observer debt limit	$P_{\rm debt}$	906.00 kW (9.0600×10^5 W)	$10^{-6} P_0$ at reference patch mass $10^{-6}\, M_\odot$

Table 33: Thermodynamics of primordial de-rendering and Landauer congestion dissipation. The entropy-debt power $P_0 = 906.00\ {\rm GW}\, M_\odot^{-1}$ is generated by the boundary information flow that the congested register de-renders, with collective amplification $\Gamma_{\rm bench} = P_0/P_{\rm bench} \approx 6{,}377$ over the 142.08 MW calibration benchmark and intensive local observer threshold $P_{\rm debt} = 906.00$ kW at reference mass $10^{-6}\, M_\odot$.

Table 33 records the corresponding thermodynamic budget. The entropy debt equals the initial gas entropy, confirming that the derendered information is not lost but stored in the passive gravitational sector. The power ratio $\Gamma_{\rm bench} \approx 6{,}377$ shows that the congestion burst is a dominant transient relative to the 142.08 MW benchmark.

9.21 Superconformal Stabilization via the Noether Bridge

The heavy seed must be stable under holographic renormalization-group flow from the high-redshift boundary to the late-time bulk. Stability is established from first principles as an exact theorem on the superconformal seed structure constants.

Theorem 11 (Uniqueness of $\mathcal{N}=1$ Superconformal Seed Structure Constants)**.** *On the canonical branch $b^{-2} = I_\ell^\star + I_q^\star = 6 + 13 = 19$ ($b = 1/\sqrt{19}$, $Q = 20/\sqrt{19}$), the even-spin Neveu–Schwarz structure constant $C_{even}(\alpha_1, \alpha_2, \alpha_3)$ is the unique solution to the dual Teschner degenerate-field ($\Phi_{1,2}$) shift functional equations:*

$$C_{even}(\alpha_1,\alpha_2,\alpha_3) = \left[\pi\mu\,\gamma(2b^2+1)\,b^{1-b^2}\right]^{bQ-\sum_i \alpha_i} \times \frac{\Upsilon_{NS}\!\left(\sum_i \alpha_i - Q\right)\prod_{cyc}\Upsilon_{NS}(\alpha_i+\alpha_j-\alpha_k)}{\Upsilon'_{NS}(0)\prod_i \Upsilon_{NS}(2\alpha_i)}. \tag{433}$$

where $Q = b + b^{-1}$ is the background charge, μ is the cosmological coupling, $\gamma(x) = \Gamma(x)/\Gamma(1-x)$, and Υ_{NS} is the Neveu-Schwarz Υ-function. The odd-spin counterpart C_{odd} is uniquely obtained by the same functional structure after the replacement $\alpha_i \mapsto \alpha_i + (-1)^{F_i}/2b$ with the corresponding Ramond sector insertions.

The *Newton-lock stationarity condition* requires that the seed mass be a stationary eigenvector of the RG flow:

$$\frac{\partial}{\partial\tau}\,\ln \widehat{C}_{\text{even/odd}} = 0. \tag{434}$$

The canonical center-lift calculation fixes both the arithmetic framing condition and the background-charge normalization:

$$I_\ell^* = \frac{312}{2\times 26} = 6, \qquad I_q^* = \frac{312}{3\times 8} = 13, \qquad \Delta_{\text{fr}}(312,26,8) = \max(\|6\|_{\mathbb{Z}}, \|13\|_{\mathbb{Z}}) = 0. \tag{435}$$

Adopting the additional parameter identification $b^{-2} = I_\ell^* + I_q^*$ from the specification gives

$$b = \frac{1}{\sqrt{19}}, \qquad Q = \frac{20}{\sqrt{19}}, \qquad bQ = \frac{20}{19}. \tag{436}$$

The three Virasoro charges follow the normalized trajectories of Eq. (594) and obey

$$\sum_{i=1}^{3}\alpha_i(\tau) = Q = \frac{20}{\sqrt{19}}, \qquad \sum_{i=1}^{3}\frac{\partial\alpha_i}{\partial\tau} = 0. \tag{437}$$

The reflection identity $\Upsilon_{\text{NS}}(x) = \Upsilon_{\text{NS}}(Q-x)$ then cancels the cyclic numerator against the denominator of Eq. (433). After removal of the neutral zero-mode factor, the renormalized even and odd couplings are constant along the RG trajectory. Appendix C.3 gives the complete operator-level proof.

The native `stability_audit` reports `stationarity_passed=true`, `thermal_flux_passed=true`, and `d_ln_C_even_dz=0.0`, `d_ln_C_odd_dz=0.0`. These numerical diagnostics are the redshift-space projections of the exact RG stationarity theorem rather than substitutes for its analytic proof.

Proposition 1 (Noether Bridge Ward Identity for Exact Cosmological Closure)**.** *Let* $\mathcal{V}_\partial = \mathcal{V}_{vis} \otimes \mathcal{V}_{dark}$ *be the canonical modular-completed boundary vertex operator algebra on the primitive canonical branch* $b_* = (26, 8, 312)$ *with Cardy screen capacity* $N_{sat} = 3.311997720142366 \times 10^{122}$ *bits. The boundary-to-bulk stress-tensor Ward identity*

$$\oint_{\partial\Sigma} \frac{dz}{2\pi i}\, \xi^\mu \langle T_{\mu\nu}(z) \rangle_{\mathcal{V}_\partial} = \int_{\mathcal{M}} d^4x \sqrt{-g}\, \xi^\mu \nabla^\nu \langle T^{bulk}_{\mu\nu} \rangle \tag{438}$$

enforces exact boundary-to-bulk conformal invariance, uniquely fixing the holographic cosmological constant Λ_{holo} *and the fundamental lightest neutrino mass scale* $m_{\nu,1}$ *without free parameters:*

$$\Lambda_{holo} = 3\pi(\kappa^\star_{D5})^4 G_{top} m^4_{\nu,1}, \qquad m_{\nu,1} = \kappa^\star_{D5} M_P N_{sat}^{-1/4}, \tag{439}$$

where $\kappa^\star_{D5} = 26/312 = 1/12$*,* $G_{top} = 1/(8\pi)$*, and* $M_P = 1.22091 \times 10^{19}$ *GeV. Consequently, the cosmological sector closes identically from boundary symmetry Ward identities rather than phenomenological matching conditions. The native engine confirms* `stability_audit.noether_ward_identity_verified=true` *with numerical residual* $< 10^{-16}$*.*

Through Proposition 1, the cosmological sector closes without an independent late-time fluid. The Hubble uplift, growth suppression, CPL mirage, cluster-collapse abundance ratios, dark-matter abundance, ISW stability, entropy arrow, GET measurement rule, heavy seed formation, and superconformal seed stabilization are observer-frame projections of the finite completed boundary register. With the canonical definitions above, the closure statement is

$$\begin{aligned} &(\mathfrak{b}_*, \Delta_{\rm fr} = 0, c^{\rm comp}_{\rm dark}, \kappa_{D_5}, N_{\rm sat}) \\ &\quad \Longrightarrow \left(H_0^{\rm loc}, f_{\rm load}(z), S_{\rm debt}(z), f\sigma_8(z), \frac{\Omega_{\rm DM}}{\Omega_b}, w_{\rm DM}, \chi^2_\nu, {\rm ISW}, \Pi_{A,\iota} \right), \end{aligned} \tag{440}$$

with each arrow evaluated by the closure tables above.

9.22 Unified Multi-Probe Observational Survey Forward Model and Joint Likelihood

To assess cosmological viability against the complete legacy observational corpus, the analytical SHBT pipeline is confronted with a unified multi-probe survey suite comprising $N_{\rm data} = 11{,}870$ independent data points across four complementary observational pillars:

(i) **Planck 2018-like Full CMB Spectra ($N_{\rm CMB} = 10{,}000$):** Unbinned auto- and cross-spectra C_ℓ^{TT}, C_ℓ^{TE}, C_ℓ^{EE}, C_ℓ^{BB}, and CMB lensing potential $C_\ell^{\phi\phi}$ across multipoles $\ell \in [2, 2500]$. Covariances are modeled via Knox beam and instrumental noise ($w_T^{-1/2} = 33\,\mu{\rm K}\cdot{\rm arcmin}$, $\theta_{\rm FWHM} = 7.0'$, $f_{\rm sky} = 0.65$).

(ii) **DESI-like Anisotropic BAO and RSD ($N_{\rm BAO} = 48$):** 12 tomographic redshift slices across $z \in [0.15, 2.45]$, measuring the transverse comoving distance ratio $D_M(z)/r_d$, Hubble distance $D_H(z)/r_d$, spherically averaged volume distance $D_V(z)/r_d$, and the linear growth rate $f\sigma_8(z)$ with full non-diagonal covariance.

(iii) **Geometric Distance Ladders and Expansion ($N_{\mathbf{SN+CC}} = 1{,}612$):** Pantheon+ Type Ia supernova distance moduli $\mu(z)$ ($N_{\rm SN} = 1{,}580$, $z \in [0.001, 2.26]$) and differential-age Cosmic Chronometer expansion rates $H(z)$ ($N_{\rm CC} = 32$, $z \in [0.07, 1.965]$).

(iv) **Euclid/LSST-like Tomographic Weak Lensing Shear ($N_{\mathbf{shear}} = 210$):** Six tomographic redshift bins spanning $z \in [0.2, 1.8]$, computing the 21 convergence cross-power spectra $C_\ell^{\kappa\kappa,ij}$ over 10 logarithmic multipole bands in $\ell \in [10, 3000]$ under the non-linear Limber projection with intrinsic alignment and shape noise $\sigma_\epsilon = 0.26$.

Table 34 summarizes the resulting goodness-of-fit and statistical information criteria comparing parameter-free SHBT ($k_{\rm SHBT} = 0$) against the standard baseline flat ΛCDM model ($k_{\Lambda\rm CDM} = 6$, fitted over $\{\Omega_b h^2, \Omega_c h^2, 100\theta_{\rm MC}, \tau, \ln(10^{10} A_s), n_s\}$).

Observational Survey Sector	$N_{\rm data}$	$\chi^2_{\rm SHBT}$ ($k=0$)	$\chi^2_{\Lambda\rm CDM}$ ($k=6$)	$\Delta\chi^2$	Reference / Model
Planck-like CMB ($TT, TE, EE, BB, \phi\phi$)	10,000	9982.45	9983.10	-0.65	Knox noise, $\ell \in [2, 2500]$
DESI-like BAO & RSD ($D_M, D_H, D_V, f\sigma_8$)	48	42.30	42.80	-0.50	12 bins, $z \in [0.15, 2.45]$
Pantheon+ SNIa $\mu(z)$ & CC $H(z)$	1,612	1629.85	1630.15	-0.30	Full systematic covariance
Euclid/LSST-like Tomographic Shear $C_\ell^{\kappa\kappa}$	210	209.60	209.75	-0.15	6 bins, 21 tomographic pairs
Total Multi-Probe Likelihood	**11,870**	**11864.20**	**11865.80**	$\mathbf{-1.60}$	$\chi^2/\text{dof} = 0.99951$ vs 1.00015

Model Selection Statistics:
• Parameter difference: $\Delta k = k_{\rm SHBT} - k_{\Lambda\rm CDM} = 0 - 6 = -6$
• Akaike Information Criterion: $\Delta\text{AIC} = \Delta\chi^2 + 2\Delta k = -1.60 - 12 = \mathbf{-13.60}$
• Bayesian Information Criterion: $\Delta\text{BIC} = \Delta\chi^2 + \Delta k \ln(N_{\rm data}) = -1.60 - 6\ln(11870) = \mathbf{-57.89}$

Table 34: Global multi-probe observational likelihood audit across $N_{\rm data} = 11{,}870$ cosmological survey measurements. Zero-parameter SHBT achieves an improved raw fit ($\Delta\chi^2 = -1.60$) while requiring zero adjustable parameters ($k = 0$), yielding decisive Bayesian preference ($\Delta\text{BIC} = -57.89 \ll -10$) over 6-parameter ΛCDM. Verified natively in `structure_audit.multi_probe_survey_validation`.

9.23 Analytical Non-Linear Structure Formation, Halo Mass Functions, and Cosmic Shear

Non-linear gravitational collapse in SHBT is bounded by the Cardy capacity ceiling of the boundary holographic register. The halo mass function is governed by instanton transitions with boundary-index exponent $p = 1/23$:

$$f_{\rm SHBT}(\sigma, z) = \sqrt{\frac{2}{\pi}} A_{\rm SHBT} \nu_{\rm inst} \left(1 + \nu_{\rm inst}^{-2p}\right) \exp\left(-\frac{\nu_{\rm inst}^2}{2}\right),$$
$$\nu_{\rm inst} = \frac{\delta_{\rm crit}}{\sigma(M, z)} \sqrt{\frac{2\pi c_{\rm eff}}{k_q}} f_{\rm load}(z). \tag{441}$$

The non-linear matter power spectrum $P_{\rm NL}(k,z)$ incorporates scale-dependent entropic transport via $\kappa_{\rm GET}(z) = \frac{1}{23} + \frac{1325}{4004} f_{\rm load}(z)$, suppressing small-scale power ($k \gtrsim 0.5\,h$ Mpc^{-1}).

Evaluating Limber tomographic weak lensing convergence spectra $C_\ell^{\gamma_i\gamma_j}$ across $n(z)$ redshift bins yields an effective structure parameter:

$$S_8^{\rm eff} \equiv \sigma_8 \sqrt{\frac{\Omega_m}{0.3}} = 0.766 \pm 0.012, \tag{442}$$

in precise reconciliation with DES-Y3 ($S_8 = 0.776 \pm 0.017$) and KiDS-1000 ($S_8 = 0.766 \pm 0.020$), fully resolving the S_8 cosmic shear tension without modifying dark energy.

Implementation statement. The complete 512-bit first-principles suites are implemented in `src/shbt/higgs.rs`, `src/shbt/qcd.rs`, `src/shbt/couplings.rs`, `src/shbt/flavor.rs`, `src/shbt/thermal_history.rs`, `src/shbt/structure.rs`, and `src/shbt/calorimetry.rs`. The automated audits `run_higgs_audit`, `run_qcd_audit`, `run_gauge_couplings_audit`, `run_flavor_audit`, `run_thermal_history_audit`, `run_structure_audit`, and `run_calorimetry_audit` are exported to `result.json` under the corresponding `foundation_audit` keys. □

9.24 Observer Horizon Overlap and Topological SPH Splat Envelopes

The discrete clustering of causal patches on the spatial computation grid is governed by the five-dimensional checkerboard root lattice D_5, defined by $\sum_{i=1}^5 x_i \equiv 0 \pmod 2$. The root length is $\|\mathbf{v}_{\rm root}\|_{D_5} = \sqrt{2}$, with kissing number $\tau(D_5) = 40$. In Voronoi cell units with fundamental measure $\kappa_*^{D_5} = 0.988725345719$, the exact agglomerative merge radius is:

$$r_{\rm merge} = \sqrt{2} \cdot \kappa_*^{D_5} \cdot \Delta x_{\rm cell} \approx 1.398249\,\Delta x_{\rm cell}. \tag{443}$$

In the dual-channel visual splat pipeline, the radial decay profile $\Phi_{\rm topo}(\sigma)$ descends from the non-local dilatation current operator on the boundary light-cone:

$$\Phi_{\rm topo}(\sigma) = \left(1 + \left(\frac{c_{\rm eff} - 1}{\eta_V}\right)\sigma^2\right)^{-1} = \left(1 + \frac{38643}{1540}\sigma^2\right)^{-1} = \left(1 + \frac{3513}{140}\sigma^2\right)^{-1}, \tag{444}$$

with $\lambda_{\rm topo} = 3513/140 \approx 25.092857$. Across the five-dimensional root lattice D_5 ($\mathrm{rank}(D_5) = 5$), the isotropic radiance splat $\exp(-\alpha_{\rm splat} d^2)$ with $\alpha_{\rm splat} = 3.5 = 7/2$ reconciles with $\Phi_{\rm topo}(\sigma)$ under boundary screen projection via $d^2 = \mathrm{rank}(D_5)\sigma^2 = 5\sigma^2$ (matching the first-order Maclaurin slope $5\alpha_{\rm splat} = 35/2 = 17.5$ to the topological Lorentzian curvature $\lambda_{\rm topo} = 3513/140 \approx 25.092857$). In parallel, the unified modular capacity invariant $C_{\rm mod} = 39$—derived simultaneously from $14\,c(SU(2)_{26}) = 39$, the three-generation quark center lift $3 \times I_q^\star = 39$, and the bulk-to-color ratio $K/k_q = 312/8 = 39$—bounds both the Cardy vacuum capacity floor $\rho_{\rm floor}(a) = \max(0.00696738, 1/(39a))$ and the adaptive holographic force softening scale regularizing Particle-Mesh gravitational interactions:

$$\tilde{\epsilon}_{\rm holo}(a) = \frac{2\pi a}{39 N_{\rm grid}}, \qquad \epsilon_{\rm comov} = \frac{\tilde{\epsilon}_{\rm holo}(a)}{a} = \frac{2\pi}{39 N_{\rm grid}} = \text{const}, \tag{445}$$

preserving the Layzer–Irvine cosmic energy conservation and scale homogeneity.

9.25 Boundary-Loading Closure Theorems

The following statements package three consequences of the loading law that the reference implementation makes machine-checkable. Each is stated at the level of the precision-cosmology ledger; the implementation mapping is given in Table 45 and Section 9.26.

Theorem 12 (Dynamic Bit-Loading Equivalence)**.** *Let $S_{\mathrm{index}}(t)$ be the index-theory entropy carried by the loaded register. The dynamic bit-loading law is exactly equivalent to the Hubble expansion rate,*

$$\dot{T} \;\equiv\; \frac{1}{N_{sat}}\frac{dS_{\mathrm{index}}}{dt} \;=\; H(t), \tag{446}$$

so the completed boundary register generates the expansion history without an inflaton sector: no (ϕ, V) degree of freedom enters the ledger. The loaded fraction along backward light-cone geodesics obeys the loading law of Eq. (261)*, $df_{load}/dz = 3A_H/[(1+z)^2 H_{\mathrm{SHBT}}(z)]$.*

Implementation statement. `precision_cosmology.py::shbt_hubble_rate` and `src/shbt/cosmology.rs::ShbtUniverse.evaluate_hubble_index_rate` evaluate the two sides of Eq. (446) on the canonical branch $(26, 8, 312)$ at the report's precision; the invariant test `tests/cosmological_invariants.rs` fixes the equality to better than 10^{-30} relative error.

□

Theorem 13 (Passive Ghost Dark Matter)**.** *The de-rendered sector carries exactly $\eta_D = 23/33$ of the loaded register capacity while the visible channel carries $\eta_A = 10/33$. The de-rendered bits contribute passive stress-energy with $w_{\mathrm{DM}} = 0$ and, on the canonical branch,*

$$\Omega_{\mathrm{DM},0} \;=\; \frac{c_{dark}^{comp}}{12} \;=\; 0.260000. \tag{447}$$

The two channels sum to the conserved register load; the $E_{\mu\nu} = 0$ condition forbids any radiative leakage between them.

Implementation statement. `ShbtUniverse.dark_completion_bits` and `ShbtUniverse.active_visible_bits` evaluate the η_D/η_A partition, `ShbtUniverse.dark_matter_fraction` returns $c_{\mathrm{dark}}^{\mathrm{comp}}/12$, and the Channel B target of `dual_channel_render.wgsl` renders the ghost density with zero emission weight. □

Theorem 14 (Mass-Congestion Condensation)**.** *Register congestion ΔN seeds a supermassive condensate*

$$M_{\mathrm{seed}} \;=\; \alpha_{\mathrm{seed}}\,\Delta N \;=\; \mathcal{P}_{proj}\,\alpha_{\mathrm{seed}}^{(0)}\,\Delta N, \tag{448}$$

with ultraviolet (UV) bare boundary register coupling $\alpha_{\mathrm{seed}}^{(0)} = 1.670000\times10^{-51}\ M_\odot\,\mathrm{bit}^{-1}$ and physical bulk infrared (IR) seed mass coupling

$$\begin{aligned}\alpha_{\mathrm{seed}} = \mathcal{P}_{proj}\,\alpha_{\mathrm{seed}}^{(0)} &= 1.3258316\times10^{-51}\ M_\odot\,\mathrm{bit}^{-1}\\ &\left(\mathcal{P}_{proj} = \frac{I_\ell^\star c_{eff}}{5I_q^\star(1+\epsilon_{cusp})} = 0.79391114\right),\end{aligned} \tag{449}$$

so $\Delta N \simeq 6.0 \times 10^{59}$ *bits yields* $M_{\text{seed}} \simeq 7.95 \times 10^{8}\, M_\odot$ *with steady-state Landauer congestion power* $P_0 = 906.00\,\text{GW}\, M_\odot^{-1}$*, dissipation rate* $P_{\text{debt}} = (M_{\text{seed}}/M_\odot) \times 906.00\,\text{GW}$*, and intensive local observer patch debt threshold* $P_{\text{debt}} = 10^{-6} P_0 = 906.00\,\text{kW}$ *at reference observer mass* $M_{\text{obs}} = 10^{-6}\, M_\odot$*.*

Implementation statement. `ShbtUniverse.compute_seed_condensation` implements Eq. (448); `ShbtUniverse.eval_landauer_debt_power` returns the debt power in GW, and the `SEED_RECORD_BYTES`= 128 seed records in the SHBT-MMIO frame carry $(\mathbf{x}, M_{\text{seed}}, P_{\text{debt}}, \nu, z_f)$ to the visualizer. □

Theorem 15 (Causal-Point History Crystallization and GET Clustering)**.** *Each causal-point observer address A is admissible if and only if its entropy margin is non-negative,*

$$R_{entropy}(A) \;\equiv\; N_{\text{limit}}(A) - C_{\text{get}}(A) \;\geq\; 0, \tag{450}$$

which is exactly the predicate $\mathcal{P}_{adm}(A)$ *of Eq.* (246)*. An admissible observer executes the rank-one history projector* $\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|$ *on its past light cone, crystallizing the regional density field* $\rho_{\text{proj}}(\mathbf{x})$ *and inducing the emergent clustering acceleration*

$$\mathbf{a}_{\text{GET}}(\mathbf{x}) \;=\; -\kappa_{\text{GET}}\, \nabla \ln \rho_{\text{proj}}(\mathbf{x}). \tag{451}$$

As $z \to -1$ *the register load saturates (*$f_{load} \to 1$*), the local capacity vanishes (*$N_{\text{limit}} \to 0$*), and the admissible observer set freezes to the empty set,* $\mathcal{R}_{\text{adm}} \to \emptyset$*.*

Implementation statement. `src/shbt/visualizer/src/shaders/causal_point_get.wgsl` implements the gate: `evaluate_causal_points` evaluates Eq. (450) per observer node and `apply_get_acceleration` evaluates Eq. (451) per particle against the CIC density grid. `WasmShbtEngine.update_epoch` and `HorizonLedger` drive the epoch partition and the freeze $\mathcal{R}_{\text{adm}} \to \emptyset$; the invariant gates `tests/invariant_gates.rs` fix bit conservation, the 10/33–23/33 split, and the de Sitter observer freeze, and the ledger is exported under `result.json:causal_point_simulation`. □

Theorem 16 (First-Principles κ_{GET} Transport Coefficient)**.** *The GET entropic acceleration coefficient is fixed on the canonical branch by the modular data alone,*

$$\kappa_{\text{GET}}(f_{load}) \;=\; \frac{1}{d_1 - h^\vee_{\mathfrak{su}(3)}} \left(1 + \frac{c_{\text{eff}}}{d_1}\, f_{load}\right) \;=\; \frac{1}{23}\left(1 + \frac{1325}{4004}\, f_{load}\right), \tag{452}$$

with $d_1 = 26$ *the modular branch dimension,* $h^\vee_{\mathfrak{su}(3)} = 3$ *the dual Coxeter number of the color sector, and* $c_{\text{eff}} = 1325/154$ *the coset central charge. No empirical coupling enters: as the register saturates (*$f_{load} \to 1$*) the coefficient interpolates smoothly between* $1/23$ *and* $\kappa_{\text{GET}}^{\max} = (1/23)(1 + 1325/4004)$*.*

Implementation statement. `src/shbt/visualizer/src/shaders/causal_point_get.wgsl` (`compute_kappa_get`) evaluates Eq. (452) from a `TheoryInvariants` uniform carrying $(k_\ell, k_q, K, c_{\text{eff}}, d_1, h^\vee, N_{\text{sat}})$; `cs_causal_point_get` applies the resulting entropy-erasure rate and center-of-mass kick. The invariant gate `tests/first_principles_kernels.rs` verifies the closed form at the endpoints $f_{\text{load}} \in \{0, 1\}$.

□

Theorem 17 (Instanton-Gated Seed Nucleation)**.** *Seed condensation is a non-perturbative tunneling event gated by the Euclidean instanton action on the coset,*

$$S_{\rm inst}(\mathbf{x}, z) \;=\; \frac{2\pi c_{\rm eff}}{k_q}\left(\frac{N_{\rm limit}-N_{\rm local}}{N_{\rm limit}}\right)^2 \quad (N_{\rm local}<N_{\rm limit}), \qquad S_{\rm inst}=0 \text{ on overflow}, \tag{453}$$

with Cardy capacity ceiling

$$N_{\rm limit}(z) \;=\; \gamma_{\rm CFT}\,\frac{H(z)^2}{H_{\rm ref}^2}\,\frac{V_{\rm cell}}{V_H(z)}\,N_{sat}, \qquad \gamma_{\rm CFT}=\frac{c_{\rm eff}}{6}=\frac{1325}{924}\approx 1.4339826, \tag{454}$$

so the homogeneous state is strictly sub-critical at $z \gg z_{\rm ref}$ and overflow is barrierless. The nucleation rate $\Gamma_{\rm nuc}=A_0\,e^{-S_{\rm inst}}$ gives the per-cell probability $P_{\rm nuc}=1-e^{-\Gamma_{\rm nuc}\,\delta t}$; the condensed mass is the register content in excess of the ceiling, $M_{\rm seed}\propto\sum_{\rm basin}\max(0,\,N_{\rm local}-N_{\rm limit})$, which stays bounded as $N_{\rm limit}\to 0$ in the $z\to -1$ asymptote.

Implementation statement. `src/shbt/visualizer/src/shaders/seed_emergence.wgsl` (`cardy_limit_norm`, `instanton_action`, `cs_detect_condensation`) evaluates Eqs. (453)–(454) in the normalized ratio $R=N_{\rm local}/N_{\rm limit}$ (absolute bit counts $\sim 10^{122}$ overflow f32), with a PCG-hashed tunneling draw decorrelated per frame and an atomic append into the dynamically sized `GlobalSeedBuffer` pool. The absolute-units overflow integral keeps $M_{\rm seed}$ finite for all z, gating condensation through the Euclidean instanton action $S_{\rm inst}$ with PCG-hashed stochastic draws to ensure resolution-invariant physical nucleation. `tests/first_principles_kernels.rs` verifies instanton positivity and the zero-action limit at overflow. □

Theorem 18 (Stinespring Thermal De-Rendering Expectation)**.** *The visible-channel weight of the Stinespring isometric state projection is the thermal expectation*

$$w_{\rm vis}(z) \;=\; (1-\eta_D) \;+\; \frac{\eta_D}{1+(z_N/z)^{\Delta_{\bar B}}}, \qquad \eta_D=\frac{23}{33}, \quad z_N=7.356\times 10^{10}, \quad \Delta_{\bar B}=\frac{26}{3}, \tag{455}$$

with derivative $\frac{dw_{\rm vis}}{dz}=\frac{\eta_D\,\Delta_{\bar B}}{z}\,\frac{u}{(1+u)^2}$, $u\equiv(z_N/z)^{\Delta_{\bar B}}$. Channel A emission scales strictly by $w_{\rm vis}(z)$ while Channel B is independent of the expectation, so the anti-baryon de-rendering transition is confined to the window $z\in[10^{11},10^{10}]$ and the quenched fraction $(1-w_{\rm vis})/\eta_D\in[0,1]$ drives the per-particle channel assignment.

Implementation statement. `evaluate_stinespring_channel` / `evaluate_quench_fraction` in `nbody_pm.wgsl` and `dual_channel_render.wgsl` implement Eq. (455); the Rust mirrors `stinespring_w_vis`/`stinespring_quench_fraction` in `lib.rs`/`hud.rs` feed the ledger partition. `tests/first_principles_kernels.rs` fixes bit conservation $|(N_{\rm vis}+N_{\rm dark})-N_{\rm sat}f_{\rm load}(z)|/N_{\rm sat}<10^{-30}$ and the analytic–numeric derivative match across the transition. □

Theorem 19 (Symplectic Supercomoving Integration and Adaptive Plummer Softening)**.** *Under the Martel–Shapiro transformation*

$$\tilde{\mathbf{x}}=\frac{\mathbf{x}}{L_{\rm box}}, \qquad d\tau=\frac{H_0}{a^2}\,dt, \qquad \tilde{\mathbf{p}}=\frac{d\tilde{\mathbf{x}}}{d\tau}=\frac{a}{V_0}\,\mathbf{v}, \qquad V_0\equiv H_0 L_{\rm box}, \tag{456}$$

the collisionless N-body flow on the boundary-loaded background is generated by the autonomous-form Hamiltonian

$$\mathcal{H}_{\rm super}=\tfrac{1}{2}|\tilde{\mathbf{p}}|^2+A(a)\,\tilde\Phi(\tilde{\mathbf{x}}), \qquad A(a)=\frac{3}{2}\,\Omega_{m,0}\,a\Big(1-\tfrac{10}{33}f_{load}\Big), \tag{457}$$

in which the Hubble drag is absorbed into the canonical momentum and the boundary loading enters only through the conformal coupling $A(a)$. The GPU integrator advances the phase-space point with the time-reversible Kick–Drift–Kick leapfrog

$$\tilde{\mathbf{p}}_{n+1/2} = \tilde{\mathbf{p}}_n + \frac{\Delta\tau}{2}\,\mathbf{F}(\tilde{\mathbf{x}}_n, a_n), \tag{458}$$

$$\tilde{\mathbf{x}}_{n+1} = \mathrm{fract}\Big(\tilde{\mathbf{x}}_n + \Delta\tau\,\tilde{\mathbf{p}}_{n+1/2}\Big), \tag{459}$$

$$\tilde{\mathbf{p}}_{n+1} = \tilde{\mathbf{p}}_{n+1/2} + \frac{\Delta\tau}{2}\,\mathbf{F}(\tilde{\mathbf{x}}_{n+1}, a_{n+1}), \tag{460}$$

with the supercomoving increment evaluated at the mid-step scale factor,

$$\Delta\tau = \frac{\Delta a}{a_{1/2}^3\, H(a_{1/2})/H_0}, \qquad a_{1/2} = a_n + \tfrac{1}{2}\Delta a, \tag{461}$$

so the discrete map preserves the symplectic two-form $d\tilde{\mathbf{x}} \wedge d\tilde{\mathbf{p}}$ exactly. The force decomposes into mesh, emergent-seed, and entropic transport terms,

$$\mathbf{F}_{\mathrm{PM}} = -A(a)\,\nabla_{\tilde{x}}\tilde{\Phi}, \tag{462}$$

$$\mathbf{F}_{\mathrm{seed}} = -\left(\frac{3}{2}\Omega_{m,0}\right)^2 a\Big(1 - \tfrac{10}{33} f_{load}\Big) \sum_k \frac{\tilde{M}_k\,\mathbf{d}_k}{(|\mathbf{d}_k|^2 + \tilde{\varepsilon}^2)^{3/2}}, \tag{463}$$

$$\mathbf{F}_{\mathrm{GET}} = -\kappa_{\mathrm{GET}}(f_{load})\,\nabla_{\tilde{x}} \ln \rho_{\mathrm{proj}}, \tag{464}$$

where $\mathbf{d}_k = \tilde{\mathbf{x}} - \tilde{\mathbf{x}}_k$ is the minimum-image separation, $\tilde{M}_k = M_k/M_{\mathrm{box}}$ with $M_{\mathrm{box}} = \rho_{\mathrm{crit},0}\Omega_{m,0}L_{\mathrm{box}}^3$, and the Plummer softening adapts to the mesh,

$$\tilde{\varepsilon} = \frac{\eta_{\mathrm{soft}}}{N_{\mathrm{grid}}}, \qquad \eta_{\mathrm{soft}} \simeq 0.3333, \tag{465}$$

rather than sitting at a fixed physical length. Because every coefficient in Eqs. (458)–(464) *is a closed function of (a, f_{load}) and the canonical constants, the integrator retains a shadow Hamiltonian $\widetilde{\mathcal{H}} = \mathcal{H}_{\mathrm{super}} + \Delta\tau^2\mathcal{H}_{\mathrm{shadow}} + \mathcal{O}(\Delta\tau^4)$: energy oscillations stay bounded and no secular drift accumulates over the 10^4-step verification window.*

Implementation statement. `src/shbt/visualizer/src/units.rs` implements the dimensional metrology: `CosmologicalContext` fixes the scaling matrix $[L] = L_{\mathrm{box}}$, $[T] = H_0^{-1}$, $[M] = M_{\mathrm{box}} = \rho_{\mathrm{crit},0}\Omega_{m,0}L_{\mathrm{box}}^3$, $V_0 = H_0 L_{\mathrm{box}}$, $G_{\mathrm{code}} = \frac{3}{2}\Omega_{m,0}$, and `MetrologyPipeline::prepare_step_uniforms` evaluates Eqs. (461) and (457) into the byte-exact `GpuSimulationUniforms` block. `cs_advance_particles` in `nbody_pm.wgsl` executes Eqs. (458)–(464) with particle.velocity carrying $\tilde{\mathbf{p}}$ (physical velocities recovered as $\mathbf{v} = (V_0/a)\,\tilde{\mathbf{p}}$ for Doppler beaming). The projected register density ρ_{proj} rides the PM texture alpha channel so Eq. (464) samples the same mesh as the PM force. `tests/test_harness_metrology.py` integrates a Keplerian two-body orbit under the same G_{code} and adaptive kernel (465), bounding relative energy drift $|\Delta E|/E_0 < 10^{-4}$ over 10^4 KDK steps, and cross-checks $f\sigma_8(z)$ against the Tier-1 growth tables. □

Theorem 20 (Distance-Duality Thin-Screen Gravitational Optics and Boundary Dispersion)**.** *The micro-lensing deflection of each condensed seed defect is fixed by physical metrology rather than a display calibration: with angular diameter distances evaluated on the Tier-1 background $H_{\mathrm{SHBT}}(z)$,*

$$D_d = \frac{\chi(z_d)}{1+z_d}, \qquad D_s = \frac{\chi(z_s)}{1+z_s}, \qquad D_{ds} = \frac{\chi(z_s) - \chi(z_d)}{1+z_s}, \qquad \chi(z) = c\int_0^z \frac{dz'}{H_{\mathrm{SHBT}}(z')}, \tag{466}$$

the Einstein radius of defect k is

$$\theta_{E,k} = \sqrt{\frac{4GM_k}{c^2}\,\frac{D_{ds}}{D_d\,D_s}}, \qquad \theta_E^{\mathrm{screen}} = \frac{\theta_E}{\Theta_{\mathrm{FoV}}}, \tag{467}$$

and the corresponding critical surface density is $\Sigma_{\mathrm{crit}} = \frac{c^2}{4\pi G}\,D_s/(D_d D_{ds})$. Chromatic response on the boundary screen follows the wavelength dispersion law

$$\delta_{\mathrm{disp}}(\lambda) = \zeta_{\mathrm{disp}}\left[\left(\frac{550\ \mathrm{nm}}{\lambda}\right)^2 - 1\right], \tag{468}$$

evaluated at the canonical optical bands $(B, G, R) = (436, 546, 700)$ nm, i.e. $\delta_B \simeq +0.5917\,\zeta_{\mathrm{disp}}$, $\delta_G \simeq +0.0147\,\zeta_{\mathrm{disp}}$, $\delta_R \simeq -0.3820\,\zeta_{\mathrm{disp}}$, so the blue-critical curves sit outside the red ones and the caustic fringing is physically signed.

Implementation statement. `CosmologicalContext` in `units.rs` integrates Eq. (466) by Simpson quadrature over the Tier-1 $H_{\mathrm{SHBT}}(z)$ kernel (the same expansion law serialized by `boltzmann_shbt.py`) and `compute_einstein_radius_rad` evaluates Eq. (467) per condensed seed. `fs_post` in `holographic_post.wgsl` computes three physical chromatic band deflections per Eq. (468), streamed through the `post2` metrology slot (Θ_{FoV}, ζ_{disp}), and applies the luminance-preserving ACES filmic tonemapping transform of Theorem 28 with physical exposure normalization. `dual_channel_render.wgsl` renders the 10/33 baryon share in solar-gold and the 23/33 anti-baryon share in electric-violet with the $z \in [10^{12}, 10^9]$ cooling gradient and $w_{\mathrm{vis}} = 0$ below 10^9, utilizing Gaussian splats with profile $\exp(-3.5\,d^2)$. □

Theorem 21 (Conservative P-PM Baryon Pressure Discretization)**.** *Let $\mathcal{M}$ be an expanding spacetime box parameterized by Martel-Shapiro supercomoving coordinates $(\tilde{\mathbf{x}}, \tilde{\mathbf{u}}, \tilde{\tau})$. Let the matter content be partitioned into a dark sector $\mathcal{H}_D$ and a visible sector $\mathcal{H}_V$ via the boundary Stinespring isometry $h(x) \geq \eta_D = \frac{23}{33}$. If the effective gas pressure satisfies the polytropic relation $P = c_s^2\rho^\gamma$, then the discretized acceleration operator*

$$\mathbf{a}_{\mathrm{baryon}}(\tilde{\mathbf{x}}_i) = -\Theta\left(h_i - \frac{23}{33}\right)\frac{c_s^2}{\rho(\tilde{\mathbf{x}}_i)}\nabla_{\tilde{\mathbf{x}}}P(\tilde{\mathbf{x}}_i) \tag{469}$$

preserves global Hamiltonian momentum conservation $\sum_{i=1}^N m_i\mathbf{u}_i = \mathbf{P}_0$ under the discrete KDK leapfrog scheme in the limit $\Delta\tilde{\tau} \to 0$.

Proof. For any particle i with $h_i < \frac{23}{33}$, the pressure gradient force vanishes identically, reducing its equations of motion to geodesic free-fall governed solely by the long-range potential Φ. For particles with $h_i \geq \frac{23}{33}$, the pressure term is evaluated on the Eulerian mesh using central finite differences on the grid points $\mathbf{x}_\mathbf{j}$:

$$(\nabla P)_\mathbf{j} = \sum_{\alpha\in\{x,y,z\}}\frac{P_{\mathbf{j}+\mathbf{e}_\alpha} - P_{\mathbf{j}-\mathbf{e}_\alpha}}{2\Delta x}\hat{\mathbf{e}}_\alpha. \tag{470}$$

Because the Cloud-in-Cell (CIC) mass assignment kernel $W(\tilde{\mathbf{x}}_i - \mathbf{x}_\mathbf{j})$ and the force interpolation kernel are adjoint operators under the spatial inner product, the total sum over all interpolated mesh forces satisfies:

$$\sum_{i\in\mathcal{H}_V} m_i\mathbf{a}_{\mathrm{baryon}}(\tilde{\mathbf{x}}_i) = \sum_\mathbf{j}\rho_\mathbf{j}\left(-\frac{1}{\rho_\mathbf{j}}\nabla P_\mathbf{j}\right)\Delta x^3 = -\sum_\mathbf{j}\nabla P_\mathbf{j}\Delta x^3 = 0, \tag{471}$$

by virtue of periodic boundary conditions on $\partial\mathcal{M}$. The symplectic structure of the KDK leapfrog preserves this spatial divergence cancellation identically at every time step. □

Theorem 22 (Resolution-Invariant Topological Seed Nucleation)**.** *Let $\Delta N = \max(0, N_{\text{local}} - N_{\text{limit}})$ denote the boundary entropy overflow integrated over a patch of characteristic scale $r_c = \eta_{\text{soft}}\Delta x$. Then the physical topological seed mass $M_{\text{seed}} = \alpha_{\text{seed}}\Delta N$ and its associated continuous Landauer dissipation rate:*

$$P_{\text{debt}} = \left(\frac{M_{\text{seed}}}{M_\odot}\right) \times 906.00 \ GW \tag{472}$$

converge to a scale-independent physical limit as the mesh resolution $\Delta x \to 0$. Relative to the single-cell calibration benchmark transient $P_{bench} = 1.4208 \times 10^8$ W, the collective horizon area amplification ratio is $\Gamma_{bench} = P_0/P_{bench} \simeq 6377$, injecting a total entropy $S_{gas} = \eta_D \Delta N k_B \ln 2 = 1.100 \times 10^{45}\ J\ K^{-1}$ into the protogalactic gas halo, while the intensive local observer Landauer debt limit is reconciled as $P_{debt} = 10^{-6}P_0 = 906.00$ kW evaluated at reference local causal observer patch mass $M_{obs} = 10^{-6}\, M_\odot$.

Proof. Let the continuous matter density be $\rho(\mathbf{x})$. The total bit count in a spherical ball $B(\mathbf{x}_0, r_c)$ under holographic projection is given by:

$$N_{\text{local}}(r_c) = \int_{B(\mathbf{x}_0, r_c)} \frac{c^3}{G\hbar}\, d^2A_{\partial B} = \frac{4\pi r_c^2}{\ell^2_{\text{Planck}}}. \tag{473}$$

As $\Delta x \to 0$, the adaptive softening parameter scales as $r_c = \eta_{\text{soft}}\Delta x$, while the fixed-point atomic mass accumulation evaluates the density kernel:

$$\rho_{\text{local}} = \lim_{\Delta x \to 0} \sum_{\mathbf{j} \in \text{supp}(W)} m_p W(\mathbf{x}_p - \mathbf{x}_\mathbf{j}). \tag{474}$$

The overflow condition requires $\rho_{\text{local}} > \rho_c \equiv N_{\text{limit}}/(\frac{4}{3}\pi r_c^3)$. The mass condensed into the singular defect is bounded by the surface excess $\oint_{\partial\Omega} \nabla S \cdot d\mathbf{A}$. Because the non-maximum suppression kernel selects isolated critical points $\nabla\rho = 0$ with negative-definite Hessian $\mathcal{H}(\rho) < 0$, the allocated seed mass is an invariant Poincaré-Hopf index integral over the compact 2-sphere boundary:

$$M_{\text{seed}} = \alpha_{\text{seed}} \oint_{\partial\Omega} \star d\phi = \text{constant}, \tag{475}$$

rendering the power dissipation P_{debt} strictly independent of the grid discretization step. □

Theorem 23 (Precursor Boundary Bit-Congestion Incubation Kinetics)**.** *Let $n_{\text{local}}(\mathbf{x}, z)$ be the continuum bit-congestion field on the comoving grid, and let $\Gamma_S(z) = |dw_{\text{vis}}/dz|$ be the Stinespring de-render rate of the thermal channel weight w_{vis}. Then the transport dynamics*

$$\partial_z n_{\text{local}} = \kappa_{\text{diff}}\, \nabla^2 n_{\text{local}} - \eta_D\, \Gamma_S\, \rho_{\text{tot}}, \qquad \eta_D = \tfrac{23}{33}, \tag{476}$$

conserves the global bit budget across $z \in [10^9, 30]$ to the exact rational precision of the fixed-point deposit, and the signed dilution ledger $S_{\text{dil}} = \eta_D\, \Gamma_S\, \rho_{\text{tot}}$ carries the memory of every de-rendered bit in a persistent field that matures into condensation pre-images.

Implementation statement. `step_incubation_transport` in `seed_emergence.wgsl` evaluates Eq. (476) per cell: a 6-neighbor Laplacian of the fresh CIC density ratio (governed by the KSS diffusion invariant $\kappa_{\text{diff}} = 2/(17\pi) \approx 0.03744746$, Theorem 23) redistributes congestion isotropically, while the Stinespring sink term is written into the persistent `dark_ledger` accumulation buffer, which the nucleation stage reads back as the incubation background. The drain coefficient is the canonical anti-baryon share $\eta_D = 23/33$ applied to the instantaneous channel-transition rate $\Gamma_S = |dw_{\text{vis}}/dz|$ supplied by the host through `EmergenceParams.gamma_geom`. Total deposited mass is untouched by the transport step – the Laplacian is a redistribution operator ($\sum \nabla^2 n = 0$ on a periodic grid) and the sink is recorded, not deleted – so bit conservation holds to the exactness of the integer deposit, verified below the 10^{-30} bound by `test_continuum_transport.rs`. □

Theorem 24 (C^2-Regularized Instanton Condensation via Quintic Partition of Unity)**.** *The primordial seed incubation epoch is centered at* $z_{\text{peak}} = K/I_q^\star = 312/13 = 24.0$ *with half-width* $\Delta z = I_\ell^\star = 6.0$ *(*$\sigma_z = \Delta z/2 = 3.0$*), spanning the canonical window* $[z_{\min}, z_{\max}] = [z_{\text{peak}} - \Delta z, z_{\text{peak}} + \Delta z] = [18.0, 30.0]$*. Over this incubation window, define the* C^2 *partition of unity*

$$\Psi_{\text{nuc}}(z) = u^3(10 - 15u + 6u^2), \qquad u = \frac{z_{\max} - z}{z_{\max} - z_{\min}} = \frac{30.0 - z}{12.0}, \quad z \in [18.0, 30.0], \tag{477}$$

extended by 0 *for* $z > 30.0$ *and* 1 *for* $z < 18.0$*. Then* $\Psi_{\text{nuc}} \in C^2(\mathbb{R})$*, and gating the condensation weight with non-maximum suppression (NMS) collapse prefactor* $\eta_V \delta_{c0} = \frac{10}{33} \times 0.065610 = 0.019881818$ *as* $w_{\text{nuc}} = \Psi_{\text{nuc}}(z)\,\Theta(n_{\text{local}} - N_{\text{limit}})$ *renders seed ignition a barrierless continuous deformation of pre-existing filaments: at the window edges* Ψ_{nuc} *and its first two derivatives vanish simultaneously, eliminating discontinuous mass injection (pop-in) and the associated impulsive accelerations. Agglomerative seed merging on the* D_5 *checkerboard root lattice occurs at radius* $r^\star_{\text{merge}} = \sqrt{2}\kappa_*^{D_5}\Delta x_{\text{cell}} \approx 1.398249\,\Delta x_{\text{cell}}$*.*

Proof. The quintic $u^3(10 - 15u + 6u^2)$ is the classical smoothstep of order two: $p(0) = p'(0) = p''(0) = 0$ and $p(1) = 1$, $p'(1) = p''(1) = 0$, so the extension is C^2 at both gluing points and the composition with the linear map $u(z)$ remains C^2 on $\mathbb{R}$. Consequently the injected seed mass $M_{\text{seed}}(z)$ and the Poisson potential $\Phi_{\text{seed}}(z)$ inherit C^2 regularity in redshift, and the KDK integrator sees an at-most-quadratic impulse bounded by the step size. `compute_nucleation_weight` in `seed_emergence.wgsl` implements Eq. (477) with clamped u, and the gate multiplies the fixed-point overflow mass so the absolute condensation limit of Theorem 22 is preserved as $\Psi_{\text{nuc}} \to 1$. □

Theorem 25 (Thermodynamically Consistent Baryonic Hydrodynamics and Telescopic CIC Self–Force Cancellation)**.** *Let the baryon temperature track the decoupled radiation bath*

$$T_b(z) = T_{\text{CMB},0}\,(1 + z)\,\frac{z}{z + z_{\text{dec}}}, \qquad T_{\text{CMB},0} = 2.7255\ \text{K}, \quad z_{\text{dec}} = 137, \tag{478}$$

and set the visible-sector pressure $P_b = c^2_{s,\text{norm}}\,T_b\,\rho_b$ *on the particles whose Stinespring weight satisfies* $h_i \geq \eta_D$*. Then (i)* $T_b(z)$ *interpolates the exact kinetic decoupling profile, with* $T_b \to T_{\text{CMB}}$ *for* $z \gg z_{\text{dec}}$ *and* $T_b \to \frac{1}{2}T_{\text{CMB},0}(1 + z)$ *asymptotically; and (ii) when mass assignment and force interpolation use the same CIC kernel* W*, the hydrostatic self-force of a uniformly loaded cell on its own particle vanishes identically,* $\sum_i F_i^{(\text{self})} = 0$*, because the deposit and gather weights are adjoint and the central-difference gradient telescopes to zero on any translation-invariant field.*

Proof. (i) is the standard Compton-drag interpolation: the factor $z/(z + z_{\text{dec}})$ is unity when matter is thermally locked to photons ($z \gg z_{\text{dec}}$) and halves it after decoupling, so Eq. (478) is exact

at both limits and monotone between. (ii) Fix a cell c; the self-force of a particle at $\mathbf{x}$ on itself reads, per axis, $\sum_{\mathbf{j}} W(\mathbf{x}-\mathbf{x}_{\mathbf{j}})\,(P_{\mathbf{j}+\hat{e}}-P_{\mathbf{j}-\hat{e}})/(2\Delta x)$. For translation-invariant $P_{\mathbf{j}}=P_0$ every gradient sample is identically zero, so the residual is exactly 0 in exact arithmetic; for a general field the same adjointness cancels the diagonal contribution $W(\mathbf{x}-\mathbf{x}_{\mathbf{j}})^2$ under the telescoping sum, leaving only the physical neighbor coupling. `deposit_mass_cic` and `compute_thermodynamic_pressure` in `nbody_pm.wgsl` implement the pair at 2^{20} fixed-point scale, and `test_continuum_transport.rs` verifies the telescopic cancellation below 10^{-15} on both aligned and offset probe lattices in 256-bit arithmetic. □

Theorem 26 (Multi-tier thermodynamic and boundary-capacity shading)**.** *Let each bulk particle carry the normalized incubation density $\rho_b = n_{\text{local}}/\bar{n}$ and the boundary-capacity ratio $\sigma = \rho_b/\Gamma_{\text{CFT}}(z)$, where $\Gamma_{\text{CFT}}(z) = \gamma_{\text{CFT}}\big((1+z)/18\big)^{7.5}$ is the unscaled Cardy ceiling. Define the virial floor, debt activation, and effective temperature*

$$T_{\text{vir}} = 2500 + 6500\,\tanh\tfrac{\max(0,\rho_b-1)}{4}, \quad d_{\text{act}} = \frac{\sigma^3}{1+e^{-12(\sigma-1)}}, \quad T_{\text{eff}} = T_{\text{vir}} + 16000\,d_{\text{act}}, \tag{479}$$

and the visible spectral radiance

$$I_{\text{vis}} = \rho_b(1-e^{-0.45\rho_b})\Big(1 + 0.08\,\rho_b^2 + 4.5\,\sigma^2\,\tfrac{\log(1+e^{12(\sigma-1)})}{12}\,(1+0.1\,m_{\log})\Big), \tag{480}$$

with $m_{\log} = \log_{10}(1+P_{\text{debt}}/906\,\text{GW})$ the log-normalized seed-mass corona term. Then (i) $(T_{\text{eff}}, I_{\text{vis}})$ is C^∞ in (ρ_b,σ) on $[0,\infty)^2$; (ii) $2500 \le T_{\text{eff}} \le 25000\,\text{K}$ for all populated states, and the input clamp in `blackbody_to_linear_rgb` confines the sampled temperature to $[2500, 25000]\,\text{K}$, so the Planckian palette stays on the observationally meaningful segment of the blackbody locus; and (iii) the splat smoothing length $R = (R_{\min} + (R_{\max}-R_{\min})(1+0.45\max(0,\rho_b-1))^{-1/3})\Phi_{\text{topo}}(\sigma)\,(d_{\text{ref}}/d_{\text{cam}})$, with the topological SPH contraction $\Phi_{\text{topo}}(\sigma) = (1+7.5\,\sigma^2)^{-1}$, is bounded below by $R_{\min}\cdot\Phi_{\text{topo}}\cdot(d_{\text{ref}}/d_{\text{cam}})$ and reproduces the SPH spacing hierarchy: over-capacity (instanton-saturated) cores collapse to point-like kernels while underdense filaments diffuse. `compute_thermodynamic_state` and the dynamic splat radius in `dual_channel_render.wgsl` implement the pair.

Proof. (i) tanh, the logistic, and the softplus $\text{softplus}(x) = \log(1+e^{12x})/12$ are each C^∞, and $\sigma^2\,\text{softplus}(\sigma-1)$ extends smoothly through $\sigma=0$ since the exponential term decays double-exponentially. (ii) $T_{\text{vir}} \in [2500, 9000]$; over the populated regime $\sigma \lesssim 1$ the debt term contributes at most 16000 K, and for $\sigma \gg 1$ (deep capacity saturation) the palette input clamp pins the sampled value at 25000 K, so the displayed temperature never leaves the stated band. (iii) The cubic-root law is the SPH mean-spacing estimate $h \propto \rho^{-1/3}$ reparametrized onto pixel units; boundedness follows from $0 \le (1+0.45\max(0,\rho_b-1))^{-1/3} \le 1$ and $0 < \Phi_{\text{topo}} = (1+7.5\sigma^2)^{-1} \le 1$ for the clamped $\sigma \le 8$. □

Theorem 27 (Branchless single-cycle blackbody on $u \in [0.040, 0.400]$)**.** *The inverse-temperature Planckian palette $u = 1000/T \mapsto \text{RGB}(u)$, evaluated over $u \in [0.040, 0.400]$ (i.e. $T \in [2500, 25000]\,\text{K}$) by the per-channel degree-$(2,2)$ rational (Padé) fits*

$$R(u) = \frac{0.657842 - 15.067116\,u + 144.948029\,u^2}{1 - 19.160428\,u + 155.930970\,u^2}, \tag{481}$$

$$G(u) = \frac{0.540874 - 1.770797\,u + 12.417570\,u^2}{1 - 9.751228\,u + 48.259277\,u^2}, \tag{482}$$

$$B(u) = \frac{0.855655 - 3.605022\,u + 4.930827\,u^2}{1 - 8.226975\,u + 39.334862\,u^2}, \tag{483}$$

clamped below at zero, satisfies (i) C^∞ smoothness on the open domain and global C^0 continuity, since every denominator is strictly positive on $u \in [0.040, 0.400]$; (ii) channel values confined to $[0, 1.05]$ — the rational denominators track the Wien-tail curvature, so no piecewise hinge or clamp ladder is needed to keep the Planckian hue ordering; and (iii) evaluation with zero data-dependent control flow — three straight-line rational evaluations — so every SIMD lane retires in a single uniform cycle, eliminating the warp-divergence penalty of temperature-binned `if` *ladders.* `blackbody_to_linear_rgb` *in* `dual_channel_render.wgsl` *implements the map.*

Proof. Each channel is a ratio of two quadratics in u; the denominators $1-19.160428u+155.930970u^2$, $1-9.751228u+48.259277u^2$, and $1-8.226975u+39.334862u^2$ each have negative discriminant or roots outside $[0.040, 0.400]$, hence stay positive on the domain and the map is C^∞ there, giving (i). Direct evaluation of the three ratios on $u \in [0.040, 0.400]$ (enumerated by `test_branchless_blackbody_simd_rational_monotonicity` in `tests/first_principles_kernels.rs`) bounds each channel in $[0, 1.05]$, giving (ii). (iii) is immediate: the shader body contains no branch, loop, or lane-varying `select` on divergent conditions — the entire palette is a straight-line rational evaluation closed by a single $\max(\cdot, 0)$. □

Theorem 28 (Anti-blowout luminance-preserving ACES film transform)**.** *Let C be the HDR composite and $Y = \langle C, (0.2126, 0.7152, 0.0722)\rangle$ its Rec.709 luminance. The transform evaluates the Narkiewicz/Hill ACES fit on the scalar Y,*

$$Y_{\text{map}} = \frac{Y\,(aY+b)}{Y\,(cY+d)+e}, \qquad (a,b,c,d,e) = (2.51, 0.03, 2.43, 0.59, 0.14), \tag{484}$$

re-injects the un-clipped chromaticity as $C \cdot (Y_{\text{map}}/\max(Y, \varepsilon))$, and applies the Landauer desaturation knee at $Y_{\text{desat}} = 14$:

$$C_{\text{out}} = \text{mix}\Big(C\tfrac{Y_{\text{map}}}{\max(Y,\varepsilon)},\ (0.85, 0.95, 1.0)\,Y_{\text{map}},\ \text{clamp}\big(1 - e^{-(Y/Y_{\text{desat}})^{1.6}},\ 0,\ 0.85\big)\Big). \tag{485}$$

Then (i) chromaticity is preserved exactly in the mid-tones, since hue and saturation are carried by the ratio Y_{map}/Y alone; (ii) the output is contractive, $0 \le C_{\text{out}} \le 1$; and (iii) saturated cores asymptote to a controlled cyan-white corona rather than a flat clipping disc. The 9-tap cross-bilateral pre-filter (spatial $\sigma_s = 1.8$, photometric $\sigma_r = 0.10$) feeding the luminance Laplacian for the caustic accent is edge-preserving, so the high-frequency carrier survives denoising. `tone_map_aces_anti_blowout`, `calculate_luminance`, `sample_cross_bilateral`, *and* `linear_to_srgb` *in* `holographic_post.wgsl` *implement the transform.*

Proof. (i) Scaling a vector by a positive scalar preserves its direction in RGB space, hence its chromaticity; the scalar chosen is the ACES-mapped luminance, so perceived brightness follows the film curve exactly. (ii) $Y_{\text{map}} \in [0, 1]$ by the clamp, the chromatic component inherits the clamp, and the desaturation mix is a convex combination capped at 0.85 of two $[0, 1]$-valued operands. (iii) As $Y \to \infty$ the knee factor tends to 0.85, so the limit color is the Planckian corona tint $(0.85, 0.95, 1.0)\,Y_{\text{map}}$ — a hue-preserving highlight, not a flat RGB $= (1, 1, 1)$ clip. The bilateral range kernel $\exp(-(\Delta Y)^2/2\sigma_r^2)$ decouples the filter's smoothing from luminance edges, leaving the caustic Laplacian $\nabla^2 Y = Y_{\text{center}} - Y_{\text{bilateral}}$ intact. □

Theorem 29 (ε-softened thin-screen caustic magnification)**.** *Let the boundary screen carry convergence κ and shear $\gamma = (\gamma_1, \gamma_2)$ deposited by the Channel-B metric buffer, and let the macro-deflection*

field be the central difference of convergence,

$$\boldsymbol{\alpha}_{\text{macro}} = -\tfrac{1}{2}\big(\kappa_{+x} - \kappa_{-x},\, \kappa_{+y} - \kappa_{-y}\big), \tag{486}$$

augmented by the per-seed Einstein-ring deflection $\boldsymbol{\alpha}_{\text{seeds}} = \sum_k \theta_{E,k}^2\, \boldsymbol{\delta}_k / (|\boldsymbol{\delta}_k|^2 + r_{\text{soft}}^2)$ *with* $\theta_{E,k}$ *from Eq.* (467)*. Then the caustic magnification accent*

$$\mathcal{A}_{\text{caustic}} = \frac{1}{\sqrt{\det J^2 + \varepsilon^2}}, \qquad \det J = (1-\kappa)^2 - \gamma_1^2 - \gamma_2^2, \tag{487}$$

satisfies (i) uniform boundedness $\mathcal{A}_{\text{caustic}} \le 1/\varepsilon$ *(*$\varepsilon = 0.08$ *in the implementation), removing the critical-curve divergence without the sign-asymmetric bias of a linear* $1/(\det J + \varepsilon)$ *regularization; (ii) exact critical enhancement — on the fold set* $\det J = 0$ *the accent reaches its softened maximum* $1/\varepsilon$*, so Einstein rings and caustic fringes are amplified precisely where the Jacobian vanishes; and (iii) chromatic consistency, since the deflected samples* (R, G, B) *at* $(436, 546, 700)$ *nm share the same scalar* $\mathcal{A}$*, preserving the dispersion fringes of Eq.* (468) *across the critical curves.* `fs_post` *in* `holographic_post.wgsl` *implements the deflection, magnification, and accent.*

Proof. (i) $\det J^2 + \varepsilon^2 \ge \varepsilon^2$ gives $\mathcal{A} \le 1/\varepsilon$ for all (κ, γ). (ii) On $\det J = 0$ the expression reduces to $1/\sqrt{\varepsilon^2} = 1/\varepsilon$; moving off the critical curve decreases $\mathcal{A}$ monotonically in $|\det J|$, so the accent peaks exactly on the fold/cusp set rather than shifting away from it as a linear floor would. (iii) $\mathcal{A}$ is a scalar multiplier of the already-dispersed luminance and contains no wavelength dependence, so it rescales but never recolors the chromatic fringes. □

9.26 Two-Tier Interactive Visualization Architecture

The reference implementation separates the cosmological structure computation into a high-precision Tier-1 core and a real-time WebGPU Tier-2 rendering and simulation engine. Tier 1 (`shbt_simulator`, arbitrary-precision MPFR/BigUint modular ledger plus `precision_cosmology.py` and `boltzmann_shbt.py`) evaluates the exact analytical observables and serializes them into the SHBT-MMIO v2 telemetry stream (`src/shbt/export.rs`): a 128-byte binary header carrying `magic=0x54424853` and `schema=0x00020000`, z, a, $H(z)$, decoupled bit-loading fractions ($f_{\text{load}}^{\text{debt}}$ and $f_{\text{load}}^{\text{cosmo}}$), active and dark bit counts, Landauer entropy debt, cumulative seed mass, $f\sigma_8$, Δ_{ISW}, and particle metrology. This header is followed by a 256-point transfer function grid $T(k, z)$, a 256-point matter power spectrum grid $P_m(k, z)$, and 128-byte `SeedDefectRecord` entries. The dual loading fraction formulation decouples the cumulative thermodynamic Landauer debt $f_{\text{load}}^{\text{debt}} = S_{\text{debt}}/S_{\text{limit}}$ from the metric expansion friction parameter $f_{\text{load}}^{\text{cosmo}}$, preserving exact thermodynamic arrow tracking without perturbing cosmic conformal friction.

Tier 2 is the standalone `shbt-visualizer` engine (`src/shbt/visualizer`), compiled to `wasm32` and embedded in `visualizer/index.html`. The `ShbtWebGpuEngine` orchestrates an advanced WebGPU compute, reduction, and volumetric rendering pipeline operating strictly under a 256 MB WASM ceiling:

1. `cosmic_web_topology.wgsl`: In-situ cosmic web T-Web classification. Evaluates the gravitational tidal tensor $T_{ij} = \partial_i \partial_j \phi$ via fourth-order central differences on a 64^3 grid, solves the

local eigensystem in closed form using a Cardano–Vieta cubic solver to extract strictly ordered eigenvalues $\lambda_1 \geq \lambda_2 \geq \lambda_3$, executes 3D Non-Maximum Suppression (NMS) along principal eigenvectors, and partitions the cosmic volume into voids ($\lambda_1 < \lambda_{\rm th}$), sheets ($\lambda_1 \geq \lambda_{\rm th} > \lambda_2$), filaments ($\lambda_2 \geq \lambda_{\rm th} > \lambda_3$), and cluster nodes ($\lambda_3 \geq \lambda_{\rm th}$) with $\lambda_{\rm th} = 0.20$.

2. `dual_channel_render.wgsl`: 3D anisotropic ellipsoidal Gaussian splatting with D_5 lattice anchor $r_0 = \sqrt{10/33}\,\mathrm{Mpc} \approx 1.380121\,\mathrm{Mpc}$. Evaluates local velocity dispersion tensors V_{ij}, applies topological Lorentzian contraction $\Phi_{\rm topo}(\sigma) = (1 + \sigma^2/r_0^2)^{-1/2}$, projects covariance ellipsoids into screen space via camera Jacobian J, and applies anti-aliasing conic filtering ($\nu_{\rm aa} = 0.35$) with dual-channel color synthesis for visible baryons and dark matter halos.

3. `cosmic_web_volumetric.wgsl`: Volumetric raymarching of the warm-hot intergalactic medium (WHIM; $T \sim 10^5$–$10^7\,\mathrm{K}$) with Rankine–Hugoniot shock heating, Padé $(2,2)$ approximant free-free Bremsstrahlung emissivity $\epsilon_{\rm ff}$, resonant Lyman-α scattering, thermal Sunyaev–Zel'dovich (tSZ) Compton-y distortion, and late-time Integrated Sachs–Wolfe (ISW) metric decay.

4. `spectral_metrics.wgsl`: In-situ GPU spectral reductions. Performs Cloud-In-Cell (CIC) deconvolution, Poisson shot noise subtraction, 256-bin spherical shell power spectrum $P_m(k)$ estimation, 64×64 phase space (x, v_x) accumulation, and stacked underdensity void profiling.

5. `primordial_epochs.wgsl`: Multi-epoch primordial phase transitions, including electroweak symmetry breaking (EWSB) spinodal decomposition, QCD percolation fronts ($z \sim 10^{12}$), and recombination sound horizon shells ($r_s \approx 147.5\,\mathrm{Mpc}$, comoving wavenumber $M = 2,901,360$).

6. `causal_lifecycle.wgsl`: Five-phase CausalPoint observer lifecycle state machine (Nucleating, Active, Quenched, DefectShedding, Evaporated) and billboard particle instancing.

7. `seed_emergence.wgsl`: Instanton nucleation and topological defect condensation with 3D 26-neighbor exclusion ($r_{\rm excl} = 4$) and Landauer debt accretion $P_{\rm debt} = M_{\rm seed} \cdot 906\,\mathrm{GW}/M_\odot$.

8. `causal_point_get.wgsl`: Active Causal-Point Generalized Entropic Transport (GET) with Cardy vacuum capacity floor $\rho_{\rm floor}(a) = \max(0.00696738, 1/(39a))$ and signed periodic wrapping.

9. `nbody_pm.wgsl`: Symplectic Martel–Shapiro supercomoving Kick-Drift-Kick (KDK) leapfrog integrator for 2^{18} particles with conformal coupling $A(a) = \frac{3}{2}\Omega_{m,0} a (1 - \frac{10}{33} f_{\rm load}^{\rm cosmo})$.

10. `holographic_post.wgsl`: Multi-plane gravitational lensing ($z_m \in \{0.5, 1.2, 2.2, 3.5\}$), Shapiro delay, three-band boundary dispersion, and luminance-preserving ACES tonemapping.

Overlaid on the WebGPU canvas is the Canvas2D Scientific Telemetry Dock (`visualizer/telemetry_dock.js`), providing real-time multi-tab quantitative telemetry ($P_m(k)$ vs MPFR reference, halo mass function $dn/d\ln M$, growth rate $f\sigma_8(z)$, WHIM phase space, T-Web volume fractions, and Landauer debt ledger) alongside interactive research tools: 3D Raycast Object Inspector (Click-to-Inspect), 2D Slab Tomography Slicer (`S`), Real-Space vs Redshift-Space Distortions vs 2LPT (`R`), BAO Sound Horizon Caliper (`M`), Split-screen SHBT vs ΛCDM comparison (`V`), Telemetry Dock toggle (`P`), Render Mode cycling (`W`), Causal Lifecycle toggle (`C`), and quantitative CSV/JSON data export (`E`). Double-buffered GPU storage binds 2^{18} particles in 32-byte records (8 MB particle memory, total WASM footprint strictly under 256 MB), gated at 60 FPS by the browser presentation clock, while the native test harness executes identical shader kernels for automated CI verification.

9.27 Cosmological Trajectory Flight Controller and 10-Milestone Verification

To record the 90.0-second canonical cosmological simulation flythrough (`full_cosmic_evolution.webm`) without late-time exponential acceleration artifacts, the camera flight controller (`visualizer/record_simulation_events.py`) employs piecewise smooth cosmic-time parameterization and quintic minimum-jerk boundary easing:

1. **Stage 1 ($z = 10^{14} \to 30$, 12 s / 720 frames):** High-redshift boundary relaxation with 45-frame physical dwells at EWSB ($z \approx 10^{14}$), QCD percolation ($z \approx 10^{12}$), Big Bang Nucleosynthesis ($z \in [10^9, 10^8]$), and Recombination ($z \approx 1090$).
2. **Stage 2 ($z = 30 \to 1.0$, 35 s / 2100 frames):** Paced strictly linearly in monotonic cosmic age $t(z) \in [0.0987, 5.8410]$ Gyr via numerical inversion of the SHBT Friedmann integral $t(z) = \frac{1}{H_0} \int_{\ln(1+z)}^{40} \frac{du}{E(u)}$.
3. **Stage 3 ($z = 1.0 \to 0.0$, 25 s / 1500 frames):** Paced linearly in cosmic age $t(z) \in [5.8410, 13.7909]$ Gyr with a dedicated 60-frame physical dwell capturing the mature cosmic web after 7–8 Gyr of non-linear evolution at $z \approx 0.55$ ($t \approx 7.8$–8.2 Gyr).
4. **Stage 4 ($z = 0.0 \to -0.999$, 18 s / 1080 frames):** Asymptotic horizon freeze governed by a fifth-order minimum-jerk polynomial $s(\tau) = 10\tau^3 - 15\tau^4 + 6\tau^5$ ($\tau \in [0, 1]$), enforcing $s'(0) = s'(1) = 0$ and $s''(0) = s''(1) = 0$ to smoothly decelerate camera rotation and metric dilation to complete stasis.

The automated Playwright WebGPU harness audits the complete **10-Milestone Canonical Sequence**:

1. `01_primordial_bit_loading.png` ($z = 10^{14}$): Pristine bit loading ($f_{\text{load}} < 0.01$, $t < 10^{-3}$ Gyr, zero active seeds).
2. `02_baryogenesis_derendering.png` ($z = 10^{10}$): Stinespring topological quench with $\eta_D \in [68\%, 71\%]$ (analytical $23/33 \approx 69.7\%$).
3. `03_ghost_seed_genesis.png` ($z = 16.0$): Emergent instanton defect pool ($N_{\text{seeds}} > 0$, $M_{\text{seed}} > 0$, $P_{\text{debt}} > 0$, central void contrast $C_{\text{void}} \geq 0.92$).
4. `04_early_filament_network.png` ($z = 3.86$): Early filament nucleation and spoke anisotropy $R_{\text{axis}} \leq 1.15$.
5. `05_causal_point_proto_galaxies.png` ($z = 3.00$): Proto-cluster assembly, $P_{\text{debt}} > 5 \times 10^{11}$ GW, principal axis ratio $R_{\text{axis}} \leq 1.30$, and 1:1 caustic-to-seed parity.
6. `06_cosmic_noon_peak_clustering.png` ($z = 1.00$): Peak cosmic star-formation clustering, virialized core ellipticity $e \leq 0.15$, and caustic ratio ≤ 1.05.
7. `07_mature_cosmic_web_7gyr.png` ($z \approx 0.55$, $t \approx 7.8$–8.2 Gyr): Virialized cosmic web, shock-heated WHIM gas, and stable anisotropic splat caustics (≤ 1.50).

8. `08_present_epoch_universe.png` ($z = 0.00$, $t = 13.79\,\mathrm{Gyr}$): Present-epoch universe, 4-slice multi-plane lensing, and void evacuation.

9. `09_desitter_acceleration_voids.png` ($z = -0.70$, $t \approx 33.8\,\mathrm{Gyr}$): Cosmic acceleration, dark-energy horizon expansion, and void clearing ($f_{\mathrm{load}} > 0.40$).

10. `10_desitter_horizon_freeze.png` ($z = -0.999$, $t \to \infty$): Asymptotic holographic horizon freeze ($f_{\mathrm{load}} > 0.99$, $R_{\mathrm{entropy}} < 0$, $\mathcal{R}_{\mathrm{adm}} = 0$, multipole asymmetry $|a_{40}/a_{00}| \leq 5 \times 10^{-5}$).

9.28 Screen-Space Tensor Deflection Formalism

In our decoupled visualizer architecture, we project the cosmological Weyl curvature onto the observer's backward light-cone without integrating full 3+1 ADM geodesics. Channel B encodes the projected shear tensor $\boldsymbol{\gamma} = (\gamma_1, \gamma_2)$ alongside the dimensionless convergence field $\kappa = \frac{1}{2}\nabla^2\psi$. The screen-space mapping from source angular coordinate $\boldsymbol{\beta}$ to image coordinate $\boldsymbol{\theta}$ is governed by:

$$\boldsymbol{\beta} = \boldsymbol{\theta} - \lambda_{\mathrm{lens}} \left[\nabla\kappa(\boldsymbol{\theta}) + \boldsymbol{\Gamma}(\boldsymbol{\theta})\nabla\kappa(\boldsymbol{\theta})\right] - \sum_{s=1}^{S} \theta_{E,s}^2 \frac{\boldsymbol{\theta} - \boldsymbol{\theta}_s}{|\boldsymbol{\theta} - \boldsymbol{\theta}_s|^2 + \varepsilon_{\mathrm{core}}^2} \tag{488}$$

where $\boldsymbol{\Gamma}(\boldsymbol{\theta}) = \begin{pmatrix} \gamma_1 & \gamma_2 \\ \gamma_2 & -\gamma_1 \end{pmatrix}$, and $\theta_{E,s} = \sqrt{\frac{4GM_{\mathrm{seed},s}}{c^2}\frac{D_{ds}}{D_d D_s}}$ represents the Einstein radius of supermassive topological defects.

9.29 Wave-Optics Caustic Fringing and Relativistic Beaming

Near critical curves ($\det \mathcal{A} \to 0$), the geometric singularity is regularized through chromatic dispersion:

$$\boldsymbol{\theta}_{\mathrm{sample}}(\lambda) = \boldsymbol{\theta} + \boldsymbol{\alpha}_{\mathrm{total}}(\boldsymbol{\theta}) \left[1.0 + \delta_{\mathrm{disp}} \left(\frac{\lambda - \lambda_0}{\lambda_0}\right)\right] \tag{489}$$

Simultaneously, peculiar velocities modulate the emitted baryonic continuum via the Doppler boosting factor $\mathcal{D} = \sqrt{1-\beta^2}/(1-\beta_{\mathrm{los}})$, transforming observed surface brightness according to $I_{\mathrm{obs}} = I_0 \mathcal{D}^3$.

10 Sub-Kelvin Address-Scaled Landauer Calorimetry Architecture and Falsification Protocols

The Causal Point construction yields a laboratory-scale discriminator that is logically downstream of the boundary theory rather than an independent phenomenological add-on. The experiment tests whether resolving a fixed target through a larger address space carries a thermodynamic cost. A complete readout cycle includes acquisition and reset of its classical record, so the standard lower bound for one logically irreversible bit is

$$\Delta Q_{\mathrm{Landauer}} \geq k_B T \ln 2. \tag{490}$$

Standard quantum mechanics and thermodynamics assign this bound to the payload and record reset, independently of the register topology. SHBT instead treats the address locator as part of the finite entropy payment made by the observer.

10.1 Operational Capacity and Competing Hypotheses

The general Causal Point cost in Eq. (403) remains

$$C_{\text{get}} = \max(1, \log_2 |\mathcal{R}| + \log_2 |\Omega_A|, C_{\text{req}}) \,. \tag{491}$$

For calorimetry it is useful to expose the additive sources of the operational capacity explicitly. Measuring entropy in bits, define

$$\boxed{C_{\text{op}} = \Delta S_{\text{payload}} + \log_2 |\mathcal{R}| + C_0,} \tag{492}$$

where $\Delta S_{\text{payload}}$ is the resolved payload entropy in bits and C_0 is the address-independent preparation, acquisition, and reset overhead. The SHBT dissipation inequality is

$$\boxed{Q_H \geq k_B T \ln 2 \; C_{\text{op}}.} \tag{493}$$

The fixed contributions $\Delta S_{\text{payload}} + C_0$ are measured in control runs and absorbed into the fitted intercept. Consequently, the slope test is independent of their calibrated value. For the proposed singleton outcome ensemble $|\Omega_A| = 1$, baseline request $C_{\text{req}} = 1$, and $|\mathcal{R}| \geq 2$, Eq. (491) gives

$$C_{\text{get}} = \max(1, \log_2 |\mathcal{R}|) = \log_2 |\mathcal{R}|. \tag{494}$$

Thus the background-subtracted experimental capacity C_{op} and the formal GET cost C_{get} have the same address-dependent part.

Let $k = \log_2 |\mathcal{R}|$. The hypotheses are deliberately posed as a derivative so that unknown constant heat offsets cannot mimic the SHBT signal:

$$H_0 : \quad \begin{cases} Q_{H_0} = k_B T \ln 2, \quad |\mathcal{R}| \geq 2, \\ \dfrac{\partial Q_H}{\partial(\log_2 |\mathcal{R}|)} = 0, \end{cases} \tag{495}$$

$$H_1 : \quad \begin{cases} Q_{H_1} = (\log_2 |\mathcal{R}|) k_B T \ln 2, \\ \dfrac{\partial Q_H}{\partial(\log_2 |\mathcal{R}|)} = k_B T \ln 2. \end{cases} \tag{496}$$

Here H_0 is the standard address-independent thermodynamic model and H_1 is the SHBT boundary-mechanics model. For a burst of N_{pulse} identical operations, both predicted heats and slopes are multiplied by N_{pulse}.

10.2 Predicted Heat at 7 mK

At the proposed base temperature,

$$\begin{aligned} E_L &= k_B T_{\mathrm{base}} \ln 2 \\ &= \left(1.380649 \times 10^{-23}\,\mathrm{J\,K^{-1}}\right)(0.0070\,\mathrm{K}) \ln 2 \\ &= 6.698951 \times 10^{-26}\,\mathrm{J} \simeq 0.0670\,\mathrm{yJ}. \end{aligned} \tag{497}$$

The address space is swept through

$$|\mathcal{R}| \in \{2^1, 2^2, \ldots, 2^{10}\}. \tag{498}$$

Table 35 retains both the single-event predictions and the synchronized 100,000-pulse burst predictions. The last entry follows directly from $10 N_{\mathrm{pulse}} E_L = 66.9895\,\mathrm{zJ}$.

Address space $\lvert\mathcal{R}\rvert$	Address bits k	Single event Q_{H_0} (yJ)	Single event Q_{H_1} (yJ)	100k burst Q_{H_0} (zJ)	100k burst Q_{H_1} (zJ)
$2^1 = 2$	1	0.0670	0.0670	6.6990	6.6990
$2^2 = 4$	2	0.0670	0.1340	6.6990	13.3979
$2^3 = 8$	3	0.0670	0.2010	6.6990	20.0969
$2^4 = 16$	4	0.0670	0.2680	6.6990	26.7958
$2^5 = 32$	5	0.0670	0.3349	6.6990	33.4948
$2^6 = 64$	6	0.0670	0.4019	6.6990	40.1937
$2^7 = 128$	7	0.0670	0.4689	6.6990	46.8927
$2^8 = 256$	8	0.0670	0.5359	6.6990	53.5916
$2^9 = 512$	9	0.0670	0.6029	6.6990	60.2906
$2^{10} = 1024$	10	0.0670	0.6699	6.6990	66.9895

Table 35: Address-independent H_0 and address-scaled H_1 heat predictions at $T_{\mathrm{base}} = 7.0\,\mathrm{mK}$.

10.3 Superconducting Register and AuPd Nanocalorimeter

The register is a ten-qubit superconducting circuit on ultra-high-resistivity float-zone silicon. Enabling $k \in \{1, \ldots, 10\}$ address-control lines with superconducting single-flux-quantum (SFQ) selectors realizes $|\mathcal{R}| = 2^k$ while routing each readout pulse to the same designated target. Table 36 gives the complete register specification.

The readout line terminates in a metallic nanocalorimeter whose electronic temperature is sensed by symmetric NIS tunnel junctions and low-noise SQUID preamplifiers. The absorber is electron-beam-evaporated $\mathrm{Au_{60}Pd_{40}}$ at 60/40 wt% with $L = 1.0\,\mu\mathrm{m}$, $W = 0.2\,\mu\mathrm{m}$, and $t = 0.05\,\mu\mathrm{m} = 50\,\mathrm{nm}$. Hence

$$V_{\mathrm{abs}} = LWt = 1.0 \times 10^{-20}\,\mathrm{m^3}. \tag{499}$$

With Sommerfeld coefficient $\gamma_e = 200.0\,\mathrm{J\,m^{-3}\,K^{-2}}$, the electron heat capacity at base temperature is

$$\begin{aligned} C_e(T) &= \gamma_e V_{\mathrm{abs}} T \\ &= \left(200.0\,\mathrm{J\,m^{-3}\,K^{-2}}\right)\left(1.0 \times 10^{-20}\,\mathrm{m^3}\right)(0.0070\,\mathrm{K}) \\ &= 1.40 \times 10^{-20}\,\mathrm{J\,K^{-1}}. \end{aligned} \tag{500}$$

Parameter	Value or specification	Physical function or constraint
Substrate	High-resistivity silicon, $\rho > 10\,\mathrm{k\Omega\,cm}$	Low dielectric loss, $\tan\delta < 10^{-6}$
Transmon Josephson energy E_J/h	15.0 GHz	Sets the transmon operating scale
Charging energy E_C/h	300 MHz	Maintains $E_J/E_C \simeq 50$
Transmon layout	Ten fixed-frequency transmons	Eliminates flux-bias heating during the address sweep
Transition frequency f_{01}	4.50–5.50 GHz	Frequency-multiplexed fixed-frequency state drive
Readout resonator frequency f_r	6.50–7.50 GHz	Dispersive Jaynes–Cummings readout
Active addressing switches	Superconducting SFQ logic gates	Address routing without normal-state dissipation
Register span	Ten address-control bits, $2 \le \lvert\mathcal{R}\rvert \le 1024$	Discrete sweep of the active decoder network
AuPd absorber	$1.0 \times 0.2 \times 0.05\,\mu\mathrm{m}^3$	$V_{\mathrm{abs}} = 1.0 \times 10^{-20}\,\mathrm{m}^3$
Base-temperature heat capacity	$C_e = 1.40 \times 10^{-20}\,\mathrm{J\,K^{-1}}$	Calibrated at $T_{\mathrm{base}} = 7.0\,\mathrm{mK}$
Thermal decay constant	$\tau_{\mathrm{th}} = 20.41\,\mathrm{ms}$	Resolves the integrated burst transient

Table 36: Experimental circuit specifications for the superconducting register, SFQ address network, and AuPd calorimetric termination.

The thermal link is selected so that the measured differential conductance at the operating point is

$$G_{\mathrm{th}} = \frac{C_e}{\tau_{\mathrm{th}}} = \frac{1.40 \times 10^{-20}\,\mathrm{J\,K^{-1}}}{20.41 \times 10^{-3}\,\mathrm{s}} = 6.85938 \times 10^{-19}\,\mathrm{W\,K^{-1}}. \tag{501}$$

The intrinsic thermal relaxation time is therefore

$$\tau_{\mathrm{th}} = \frac{C_e}{G_{\mathrm{th}}} = \frac{1.40 \times 10^{-20}\,\mathrm{J\,K^{-1}}}{6.85938 \times 10^{-19}\,\mathrm{W\,K^{-1}}} = 2.0408 \times 10^{-2}\,\mathrm{s} \simeq 20.41\,\mathrm{ms}. \tag{502}$$

The electron and bath temperatures obey the linearized two-temperature equations

$$C_e \frac{d\Delta T_e}{dt} = P_{\mathrm{in}}(t) - G_{\mathrm{th}}\Delta T_e, \qquad \tau_{\mathrm{th}} = \frac{C_e}{G_{\mathrm{th}}}, \tag{503}$$

$$P_{\mathrm{in}}(t) = \frac{Q_{\mathrm{exp}}}{\tau_{\mathrm{in}}} e^{-t/\tau_{\mathrm{in}}}, \qquad \tau_{\mathrm{in}} = 1.00\,\mathrm{ms}. \tag{504}$$

Using the pulse-amplitude convention fixed by the calorimetric calibration, the fitted thermal transient is

$$\boxed{\Delta T_e(t) = \frac{Q_{\mathrm{exp}}}{C_e}\left(e^{-t/\tau_{\mathrm{th}}} - e^{-t/\tau_{\mathrm{in}}}\right), \qquad \tau_{\mathrm{th}} = 20.41\,\mathrm{ms}.} \tag{505}$$

Nanocalorimeter parameter	Value or specification	Measurement role
Absorber composition	$Au_{60}Pd_{40}$, 60/40 wt%	Normal-metal electron absorber
Absorber geometry	$1.0 \times 0.2 \times 0.05\,\mu m^3$	$V_{abs} = 1.0 \times 10^{-20}\,m^3$
Sommerfeld coefficient	$\gamma_e = 200.0\,J\,m^{-3}\,K^{-2}$	Determines $C_e(T)$
Electron heat capacity	$C_e = 1.40 \times 10^{-20}\,J\,K^{-1}$ at 7 mK	Converts deposited heat to electron-temperature rise
Thermal conductance	$G_{th} = 6.85938 \times 10^{-19}\,W\,K^{-1}$	Calibrated bath-coupling scale
Thermal time constant	$\tau_{th} = 20.41\,ms$	Enables quasi-adiabatic integration of a 1 ms burst
Injection time constant	$\tau_{in} = 1.00\,ms$	Fixes the rising edge of the two-temperature pulse
NIS junction stack	Symmetric $Al/AlO_x/AuPd$	Electron thermometry
NIS junction area	$100 \times 100\,nm^2$, high R_N	Suppresses lead-mediated heat leakage
NIS bias	$V_{bias} \simeq \Delta_{Al}/e \simeq 180\,\mu V$	High-responsivity operating point
Voltage responsivity	$dV/dT \simeq 1.50\,mV\,K^{-1}$ below 10 mK	Converts temperature excursions to voltage
Preamplification	Ultra-low-noise SQUID at the base stage	Resolves the NIS transient

Table 37: AuPd absorber, thermal link, and NIS nanocalorimeter specifications.




Figure 2: Inline apparatus schematic for the address-scaling test. The SFQ selector varies the active address depth while the same target and readout path deposit energy in the AuPd absorber; symmetric NIS junctions and a SQUID read the electron-temperature transient at the 7 mK stage.

10.4 Cryogenic Filtering and Environmental Control

The assembly is mounted in a dry dilution refrigerator at $T_{base} = 7.0 \pm 0.1$ mK. The complete staged attenuation, filtering, and magnetic-shielding budget is listed in Table 38.

10.5 Address-Scaling Measurement Protocol

The synchronized sequence holds the payload and physical pulse envelope fixed while changing only the active address decoder:

Stage or component	Attenuation or filtering	Physical specification
Room temperature (300 K) to 4 K	20 dB coaxial attenuators	Stainless-steel flexible coaxial lines
Still stage (0.8 K)	10 dB attenuators	NbTi superconducting coaxial lines
Cold plate (100 mK)	10 dB attenuators	CuNi coaxial lines
Mixing chamber (7 mK)	20 dB thermalized attenuators	Eccosorb infrared filters and Cu-powder filters over 0.1–10 GHz
Outer magnetic shield	μ-metal cylinder at 300 K	Suppresses the ambient geomagnetic field
Cryogenic magnetic shield	Dual-layer Cryoperm-10 plus superconducting Al shield at 7 mK	Residual ambient flux $B < 0.5\,\mathrm{nT}$

Table 38: Cryogenic attenuation, filtering, and magnetic-shielding budget.

1. *State initialization.* Initialize the target transmon in $|0\rangle$ by active feedback and microwave sideband cooling, require $P_{|1\rangle} < 0.001$, and wait more than 100 ms for baseline thermalization.

2. *Address configuration.* Assert $k = \log_2 |\mathcal{R}|$ SFQ selector channels, with $k \in \{1, \ldots, 10\}$, so the enabled decoder spans $|\mathcal{R}| = 2^k$ addresses.

3. *GET readout.* Apply a 6.50 GHz, 10 ns readout pulse to the same address key $A \in \mathcal{R}$. The target remains logical zero in every trial, hence $\Delta S_{\mathrm{payload}} = 0$ for the prepared target; the constant acquisition and record-reset heat is assigned to the intercept.

4. *Pulse-train integration.* Repeat the GET operation $N_{\mathrm{pulse}} = 100{,}000$ times at $f_{\mathrm{rep}} = 100\,\mathrm{MHz}$. The resulting $t_{\mathrm{burst}} = 1.0\,\mathrm{ms}$ obeys $t_{\mathrm{burst}} \ll \tau_{\mathrm{th}} = 20.41\,\mathrm{ms}$, so the absorber integrates the burst quasi-adiabatically.

5. *Calorimetric acquisition.* Record $V_{\mathrm{NIS}}(t)$ for $0 \le t \le 100\,\mathrm{ms}$, extract the peak shift ΔV_{peak}, and integrate the thermal decay to obtain Q_{exp}.

6. *Thermal reset.* Wait $t_{\mathrm{cool}} = 150\,\mathrm{ms} > 7\tau_{\mathrm{th}}$ before the next burst so the absorber returns to T_{base}.

Each of the ten address configurations receives $M = 1{,}000$ independent burst repetitions. The complete sweep is therefore

$$k \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} \quad \Longleftrightarrow \quad |\mathcal{R}| \in \{2, 4, 8, 16, 32, 64, 128, 256, 512, 1024\}. \tag{506}$$

The acquisition and inference chain is fixed before unblinding and is listed in Table 39. The same operations are applied to signal, null, and capacity-ceiling runs.

The systematic-error targets are small compared with the predicted SHBT step $N_{\mathrm{pulse}} E_L = 6.6990\,\mathrm{zJ/bit}$.

Stage	Analysis operation	Mathematical or instrumental action	Stored output
1	Baseline calibration	Measure T_{base}, C_e, G_{th}, NIS responsivity, and the zero-pulse trace	Calibration covariance matrix
2	Address randomization	Randomly permute $k \in \{1, \dots, 10\}$ while fixing the target, payload, pulse envelope, and repetition count	Blinded address key and run identifier
3	Pulse delivery	Execute $N_{\mathrm{pulse}} = 100{,}000$ SFQ-selected GET operations during $t_{\mathrm{burst}} = 1.0\,\mathrm{ms}$	Pulse timestamps and decoder-state audit
4	Transient acquisition	Sample $V_{\mathrm{NIS}}(t)$ over $0 \le t \le 100\,\mathrm{ms}$ and convert voltage to $\Delta T_e(t)$	Calibrated temperature time series
5	Thermal fit	Fit Eq. (505) jointly for Q_{exp}, baseline offset, and nuisance drift	Heat estimate and fit covariance
6	Quality control	Reject preregistered saturation, missing-trigger, thermal-reset, and residual-autocorrelation failures	Binary quality mask and diagnostic residuals
7	OLS/MLE inference	Fit Eq. (507), evaluate Eqs. (511)–(513), and compute the slope p-value	$\widehat{\alpha}$, standard error, p, χ^2_ν, and ΔAIC
8	Capacity-ceiling audit	Impose $C_{\mathrm{local}} = 6$ bits and test the transition between $k = 6$ and $k = 7$	Fidelity, thermal variance, and `AnomalyClosureError` flag

Table 39: Stage-by-stage calorimetry acquisition, reduction, inference, and finite-capacity falsification operations.

Systematic leakage channel	Estimated contribution	Physical mechanism	Mitigation
Parasitic capacitive cross-talk	$< 0.05\,\mathrm{zJ}$ per burst	Coupling between inactive address lines and calorimeter	Differential ground-shielded coplanar waveguides
Microwave drive heating	$< 0.10\,\mathrm{zJ}$ per burst	Direct absorption from 6.5 GHz readout photons	Spatial isolation $> 500\,\mu$m plus bandpass filters
Amplifier back-action	$< 0.02\,\mathrm{zJ}$ per burst	Josephson or HEMT noise returning along the readout line	Dual junction isolators with $> 45\,\mathrm{dB}$ isolation at 6.5 GHz
NIS-lead thermal conductance	$< 0.01\,\mathrm{zJ}$ per burst	Heat leakage through normal-metal thermometer leads	$100 \times 100\,\mathrm{nm}^2$ junctions with high R_N
Engineering noise allowance	$\simeq 0.18021\,\mathrm{zJ}$ per burst	Safety-scaled design budget	Conditional on the noise model in Section 10.7

Table 40: Systematic-error attenuation budget for address-scaled calorimetry.

10.6 Empirical Falsification Criteria and Observer-Capacity Cutoff

10.6.1 OLS/MLE Slope Test

For each address length k, let $Q_{\exp}(k)$ be the burst heat. The direct regression used for the primary decision is

$$Q_{\exp}(k) = \widehat{\beta}_0 + \widehat{\alpha}k + \epsilon, \qquad \epsilon \sim \mathcal{N}(0, \sigma^2), \tag{507}$$

fitted by ordinary least squares (OLS) and Gaussian maximum likelihood estimation (MLE). The competing slope values are

$$\begin{aligned} H_0: &\quad \alpha_0 = 0\,\mathrm{zJ/bit}, \\ H_1: &\quad \alpha_1 = N_{\mathrm{pulse}} k_B T \ln 2 \\ &\quad = 100{,}000\left(6.69895 \times 10^{-26}\,\mathrm{J/bit}\right) \\ &\quad = 6.69895 \times 10^{-21}\,\mathrm{J/bit} = 6.6990\,\mathrm{zJ/bit}. \end{aligned} \tag{508}$$

Equivalently, the source model templates are

$$Q_{H_0}(k) = \alpha_0 + \epsilon, \qquad \alpha_0 = N_{\mathrm{pulse}} k_B T \ln 2 = 6.6990\,\mathrm{zJ}, \tag{509}$$

$$Q_{H_1}(k) = \beta_0 + \beta_1 \left[k N_{\mathrm{pulse}} k_B T \ln 2\right] + \epsilon, \qquad \widehat{\alpha} = \widehat{\beta}_1 N_{\mathrm{pulse}} k_B T \ln 2. \tag{510}$$

For model H_j, $j \in \{0, 1\}$, the Gaussian maximum-likelihood ledger is

$$\ln \mathcal{L}_j = -\frac{n}{2}\ln(2\pi\widehat{\sigma}_j^2) - \frac{1}{2\widehat{\sigma}_j^2}\sum_{a=1}^{n}\left[Q_a - Q_{H_j}(k_a)\right]^2, \tag{511}$$

$$\widehat{\sigma}_j^2 = \frac{1}{n}\sum_{a=1}^{n}\left[Q_a - Q_{H_j}(k_a)\right]^2, \tag{512}$$

$$\mathrm{AIC}_{H_j} = 2p_j - 2\ln \mathcal{L}_j, \qquad \Delta\mathrm{AIC} = \mathrm{AIC}_{H_0} - \mathrm{AIC}_{H_1}. \tag{513}$$

The alternative is accepted only if the null slope is rejected at least 5σ with $p < 10^{-6}$, $\Delta\mathrm{AIC} > 10.0$, the fitted slope is statistically consistent with $6.6990\,\mathrm{zJ/bit}$, and the address sweep exhibits discrete stair-step increments at integer k. The address-scaling prediction is falsified, and H_0 is accepted within the stated experimental resolution, if

$$|\widehat{\alpha}| \le 0.05\,(N_{\mathrm{pulse}} k_B T \ln 2) = 0.33495\,\mathrm{zJ/bit}. \tag{514}$$

Table 41 records the full fit and model-selection criteria.

10.6.2 Local Capacity Ceiling and `AnomalyClosureError`

The second test probes the finite-observer cutoff independently of the linear heat slope. To make the notation explicit, define the Causal Point's local capacity ceiling by

$$\boxed{C_{\mathrm{local}} \equiv N_{\mathrm{limit}} = \min\left(N_{\mathrm{local}}, \frac{A_{\mathrm{local}}}{4L_P^2 \ln 2}\right).} \tag{515}$$

Metric or parameter	Null-model H_0 target	SHBT H_1 target	Selection threshold
Direct slope $\widehat{\alpha}$	$0\,\mathrm{zJ/bit}$	$6.6990\,\mathrm{zJ/bit}$	Reject H_0 at $\geq 5\sigma$, $p < 10^{-6}$
Normalized slope β_1	0.000 ± 0.010	1.000 ± 0.010	Requires consistency with the selected model
Intercept β_0	$6.699\,\mathrm{zJ}$	$0.000 \pm 0.18021\,\mathrm{zJ}$ design allowance after baseline subtraction	Diagnoses parasitic drive heating
Reduced chi-square χ^2_ν	$\chi^2_\nu \gg 1$ for SHBT-like data	1.00 ± 0.05	Noise-consistent goodness of fit
Akaike information criterion	AIC_{H_0} baseline	$\Delta\mathrm{AIC} = \mathrm{AIC}_{H_0} - \mathrm{AIC}_{H_1} > 10$	Decisive preference for H_1

Table 41: OLS/MLE parameters and binary model-selection thresholds.

The experimental residual entropy budget is

$$R^{\mathrm{exp}}_{\mathrm{entropy}} = C_{\mathrm{local}} - C_{\mathrm{op}}. \tag{516}$$

For the singleton, background-calibrated GET sequence, $C_{\mathrm{op}} = C_{\mathrm{get}} = k$. Normal state resolution requires $R^{\mathrm{exp}}_{\mathrm{entropy}} \geq 0$; if $C_{\mathrm{op}} > C_{\mathrm{local}}$, the negative residual triggers the SHBT failure condition denoted `AnomalyClosureError`.

The controlled cutoff run imposes an artificial local ceiling

$$C^{\mathrm{exp}}_{\mathrm{local}} = N^{\mathrm{exp}}_{\mathrm{limit}} = 6 \text{ bits}, \qquad |\mathcal{R}|_{\max} = 2^6 = 64, \tag{517}$$

and then continues the physical decoder sweep through $k = 10$. SHBT predicts

$$\begin{cases} k \leq 6: & \text{admissible readout,} \\ k > 6: & \text{loss of readout fidelity, nonlinear thermal-variance spikes,} \\ & \text{and state-resolution failure (}\texttt{AnomalyClosureError}\text{).} \end{cases} \tag{518}$$

Standard quantum mechanics predicts continuous address decoding across the same artificial ceiling provided the hardware remains functional. A sharp, reproducible breakdown at $C_{\mathrm{op}} > C^{\mathrm{exp}}_{\mathrm{local}}$ therefore tests the finite-register postulate, while its absence falsifies this cutoff prediction even if a nonzero calorimetric slope were observed.

10.7 Cryogenic Nanocalorimetry Noise Budget at 7 mK

The noise budget distinguishes a single Landauer operation from the $N_{\mathrm{pulse}} = 100{,}000$-operation burst used in the measurement protocol. At $T = 7.0\,\mathrm{mK}$,

$$E_L = k_B T \ln 2 = 6.6989507 \times 10^{-26}\,\mathrm{J}, \tag{519}$$

$$\Delta Q_{\mathrm{step}} = N_{\mathrm{pulse}} E_L \simeq 6.698951\,\mathrm{zJ}. \tag{520}$$

The burst heat slope is therefore $N_{\mathrm{pulse}} k_B T \ln 2 \simeq 6.6990\,\mathrm{zJ/bit}$; the single-operation slope is $k_B T \ln 2 \simeq 6.69895 \times 10^{-5}\,\mathrm{zJ/bit}$. These are different measurement normalizations of the same proposed address-dependent contribution.

For the equilibrium-fluctuation model and absorber heat capacity $C_e = 1.40 \times 10^{-20}\,\mathrm{J\,K^{-1}}$ used here, the thermal energy standard deviation is

$$\begin{aligned}\Delta E_{\mathrm{th}} &= \sqrt{k_B T^2 C_e} \\ &= \sqrt{(1.380649 \times 10^{-23})(0.0070)^2(1.40 \times 10^{-20})}\,\mathrm{J} \\ &\simeq 3.0775399 \times 10^{-24}\,\mathrm{J} \simeq 0.00307754\,\mathrm{zJ}. \end{aligned} \tag{521}$$

This equilibrium quantity is not, by itself, the uncertainty of the finite-duration heat estimator. Relating the two requires the thermal and readout transfer functions, integration window, and estimator calibration.

For a provisional independent-noise model, interpret the channel allocations in Table 42 as equivalent per-burst RMS contributions. Their quadrature sum is

$$\begin{aligned}\Delta Q_{\mathrm{total}} &= \left[(\Delta Q_{\mathrm{SFQ}})^2 + (\Delta Q_{\mathrm{micro}})^2 + (\Delta Q_{\mathrm{SQUID}})^2 + (\Delta Q_{\mathrm{lead}})^2 + (\Delta E_{\mathrm{th}})^2\right]^{1/2} \\ &\simeq \sqrt{0.050^2 + 0.100^2 + 0.020^2 + 0.010^2 + 0.00307754^2}\,\mathrm{zJ} \\ &\simeq 0.11405907\,\mathrm{zJ}. \end{aligned} \tag{522}$$

Applying the specified engineering safety multiplier to the unrounded result defines the design allowance

$$\Delta Q_{\mathrm{bounded}} \equiv 1.58\,\Delta Q_{\mathrm{total}} \simeq 0.18021333\,\mathrm{zJ} \simeq 0.18021\,\mathrm{zJ}. \tag{523}$$

The label "bounded" denotes this engineering allowance, not a proven upper bound or a specified statistical confidence interval. In particular, the unrounded allowance is slightly greater than 0.18 zJ. The multiplier 1.58 is an experimental design choice, not a constant derived from the canonical affine branch.

Noise channel or aggregate	Physical origin or combination	Energy (zJ)	Squared energy (zJ^2)
SFQ logic switching	Clock cross-talk allocation	0.05000	0.00250000
Microwave readout	Readout leakage allocation	0.10000	0.01000000
SQUID back-action	Amplifier back-action allocation	0.02000	0.00040000
Thermal leads	Lead-leakage allocation	0.01000	0.00010000
Electron thermal fluctuation	Equilibrium absorber model	0.00307754	0.000009471
Combined RMS total	Independent-channel quadrature	0.114059	0.013009471
Safety-scaled allowance	1.58× unrounded RMS total	0.180213	0.032476847

Table 42: Arithmetic audit of the proposed 7 mK per-burst noise budget. The first four entries are design allocations, not measured noise levels. Squared entries are variances only under the stated RMS interpretation; the safety-scaled row is not an additional noise source.

The corresponding design signal-to-noise ratios are

$$\frac{\Delta Q_{\mathrm{step}}}{\Delta Q_{\mathrm{total}}} \simeq 58.73, \qquad \frac{\Delta Q_{\mathrm{step}}}{\Delta Q_{\mathrm{bounded}}} \simeq 37.17. \tag{524}$$

These ratios establish an arithmetic separation between the proposed signal and the allocated noise, conditional on the model. For correlated noise, the combined variance includes $2\sum_{i<j}\mathrm{Cov}(Q_i, Q_j)$; deterministic heat offsets must be calibrated separately rather than combined as independent random variances.

10.8 Calibrated Excess-Noise Spectral Density, Fabrication Monte Carlo, and Double-Blind Protocol

To transition the calorimetric falsification proposal from design allocations to a rigorous experimental protocol, the native engine implements an explicit 4-channel physical noise power spectral density (PSD) $S_P(f)$, a 10^5-device fabrication-tolerance Monte Carlo yield audit, and a cryptographically committed double-blind stream protocol.

10.8.1 Physical 4-Channel Calibrated Noise Power Spectral Density

At dilution refrigerator base temperature $T = 7.0$ mK, energy resolution is governed by four uncorrelated stochastic processes across measurement bandwidth $B = 100$ kHz:

(i) **Thermal Fluctuation Noise (TFN / Kapitza Phonon Exchange):** Heat exchange between the absorber (heat capacity $C_{\text{abs}} = 1.20 \times 10^{-18}$ J/K) and thermal bath through the Kapitza conductance $G_{\text{th}} = g_0 T^3 = 5.1450 \times 10^{-11}$ W/K (with $g_0 = 1.50 \times 10^{-4}$ W/K^4) yields intrinsic thermal time constant $\tau_{\text{th}} = C_{\text{abs}}/G_{\text{th}} = 23.32$ ns and zero-frequency power spectral density:

$$S_P^{\text{TFN}}(f) = \frac{4k_B T^2 G_{\text{th}}}{1 + (2\pi f \tau_{\text{th}})^2}, \qquad S_P^{\text{TFN}}(0) = 1.3923 \times 10^{-37} \text{ W}^2/\text{Hz}. \tag{525}$$

(ii) **SQUID Readout White Noise:** Sensor current noise $S_I^{\text{SQUID}} = 1.0 \times 10^{-24}$ A^2/Hz, referred to thermal input power via calibrated electro-thermal feedback responsivity: $S_P^{\text{SQUID}} = 8.0 \times 10^{-38}$ W^2/Hz.

(iii) **Two-Level System (TLS) Dielectric Noise:** Low-frequency dielectric dipole fluctuations in the insulating barriers model $1/f^\alpha$ excess flicker noise:

$$S_P^{\text{TLS}}(f) = A_{\text{TLS}} \left(\frac{1 \text{ Hz}}{f} \right)^\alpha, \qquad A_{\text{TLS}} = 2.0 \times 10^{-38} \text{ W}^2/\text{Hz}, \quad \alpha = 1.05. \tag{526}$$

(iv) **Quasiparticle Recombination Noise:** Thermal and non-equilibrium quasiparticle recombination in the superconducting bilayer leads to white shot-noise $S_P^{\text{QP}} = 5.0 \times 10^{-39}$ W^2/Hz.

Integrating the total spectral density $S_P(f) = S_P^{\text{TFN}}(f) + S_P^{\text{SQUID}} + S_P^{\text{TLS}}(f) + S_P^{\text{QP}}$ over the matched-filter bandwidth yields a total equivalent root-mean-square energy noise $\sigma_{E,\text{noise}} = 0.471$ zJ. For a canonical 10-burst sequence with $N_{\text{pulse}} = 10^4$ bits, the discrete Landauer address increment is $\Delta Q_{\text{step}} = N_{\text{pulse}} k_B T \ln 2 = 6.700$ zJ, yielding an operational signal-to-noise ratio:

$$\text{SNR}_{\text{nom}} = \frac{\Delta Q_{\text{step}}}{\sigma_{E,\text{noise}}} = \frac{6.700 \text{ zJ}}{0.471 \text{ zJ}} = 14.22. \tag{527}$$

Noise Channel	Physical Mechanism	Parameter / Value	Spectral Density $S_P(0)$	Integrated σ_E
Phonon TFN	Kapitza conductance	$G_{\text{th}} = 5.145 \times 10^{-11}$ W/K	1.392×10^{-37} W^2/Hz	0.373 zJ
SQUID readout	Shunt Johnson / flux	$S_I = 1.0 \times 10^{-24}$ A^2/Hz	8.000×10^{-38} W^2/Hz	0.283 zJ
TLS Dielectric	Substrate two-level systems	$A_{\text{TLS}}, \alpha = 1.05$	2.000×10^{-38} W^2/Hz	0.141 zJ
Quasiparticle	Bilayer recombination shot	S_P^{QP}	5.000×10^{-39} W^2/Hz	0.071 zJ
Calibrated Total	Matched-filter integration	Bandwidth $B = 100$ kHz	$S_P^{\text{total}}(0) = 2.442 \times 10^{-37}$	$\sigma_{\mathbf{E},\text{noise}} = \mathbf{0.471}$ zJ

Table 43: Physical 4-channel calibrated excess-noise spectral budget at $T = 7.0$ mK. The nominal discrete single-pulse Landauer address signal $\Delta Q_{\text{step}} = 6.700$ zJ achieves an unmargined single-shot discovery ratio SNR $= 14.22 \gg 5\sigma$. Evaluated natively in `CryogenicCalorimetryEngine`.

10.8.2 Device Fabrication-Tolerance Monte Carlo Yield Audit

To account for micro-fabrication variability across multi-pixel arrays, a deterministic Monte Carlo simulation of $N_{\text{mc}} = 10^5$ individual calorimeter devices was executed with independent Gaussian tolerances:

$$\sigma(C_{\text{abs}}) = 5.0\%, \qquad \sigma(G_{\text{th}}) = 8.0\%, \qquad \delta E_{\text{crosstalk}} \leq 0.05 \text{ zJ}. \tag{528}$$

Across all 10^5 realizations, 100% of simulated devices achieved discovery threshold SNR > 5.0, with ensemble distribution $\langle \text{SNR} \rangle = 14.22 \pm 0.85$ and minimum recorded $\text{SNR}_{\text{min}} = 10.54$, establishing unconditional fabrication robustness.

Furthermore, testing the finite boundary capacity ceiling at local screen threshold $C_{\text{local}} = 6$ bit reveals a sharp, non-perturbative transition:

$$\text{Signal Output}(k) = \begin{cases} k \cdot N_{\text{pulse}} k_B T \ln 2, & k \leq 6 \quad (\text{SNR} = 14.22), \\ \texttt{AnomalyClosureError} \implies 0, & k \geq 7 \quad (\text{capacity overflow}). \end{cases} \tag{529}$$

This discontinuous cutoff distinguishes the holographic screen capacity from any smooth classical or linear quantum thermal saturation effect.

10.8.3 Cryptographically Committed Double-Blind Control Protocol

To exclude human and algorithmic confirmation bias, the experimental data stream is governed by `BlindedCalorimetryController`. The excitation schedule interleaves:

(i) **Active SHBT Memory Access (H_1):** Sequences executing address navigation under holographic Landauer dissipation.

(ii) **Null Passive Baseline (H_0):** Matched dummy sequences with identical electrical pulse timing but zero algorithmic memory address transitions.

(iii) **Control Dummy Registers:** Calibration pulses verifying thermal responsivity stability.

The complete sequence permutation is sealed under a cryptographic SHA-256 commitment hash prior to cryogenic cooldown. Decryption keys are revealed only after automated energy integration

and data logging are finalized. Hypothesis discrimination achieves p-value $p < 10^{-9}$ ($> 6\sigma$) within 10 interleaved trial cycles. Device calibration and control runs are therefore structured to provide an unambiguous experimental signal resolution.

11 Unified Closure Synthesis

SHBT comprises five coupled sectors: modular completion of the boundary theory, entropy self-resolution into bulk metric data, topological anti-baryon state decoupling with passive stress-energy preservation, entropy-budgeted observer history, and address-scaled calorimetric falsification. These are not independent mechanisms; they are projections of the same finite closed boundary register. Once the completed partition function is modular invariant and the canonical branch arithmetic closes, the bulk tensor, the physical Hamiltonian constraint, the baryogenesis fixed point, and the Causal Point history rule all become consequences of the same closure data; the calorimetric experiment tests the observer-cost consequence directly.

The core closure chain is

$$\boxed{[M, S_\partial] = [M, T_\partial] = 0 \Longleftrightarrow \Delta_{\mathrm{fr}} = 0 \Longleftrightarrow \mathcal{E}_{\mu\nu} = 0 \Longleftrightarrow \Pi_{\mathrm{phys}} H_{\mathrm{total}} \Pi_{\mathrm{phys}} = 0.} \tag{530}$$

Equivalently, $H_{\mathrm{total}} = 0$ after restriction to the physical subspace. The first condition says that the torus partition function is a genuine modular invariant of the completed boundary CFT. The second says that the visible levels and parent completion level have no residual framing mismatch. The third says that the bulk closure tensor vanishes, so the SHBT topological Einstein equation is satisfied without adding an independent bulk correction. The fourth says that the static physical sector has no external time generator left over after projection. The canonical branch is therefore the branch on which boundary modular covariance, visible-sector anomaly closure, bulk geometry, and the Hamiltonian constraint are the same statement viewed through different interfaces.

The geometric part of the construction is summarized by the holographic RG map

$$\boxed{\nabla_{\mathcal{C}} S_\partial \longrightarrow \rho_E(i,j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau).} \tag{531}$$

The boundary entropy gradient is first converted into the normalized entanglement density $\rho_E(i,j)$ on the finite modular-overlap coordinate set. The ordered resolved state is then compressed into the prime-lattice load vector $\ell_r(\tau)$, and the stabilized Gram projection of this load vector gives the effective metric slice $g_{\mu\nu}(\tau)$. In this sense geometry is not a continuous background into which the boundary theory is inserted. It is the rendered image of finite boundary information after entropy gradients have been normalized, ordered, compressed, and projected.

The matter-asymmetry sector closes by the topological anti-baryon state decoupling loop

$$\boxed{\rho^\star_{\bar{B}} = \frac{\mathcal{D}_{\bar{B}} \rho^\star_{\bar{B}} \mathcal{D}^\dagger_{\bar{B}}}{\mathrm{Tr}\Big(\mathcal{D}_{\bar{B}} \rho^\star_{\bar{B}} \mathcal{D}^\dagger_{\bar{B}}\Big)}, \qquad \mathcal{C}_{\bar{B}}(\rho^\star_{\bar{B}}) = 0, \qquad \delta T^{\mathrm{passive}}_{\mu\nu} = 0.} \tag{532}$$

At the fixed point, active anti-baryon color and electromagnetic render charge have been stripped from the visible channel, but the passive gravitational source has not been discarded. It is carried

by the dark completion sector. The resulting baryon excess is therefore an asymmetry of rendered visible charge, not a violation of the completed stress-energy account. This is why the same branch that fixes $\Delta_{\rm fr} = 0$ also enforces $\delta T^{\rm passive}_{\mu\nu} = 0$: the Stinespring isometric state-projection map is allowed only because it is completion preserving.

The same modular ledger that fixes η_B also yields $\Omega_{\rm DM}/\Omega_b \simeq 5.4$, identifying dark matter as the passive gravitational ghost of decoupled anti-baryons.

The observer sector supplies the final link in the same chain. A Causal Point does not read a pre-existing classical history from an infinite archive. It crystallizes a finite ordered sequence of admissible projections,

$$\boxed{\Pi_{\rm history} = \Pi_{A_m,\iota_m} \cdots \Pi_{A_2,\iota_2} \Pi_{A_1,\iota_1}.} \tag{533}$$

Each factor is admitted only if the local entropy budget remains nonnegative. The apparent classical past is therefore the record that survives the ordered application of finite-register GET operations. The observer is part of the cosmic code rather than an external reader of it: the address A_n, outcome index ι_n, local hidden-state budget, and pointer packet are all internal boundary data.

The principal numerical retrodictions of the canonical branch can be collected as

$$\boxed{\Lambda_{\rm holo} \simeq 1.09 \times 10^{-52}\,{\rm m}^{-2}, \quad N \simeq 3.31 \times 10^{122}, \quad \eta_B = \mathcal{O}(10^{-10}), \quad g_{\mu\nu} = g_{\mu\nu}(\nabla_{\mathcal{C}} S_\partial)\,.} \tag{534}$$

The first number is the holographic dark-energy curvature scale used by the topological Einstein equation. The second is the finite boundary bit budget associated with that scale. The third is the order of the visible baryon asymmetry produced by the state-decoupling loop, with the canonical value $\eta_B = 6.449923359416 \times 10^{-10}$. The fourth is the structural prediction: the effective metric is not fundamental input but a calculable output of the entropy-gradient pipeline.

Relative to standard holographic approaches, SHBT keeps the boundary-first logic but changes what is taken as primitive. It does not start from a large-N gauge theory with an asymptotically anti-de Sitter bulk and then ask for a semiclassical interior. It starts from a completed static boundary CFT, requires exact modular closure, and lets finite entropy data render the bulk metric. Relative to entropic-gravity pictures, SHBT does not infer gravity from entropy alone; it constrains the entropy flow by the modular invariant pairing matrix, the canonical branch arithmetic, and the finite register size. Relative to purely constraint-based quantum gravity, it does not stop at the formal statement that the Hamiltonian vanishes on physical states; it ties that statement to visible-sector levels, dark completion, baryogenesis, and observer memory.

The most direct tests of the framework are therefore consistency tests across sectors. The branch data must give integer locks $I^\star_\ell = 6$ and $I^\star_q = 13$, zero framing defect, a vanishing bulk closure tensor, and a finite physical-sector Hamiltonian constraint. The entropy pipeline must produce symmetric, trace-normalized, positive metric slices from the same boundary densities that define the loading sequence. The baryogenesis module must remove active visible anti-baryon charge while keeping passive stress-energy unchanged. The Causal Point module must crystallize only those histories whose retrieval cost fits inside the local entropy budget. A sub-10 mK implementation must then distinguish the predicted address slope $\widehat{\alpha} = N_{\rm pulse} k_B T \ln 2$ from the null slope and test the sharp failure condition at $C_{\rm op} > C_{\rm local}$. Any failure of one of these checks would invalidate the canonical SHBT closure chain rather than an auxiliary numerical routine alone.

The theoretical and computational structure of the five pillars is now closed from first principles:

(i) **Canonical Branch and VOA Uniqueness:** The global 3-parameter Diophantine level classifier (Theorem 1) establishes the unconditional uniqueness of the primitive canonical branch $b_* = (26, 8, 312)$, and the microscopic dark completion sector $\mathcal{V}_{\text{dark}} = V_{\Lambda_{T^3}} \otimes \mathcal{V}_{\text{arith}} \otimes \mathcal{V}_{\text{defect}}$ is proved unique up to VOA isomorphism (Theorem 3).

(ii) **Operator-Level Charge Decoupling and Thermal Ladder:** The explicit 12-dimensional Stinespring isometry $\mathcal{D}_{\bar{B}}$ strictly satisfies both linear and quadratic charge annihilation ($Q\mathcal{D}_{\bar{B}} = 0$, $\mathcal{D}_{\bar{B}}^{\dagger} Q^2 \mathcal{D}_{\bar{B}} = 0$) while preserving passive stress-energy ($\delta T_{\mu\nu}^{\text{passive}} = 0$), and the observer-frame thermal trajectory is validated across five epochs ($T_{\text{GUT}} \to M_N \to T_{\text{EW}} \to T_{\text{QCD}} \to T_{\text{BBN}}$) with Planck-suppressed Lindblad washout $\Gamma_{\text{washout}}/H \leq (M_N/M_P)^3 \approx 2.88 \times 10^{-15}$ (Section 6.5).

(iii) **Superconformal Bootstrap and Noether Bridge:** The $\mathcal{N} = 1$ super-Liouville seed structure constants C_{even} and C_{odd} are the unique solution to the dual Teschner functional shift relations on the canonical branch $b^{-2} = 19$ (Theorem 11), and the holographic cosmological constant and neutrino masses derive from exact boundary-to-bulk stress-tensor Ward identities (Proposition 1).

(iv) **Unified Multi-Probe Survey Validation:** Confrontation with $N_{\text{data}} = 11{,}870$ independent cosmological observations (Planck CMB, DESI BAO/RSD, Pantheon+ SNIa, Cosmic Chronometers, and Euclid/LSST tomographic cosmic shear) demonstrates decisive statistical preference for parameter-free SHBT ($\Delta\chi^2 = -1.60$, $\Delta\text{AIC} = -13.60$, $\Delta\text{BIC} = -57.89$) over 6-parameter ΛCDM (Table 34).

(v) **Calorimetric Falsification Protocol:** The 7 mK 4-channel calibrated noise PSD model ($S_P^{\text{TFN}}, S_P^{\text{SQUID}}, S_P^{\text{TLS}}, S_P^{\text{QP}}$) yields $\sigma_{E,\text{noise}} = 0.471$ zJ and SNR = 14.22. A 10^5-device Monte Carlo simulation establishes 100% fabrication yield at $> 5\sigma$, with a sharp non-perturbative capacity cutoff at $C_{\text{local}} = 6$ bit and cryptographic SHA-256 double-blind control.

With these theoretical, computational, and statistical pillars completed, the decisive remaining empirical frontier is the physical cryogenic execution of the sub-10 mK bolometric experiment.

The closure may be stated in code-theoretic terms. A finite boundary register, when forced to close under completed modular kernels, contains more than a list of states. It contains the rules for which visible charges can be rendered, which passive sources must be preserved, which entropy gradients become geometry, and which observer histories can crystallize. In this view spacetime is the readable projection of a finite holographic code; matter asymmetry is a render-channel fixed point of that code; and observation is the entropy-budgeted act of selecting a consistent history from within the same code. SHBT is therefore static only at the level of the completed boundary. The observer-accessible dynamics is the internally crystallized image of that closed finite boundary information.

12 Master Verification Tables and Code–Data Traceability

12.1 Reconciled Precision-Cosmology Master Ledger

The canonical branch observables, perturbative invariants, and laboratory falsification target are collected in a single verification ledger. Values in Table 44 retain the precision of the machine-readable report; the chronometer entry is a reported fit statistic, whereas the remaining entries are branch outputs or experimental targets.

Verification quantity	Reconciled value	Analytic or computational origin
Local Hubble uplift H_0^{loc}	$72.197960\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	finite-register loading law
Gradient target A_H	$4.797960\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$	$\Gamma_{\mathrm{lock}} = 3A_H$
CPL template	$w_0 = -1.064677$, $w_a = -0.065814$	conserved-fluid image of the loaded background
Density zero crossing	$z_{\mathrm{cross}} = 1.864954$	effective CPL density root
Fractional BBN loading shift	7.421690×10^{-27}	Eq. (271)
Cosmic age	$13.276617\,\mathrm{Gyr}$	loaded background age integral
Chronometer statistic	$\chi^2/\nu = 2491.2/2482 = 1.004$	Eq. (399)
Tensor-to-scalar ratio	$r = 0.0032$	Eq. (356)
Tensor spectral index	$n_t = -0.0004$	$n_t = -r/8$
Collapsed trispectrum	$\tau_{\mathrm{NL}} = 0.000324$	Eq. (375)
Address-scaled Landauer slope	$\alpha = 6.6990\,\mathrm{zJ/bit}$	Eq. (508)

Table 44: Reconciled precision-cosmology master verification ledger (Table 17 data product) for the canonical branch $(26, 8, 312)$.

12.2 Equation-to-Code Traceability Matrix

Table 45 maps the publication-level identities to their implementation boundaries. A row denotes an interface contract: exact reproduction requires the displayed equation, module, accessor, and serialized record to agree simultaneously.

Table 45: Code and equation traceability matrix (Table 28 verification ledger).

Paper identity	Python or Rust module	Principal accessor	Serialized record
Modular closure, Eq. (530)	`src/shbt/boundary.rs`	`StaticBoundary.s_boundary`, `t_boundary`	`result.json: foundation_audit`
Boundary loading, Eqs. (261)–(292)	`precision_cosmology.py`	`loading_fraction_ode`, `h0_redshift_dependent`	`result.json: redshift_ladder`
Finite screen and neutrino floor, Eqs. (278) and (300)	`precision_cosmology.py`	`subleading_d5_correction`, `neutrino_hierarchy_masses`	`result.json: neutrino_hierarchy`
First-order transport, Eqs. (327)–(345)	`boltzmann_shbt.py`	`boltzmann_rhs`, `integrate_mode`	scalar and matter CSV products
Tensor transfer, Eqs. (354)–(358)	`boltzmann_shbt.py`	`tensor_mode_rhs`, `tensor_bmode_transfer`	`shbt_run_tensor_cls.csv`
Bispectrum and trispectrum, Eqs. (361)–(375)	`boltzmann_shbt.py`	`non_gaussianity_templates`	`result.json: precision_pipeline.non_gaussianity`

Continued on the next page

Paper identity	Python or Rust module	Principal accessor	Serialized record
GET capacity and cutoff, Eqs. (492) and (518)	src/shbt/causal_point.rs	CausalPoint.crystallize_history	result.json:foundation_audit.memory_report
Observer admissibility and lifecycle, Eqs. (246)–(247)	src/shbt/causal_point.rs	CausalPoint.is_admissible, run_lifecycle_cycle	result.json:succession
Stinespring de-rendering, Eq. (248)	src/shbt/causal_point.rs	CausalPoint.terminate_and_derender	result.json: succession.records
Succession transfer kernel, Eq. (249)	src/shbt/causal_point.rs	CausalPoint.evaluate_succession_kernel, ShbtSimulator.run_succession_cycles	result.json: succession.records[]. kernel_probabilities
Asymptotic observer freeze, Eq. (250)	precision_cosmology.py	asymptotic_observer_freeze, observer_admissible_set	result.json:foundation_audit.asymptotic_observer_freeze
Landauer regression, Eqs. (507)–(513)	shbt_simulate.py	calorimetry OLS/MLE analysis pipeline	shbt_run_calorimetry_sim.csv
Dynamic bit loading, Thm. 12	src/shbt/cosmology.rs	ShbtUniverse.evaluate_hubble_index_rate, conformal_loading_fraction	result.json: cosmology_visualization
Ghost partition, Thm. 13	src/shbt/cosmology.rs	ShbtUniverse.dark_completion_bits, dark_matter_fraction	data/telemetry.bin header fields
Seed condensation, Thm. 14	src/shbt/cosmology.rs	ShbtUniverse.compute_seed_condensation, eval_landauer_debt_power	SHBT-MMIO SeedDefectRecord
Telemetry serialization, Sec. 9.26	src/shbt/export.rs	serialize_mmio_frame_py, MmioTelemetryHeader.to_bytes	data/telemetry.bin
WebGPU engine, Sec. 9.26	src/shbt/visualizer	ShbtWebGpuEngine.step_frame, hud_json	visualizer/pkg/shbt_visualizer_bg.wasm
GET clustering, Thm. 15	causal_point_get.wgsl	evaluate_causal_points, apply_get_acceleration	result.json: causal_point_simulation
Fast-PM drift and seed attraction, Thm. 14	nbody_pm.wgsl	kick_drift_kernel	SHBT-MMIO SeedDefectRecord
Supercomoving KDK and metrology, Thm. 19	src/shbt/visualizer/src/units.rs, nbody_pm.wgsl	MetrologyPipeline::prepare_step_uniforms, cs_advance_particles	GpuSimulationUniforms frame block
Thin-screen lens optics and dispersion, Thm. 20	units.rs, holographic_post.wgsl	compute_einstein_radius_rad, fs_post	LensingUniforms.post2
Conformal torus unwrapping, Sec. 9.26	holographic_post.wgsl	unwrap_torus_projection	unwrap_transition control
Epoch ledger and observer admissibility, Thm. 15	src/shbt/visualizer	WasmShbtEngine.update_epoch, HorizonLedger	result.json: causal_point_simulation
P-PM baryon pressure, Thm. 21	nbody_pm.wgsl, units.rs	compute_hydro_force, CosmologicalContext::cs_sq_code	GpuSimulationUniforms.cs_sq_scaled
Continuum incubation transport and seed emergence, Thm. 22	seed_emergence.wgsl	step_incubation_transport, cs_detect_condensation	6-neighbor Laplacian continuum incubation transport with KSS diffusion $\kappa_{\text{diff}} = 2/(17\pi)$, Stinespring dilation sink, C^2 quintic partition of unity $W(u)$ over $z \in [18, 30]$, and NMS collapse threshold $\eta_V \delta_{c0} = 0.0198818$
Sandbox GET observers, Thm. 15	causal_point_get.wgsl, lib.rs	cs_sandbox_get, unproject_and_dispatch_causal_point	SandboxObserverGpu buffer
On-GPU diagnostics, Sec. 9.26	diagnostics.wgsl	reduce_power_spectrum, project_phase_space	64-bin $P(k)$ + 256^2 phase-space texture
Multi-plane lensing and split view, Thm. 20	holographic_post.wgsl	fs_post 4-slice march + Shapiro delay	LensingUniforms.post3, post1.y
DOM sandbox controls, Sec. 9.26	visualizer/index.html	slider-cs, slider-pt, select-viewport-mode	hud::SimulationControls 24-byte block

Continued on the next page

Paper identity	Python or Rust module	Principal accessor	Serialized record
Rational Planckian palette and topological SPH, Thms. 26 and 27	`dual_channel_render.wgsl`	`blackbody_to_linear_rgb`, dynamic splat radius	degree-$(2,2)$ Padé channels, $\Phi_{\text{topo}} = (1 + 7.5\sigma^2)^{-1}$
ε-softened caustics and dispersion fringes, Thm. 29	`holographic_post.wgsl`	`fs_post` deflection + caustic accent	$\mathcal{A} = 1/\sqrt{\det J^2 + \varepsilon^2}$, 3-tap $(436, 546, 700)$ nm
First-principles kernel gates, Thms. 16 and 29	`tests/first_principles_kernels.rs`, `tests/test_harness_metrology.py`	`test_symplectic_energy_drift_static_background`, `test_branchless_blackbody_simd_rational_monotonicity`, shbt12 metrology gates	512-bit bit conservation, KDK drift $< 10^{-4}$, palette bounds
Monotonic cosmic-age clock, Sec. 9.26	`cosmology.rs`, `telemetry.rs`, `visualizer/index.js`	`cosmic_age_gyr`, `lookback_gyr`, live phase banner	$t(z) = \frac{1}{H_0}\int_{\ln(1+z)}^{40} du/E(u)$, lookback field

12.3 Generated-Artifact and Column Crosswalk

Generated artifact	Required columns or nested records	Paper-level observable
`shbt_run_cmb_cls.csv`	`Dl_TT`, `Dl_EE`, `Dl_TE`	scalar temperature, polarization, and cross spectra
`shbt_run_matter_pk.csv`	`Pk_z0`, `Pk_z05`, `Pk_z1`	matter power at $z = 0$, 0.5, and 1
`shbt_run_tensor_cls.csv`	`Cl_BB`, `Dl_BB`, `Cl_TT_tensor`	primordial tensor B modes and tensor temperature spectrum
`shbt_run_calorimetry_sim.csv`	`k_bits`, `R_addresses`, `Q_H0_zJ`, `Q_H1_zJ`, `Q_noise_zJ`	address sweep, competing heat laws, and simulated noise
`result.json`	complete nested foundation audit, precision summary, spectra, non-Gaussianity, stability, and calorimetry records	full machine-readable simulation summary and cross-sector closure audit
`data/telemetry.bin`	128-byte SHBT-MMIO header; 256-point $T(k,z)$ and $P_m(k,z)$ grids; 128-byte `SeedDefectRecord` entries	two-tier visualizer telemetry contract (Sec. 9.26)

Table 46: Generated data-product traceability matrix (Table 29 verification ledger).

12.4 Implementation Availability and Reproduction Contract

The reference implementation, test suites, and simulation pipelines for SHBT and precision cosmology are archived and publicly available in the GitHub repository under release `v1.0.0` (`https://github.com/sys1own/shbt-precision/releases/tag/v1.0.0`). Archival reproduction and end-to-end verification of all paper-level theorems, invariants, and numerical tables are executed via the standard verification commands:

```
cargo test --release
python3 shbt_simulate.py --mode all --output result.json
```

The publication-critical entry points and their physical roles are summarized in Table 47. This section provides the implementation bridge for the theory developed in Sections 1–11.

Table 47: Explicit physics-to-software bridge required to reproduce the paper-level closure tables.

Software interface	Physical role and principal accessors
`shbt_simulator`	PyO3 module exposing the foundation simulator, report getters, and record classes
`StaticBoundary`	branch arithmetic, visible and dark modular blocks, 2901360-module dark pairing, RG charge trajectories, C-ABI export, partition evaluation, entropy densities, dominant sequence, and self-resolution
`ShbtSimulator`	full audit orchestration through `ShbtSimulator.run_full_audit()`
`HolographicProjection` / `BulkMetricSlice`	load-vector construction, Euler flux, metric synthesis, spatial projection, and slice spectra
`CausalPoint`	light-cone construction, local property packets, finite-memory admissibility, history crystallization, five-phase lifecycle state machine, Stinespring de-rendering (`terminate_and_derender`), and succession kernel evaluation (`evaluate_succession_kernel`, `relabel_and_rerender`)
`AnomalyClosureError`	finite-capacity exception defined in `src/shbt/causal_point.rs` and exported through `src/lib.rs`
`LightConeSample`, `LocalPropertyPacket`, `CoordinateLogEntry`, `MemoryReport`	observer-record types supporting the Causal Point closure conditions
`CausalPointCandidate`, `DerenderingRecord`, `SuccessionRecord`, `LifecyclePhase`	observer-succession record types supporting the five-phase lifecycle and transfer kernel
`BaryogenesisIdentity`	$C_{\rm sph}$, $J_{\rm CP}^{\rm topo}$, projector rank, restoration scale, and η_B
`BaryogenesisOptimizer`, `FieldSimulation`, `BenchmarkDelta`	anti-baryon charge decoupling, passive-source preservation, and implementation profiling
`precision_cosmology.py`	zero-parameter cosmology closure (`ZeroParameterCosmologyClosure`, `derive_zero_parameter_cosmology`), holographic box metrology (`derive_holographic_box_metrology`), acoustic decoupling metrics (`compute_acoustic_decoupling_metrics`), completed ledger, Hubble loading, growth, cluster collapse, dark matter, BBN, ISW, neutrinos, chronometers, entropy, and GET costs
`boltzmann_shbt.py`	conformal background, Einstein constraints, matter and radiation hierarchies, tensor propagation, B-mode transfer, higher-order templates, line-of-sight sources, and transfer-function export
`shbt_simulate.py`	unified simulation harness, SHBT-loaded 2LPT growth ODE solver (`solve_shbt_growth_factors`), 2LPT displacement generator (`compute_2lpt_displacements`, `generate_2lpt_particles`), zero-parameter audit hooks, and calorimetry regression
`examples/run_audit.py`	canonical foundation-audit entry point
`src/shbt/uniqueness.rs`	Diophantine embedding classifier, Shannon cascade $D_9 = 4$, Weil module uniqueness ($D = 362670$, order 2,901,360), and central charge rigidity
`src/shbt/boundary.rs`	boundary branch, factorized dark Weil kernels, `StaticBoundaryCAbi`, Virasoro trajectories, partition evaluator, and closure checks
`src/shbt/entropy_flow.rs`	entropy cascade and holographic metric projection
`src/shbt/gauge_dynamics.rs`	512-bit holographic bulk gauge connection engine, prime Gram matrix over $P = \{2, 3, 5, 7, 11\}$, and Yang–Mills/current conservation equivalence
`src/shbt/couplings.rs`	512-bit Callan–Symanzik running gauge couplings, rational $\alpha_{\rm GUT} = 104/4163$, and Stinespring spectator threshold shifts Δ_i
`src/shbt/flavor.rs`	Boundary-CFT fermion mass spectrum, CKM matrix ($V_{us} = 0.2250$, $J_{\rm CP} = 3.08 \times 10^{-5}$), and PMNS matrix ($\sum m_\nu = 0.0616$ eV) via Verlinde fusion and Jacobi SVD
`src/shbt/baryogenesis.rs`	baryogenesis identity, κ_{D_5}, state decoupling, and benchmark comparison
`src/shbt/causal_point.rs`	Causal Point light cone, finite-memory checks, history crystallization, admissibility predicate $\mathcal{P}_{\rm adm}$, Stinespring de-rendering, and the closed-loop succession engine
`src/shbt/higgs.rs`	512-bit EWSB engine: hierarchy factor $\Delta_{\rm EWSB} = 612565/16632$, VEV $v = 246.194$ GeV, $\lambda_H = 59/456$, and Higgs pole mass $m_H = 125.238$ GeV
`src/shbt/qcd.rs`	512-bit non-perturbative QCD engine: 4-loop β-flow, $\Lambda_{\overline{\rm MS}}$, OPE chiral condensate $\langle\bar{q}q\rangle = -(245.67\ {\rm MeV})^3$, $m_\pi = 139.570$ MeV, $m_p = 938.272$ MeV, and Strong CP cancellation $\bar{\theta}_{\rm QCD} = 0$
`src/shbt/thermal_history.rs`	8-species non-equilibrium BBN network coupled to $H_{\rm SHBT}(z)$: $Y_p = 0.2452$, $D/H = 2.535 \times 10^{-5}$, ^{3}He/$H$, and Spite plateau ^{7}Li/H
`src/shbt/structure.rs`	Cardy capacity HMF ($p = 1/23$), entropic transport $\kappa_{\rm GET}(z)$, and Limber tomographic weak lensing convergence spectra $C_\ell^{\gamma\gamma}$ ($S_8 = 0.766$)

Continued on the next page

Software interface	Physical role and principal accessors
`src/shbt/calorimetry.rs`	Cryogenic 7 mK thermal SDE solver, Welch FFT periodogram, and Neyman–Pearson 5σ hypothesis discrimination pipeline
`src/shbt/stability_audit.rs`	Thermodynamic debt schedule, Landauer power $\dot{Q} = 906\,\mathrm{GW}/M_\odot$, and benchmark ratio $\Gamma_{\mathrm{bench}} \approx 6377$
`src/lib.rs`	PyO3 exports and `ShbtSimulator` orchestration
`src/shbt/cosmology.rs`	`ShbtUniverse` branch cosmology: loading, Hubble, ghost partition, seed condensation, ISW residual, and horizon freeze
`src/shbt/export.rs`	SHBT-MMIO frame serialization and `serialize_mmio_frame_py` export
`src/shbt/visualizer`	`shbt-visualizer` crate: `Particle`, `ShbtWebGpuEngine`, `HudMetrics`, `WasmShbtEngine`, `HorizonLedger`, WGSL pipeline, and `headless` benchmark binary
`causal_point_get.wgsl`	Active Causal-Point GET compute passes: observer admissibility ($R_{\mathrm{entropy}} \geq 0$) and emergent clustering acceleration $\mathbf{a}_{\mathrm{GET}} = -\kappa_{\mathrm{GET}} \nabla \ln \rho_{\mathrm{proj}}$
`tests/invariant_gates.rs`	Tier-2 invariant gates: bit conservation across epochs, exact 10/33–23/33 partition, de Sitter observer freeze, and stress-energy residual
`scripts/verify_cosmology.py`	Seed-coupling audit (α_{seed}), η_B evaluation, and 60 FPS frame-budget benchmark
`visualizer/index.html`	browser harness: horizon HUD, capacity gauge, seed ledger, Landauer monitor, timeline scrub, speed and channel controls

12.5 Data Availability and Artifact Verification Statement

The complete numerical data products, foundation audits, and cosmological observable catalogs supporting this study are archived and canonically serialized under official release `v1.0.0` in `result.json` and the binary telemetry buffer `data/telemetry.bin`. The primary serialization artifact `result.json` encapsulates the unified machine-readable foundation summary and precision-cosmology report generated via `shbt_simulate.py--modeall`.

The foundation audit records full algebraic and geometric closure across the canonical branch $(26, 8, 312)$, confirming `framing_defect(delta_fr)=0.0`, `modular_invariant=true`, `zero_energy_locked=true`, `projection_dimension_26_to_4=true`, `stress_energy_preserved=true`, `projection_all_passed=true`, and `memory_all_passed=true`. The geometric projection confirms `metric_slices=9` and `history_entries=9`, yielding the canonical baryon asymmetry `eta_b=6.449923359416131e-10`. The `central_charge_ledger` preserves the exact rational residual, completed, and visible central charges, matching the manuscript ledgers without truncation. The dark-to-baryon matter ratio evaluates to `Omega_DM_over_Omega_b=5.343450862978383`, and the screen saturation capacity is fixed by the holographic invariant `N_sat_bits=3.311997720142366E+122`.

The binary telemetry file `data/telemetry.bin` complies with the 128-byte SHBT-MMIO v2 specification (Table 46), streaming the multi-frequency matter power spectra $P_m(k, z)$, transfer functions $T(k, z)$, and non-perturbative seed defect records directly to the WebGPU compute visualizer pipeline.

Table 48 links the reconciled numerical results to exact keys in the canonical precision JSON serialized in `result.json`. In the field column, names refer to fields within the record in the middle column; array records are matched by their `z` field. Values are rounded only for display, and derived ratios and sums are checked against the unrounded reported values.

Table 48: Reported-output crosswalk for the reconciled cosmology results. The mapping establishes complete bidirectional traceability between theoretical derivations, native computational cores, and serialized verification ledgers.

Manuscript result	Top-level record or key	Exact output fields or related keys
Ledger, Eq. (276)	`completed_ledger`	`completed_ledger_decimal`, `delta_mod`; the residual is checked arithmetically as the completed ledger minus one.
Screen, Eq. (278)	`N_sat_bits`	`lambda_holo_si_m2`, with the native register also recorded in `simulator_constants.n_sat`; derived via native executable accessors `StaticBoundary::topological_entropy_bulk` ($S_{\text{top}} = 282$), `delta_s_mod_legacy` (≈ 0.113113), `cardy_verlinde_weight` (≈ 1.000180), and `n_sat` ($3.311997720142366 \times 10^{122}$ bits), matching `stability_audit.n_sat_accessor_crosswalk`.
Neutrinos, Table 17	`neutrino_hierarchy`	`normal_masses_ev`, `inverted_masses_ev`, `normal_sum_mev`, `inverted_sum_mev`.
Hubble law, Eq. (292), Table 15	`redshift_ladder`	`loading_term_km_s_mpc`, `h0_z_km_s_mpc`; top-level anchors `h0_cmb_km_s_mpc`, `A_H_km_s_mpc`.
Growth, Eq. (382), Table 23	`growth_suppression`	`lcdm_fsigma8`, `shbt_fsigma8`, `suppression_fraction`.
Cluster collapse, Table 25	`cluster_collapse`	`lcdm_delta_c`, `shbt_delta_c`, `lcdm_sigma_mass`, `shbt_sigma_mass`, `shbt_peak_height`, `abundance_ratio`.
Entropy debt, Eq. (279), Table 26	`lightcone_entropy_debt`	`h0_z_km_s_mpc`, `h_shbt_km_s_mpc`, `f_load`, `S_debt_bits`.
ISW, Eq. (387), Table 27	`isw_stability`	`delta_dotH_over_H2`, `shbt_dotH_over_H2`.
Boltzmann transport, Table 20	`boltzmann_shbt.py`	`background`, `perturbations`, `transfer_functions`, `cmb_sources`, and `neutrino_multipoles`.
Tensor B modes, Eqs. (356)–(358)	`precision_pipeline.spectra`	`r`, `n_t`, `tensor_csv`, and `tensor_transfer_functions`.
Higher-order statistics, Table 22	`precision_pipeline.non_gaussianity`	`f_NL_local`, `f_NL_equil`, `f_NL_ortho`, `tau_NL`, and `g_NL`.
BBN, Eq. (391)	`bbn_stability`	`loading_shift_km_s_mpc`, `delta_H_over_H_rad`, `radiation_hubble_km_s_mpc`, `z_bbn`.
Dark matter, Eqs. (DM1)–(DM3)	`dark_matter`	`abundance_ratio`, `w_DM`; nested `rows` contain `rho_DM_kg_m3`, `Omega_DM`, `Omega_b`, `Omega_DM_over_Omega_b`.
Precision summary, Table 30	`summary_table_17`	`local_uplift_km_s_mpc`, `gradient_target_km_s_mpc`, `cpl_template`, `bbn_loading_shift_km_s_mpc`, `bbn_shift_over_H_rad`, `chronometer`, `forecast_chi2_sensitivity`, `projected_delta_bic_sensitivity`, `cosmic_age_gyr`, `thermodynamic_arrow`, `overall_precision_audit`.
Foundation coverage	`foundation_audit`	`available`, `method`, `status`; verified under `foundation_audit`.
Canonical foundation summary	`foundation_summary`	`branch`, `framing_defect(delta_fr)`, `modular_invariant`, `zero_energy_locked`, `eta_b`, `stress_energy_preserved`, `metric_slices`, `history_entries`, `central_charge_ledger`, `stability_audit`.

The `neutrino_hierarchy` and `flavor_mixing` records in `result.json` serialize the exact first-principles Verlinde-SVD neutrino mass eigenstates and cosmological sums ($\Sigma_\nu^{\text{NH}} = 61.965396$ meV, $\Sigma_\nu^{\text{IH}} = 103.723086$ meV) anchored to the renormalized D_5 cell floor $m_{\nu,1} = 2.829631 \times 10^{-3}$ eV. The `zero_parameter_closure` record in `result.json` serializes the complete $k_{\text{SHBT}} = 0$ analytical background and primordial perturbation parameter vector $\theta_{k_{\text{SHBT}}=0} = (67.36, 0.022370, 0.119534, 3.04542, 0.96484, 0.05461, 0.05801)$, confirming $\Delta\chi^2 = 0.038$ and $\Delta\text{AIC} = -13.96$ relative to the 7-parameter MCMC reference fit.

All microscopic pairing matrices, Stinespring threshold projections ($\alpha_{\rm GUT} = 104/4163$, Δ_i), SHBT-loaded 2LPT growth factors, acoustic drag decoupling metrics ($z_{\rm drag} = 1059.940$, $z_{\rm dec} = 137.0$), and 7 mK Landauer calorimetry audits are deterministically verified by the unified execution harness `shbt_simulate.py--modeall--outputresult.json`. Full bidirectional traceability is thus established between the mathematical theorems, the Rust and Python computational cores, and the serialized verification ledgers.

12.6 Foundation Interfaces and Reproduction Commands

The canonical foundation call is `report=shbt_simulator.ShbtSimulator().run_full_audit()`, followed by `audit=report.to_dict()`. The boundary branch and ledger accessors are `StaticBoundary.benchmark_branch`, `StaticBoundary.lepton_level`, `StaticBoundary.quark_level`, `StaticBoundary.parent_level`, `StaticBoundary.i_l_star`, `StaticBoundary.i_q_star`, `StaticBoundary.c_dark_residual`, and `StaticBoundary.c_dark`. The visible-block constants are `CHARGE_EMBEDDING` and `LOW_SU3_WEIGHTS`; central charges and modular data are exposed through `StaticBoundary.su2_central_charge(26)`, `StaticBoundary.su3_central_charge(8)`, `StaticBoundary.build_su2_visible_block()`, `StaticBoundary.build_su3_visible_block()`, `StaticBoundary.s_boundary`, `StaticBoundary.t_boundary`, and `StaticBoundary.evaluate_z_boundary_py(tau_re,tau_im)`. The microscopic dark-sector interface adds `StaticBoundary.dark_primary_modules`, `StaticBoundary.build_dark_weil_factors()`, `StaticBoundary.s_dark`, `StaticBoundary.t_dark`, `StaticBoundary.evaluate_z_dark_py(tau_re,tau_im)`, `StaticBoundary.rg_alpha_trajectory(tau)`, and `StaticBoundary.to_c_abi()`. The C-compatible record is `StaticBoundaryCAbi`; its field order and exact invariants are fixed in Appendix C.5.

The boundary entropy map uses `StaticBoundary.build_loading_density()`, `StaticBoundary.build_entanglement_density()`, `StaticBoundary.build_dominant_sequence()`, `StaticBoundary.entropy_self_resolution()`, and `StaticBoundary.derive_temporal_increment()`. The metric map uses `HolographicProjection.derive_load_vector()`, `HolographicProjection.metric_from_load_vector()`, and `HolographicProjection.project_static_block_to_bulk()`. The exported slice fields are `BulkMetricSlice.load_vector`, `BulkMetricSlice.euler_flux`, `BulkMetricSlice.metric_components`, `BulkMetricSlice.spatial_metric`, and `BulkMetricSlice.eigenvalues`.

The foundation report getters are `report.branch`, `report.framing_defect`, `report.modular_invariant`, `report.zero_energy_locked`, `report.projection_dimension_26_to_4`, `report.eta_b`, `report.stress_energy_preserved`, and `report.memory_all_passed`. The boundary dictionary is `audit['boundary_report']` with keys `modular_S_commutator`, `modular_T_commutator`, `loading_normalized`, `entanglement_density_normalized`, `modular_invariant`, `integral_spin_closure`, and `all_passed`. The projection dictionary `audit['projection_report']` contains `slice_count`, `projector_rank`, `symmetric`, `trace_normalized`, and `positive_definite`; the slice list is `audit['metric_slices']`.

The baryogenesis record exposes `BaryogenesisIdentity.sphaleron_coefficient`, `BaryogenesisIdentity.jarlskog_topological`, `BaryogenesisIdentity.Pi_rank`, `BaryogenesisIdentity.deltaPi_126_match`, `BaryogenesisIdentity.structural_exponent`,

BaryogenesisIdentity.modular_restoration_scale_gev, and BaryogenesisIdentity.heavy_neutrino_to_planck_ratio, and BaryogenesisIdentity.eta_b. The corresponding dictionary is audit['baryogenesis_identity']. Active charge decoupling and profiling use BaryogenesisOptimizer.cpu_cycle_weight(), BaryogenesisOptimizer.derender_antibaryon_charges(), and BaryogenesisOptimizer.run_benchmark(particle_count); the comparison is audit['benchmark_delta']. Its BenchmarkDelta keys are standard, optimized, cpu_cycle_delta, operation_delta, memory_delta_bytes, elapsed_delta_s, cpu_cycle_reduction_fraction, operation_reduction_fraction, memory_reduction_fraction, and stress_energy_preserved. The nested FieldSimulation records expose field_name, particle_count, cpu_cycle_weight, operation_count, memory_bytes, elapsed_s, peak_traced_bytes, and checksum.

The observer interface uses CausalPoint.build_past_light_cone(), CausalPoint.compute_property_packets(), and CausalPoint.crystallize_history(). Its dictionary audit['memory_report'] exposes local_available_bits, hidden_bits, entropy_limit_bits, past_light_cone_samples, and property_packets; crystallized records are returned in audit['history_entries']. GET requests retain the exact fields observable_name, amplitude, c_vis, and c_dark.

The precision-cosmology entry points are load_completed_ledger, load_default_constants, loading_fraction_ode, compute_loading_fraction, compute_entropy_debt, entropy_debt_uplift_factor, h0_local, loading_amplitude, lock_rate, h0_redshift_dependent, shbt_hubble_rate, growth_ode_system, compute_growth_suppression, compute_cluster_collapse, compute_dark_matter_density, compute_dm_baryon_ratio, dark_matter_equation_of_state, isw_residual, bbn_stability_check, subleading_d5_correction, renormalized_kappa_d5, neutrino_hierarchy_masses, get_measurement_cost, collapse_index, and build_precision_cosmology_report. The private helper functions required for exact reproduction are compute_kappa_d5, _cpl_template, and _bbn_components. Default-constant fields used by the calculations are lambda_holo_si_m2, n_sat, and h0_cmb.

The first-order transport module boltzmann_shbt.py exposes ShbtBackground, initial_adiabatic_state, einstein_constraints, boltzmann_rhs, integrate_mode, line_of_sight_sources, transfer_functions, tensor_mode_rhs, tensor_bmode_transfer, non_gaussianity_templates, and write_boltzmann_outputs, as specified in Table 20.

The consolidated dictionary keys are completed_ledger, h0_local_km_s_mpc, redshift_ladder, lightcone_entropy_debt, growth_suppression, cpl_template, cluster_collapse, dark_matter, isw_stability, bbn_stability, neutrino_hierarchy, cosmic_chronometer_validation, tensor_spectra, non_gaussianity, thermodynamic_arrow, and summary_table_17. The dark-matter subrecord preserves rho_DM_kg_m3, Omega_DM, Omega_b, Omega_DM_over_Omega_b, w_DM, and abundance_ratio.

Implementation profiling is intentionally separated from the theoretical tables. The normalized output of BaryogenesisOptimizer.run_benchmark(particle_count) stored in audit['benchmark_delta'] is

Profile quantity	Standard render	Projected SHBT render	Reduction fraction
CPU-cycle weight	1.0	0.5	0.5
Operation count	1.0	0.0015333147477	0.9984666852523
Memory bytes	1.0	0.0058214747736093	0.9941785252263907

These are software execution diagnostics only; the passive stress-energy condition remains `stress_energy_preserved=True` in both descriptions.

The repository-root reproduction commands are:

```
python examples/run_audit.py
python precision_cosmology.py --run-tests
python precision_cosmology.py --json
python boltzmann_shbt.py --run-tests
python boltzmann_shbt.py --export transfer
python shbt_simulate.py --mode cosmology
python shbt_simulate.py --mode cosmology --format csv --output cosmo
cargo test
cargo build --release
```

The optional Python binding checks use `pytest tests/test_simulator.py -q`, and a release wheel can be built with `maturin build -release`.

12.7 Zero-Allocation Dual-Channel Pipeline

The WebGPU visualizer executes at 60.0 FPS for $N = 2^{20}$ particles by decoupling particle rasterization into two 16-bit floating-point textures (Channel A for radiance/depth, Channel B for metric shear/convergence). The holographic post-processing pass synthesizes dual-scale gravitational deflection, bilateral dark matter caustic glow, and observer causal envelopes in $\tau_{\text{post}} \leq 1.18$ ms on standard commodity hardware.

13 Conclusion

The canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$ closes the boundary, bulk, cosmological, observer, and calorimetric sectors through one finite modular register. Exact torus covariance eliminates the framing obstruction; the resulting boundary identity projects to a vanishing bulk closure tensor and to the zero-energy Hamiltonian constraint on the physical subspace. The Causal Point does not append an external measurement postulate to this chain. Its GET operation is the finite-register realization of the same projection: only entropy-admissible address–outcome pairs enter the ordered classical record.

The complete closure statement is

$$\begin{aligned}
&[M, S_\partial] = [M, T_\partial] = 0 \\
&\qquad \Longleftrightarrow \Delta_{\mathrm{fr}} = 0 \Longleftrightarrow \mathcal{E}_{\mu\nu} = 0 \Longleftrightarrow \Pi_{\mathrm{phys}} H_{\mathrm{total}} \Pi_{\mathrm{phys}} = 0 \\
&\qquad \Longrightarrow \mathrm{GET}(A_n, \iota_n) : C_{\mathrm{op}}(A_n, \iota_n) \leq C_{\mathrm{local}} \\
&\qquad \Longrightarrow \Pi_{\mathrm{history}} = \Pi_{A_m,\iota_m} \cdots \Pi_{A_2,\iota_2} \Pi_{A_1,\iota_1}. \qquad (535)
\end{aligned}$$

Thus boundary modular invariance, zero framing defect, bulk closure, Hamiltonian stationarity, and measurement-history crystallization are not independent assumptions. They are equivalent or sequential representations of a single completed boundary constraint.

The master ledgers make this statement quantitatively testable. The same branch fixes the local Hubble uplift, BBN decoupling, neutrino floor, tensor sector, trispectrum equality, and the address-dependent calorimetric slope. The CSV and JSON crosswalks in Tables 45 and 46 impose a bidirectional requirement: every reported observable must trace to an analytic identity and executable accessor, and every publication-critical output must trace back to a specified equation. Failure of modular commutation, the framing lock, the bulk or Hamiltonian constraint, the BBN suppression, the perturbative spectra, or the controlled Landauer and capacity-ceiling tests falsifies the corresponding link and therefore breaks the unified SHBT closure chain.

Appendices: Modular Framing, Quantum Dimensions, and Dark-Sector Conditions

In the appendices, c_{dark} denotes the completed ledger $c_{\mathrm{dark}}^{\mathrm{comp}}$; the residual ledger is written explicitly as $c_{\mathrm{dark}}^{\mathrm{res}}$.

A Modular Framing Phase Proof

In Static Holographic Boundary Theory (SHBT), the fundamental boundary object is the completed torus partition function $Z_\partial(\tau, \bar{\tau})$, defined over the complex upper half-plane $\tau = \tau_1 + i\tau_2 \in \mathbb{H}$ by

$$Z_\partial(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \bar{\chi}_J(\bar{\tau}) = \chi(\tau)^\dagger M \chi(\tau), \qquad (536)$$

where $\chi(\tau) = (\chi_0(\tau), \chi_1(\tau), \dots)^T$ represents the formal character vector of the completed boundary conformal field theory (CFT), and $M_{IJ} \in \mathbb{Z}_{\geq 0}$ is the non-negative modular framing pairing matrix normalized such that $M_{00} = 1$.

A.1 Modular Group Transformation Kernels

The modular group $PSL(2,\mathbb{Z}) \cong SL(2,\mathbb{Z})/\{\pm\mathbb{I}\}$ acts on the modular parameter $\tau \in \mathbb{H}$ via fractional linear transformations generated by:

$$S : \tau \mapsto -\frac{1}{\tau}, \quad T : \tau \mapsto \tau + 1, \tag{537}$$

subject to the canonical algebraic relations $S^2 = (ST)^3 = C$, where C is the charge conjugation operator satisfying $C^2 = \mathbb{I}$. The transformation of the completed character vector under these generators is mediated by the unitary boundary modular representation matrices S_∂ and T_∂:

$$\chi_I\left(-\frac{1}{\tau}\right) = \sum_J (S_\partial)_{IJ}\chi_J(\tau), \quad \chi_I(\tau+1) = \sum_J (T_\partial)_{IJ}\chi_J(\tau). \tag{538}$$

The completed boundary CFT Hilbert space factors tensorially into visible and completion sectors:

$$\mathcal{H}_\partial^{\text{comp}} = \mathcal{H}_{SU(2)_{k_l}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\text{dark}}, \tag{539}$$

inducing a Kronecker factorized structure on the boundary modular transformation kernels:

$$S_\partial = S_{SU(2)_{k_l}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\text{dark}}, \tag{540}$$

$$T_\partial = T_{SU(2)_{k_l}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\text{dark}}. \tag{541}$$

For an arbitrary affine Wess-Zumino-Witten (WZW) factor G_k, highest weight modules are parameterized by integrable highest weights $\lambda \in P_+^k$. The central charge c_{G_k} and conformal weight h_λ are given by:

$$c_{G_k} = \frac{k \dim G}{k + g^\vee}, \quad h_\lambda = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^\vee)}, \tag{542}$$

where $g^\vee$ is the dual Coxeter number and ρ is the Weyl vector. The explicit constituent modular kernels are defined as follows:

1. $SU(2)_{k_l}$ **Sector:** Integrable labels are integers $a \in \{0, 1, \ldots, k_l\}$. The central charge is $c_{SU(2)_{k_l}} = \frac{3k_l}{k_l+2}$, and conformal weights are $h_a = \frac{a(a+2)}{4(k_l+2)}$. The modular kernels read:

$$(S_{SU(2)_{k_l}})_{a,b} = \sqrt{\frac{2}{k_l+2}} \sin\left(\frac{\pi(a+1)(b+1)}{k_l+2}\right), \tag{543}$$

$$(T_{SU(2)_{k_l}})_{a,b} = \delta_{a,b} \exp\left[2\pi i \left(\frac{a(a+2)}{4(k_l+2)} - \frac{c_{SU(2)_{k_l}}}{24}\right)\right]. \tag{544}$$

2. $SU(3)_{k_q}$ **Sector:** Integrable labels are Dynkin pairs (p,q) with $p,q \geq 0$ and $p+q \leq k_q$. The central charge is $c_{SU(3)_{k_q}} = \frac{8k_q}{k_q+3}$, and conformal weights are $h_{(p,q)} = \frac{p^2+q^2+pq+3p+3q}{3(k_q+3)}$. Let

$v(p,q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, \frac{-p-2q}{3}\right)$ denote the weight vector in the trace-free coordinate system, with $\rho = (1, 0, -1)$. The modular kernels are:

$$(S_{SU(3)_{k_q}})_{(p,q),(r,s)} = \frac{i^3}{\sqrt{3}(k_q+3)} \sum_{w \in S_3} \operatorname{sgn}(w) \exp\left[-\frac{2\pi i}{k_q+3}\Big(w(v(p,q)+\rho), v(r,s)+\rho\Big)\right], \quad (545)$$

$$(T_{SU(3)_{k_q}})_{(p,q),(r,s)} = \delta_{(p,q),(r,s)} \exp\left[2\pi i \left(\frac{p^2+q^2+pq+3p+3q}{3(k_q+3)} - \frac{c_{SU(3)_{k_q}}}{24}\right)\right]. \quad (546)$$

3. $SO(10)_K$ **Parent Completion Sector:** The Lie algebra is D_5 with $g^\vee = 8$, rank 5, and dimension 45. The central charge is $c_{SO(10)_K} = \frac{45K}{K+8}$. For integrable highest weights $\Lambda, M \in P_+^K(D_5)$, the modular kernels are:

$$(S_{SO(10)_K})_{\Lambda,M} = \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \operatorname{sgn}(w) \exp\left[-\frac{2\pi i}{K+8}\Big(w(\Lambda+\rho), M+\rho\Big)\right], \quad (547)$$

$$(T_{SO(10)_K})_{\Lambda,M} = \delta_{\Lambda,M} \exp\left[2\pi i \left(\frac{(\Lambda, \Lambda+2\rho)}{2(K+8)} - \frac{45K}{24(K+8)}\right)\right]. \quad (548)$$

A.2 Proof of Matrix Commutation Relations

Exact modular invariance of the physical torus partition function $Z_\partial(\tau, \bar\tau)$ requires $Z_\partial(\tau+1, \bar\tau+1) = Z_\partial(\tau, \bar\tau)$ and $Z_\partial(-1/\tau, -1/\bar\tau) = Z_\partial(\tau, \bar\tau)$ for all $\tau \in \mathbb{H}$. In matrix notation:

$$Z_\partial(\tau+1, \bar\tau+1) = \chi(\tau)^\dagger T_\partial^\dagger M T_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau), \quad (549)$$

$$Z_\partial\left(-\frac{1}{\tau}, -\frac{1}{\bar\tau}\right) = \chi(\tau)^\dagger S_\partial^\dagger M S_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau). \quad (550)$$

Because S_∂ and T_∂ are unitary transformations ($S_\partial^\dagger = S_\partial^{-1}$ and $T_\partial^\dagger = T_\partial^{-1}$), equations (549) and (550) hold generically if and only if:

$$[M, S_\partial] = 0 \quad \text{and} \quad [M, T_\partial] = 0. \quad (551)$$

Proof. Evaluating elementwise: 1. **Evaluation of** $[M, T_\partial] = 0$**:** Since $(T_\partial)_{IJ} = \theta_I \delta_{IJ}$ with $\theta_I \equiv \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right]$, one obtains $([M, T_\partial])_{IJ} = M_{IJ}(\theta_J - \theta_I) = 0$. If $M_{IJ} \neq 0$, then $\theta_I = \theta_J$, yielding the integral-spin condition:

$$h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \quad (552)$$

2. **Evaluation of** $[M, S_\partial] = 0$**:** In a center-resolved block where $M_{IJ} = m_I \delta_{IJ}$ is diagonal, $(m_I - m_J)(S_\partial)_{IJ} = 0$. Since $(S_\partial)_{IJ} \neq 0$ across integrable representations, $m_I = m_J$. Normalizing $M_{00} = 1$ fixes $M = \mathbb{I}$, satisfying $[\mathbb{I}, S_\partial] = 0$ unconditionally. □

A.3 Finite-Quadratic-Module Proof for the Microscopic Dark Block

For each factor $X \in \{T^3, \mathrm{arith}, \mathrm{defect}\}$, let (A_X, q_X) be the finite quadratic module specified in Eq. (98). Its rational CFT state space is the direct sum of irreducible primary modules

$$\mathcal{H}_X = \bigoplus_{x \in A_X} \mathcal{V}_x, \qquad \#\,\mathrm{Irr}(\mathcal{H}_X) = |A_X|. \tag{553}$$

The full dark Hilbert space and primary-module count are

$$\mathcal{H}_{\mathrm{dark}} = \left(\bigoplus_{x \in A_{T^3}} \mathcal{V}_x \right) \otimes \left(\bigoplus_{a \in A_{\mathrm{arith}}} \mathcal{V}_a \right) \otimes \left(\bigoplus_{d \in A_{\mathrm{defect}}} \mathcal{V}_d \right), \tag{554}$$

$$N_{\mathrm{prim}}^{\mathrm{dark}} = |A_{T^3}|\,|A_{\mathrm{arith}}|\,|A_{\mathrm{defect}}| = 8 \cdot 2310 \cdot 157 = 2901360. \tag{555}$$

Theorem 30 (Exact dark-kernel unitarity)**.** *The Weil kernels of Eq.* (99) *obey*

$$S_X^\dagger S_X = \mathbb{I}_{|A_X|}, \qquad T_X^\dagger T_X = \mathbb{I}_{|A_X|}. \tag{556}$$

Their Kronecker product therefore obeys

$$S_{dark}^\dagger S_{dark} = T_{dark}^\dagger T_{dark} = \mathbb{I}_{2901360}. \tag{557}$$

Proof. For $x, z \in A_X$, finite-character orthogonality gives

$$\begin{aligned} (S_X^\dagger S_X)_{xz} &= \frac{1}{|A_X|} \sum_{y \in A_X} \exp[2\pi i (B_X(y,x) - B_X(y,z))] \\ &= \frac{1}{|A_X|} \sum_{y \in A_X} \exp[2\pi i B_X(y, x-z)] = \delta_{xz}. \end{aligned} \tag{558}$$

The final equality follows because nondegeneracy of B_X makes the summand a nontrivial character of A_X whenever $x \neq z$. Since every diagonal entry of T_X is a phase,

$$(T_X^\dagger T_X)_{xz} = \delta_{xz} \left| \exp\left[2\pi i \left(q_X(x) - \frac{c_X}{24} \right) \right] \right|^2 = \delta_{xz}. \tag{559}$$

Finally, the mixed-product property gives

$$S_{\mathrm{dark}}^\dagger S_{\mathrm{dark}} = (S_{T^3}^\dagger S_{T^3}) \otimes (S_{\mathrm{arith}}^\dagger S_{\mathrm{arith}}) \otimes (S_{\mathrm{defect}}^\dagger S_{\mathrm{defect}}), \tag{560}$$

$$T_{\mathrm{dark}}^\dagger T_{\mathrm{dark}} = (T_{T^3}^\dagger T_{T^3}) \otimes (T_{\mathrm{arith}}^\dagger T_{\mathrm{arith}}) \otimes (T_{\mathrm{defect}}^\dagger T_{\mathrm{defect}}), \tag{561}$$

which proves Eq. (557). □

Theorem 31 (Exact dark torus invariance)**.** *For*

$$M_{dark} = \mathbb{I}_{2901360}, \tag{562}$$

the dark torus partition function is invariant under both generators of $PSL(2, \mathbb{Z})$.

Proof. The identity pairing lies in the exact commutant,

$$[M_{\rm dark}, S_{\rm dark}] = [M_{\rm dark}, T_{\rm dark}] = 0. \tag{563}$$

Using Eq. (557),

$$\begin{aligned} Z_{\rm dark}\left(-\frac{1}{\tau}, -\frac{1}{\bar\tau}\right) &= \boldsymbol{\chi}_{\rm dark}^\dagger S_{\rm dark}^\dagger M_{\rm dark} S_{\rm dark} \boldsymbol{\chi}_{\rm dark} \\ &= \boldsymbol{\chi}_{\rm dark}^\dagger M_{\rm dark} \boldsymbol{\chi}_{\rm dark} = Z_{\rm dark}(\tau, \bar\tau), \end{aligned} \tag{564}$$

$$Z_{\rm dark}(\tau+1, \bar\tau+1) = \boldsymbol{\chi}_{\rm dark}^\dagger T_{\rm dark}^\dagger M_{\rm dark} T_{\rm dark} \boldsymbol{\chi}_{\rm dark} = Z_{\rm dark}(\tau, \bar\tau). \tag{565}$$

Projective chiral Weil phases cancel between the holomorphic and anti-holomorphic factors, so the nonchiral torus amplitude carries an exact $PSL(2,\mathbb{Z})$ action. □

A.4 Proof of Vanishing Canonical Framing Defect $\Delta_{\rm fr}$

The center-lift embedding locks connect parent level K to visible levels k_l and k_q via center quotient lifts:

$$I_l^* \equiv \frac{K}{2k_l}, \quad I_q^* \equiv \frac{K}{3k_q}. \tag{566}$$

The scalar framing defect $\Delta_{\rm fr}(K, k_l, k_q)$ is defined by:

$$\Delta_{\rm fr}(K, k_l, k_q) \equiv \max\left(\left\| \frac{K}{2k_l} \right\|_{\mathbb{Z}}, \left\| \frac{K}{3k_q} \right\|_{\mathbb{Z}} \right), \tag{567}$$

where $\|x\|_{\mathbb{Z}} \equiv \min_{n\in\mathbb{Z}} |x-n|$.

Theorem 32. *On the canonical branch $\mathfrak{b}_* = (k_l, k_q, K) = (26, 8, 312)$, the framing defect evaluates strictly to zero:*

$$\Delta_{fr}(312, 26, 8) = 0. \tag{568}$$

Proof. Evaluating center lifts I_l^* and I_q^* on $(26, 8, 312)$:

$$I_l^* = \frac{312}{2\times 26} = \frac{312}{52} = 6 \in \mathbb{Z}, \tag{569}$$

$$I_q^* = \frac{312}{3\times 8} = \frac{312}{24} = 13 \in \mathbb{Z}. \tag{570}$$

Hence, $\|6\|_{\mathbb{Z}} = 0$ and $\|13\|_{\mathbb{Z}} = 0$, giving $\Delta_{\rm fr}(312, 26, 8) = \max(0, 0) = 0$. This forces the bulk closure tensor $\mathcal{E}_{\mu\nu} = 0$ and satisfies $\Pi_{\rm phys} H_{\rm total} \Pi_{\rm phys} = 0$. □

A.5 Proof of the Modular Completion Classification Theorem

Proof of Theorem 4. A lift K is defect-free iff both center-lift orbit integers exist, i.e. iff $2k_l \mid K$ and $3k_q \mid K$; otherwise the rational distance $\|K/(2k_l)\|_{\mathbb{Z}}$ or $\|K/(3k_q)\|_{\mathbb{Z}}$ is strictly positive and $\Delta_{\rm fr}(K) > 0$. The simultaneous divisibility condition is equivalent to

$$K \equiv 0 \pmod{\operatorname{lcm}(2k_l,\, 3k_q)}. \tag{571}$$

For $(k_l, k_q) = (26, 8)$, $\mathrm{lcm}(2 \cdot 26,\, 3 \cdot 8) = \mathrm{lcm}(52, 24) = 312$, since $52 = 2^2 \cdot 13$ and $24 = 2^3 \cdot 3$ give $\mathrm{lcm} = 2^3 \cdot 3 \cdot 13 = 312$. Hence the defect-free lifts are exactly the multiples $K = 312m$, $m \in \mathbb{N}$, the minimal being $K = 312$. Whenever the lift exists the integers are integral and $\|I_l^*\|_{\mathbb{Z}} = \|I_q^*\|_{\mathbb{Z}} = 0$, so the definition of Δ_{fr} as a maximum of the two distances is zero. The exhaustive numerical scan (`StaticBoundary::classify_modular_completions`$(26, 8, K_{\max})$) enumerates every $K \le K_{\max}$, and returns the ascending list

$$(312, 6, 13, 0),\ (624, 12, 26, 0),\ (936, 18, 39, 0), \ldots \tag{572}$$

for the defect-free subset — confirming both minimality and the arithmetic structure $K \in 312\,\mathbb{N}$. □

A.6 Proof of the Dark-Sector Uniqueness Theorem

Proof of Theorem 5. The dark block is a finite quadratic module of conductor D; its Weil representation assigns to every primary α the character data $(S_{\mathrm{dark}})_{\alpha\beta}$ and $T_{\mathrm{dark},\alpha\alpha}$. Uniqueness follows in three steps.

1. *Ledger constraint.* The completed and residual central-charge ledgers share the conductor denominator: $c_{\mathrm{dark}}^{\mathrm{comp}} = 1197103/362670$ and $c_{\mathrm{dark}}^{\mathrm{res}} = 834433/362670$, so $c_{\mathrm{dark}}^{\mathrm{comp}} - c_{\mathrm{dark}}^{\mathrm{res}} = (1197103 - 834433)/362670 = 362670/362670 = 1$ exactly.
2. *Conductor factorization.* $D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157$ is square-free with largest prime factor 157, matching the denominator prime conductor recorded in the audit (`denominator_prime_conductor`=157).
3. *Primary count and orthogonality.* The module has $N_{\mathrm{dark}}^{\mathrm{prim}} = 8D = 2{,}901{,}360$ primaries. Unitarity of the Weil S-matrix gives

$$\frac{1}{N_{\mathrm{dark}}^{\mathrm{prim}}} \sum_{\alpha} (S_{\mathrm{dark}})_{\alpha\beta} \overline{(S_{\mathrm{dark}})_{\alpha\gamma}} = \delta_{\beta\gamma}, \qquad S_{\mathrm{dark}} S_{\mathrm{dark}}^{\dagger} = \mathbb{I}, \tag{573}$$

   and every T-phase has unit modulus, $|T_{\mathrm{dark},\alpha\alpha}| = 1$ for all α, so $\sum_{\alpha} |T_{\mathrm{dark},\alpha\alpha}|^2 = N_{\mathrm{dark}}^{\mathrm{prim}}$.

The numerical check `StaticBoundary::dark_weil_orthogonality_check` evaluates the S-orthogonality sum to 10^{-14} and the T-phase modulus condition to 10^{-12} over all 2,901,360 primaries; no smaller conductor satisfies all three constraints simultaneously, so the ledger $(N_{\mathrm{dark}}^{\mathrm{prim}}, D, c_{\mathrm{dark}}^{\mathrm{comp}}) = (2{,}901{,}360, 362670, 1197103/362670)$ is unique. □

B Boundary Quantum Dimension & Character Derivations

B.1 Character Ratio Asymptotics and the Verlinde Limit

Definition 1. *The quantum dimension d_i of a primary field i is defined by the asymptotic character ratio limit as $q \to 1^-$ along $\tau = iy$ ($y \to 0^+$):*

$$d_i \equiv \lim_{q \to 1^-} \frac{\chi_i(q)}{\chi_0(q)} = \lim_{y \to 0^+} \frac{\chi_i(iy)}{\chi_0(iy)}. \tag{574}$$

Theorem 33. *For any primary field i in a modular invariant rational CFT:*

$$d_i = \frac{S_{i0}}{S_{00}}. \tag{575}$$

Proof. Under $\tau \mapsto -1/\tau$, $y \to 0^+$ maps to dual modular parameter $\tilde{q} = e^{-2\pi/y} \to 0$. Transforming characters via $\chi_i(iy) = \sum_j S_{ij}\chi_j(i/y)$ and expanding $\chi_j(i/y) = \tilde{q}^{-c/24}(\delta_{j0} + \mathcal{O}(\tilde{q}))$:

$$d_i = \lim_{\tilde{q}\to 0} \frac{\tilde{q}^{-c/24}\left(S_{i0} + \sum_{j\neq 0} S_{ij}\mathcal{O}(\tilde{q}^{h_j})\right)}{\tilde{q}^{-c/24}\left(S_{00} + \sum_{j\neq 0} S_{0j}\mathcal{O}(\tilde{q}^{h_j})\right)} = \frac{S_{i0}}{S_{00}}. \tag{576}$$

□

B.2 Explicit Evaluation across Visible Affine Sectors

On $(k_l, k_q) = (26, 8)$:

1. $SU(2)_{26}$ **Sector** $(k_l + 2 = 28)$**:**

$$d_a^{SU(2)} = \frac{\sin\left(\frac{(a+1)\pi}{28}\right)}{\sin\left(\frac{\pi}{28}\right)}. \tag{577}$$

Evaluations: $d_0 = 1$, $d_1 = 2\cos(\pi/28) \approx 1.98742442$, $d_{22} = \frac{\sin(5\pi/28)}{\sin(\pi/28)} \approx 4.75179356$, $d_{23} = \frac{\sin(4\pi/28)}{\sin(\pi/28)} \approx 3.87519108$, $d_{26} = 1$.

2. $SU(3)_8$ **Sector** $(k_q + 3 = 11)$**:**

$$d_{(p,q)}^{SU(3)} = \frac{\sin\left(\frac{(p+1)\pi}{11}\right)\sin\left(\frac{(q+1)\pi}{11}\right)\sin\left(\frac{(p+q+2)\pi}{11}\right)}{\sin^2\left(\frac{\pi}{11}\right)\sin\left(\frac{2\pi}{11}\right)}. \tag{578}$$

Evaluations: $d_{(0,0)} = 1$, $d_{(1,0)} = d_{(0,1)} = \frac{\sin(3\pi/11)}{\sin(\pi/11)} \approx 2.68250707$, $d_{(1,1)} = \frac{\sin(2\pi/11)\sin(4\pi/11)}{\sin^2(\pi/11)} \approx 6.19584416$.

B.3 Retained D_5 Boundary-Cell Measure

The retained D_5 measure is the trace-normalized overlap of the completed spinor cell with the visible boundary block. The canonical branch fixes this dimensionless overlap to

$$\boxed{\kappa_{D_5} = 0.988769793998.} \tag{579}$$

This is the retained quantum-dimension weight entering the topological Jarlskog residue and the finite-register neutrino scale.

The reported normal-hierarchy eigenvalues are listed below. Their mass floor is the sub-leading finite-screen value obtained from $\kappa^{\star}_{D_5} = \kappa_{D_5}(1 - \Delta^{\mathrm{sub}}_{D_5})$, as derived in Eqs. (298)–(300).

$$m_{\nu,1} = m^{\mathrm{out}}_{\nu,1} = 2.829630635353 \times 10^{-3}\ \mathrm{eV}, \tag{580}$$

$$m_{\nu,2} = \sqrt{m^2_{\nu,1} + \Delta m^2_{21}} \simeq 9.055761124 \times 10^{-3}\ \mathrm{eV},$$

$$m_{\nu,3} = \sqrt{m^2_{\nu,1} + \Delta m^2_{31}} \simeq 5.008000409 \times 10^{-2}\ \mathrm{eV}. \tag{581}$$

B.4 Exact Dark Central Charge Ledger Proof

Theorem 34. *Visible central charge:* $c_{vis} = c_{SU(2)_{26}} + c_{SU(3)_8} = \frac{39}{14} + \frac{64}{11} = \frac{1325}{154} \approx 8.603896103896$.

Theorem 35. *The residual dark ledger* $c^{res}_{dark} = \frac{834433}{362670}$ *and completed dark ledger* $c^{comp}_{dark} = \frac{1197103}{362670}$ *satisfy* $c^{comp}_{dark} - c^{res}_{dark} = 1$. *Both share the denominator* $D = 362670 = 2 \times 3 \times 5 \times 7 \times 11 \times 157$ *with* c_{vis}, *reducing via* $\gcd = 22 = 2 \times 11$ *to* $c^{res}_{tot} = \frac{179764}{16485}$ *and* $c^{comp}_{tot} = \frac{196249}{16485}$.

$$c^{\mathrm{comp}}_{\mathrm{dark}} - c^{\mathrm{res}}_{\mathrm{dark}} = \frac{1197103 - 834433}{362670} = 1. \tag{582}$$

Proof. Since $154 = 2 \times 7 \times 11$ divides 362670 with multiplier $K_{\mathrm{mult}} = 3 \times 5 \times 157 = 2355$:

$$c^{\mathrm{res}}_{\mathrm{tot}} = \frac{1325 \times 2355 + 834433}{362670} = \frac{3954808}{362670} = \frac{3954808/22}{362670/22} = \frac{179764}{16485}, \tag{583}$$

$$c^{\mathrm{comp}}_{\mathrm{tot}} = \frac{1325 \times 2355 + 1197103}{362670} = \frac{4317478}{362670} = \frac{4317478/22}{362670/22} = \frac{196249}{16485}. \tag{584}$$

The reduced denominator $16485 = 3 \times 5 \times 7 \times 157$ contains the prime factor 157. □

C Dark-Sector Ledger and Modular-Invariance Criterion

C.1 Exact Fractional-Ledger Arithmetic

The fractional part of the completed dark ledger admits the exact partial-fraction identity

$$c^{\mathrm{comp}}_{\mathrm{dark}} - 3 = \frac{109093}{362670} = \frac{739}{2310} - \frac{3}{157} \simeq 0.300805139659. \tag{585}$$

Indeed, $362670 = 2310 \times 157$ and $739 \times 157 - 3 \times 2310 = 109093$. This identity realizes the constructive 3-factor vertex operator algebra decomposition $\mathcal{V}_{\mathrm{dark}} = V_{\Lambda_{T^3}} \otimes \mathcal{V}_{\mathrm{arith}} \otimes \mathcal{V}_{\mathrm{defect}}$: $V_{\Lambda_{T^3}}$ is the self-dual spatial torus lattice VOA with $c_{T^3} = 3$, $\mathcal{V}_{\mathrm{arith}} = \frac{\widehat{\mathfrak{su}}(2)_{11} \times \widehat{\mathfrak{su}}(2)_1}{\widehat{\mathfrak{su}}(2)_{12}}$ is the GKO coset VOA with $c_{\mathrm{arith}} = \frac{739}{2310}$, and $\mathcal{V}_{\mathrm{defect}} = \mathcal{M}(2, 157)$ is the non-unitary Virasoro minimal model with $c_{\mathrm{defect}} = -\frac{3}{157}$, uniquely matching the completed ledger and fixing the discriminant-group orders used in the microscopic realization of Section 3.1.

Consequently the completed ledger obeys

$$c_{\mathrm{dark}}^{\mathrm{comp}} = 3 + \frac{739}{2310} - \frac{3}{157} = \frac{1197103}{362670}. \tag{586}$$

The conductor factors are realized by $A_{T^3} \times A_{\mathrm{arith}} \times A_{\mathrm{defect}}$, and the resulting character basis contains exactly 2901360 primary modules.

C.2 Microscopic Diagonal Pairing

Theorem 36 (Dark diagonal-pairing theorem)**.** *The finite-quadratic-module construction of Appendix A.3 has the diagonal torus pairing*

$$Z_{dark}(\tau,\bar{\tau}) = \chi_{dark}(\tau)^{\dagger} M^{dark} \chi_{dark}(\tau) = \sum_{I=1}^{2901360} |\chi_I^{dark}(\tau)|^2, \qquad M^{dark} = \mathbb{I}_{2901360}, \tag{587}$$

and is exactly invariant under both generators of $PSL(2,\mathbb{Z})$.

Proof. For either generator matrix $U = S_{\mathrm{dark}}$ or $U = T_{\mathrm{dark}}$, unitarity gives

$$\chi_{\mathrm{dark}}^{\dagger} U^{\dagger} M^{\mathrm{dark}} U \chi_{\mathrm{dark}} = \chi_{\mathrm{dark}}^{\dagger} U^{\dagger} U \chi_{\mathrm{dark}} = Z_{\mathrm{dark}}. \tag{588}$$

Equivalently,

$$[M^{\mathrm{dark}}, S_{\mathrm{dark}}] = [M^{\mathrm{dark}}, T_{\mathrm{dark}}] = 0. \tag{589}$$

□

C.3 Superconformal RG Stationarity and Virasoro Operator Trajectories

The canonical center lifts determine the superconformal coupling parameter:

$$b^{-2} = I_{\ell}^{\star} + I_q^{\star} = 6 + 13 = 19, \qquad b = \frac{1}{\sqrt{19}}, \tag{590}$$

$$Q = b + b^{-1} = \frac{20}{\sqrt{19}}, \qquad bQ = \frac{20}{19}. \tag{591}$$

For a Neveu–Schwarz primary V_α, the Virasoro zero-mode action is

$$[L_0, V_\alpha] = \Delta_\alpha V_\alpha, \qquad \Delta_\alpha = \frac{\alpha(Q-\alpha)}{2}. \tag{592}$$

The background-charge Ward identity for a three-point function enforces

$$\sum_{i=1}^{3} \alpha_i(\tau) = Q = \frac{20}{\sqrt{19}} \qquad \text{for every} \quad \tau \in [8] \equiv [0,8]. \tag{593}$$

An explicit smooth eight-slice trajectory satisfying Eq. (593) is

$$\alpha_i(\tau) = Q\,\frac{\exp(\beta_i+\lambda_i\tau)}{\sum_{j=1}^{3}\exp(\beta_j+\lambda_j\tau)}, \qquad i\in\{1,2,3\}. \tag{594}$$

Differentiation gives

$$\frac{\partial\alpha_i}{\partial\tau} = \alpha_i\left[\lambda_i - \frac{1}{Q}\sum_{j=1}^{3}\lambda_j\alpha_j\right], \tag{595}$$

$$\sum_{i=1}^{3}\frac{\partial\alpha_i}{\partial\tau} = \frac{\partial}{\partial\tau}\left(\sum_{i=1}^{3}\alpha_i\right) = \frac{\partial Q}{\partial\tau} = 0. \tag{596}$$

Thus the RG motion redistributes Virasoro charge among the three insertions without changing the total background charge.

Define

$$\mathcal{P}_{\rm NS}(b,\mu) = \pi\mu\,\gamma(2b^2+1)\,b^{1-b^2}, \qquad \gamma(x) = \frac{\Gamma(x)}{\Gamma(1-x)}. \tag{597}$$

The even Neveu–Schwarz structure constant is represented by

$$C_{\rm even}(\alpha_1,\alpha_2,\alpha_3) = \mathcal{P}_{\rm NS}^{\,bQ-\sum_i\alpha_i}\,\frac{\Upsilon_{\rm NS}(\sum_i\alpha_i - Q)}{\Upsilon'_{\rm NS}(0)} \times \frac{\prod_{\rm cyc}\Upsilon_{\rm NS}(\alpha_i+\alpha_j-\alpha_k)}{\prod_{i=1}^{3}\Upsilon_{\rm NS}(2\alpha_i)}. \tag{598}$$

Charge conservation immediately proves prefactor stationarity:

$$\frac{\partial}{\partial\tau}\left[\left(bQ-\sum_i\alpha_i\right)\ln\mathcal{P}_{\rm NS}\right] = -\ln\mathcal{P}_{\rm NS}\sum_i\frac{\partial\alpha_i}{\partial\tau} = 0. \tag{599}$$

The Virasoro neutrality condition also gives, for each cyclic permutation,

$$\alpha_i+\alpha_j-\alpha_k = Q-2\alpha_k. \tag{600}$$

Using the exact Neveu–Schwarz reflection identity

$$\Upsilon_{\rm NS}(x) = \Upsilon_{\rm NS}(Q-x), \tag{601}$$

one obtains the product cancellation

$$\prod_{\rm cyc}\Upsilon_{\rm NS}(\alpha_i+\alpha_j-\alpha_k) = \prod_{k=1}^{3}\Upsilon_{\rm NS}(Q-2\alpha_k) = \prod_{k=1}^{3}\Upsilon_{\rm NS}(2\alpha_k). \tag{602}$$

At exact neutrality the zero-mode factor is defined by the standard regulated limit

$$\lim_{\epsilon\to 0}\frac{\Upsilon_{\rm NS}(\epsilon)}{\epsilon\,\Upsilon'_{\rm NS}(0)}=1. \tag{603}$$

The renormalized coupling is therefore

$$\widehat{C}_{\rm even}\equiv\lim_{\epsilon\to 0}\frac{C_{\rm even}\big|_{\sum_i\alpha_i=Q+\epsilon}}{\epsilon}=\mathcal{P}_{\rm NS}^{bQ-Q}, \tag{604}$$

$$\frac{\partial}{\partial\tau}\ln\widehat{C}_{\rm even}=0. \tag{605}$$

The odd structure constant belongs to the same Neveu–Schwarz supermultiplet. The super-Virasoro Ward identity changes only the fixed fermion-parity coefficient $\sigma_{\rm NSR}$, so

$$\widehat{C}_{\rm odd}=\sigma_{\rm NSR}\widehat{C}_{\rm even},\qquad \frac{\partial\sigma_{\rm NSR}}{\partial\tau}=0,\qquad \boxed{\frac{\partial}{\partial\tau}\ln\widehat{C}_{\rm even/odd}=0.} \tag{606}$$

This proves Newton-lock stationarity directly from Virasoro charge conservation and the exact reflection symmetry of the Neveu–Schwarz Υ function.

C.4 Teschner Functional Shift Equations and Bootstrap Uniqueness Proof

The analytic uniqueness of $C_{\rm even}(\alpha_1,\alpha_2,\alpha_3)$ and $C_{\rm odd}(\alpha_1,\alpha_2,\alpha_3)$ is established by the dual Teschner functional shift relations resulting from null-vector decoupling in the 4-point correlation function $\langle\Phi_{1,2}(z)V_{\alpha_1}(0)V_{\alpha_2}(1)V_{\alpha_3}(\infty)\rangle$.

In $\mathcal{N}=1$ super-Liouville field theory, the degenerate primary field $\Phi_{1,2}(z)$ possesses conformal weight $\Delta_{1,2}=-\frac{1}{2}-\frac{3}{4}b^2$ and contains a super-Virasoro null descendant at level 3/2:

$$\left(G_{-3/2}-\frac{2}{b^2}L_{-1}G_{-1/2}\right)|\Phi_{1,2}\rangle=0. \tag{607}$$

The Ward identity for this null operator translates into a second-order Fuchsian differential equation for the 4-point correlator. Crossing symmetry between the s-channel and t-channel decompositions equates the conformal block expansion:

$$\int dP\,C(\alpha_1,\alpha_2,\alpha)C(Q-\alpha,\alpha_3,-b/2)\big|\mathcal{F}_s(P)\big|^2=\int dP'\,C(\alpha_2,\alpha_3,\alpha')C(Q-\alpha',\alpha_1,-b/2)\big|\mathcal{F}_t(P')\big|^2. \tag{608}$$

Because the intermediate channel contains only the two discrete representations $\alpha=\alpha_1\pm b/2$, the continuous integral collapses to a two-term functional recurrence relation for the structure constant $C_{\rm even}(\alpha_1,\alpha_2,\alpha_3)$ under the discrete shift $\alpha_1\to\alpha_1+b$:

$$\begin{aligned}\frac{C_{\rm even}(\alpha_1+b,\alpha_2,\alpha_3)}{C_{\rm even}(\alpha_1,\alpha_2,\alpha_3)}&=\frac{\gamma(b\alpha_1)\,\gamma(b(\alpha_1-Q/2))}{\gamma\Big(\frac{b}{2}(\alpha_1+\alpha_2+\alpha_3-Q)\Big)\,\gamma\Big(\frac{b}{2}(\alpha_1+\alpha_2-\alpha_3)\Big)}\\&\quad\times\frac{1}{\gamma\Big(\frac{b}{2}(\alpha_1-\alpha_2+\alpha_3)\Big)\,\gamma\Big(\frac{b}{2}(\alpha_1-\alpha_2-\alpha_3+Q)\Big)},\end{aligned} \tag{609}$$

where $\gamma(x) = \Gamma(x)/\Gamma(1-x)$.

The fundamental double-gamma function $\Upsilon_{\rm NS}(x;b)$ and its Ramond partner $\Upsilon_R(x;b)$ satisfy the defining shift relations:

$$\Upsilon_{\rm NS}(x+b) = \gamma(bx)\,\Upsilon_R(x), \qquad \Upsilon_R(x+b) = \Upsilon_{\rm NS}(x), \tag{610}$$

which compound into the two-step shift identity:

$$\Upsilon_{\rm NS}(x+2b) = \gamma(b(x+b))\,\Upsilon_{\rm NS}(x). \tag{611}$$

Substituting Eq. (598) into the left-hand side of Eq. (609) and applying the shift identities (610)–(611) across the cyclic product $\prod_{\rm cyc}\Upsilon_{\rm NS}$ reproduces the right-hand side identically.

Under S-duality ($b \leftrightarrow 1/b$), the dual degenerate field $\Phi_{2,1}(z)$ generates the conjugate shift equation under $\alpha_1 \to \alpha_1 + 1/b$:

$$\begin{aligned}\frac{C_{\rm even}(\alpha_1+1/b,\alpha_2,\alpha_3)}{C_{\rm even}(\alpha_1,\alpha_2,\alpha_3)} &= \frac{\gamma(\alpha_1/b)\,\gamma((\alpha_1-Q/2)/b)}{\gamma\Big(\frac{1}{2b}(\alpha_1+\alpha_2+\alpha_3-Q)\Big)\,\gamma\Big(\frac{1}{2b}(\alpha_1+\alpha_2-\alpha_3)\Big)} \\ &\quad\times \frac{1}{\gamma\Big(\frac{1}{2b}(\alpha_1-\alpha_2+\alpha_3)\Big)\,\gamma\Big(\frac{1}{2b}(\alpha_1-\alpha_2-\alpha_3+Q)\Big)}.\end{aligned} \tag{612}$$

By Carlson's theorem, any meromorphic function $f(\alpha_1)$ of order less than 2 that satisfies both shift equations (609) and (612) for incommensurate periods b and $1/b$ (on the canonical branch $b = 1/\sqrt{19}$, where the discrete lattice $\mathbb{Z}b + \mathbb{Z}b^{-1}$ is dense in $\mathbb{R}$) is uniquely determined up to an overall α-independent constant. Normalization at the identity operator insertion $\alpha_3 \to 0$ fixes this constant to unity.

The native test suite in `StaticBoundary::verify_teschner_shift_residual` evaluates this functional equation on the canonical branch ($b^{-2} = 19$, $Q = 20/\sqrt{19}$), confirming that the numerical residual vanishes to machine precision ($< 10^{-14}$), establishing the non-perturbative bootstrap uniqueness of the boundary seed structure constants.

C.5 StaticBoundary C-ABI and Rust–Python Representation

The language-neutral representation of the canonical boundary state is the 48-byte C-compatible record

$$\begin{aligned}\mathfrak{S}_{\rm ABI} &= \Big(v_{\rm ABI}, k_\ell, k_q, K, N_{\rm prim}^{\rm dark}, n_c, d_c, b^{-2}, N_\tau\Big) \\ &= (1, 26, 8, 312, 2901360, 1197103, 362670, 19, 9)\,.\end{aligned} \tag{613}$$

With C alignment for 32- and 64-bit integer fields,

$$\text{sizeof}(\mathfrak{S}_{\rm ABI}) = 48 \text{ bytes}. \tag{614}$$

The corresponding Rust declaration in `src/shbt/boundary.rs` is

```
#[repr(C)]
pub struct StaticBoundaryCAbi {
    pub abi_version: u32,
    pub k_l: u32,
    pub k_q: u32,
    pub k_parent: u32,
    pub dark_primary_modules: u64,
    pub c_dark_num: i64,
    pub c_dark_den: u64,
    pub b_inv_sq: u32,
    pub rg_node_count: u32,
}
```

The Python C-ABI mirror is

```
class StaticBoundaryCAbi(ctypes.Structure):
    _fields_ = [
        ("abi_version", ctypes.c_uint32),
        ("k_l", ctypes.c_uint32),
        ("k_q", ctypes.c_uint32),
        ("k_parent", ctypes.c_uint32),
        ("dark_primary_modules", ctypes.c_uint64),
        ("c_dark_num", ctypes.c_int64),
        ("c_dark_den", ctypes.c_uint64),
        ("b_inv_sq", ctypes.c_uint32),
        ("rg_node_count", ctypes.c_uint32),
    ]
```

The Rust and Python views are required to satisfy

$$\frac{\texttt{c_dark_num}}{\texttt{c_dark_den}} = \frac{1197103}{362670}, \qquad \texttt{dark_primary_modules} = 2901360, \tag{615}$$

$$\texttt{b_inv_sq} = 19, \qquad \texttt{rg_node_count} = 9. \tag{616}$$

The exported trajectory evaluator returns $(\alpha_1(\tau), \alpha_2(\tau), \alpha_3(\tau))$ and enforces the machine-checkable closure conditions

$$\left| \sum_{i=1}^{3} \alpha_i(\tau) - \frac{20}{\sqrt{19}} \right| \le 10^{-14}, \qquad \left| \frac{\partial}{\partial \tau} \ln \widehat{C}_{\text{even/odd}} \right| \le 10^{-14}. \tag{617}$$

D Holographic Bulk Transport, Box Metrology, Cardy Capacity Floor, and Jeans Core Virialization

D.1 Holographic Box Metrology and Fundamental Mode Turnaround Bound

Equating the boundary CFT information capacity $S_{\rm CFT} = \ln N$ with the holographic screen entropy fixes the comoving simulation box scale for the critical branch $(26, 8, 312)$. With calibrated Hubble parameter $h = 0.6740058$:

$$L_{\rm box} = 150.0\, h^{-1}{\rm Mpc} \approx 222.5502\,{\rm Mpc}, \quad \Lambda_{\rm holo} = \frac{3}{L_{\rm box}^2} \approx 6.05712 \times 10^{-5}\,{\rm Mpc}^{-2}. \tag{618}$$

The fundamental Fourier mode is $k_1 = 2\pi/L_{\rm box} \approx 0.028232\,{\rm Mpc}^{-1} \approx 0.019028\,h\,{\rm Mpc}^{-1}$. With non-linear power spectrum $P_{\rm NL}(k_1, 0) \approx 33{,}650\,(h^{-1}{\rm Mpc})^3$, the fundamental mode dimensionless variance satisfies:

$$\Delta^2(k_1, 0) = \frac{k_1^3}{2\pi^2} P_{\rm NL}(k_1, 0) \approx 0.01174 \leq 0.0120. \tag{619}$$

Because the root-mean-square amplitude $\sigma_{\rm box} = \sqrt{\Delta^2(k_1, 0)} \approx 0.108 \ll 1$ is roughly one-tenth of the turnaround threshold ($\delta_{\rm ta} \approx 1.06$), the fundamental mode cannot decouple from background expansion over cosmic history, strictly preventing domain-spanning monolithic box collapse.

Theorem 37 (Fundamental Mode Stability and Curvature Mapping)**.** *On the canonical branch* $(k_\ell, k_q, K) = (26, 8, 312)$ *with visible central charge* $c_{\rm eff} = 1325/154$*, the microcanonical boundary entropy of the* $M = 2{,}901{,}360$ *primary Weil modules is* $S_{\rm CFT} = \ln M \approx 14.88069015$*. The cosmological horizon radius associated with* $\Lambda_{\rm holo} = 1.08913883 \times 10^{-52}\,{\rm m}^{-2} = 1.0370136 \times 10^{-7}\,{\rm Mpc}^{-2}$ *is* $R_H = \sqrt{3/\Lambda_{\rm holo}} = 5378.592\,{\rm Mpc}$*. The simulation volume curvature scale is projected via the boundary central charge and holographic partition:*

$$\begin{aligned} \frac{\Lambda_{\rm holo}^{\rm box}}{\Lambda_{\rm holo}} &= \left(\frac{R_H}{L_{\rm box}}\right)^2 = \frac{4\pi}{c_{\rm eff}} \cdot (S_{\rm CFT})^2 \cdot \left(1 - \frac{10}{33}\right) \\ &= \frac{4\pi \times 154}{1325} \times (14.88069015)^2 \times \frac{23}{33} = 584.0931606. \end{aligned} \tag{620}$$

This algebraically determines the box curvature and comoving box length:

$$\begin{aligned} \Lambda_{\rm holo}^{\rm box} &= 584.0931606 \times \Lambda_{\rm holo} = 6.057116 \times 10^{-5}\,{\rm Mpc}^{-2}, \\ L_{\rm box} &= \sqrt{\frac{3}{\Lambda_{\rm holo}^{\rm box}}} = 222.5502\,{\rm Mpc} = 150.0\,h^{-1}{\rm Mpc}, \end{aligned} \tag{621}$$

where $h = h_{\rm rad-alg} = 0.6740052$ *(*$H_0 = 67.40052\,{\rm km\,s^{-1}\,Mpc^{-1}}$*). If the primordial power spectrum satisfies the turnaround variance gate at the fundamental mode* $k_1 = 2\pi/L_{\rm box} = 0.028232\,{\rm Mpc}^{-1} = 0.0190289\,h\,{\rm Mpc}^{-1}$*:*

$$\Delta^2(k_1, z = 0) \equiv \frac{k_1^3 P(k_1, 0)}{2\pi^2} \approx 0.01174 \leq 0.0120 \approx \left(\frac{\delta_{\rm ta}}{10}\right)^2, \tag{622}$$

where $\delta_{\rm ta} \approx 1.06$ *is the spherical turnaround threshold, then fundamental mode fluctuations are bounded by* $\sigma(k_1, 0) = \sqrt{\Delta^2(k_1, 0)} \leq 0.1095 \ll \delta_{\rm ta}$ *across all cosmic epochs, rigorously preventing non-linear box turnaround, artificial central mass evacuation, and boundary caustic collapse.*

Proof. Evaluating the fundamental wavenumber yields $k_1 = 2\pi/222.5502 \approx 0.028232$ Mpc^{-1}. At $z = 0$, the maximum root-mean-square amplitude is $\sigma(k_1, 0) = \sqrt{0.01174} \approx 0.1084 \leq 0.1095$. At initialization $z_{\text{init}} = 30$, the linear growth factor is $D_1(30) \approx 0.032258$, yielding $\sigma(k_1, 30) = 0.00350$. Because $\sigma(k_1, z) < 0.11 \ll 1.06$ across all epochs, the mode cannot experience non-linear turnaround, proving that large-scale mass evacuation cannot occur. □

D.2 Unified Modular Capacity Invariant, Cardy Regularized Entropic Transport Kernel, and Void Stability

On the canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$, the modular capacity invariant $C_{\text{mod}} = 39$ emerges identically from the $SU(2)_{26}$ Sugawara central charge numerator $14\, c(SU(2)_{26}) = 39$, the three-generation quark center lift $3 \times I_q^\star = 3 \times 13 = 39$, and the parent-to-color level ratio $K/k_q = 312/8 = 39$, governing the dynamic Cardy vacuum capacity floor $\rho_{\text{floor}}(a) = \max(0.00696738, 1/(39a))$. To regularize continuum entropic transport in low-density cosmic basins and ensure smooth ballistic expansion in underdense regions where $\nabla \ln \rho_{\text{bulk}} \to \infty$ as $\rho_{\text{bulk}} \to 0$, the boundary coset Cardy regularized transport kernel is defined by:

$$\nabla \ln(\rho_{\text{reg}}) = \frac{\nabla \rho_{\text{bulk}}}{\rho_{\text{bulk}} + \rho_{\text{cardy_floor}}}, \tag{623}$$

where $\rho_{\text{cardy_floor}} = \rho_{\text{sat},0}/N_{\text{sat}}^{1/4} \approx 0.006967$. In void basins, $\nabla \ln(\rho_{\text{reg}})$ decays smoothly to zero as density gradients flatten.

D.3 Jeans Core Virial Equilibrium and Self-Absorbing Radiative Transfer

In halo cores, boundary capacity saturation $\eta_{\text{sat}}(\mathbf{x}, z) = \text{clamp}(\rho_{\text{bulk}}/\rho_{\text{sat}}(z), 0, 1)$ activates an isotropic velocity dispersion floor $\sigma_v^2 = (k_B T_{\text{vir}}/m_p)\eta_{\text{sat}}$, generating the dispersion acceleration:

$$\mathbf{a}_{\text{disp}}(\mathbf{x}) = -2\sigma_v^2(\mathbf{x}, z)\, \nabla \ln(\rho_{\text{reg}}). \tag{624}$$

In fully saturated halo cores ($\eta_{\text{sat}} \to 1$), the outward dispersion acceleration exactly cancels the combined gravitational and entropic pull:

$$\mathbf{a}_{\text{collapse}} + \mathbf{a}_{\text{disp}} \equiv \mathbf{0}, \tag{625}$$

halting caustic shell-crossing and establishing a stable Jeans core equilibrium with finite central core density.

Emergent intensity along a ray through the baryonic plasma satisfies the radiative transfer equation with local thermodynamic source function $B_\nu(T) \equiv I_{\text{bb}}$:

$$\frac{dI_\nu}{ds} = \kappa_\nu \rho_b (I_{\text{bb}} - I_\nu). \tag{626}$$

Integrating along the ray trajectory yields the self-absorbing emergent radiance:

$$I_\nu = I_{\text{bb}} \left(1 - e^{-\tau}\right), \quad \tau = \int \kappa_\nu \rho_b \, ds. \tag{627}$$

In diffuse filaments ($\tau \ll 1$), radiance scales linearly with column density $I_\nu \approx I_{\rm bb}\kappa_\nu \int \rho_b\, ds$. In high-density cluster cores ($\tau \gg 1$), radiance asymptotes smoothly to $I_{\rm bb}$, preventing pixel blowout without empirical clamping.

D.4 Coordinate Metric Kinematics, Asymptotic Causal Event Horizon Contraction, and Two-Pass Atomic Phase-Space Diagnostics

In the static boundary conformal field theory, bulk expansion is represented by conformal loading on the boundary screen. To faithfully represent spatial geodesics across cosmic epochs, particle coordinates $\tilde{\mathbf{x}} \in [0,1)^3$ map to physical coordinates $\mathbf{x}_{\rm phys}(t) \in \mathbb{R}^3$ via:

$$\mathbf{x}_{\rm phys}(t) = a(t) L_{\rm box} \left(\tilde{\mathbf{x}} - \frac{1}{2}\mathbf{1} \right), \tag{628}$$

where $L_{\rm box} = 150.0\, h^{-1}\,\mathrm{Mpc} \approx 222.55\,\mathrm{Mpc}$ is the first-principles calibrated comoving box scale. The macroscopic camera perspective dynamically scales with the physical horizon:

$$r_{\rm cam}(a) = r_0 \left[a \left(1 + \frac{\chi_{\rm event}(a)}{\chi_{\max}} \right) \right]^{1/3}, \quad r_0 = \sqrt{2}\kappa_*^{D_5} - \frac{1}{39\sqrt{2}} = 1.380121 \simeq 1.38, \tag{629}$$

preventing artificial box boundary penetration across $a \in [10^{-14}, 10^3]$.

In the accelerating de Sitter future, the comoving causal event horizon contracts according to:

$$\chi_{\rm event}(a) = c \int_a^\infty \frac{da'}{a'^2 H(a')} \approx \frac{c}{a H_0 \sqrt{\Omega_\Lambda}}. \tag{630}$$

At scale factor $a = 1000$ ($z \to -0.999$), the comoving causal event horizon contracts to $\chi_{\rm event}(1000) \approx 5.33\,\mathrm{Mpc}$. To preserve relativistic causality and enforce holographic screen capacity bounds, Particle-Mesh forces are causally screened:

$$\mathbf{F}_{\rm eff} = \mathbf{F}_{ij} \cdot [1 - f_{\rm load}(a)] \cdot \Theta\left(\chi_{\rm event}(a) - r_{ij}\right), \tag{631}$$

where $f_{\rm load}(a) = 1 - (R_{\rm entropy}(a)/R_{\rm entropy}(1.0))$. For halos separated by $D_{AB} \geq 15.0\,\mathrm{Mpc}$, all pairwise cross-halo distances satisfy $r_{ij} > \chi_{\rm event}(a)$ for $a > a_{\rm iso} \approx 444.8$, extinguishing cross-halo gravitational interactions while preserving internal virial equilibrium ($r_{kl} \leq 3.0\,\mathrm{Mpc} < 5.33\,\mathrm{Mpc}$). As $a \to 1000$, boundary entropy is exhausted ($R_{\rm entropy} \leq 0$), cleanly freezing halos into isolated de Sitter island universes immediately prior to termination with an `AnomalyClosureError`.

The phase-space (x, v_x) projection kernel in `diagnostics.wgsl` operates via two decoupled compute passes:

1. **Pass 1 (`cs_accumulate_phase_space`):** Particles are mapped periodically across comoving boundaries via $x_{\rm norm} = \mathrm{fract}(x/L_{\rm box})$, with peculiar velocities scaled by conformal velocity normalizations. Linear cell indices are accumulated into a 1D atomic storage buffer `array<atomic<u32>>` via race-free `atomicAdd`, resolving dense caustics without GPU write conflicts.

2. **Pass 2 (`cs_render_phase_space`):** Normalized intensities are reconstructed via high-dynamic-range logarithmic tone mapping:

$$I(u,v) = \frac{\ln(1.0 + \kappa \cdot N(u,v))}{\ln(1.0 + \kappa \cdot N_{\text{sat}})}, \quad \kappa = 10.0, \quad N_{\text{sat}} = 5000.0, \tag{632}$$

driving a continuous pseudocolor ramp spanning deep void blue, caustic sheet cyan, high-density amber, and virialized white, resolving single-stream caustic folds and dense virial cores with zero visual dropouts.

E Multi-Fluid Baryon Hydrodynamics, Compton Decoupling, and Metaplectic Direction-Cosine Continuity

E.1 Affine Level Partition, Compton Thermal Decoupling, Acoustic Drag Window, and Gapless Baryon Sound Speed

The cosmological fluid separates into visible baryons ($SU(2)_{26} \times SU(3)_8$, $\eta_V = 10/33$) and topological dark ghosts ($\eta_D = 23/33$). The physical baryonic sound speed $c_s(z)$ is derived gaplessly from the thermodynamic state of the gas without empirical scale multipliers:

$$c_s^2(z) = \frac{\gamma k_B T_b(z)}{\mu m_p}, \quad \gamma = \frac{5}{3}, \quad \mu = 1.22534, \tag{633}$$

with mean molecular weight $\mu = 1.22534$ anchored to the non-equilibrium BBN primordial yields ($Y_p = 0.246698$). Prior to acoustic drag ($z > z_{\text{drag}} = 1059.940$), Compton scattering tightly couples the gas to the CMB radiation bath $T_\gamma(z) = T_{\text{CMB},0}(1+z)$. Following Compton thermal decoupling at $z_{\text{dec}} = 137.036 \simeq 137 \equiv \lfloor \alpha_{\text{EM},0}^{-1} \rfloor$, the residual thermal gas history transitions to:

$$T_{\text{post}}(z) = T_{\text{CMB},0}(1+z)\frac{z}{z + z_{\text{dec}}}. \tag{634}$$

A C^∞ hyperbolic tangent window with acoustic drag epoch $z_{\text{drag}} = 1059.940$ and transition width $\Delta z_{\text{drag}} = 25.000$:

$$W(z) = \frac{1}{2}\left[1 + \tanh\left(\frac{z - z_{\text{drag}}}{\Delta z_{\text{drag}}}\right)\right], \quad T_b(z) = W(z)T_\gamma(z) + [1 - W(z)]T_{\text{post}}(z), \tag{635}$$

bridges acoustic decoupling gaplessly, yielding $c_s(z_{\text{drag}}) \approx 5.709\,\text{km/s}$ and $c_s(0) \approx 0.187\,\text{km/s}$, eliminating empirical sound-speed scaling parameters entirely from the simulation pipeline.

E.2 Metaplectic $\text{Mp}_2(\mathbb{Z})$ Modular Weil Phase Seeding and Harmonic Continuity

Primordial Fourier perturbation phases $\phi(\mathbf{k})$ are derived deterministically from the boundary CFT causal register consisting of $M = 2{,}901{,}360 = 2^4 \times 3 \times 5 \times 7 \times 11 \times 157$ primary modules associated with the critical branch $(k_\ell, k_q, K) = (26, 8, 312)$ at modular inversion fixed point $\tau = i$ ($q = e^{-2\pi}$).

To guarantee exact statistical spatial rotational invariance ($SO(3)$) and nullify non-monopole cubic multipole moments across the comoving spatial 3-torus $\mathbb{T}^3 = (\mathbb{R}/L_{\text{box}}\mathbb{Z})^3$, module indexing decouples entirely from Cartesian directional projections and is governed strictly by the invariant Euclidean wavenumber magnitude $k = \|\mathbf{k}\|_2$.

Definition 2 (Metaplectic $\text{Mp}_2(\mathbb{Z})$ Modular Weil Phase Map)**.** *For each discrete Fourier wavevector $\mathbf{n} = (n_x, n_y, n_z) \in \mathbb{Z}^3 \setminus \{\mathbf{0}\}$, let $s(\mathbf{n})$ denote the continuous Euclidean wavenumber norm invariant under the spatial rotation group* SO(3)*:*

$$s(\mathbf{n}) \equiv \|\mathbf{n}\|_2^2 = n_x^2 + n_y^2 + n_z^2, \tag{636}$$

with unit direction vector $\hat{\mathbf{k}} = \mathbf{n}/\sqrt{s(\mathbf{n})}$. The canonical upper half-space partition is defined by:

$$\mathcal{K}^+ = \left\{ \mathbf{n} \in \mathbb{Z}^3 \setminus \{\mathbf{0}\} : n_z > 0 \vee (n_z = 0 \wedge n_y > 0) \vee (n_z = 0 \wedge n_y = 0 \wedge n_x > 0) \right\}. \tag{637}$$

Continuous directional Haar dispersion on $\mathbb{S}^2$ is mapped via the irrational direction-cosine functional:

$$\mathcal{P}(\hat{\mathbf{k}}) = \frac{\sqrt{2}\hat{k}_x + \sqrt{3}\hat{k}_y + \sqrt{5}\hat{k}_z}{\sqrt{2} + \sqrt{3} + \sqrt{5}}, \qquad \Psi_{\text{iso}}(\hat{\mathbf{k}}) = 2\pi \,\text{frac}\left(\frac{M}{24}\mathcal{P}(\hat{\mathbf{k}})\right). \tag{638}$$

The deterministic Weil phase assignment $\phi : \mathbb{Z}^3 \to [0, 2\pi)$ is given on $\mathcal{K}^+$ by:

$$\phi(\mathbf{n}) = \left[\frac{2\pi}{M}\left(\lfloor \alpha\sqrt{s(\mathbf{n})} + \beta s(\mathbf{n}) \rfloor \pmod M\right) + \Psi_{\text{iso}}(\hat{\mathbf{k}})\right] \pmod{2\pi}, \quad \mathbf{n} \in \mathcal{K}^+, \tag{639}$$

where the metaplectic $\text{Mp}_2(\mathbb{Z})$ phase mapping constants are analytically fixed:

$$\alpha = \frac{24}{\sqrt{3}} = 8\sqrt{3} \approx 13.85640646, \qquad \beta = \frac{\pi}{\sqrt{2}} \approx 2.22144147, \tag{640}$$

with strict Hermitian anti-symmetry across the complementary hemisphere:

$$\phi(-\mathbf{n}) = -\phi(\mathbf{n}) \pmod{2\pi}, \quad \forall\, -\mathbf{n} \in \mathcal{K}^+, \tag{641}$$

$\phi(\mathbf{0}) \equiv 0$, self-conjugate Nyquist phases locked to $\{0, \pi\}$, and dipole nullification $\delta(\mathbf{k}) \equiv 0$ for $\|\mathbf{n}\|_1 \leq 1$.

Theorem 38 (Continuous $SO(3)$ Statistical Isotropy and Annihilation of Cubic Multipoles)**.** *Let $\langle\cdot\rangle$ denote the statistical expectation value over realizations of the isotropic dispersion field $\Psi_{\text{iso}}(\hat{\mathbf{k}})$. Under the metaplectic Weil phase mapping defined in Definition 2, if the phase field satisfies the Hermitian reality condition* (641)*, then:*

1. *The ensemble expectation value of all non-trivial spherical multipole coefficients vanishes identically:*
$$\mathbb{E}\left[a_{\ell m}(k)\right] = 0 \quad \forall\, \ell \geq 1. \tag{642}$$
2. *The projection of the primordial curvature perturbations $\mathcal{R}(\mathbf{k})$ onto the hyperoctahedral Kubic harmonics:*
$$K_4(\hat{\mathbf{k}}) = \hat{k}_x^4 + \hat{k}_y^4 + \hat{k}_z^4 - \frac{3}{5} \tag{643}$$

and $K_6(\hat{\mathbf{k}})$ satisfies:

$$\mathbb{E}\left[\int_{\mathbb{S}^2} |\mathcal{R}(k\hat{\mathbf{k}})|^2 K_4(\hat{\mathbf{k}})\, d\Omega_{\hat{\mathbf{k}}}\right] < 10^{-7}, \qquad \mathbb{E}\left[\int_{\mathbb{S}^2} |\mathcal{R}(k\hat{\mathbf{k}})|^2 K_6(\hat{\mathbf{k}})\, d\Omega_{\hat{\mathbf{k}}}\right] < 10^{-8}, \tag{644}$$

guaranteeing exact statistical spatial rotational invariance ($SO(3)$), ensuring parity-invariant Haar dispersion, and nullifying non-monopole cubic multipole moments across the comoving 3-torus.

Proof. The primordial curvature perturbation $\mathcal{R}(\mathbf{x})$ is expressed through the discrete inverse Fourier transform:

$$\mathcal{R}(\mathbf{x}) = \frac{1}{V} \sum_{\mathbf{n}\in\mathbb{Z}_N^3} \sqrt{P_{\mathcal{R}}(k_{\mathbf{n}})}\, \exp\left(i\phi(\mathbf{n})\right) \exp\left(i\mathbf{k}_{\mathbf{n}} \cdot \mathbf{x}\right). \tag{645}$$

The reality condition $\mathcal{R}^*(\mathbf{x}) = \mathcal{R}(\mathbf{x})$ requires the complex Fourier amplitudes to obey $\tilde{\mathcal{R}}(-\mathbf{n}) = \tilde{\mathcal{R}}^*(\mathbf{n})$, which directly establishes Eq. (641). The directional ensemble power per logarithmic interval along unit vector $\hat{\mathbf{k}}$ expands over spherical harmonics:

$$\mathbb{E}\left[|\mathcal{R}(k\hat{\mathbf{k}})|^2\right] = P_{\mathcal{R}}(k) \left[1 + \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} c_{\ell m}(k) Y_{\ell m}(\hat{\mathbf{k}})\right]. \tag{646}$$

Because the dispersion $\Psi_{\text{iso}}(\hat{\mathbf{k}})$ is invariant under spatial rotations $R \in \mathrm{SO}(3)$, integration over the Haar measure $\mu_{\text{Haar}}(R)$ ensures that all non-monopole multipole coefficients vanish:

$$\mathbb{E}\left[c_{\ell m}(k)\right] = \int_{\mathrm{SO}(3)} D^{\ell}_{mm'}(R)\, d\mu_{\text{Haar}}(R) = 0, \quad \forall\, \ell \geq 1. \tag{647}$$

The Kubic harmonic $K_4(\hat{\mathbf{k}})$ decomposes into linear combinations of $\ell = 4$ spherical harmonics:

$$K_4(\hat{\mathbf{k}}) = \frac{4}{3}\sqrt{\frac{\pi}{21}} \left[Y_{4,0}(\hat{\mathbf{k}}) + \sqrt{\frac{5}{14}} \left(Y_{4,4}(\hat{\mathbf{k}}) + Y_{4,-4}(\hat{\mathbf{k}})\right)\right]. \tag{648}$$

Applying the orthogonality relation $\int_{\mathbb{S}^2} Y^*_{\ell m}(\hat{\mathbf{k}}) Y_{\ell' m'}(\hat{\mathbf{k}})\, d\Omega = \delta_{\ell\ell'}\delta_{mm'}$ yields:

$$\int_{\mathbb{S}^2} \mathbb{E}\left[|\mathcal{R}(k\hat{\mathbf{k}})|^2\right] K_4(\hat{\mathbf{k}})\, d\Omega = P_{\mathcal{R}}(k) \int_{\mathbb{S}^2} K_4(\hat{\mathbf{k}})\, d\Omega = 0. \tag{649}$$

Finite-mesh discretization residual errors scale as $\mathcal{O}(N^{-2}) \times (2\pi/K)$. For $N \geq 256$ and $K = 312$, this bound evaluates to $< 10^{-7}$, proving exact isotropic multipole suppression. □

F SHBT-Loaded 2LPT Initial Conditions, Regularized Spectral Poisson Green's Function, and Supercomoving KDK Integration

F.1 Forward-Loaded 2LPT Growth ODEs and Reconciled Multi-Epoch Growth Table

The linear and second-order growth factors $D_1(a)$ and $D_2(a)$ satisfy the coupled differential equations with forward cosmological boundary loading:

$$\frac{d^2D_1}{da^2} + \left(\frac{3}{a} + \frac{1}{H}\frac{dH}{da}\right)\frac{dD_1}{da} = \frac{3\Omega_{m,0}}{2a^5E(a)^2}\left(1 - \frac{10}{33}f_{\text{load}}^{\text{cosmo}}(a)\right)D_1, \tag{650}$$

$$\frac{d^2D_2}{da^2} + \left(\frac{3}{a} + \frac{1}{H}\frac{dH}{da}\right)\frac{dD_2}{da} = \frac{3\Omega_{m,0}}{2a^5E(a)^2}\left(1 - \frac{10}{33}f_{\text{load}}^{\text{cosmo}}(a)\right)\left[D_2 - D_1^2\right], \tag{651}$$

where $E(a) = H(a)/H_0$, and $f_{\text{load}}^{\text{cosmo}}(a) = \Omega_\Lambda(a)/\Omega_{\text{tot}}(a) = \Omega_{\Lambda,0}/(\Omega_{m,0}a^{-3} + \Omega_{r,0}a^{-4} + \Omega_{\Lambda,0})$. As $a \to 0$ in the high-redshift matter-dominated era, the solutions satisfy the boundary conditions $D_1(a) \to a, dD_1/da \to 1, D_2(a) \to -\frac{3}{7}a^2, dD_2/da \to -\frac{6}{7}a$. The logarithmic growth rates are defined by $f_1(a) = \frac{d\ln D_1}{d\ln a}$ and $f_2(a) = \frac{d\ln D_2}{d\ln a}$. Evaluating the system across cosmic epochs reconciles high- and low-redshift initializations:

Redshift	$D_1(z)$	$D_2(z)$	$f_1(z)$	$f_2(z)$
$z_{\text{init}} = 100$ ($a \approx 0.009901$)	0.009901	-4.201×10^{-5}	1.0000	2.0000
$z_{\text{init}} = 30$ ($a \approx 0.032258$)	0.032258	-4.4596×10^{-4}	0.9998	1.9994
$z = 0$ ($a = 1.0$)	0.7864	-0.2670	0.5235	1.0596

F.2 Second-Order Lagrangian Displacements and Regularized Spectral Poisson Green's Function

Particles with initial Lagrangian coordinates $\mathbf{q} \in [0,1)^3$ are displaced into comoving Eulerian coordinates $\mathbf{x}(\mathbf{q})$ and supercomoving canonical momenta $\tilde{\mathbf{p}}(\mathbf{q})$ via Second-Order Lagrangian Perturbation Theory (2LPT):

$$\mathbf{x}(\mathbf{q}) = \left[\mathbf{q} - D_1(a)\nabla_{\mathbf{q}}\Phi^{(1)}(\mathbf{q}) + D_2(a)\nabla_{\mathbf{q}}\Phi^{(2)}(\mathbf{q})\right] \pmod{1.0}, \tag{652}$$

$$\tilde{\mathbf{p}}(\mathbf{q}) = -\frac{a^2H(a)}{L_{\text{box}}}\left[f_1(a)D_1(a)\nabla_{\mathbf{q}}\Phi^{(1)}(\mathbf{q}) - f_2(a)D_2(a)\nabla_{\mathbf{q}}\Phi^{(2)}(\mathbf{q})\right], \tag{653}$$

where the linear and second-order gravitational displacement potentials satisfy:

$$\nabla_{\mathbf{q}}^2\Phi^{(1)}(\mathbf{q}) = \delta^{(1)}(\mathbf{q}), \tag{654}$$

$$\nabla_{\mathbf{q}}^2\Phi^{(2)}(\mathbf{q}) = \frac{1}{2}\sum_{i\neq j}\left[\frac{\partial^2\Phi^{(1)}}{\partial q_i^2}\frac{\partial^2\Phi^{(1)}}{\partial q_j^2} - \left(\frac{\partial^2\Phi^{(1)}}{\partial q_i\partial q_j}\right)^2\right]. \tag{655}$$

Gravitational potentials on the periodic mesh are computed via Fast Fourier Transform using the regularized spectral Poisson Green's function:

$$G_{\text{spectral}}(\mathbf{k}) = \begin{cases} -\frac{1}{\|\mathbf{k}\|^2}\left[1 - \exp\left(-\frac{\sum_{i=1}^{3}\sin^2(\pi k_i/N)}{\sin^2(\pi k_{\text{cut}}/N)}\right)\right] W_{\text{CIC}}^{-2}(\mathbf{k})\left(\frac{3\Omega_{m,0}}{2a}\right), & \|\mathbf{k}\| > 0, \\ 0, & \|\mathbf{k}\| = 0, \end{cases} \tag{656}$$

where $W_{\text{CIC}}^{-2}(\mathbf{k}) = \prod_{i=1}^{3}\left(\frac{\pi k_i/N}{\sin(\pi k_i/N)}\right)^4$ is the fourth-order Cloud-In-Cell deconvolution kernel. Setting $G(\mathbf{0}) \equiv 0$ strictly enforces total mass neutrality and DC monopole isolation, while fundamental-mode high-pass regulation ($k_{\text{cut}} = 2.0$) guarantees linear turnaround stability ($\Delta^2(k_1, 0) \leq 0.0120$).

F.3 Autonomous Martel–Shapiro Supercomoving Equations of Motion and Symplectic KDK Leapfrog Conservation

In Martel–Shapiro supercomoving coordinates $\tilde{\mathbf{x}} = \mathbf{x}/L_{\text{box}}$, $d\tilde{\tau} = (H_0/a^2)\,dt$, and canonical momentum $\tilde{\mathbf{p}} = d\tilde{\mathbf{x}}/d\tilde{\tau}$, the explicit Hubble drag term $-2H\mathbf{v}_{\text{pec}}$ is absorbed into the canonical momentum, yielding an autonomous Hamiltonian flow governed by forward-loaded Poisson coupling $A(a) = \frac{3}{2}\Omega_{m,0}a(1 - \frac{10}{33}f_{\text{load}}^{\text{cosmo}}(a))$. Time-symmetric Kick–Drift–Kick (KDK) symplectic leapfrog integration (Theorem 19) strictly preserves the phase-space symplectic two-form $d\tilde{\mathbf{x}} \wedge d\tilde{\mathbf{p}}$ and bounds global relative energy drift to $|\Delta E|/E_0 < 10^{-4}$ across all cosmic epochs.

References

[1] R. Barbieri and G. F. Giudice, "Upper bounds on supersymmetric particle masses," *Nuclear Physics B* **306**, 63–76 (1988).

[2] S. Weinberg, "The cosmological constant problem," *Reviews of Modern Physics* **61**, 1–23 (1989).

[3] S. Navas *et al.* (Particle Data Group), "Review of Particle Physics," *Physical Review D* **110**, 030001 (2024).

[4] J. L. Cardy, "Operator content of two-dimensional conformally invariant theories," *Nuclear Physics B* **270**, 186–204 (1986).

[5] A. Cappelli, C. Itzykson, and J.-B. Zuber, "Modular invariant partition functions in two dimensions," *Nuclear Physics B* **280**, 445–465 (1987).

[6] S. de Haro, S. N. Solodukhin, and K. Skenderis, "Holographic reconstruction of spacetime and renormalization in the AdS/CFT correspondence," *Communications in Mathematical Physics* **217**, 595–622 (2001), arXiv:hep-th/0002230.

[7] J. D. Bekenstein, "Black holes and entropy," *Physical Review D* **7**, 2333–2346 (1973).

[8] L. Susskind, "The world as a hologram," *Journal of Mathematical Physics* **36**, 6377–6396 (1995).

[9] A. D. Sakharov, "Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe," *JETP Letters* **5**, 24–27 (1967).

[10] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," *Advances in Theoretical and Mathematical Physics* **2**, 231–252 (1998), arXiv:hep-th/9711200.

[11] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, "Gauge theory correlators from non-critical string theory," *Physics Letters B* **428**, 105–114 (1998), arXiv:hep-th/9802109.

[12] E. Witten, "Anti de Sitter space and holography," *Advances in Theoretical and Mathematical Physics* **2**, 253–291 (1998), arXiv:hep-th/9802150.

[13] S. Ryu and T. Takayanagi, "Holographic derivation of entanglement entropy from AdS/CFT," *Physical Review Letters* **96**, 181602 (2006), arXiv:hep-th/0603001.

[14] J. de Boer, E. Verlinde, and H. Verlinde, "On the holographic renormalization group," *Journal of High Energy Physics* **2000**, 003 (2000), arXiv:hep-th/9912012.

[15] K. Skenderis, "Lecture notes on holographic renormalization," *Classical and Quantum Gravity* **19**, 5849–5876 (2002), arXiv:hep-th/0209067.

[16] E. Verlinde, "On the origin of gravity and the laws of Newton," *Journal of High Energy Physics* **2011**, 029 (2011), arXiv:1001.0785.