

Static Holographic Boundary Theory and Precision Cosmology

July 16, 2026

Abstract

Static Holographic Boundary Theory (SHBT) is a boundary-first closure program: the primitive object is a completed boundary conformal field theory whose torus partition function is exactly modular invariant, not a pre-existing bulk spacetime. The canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$ locks the weak and color center lifts to $I_\ell^* = 6$ and $I_q^* = 13$, so the framing defect Δ_{fr} vanishes. On this branch, modular closure is equivalent to a vanishing bulk closure tensor and to the projected Hamiltonian constraint $\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0$.

This unified manuscript combines the foundational SHBT simulator paper with the finite-capacity precision-cosmology manuscript. The common data determine the visible modular arithmetic, the residual and completed dark ledgers $c_{\text{dark}}^{\text{res}} = 834433/362670$ and $c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = 1197103/362670$, the finite horizon register, holographic RG metric slices, baryogenesis by anti-baryon de-rendering, and Causal Point observer histories. The cosmology part uses the completed ledger to define an entropy-debt variable $\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$ and an observer-frame loading law that interpolates between the Planck CMB intercept and the local Hubble value while decaying as $(1+z)^{-1}$ toward the early universe.

The resulting paper presents both the closure theory and its audit trail. The final integration makes explicit the Fefferman–Graham/topological-Einstein bridge, the entropy-debt redshift ODE, the Hubble-loading amplitude A_H , the lock rate $\Gamma_{\text{lock}} = 3A_H$, the structure-growth and ISW residual equations, BBN decoupling, chronometer covariance, and the finite GET measurement rule. The simulator checks exact branch closure, nine positive trace-normalized metric slices, nine finite-memory history entries, stress-energy-preserving anti-baryon de-rendering with $\eta_B = 6.449923359416 \times 10^{-10}$, Newton-lock stationarity, the unity-of-scale identity, Hubble-ladder tracking, growth suppression, cluster and ISW stability, and the thermodynamic light-cone entropy ledger.

1 Introduction

Quantum gravity and Standard Model phenomenology are usually formulated in opposite directions. Perturbative quantum field theory starts from local particle multiplets, gauge charges, and renormalization-group data, and then asks how this successful low-energy structure survives coupling to gravity. Holography starts from boundary data and asks how local bulk spacetime and effective fields emerge. A realistic theory must make these two directions meet: it must account for the observed particle content, vacuum response, baryon asymmetry, and cosmological scales without repeatedly inserting small numbers by hand. This is the naturalness problem in its broadest

form. The Standard Model is phenomenologically successful, but the gravitational and vacuum sectors appear highly sensitive to ultraviolet completion unless a deeper closure principle fixes the allowed data [1, 2, 3].

This manuscript unifies two previously separate presentations. Sections 2–8 develop the static boundary theory, while Section 9 adds the finite-capacity cosmological implications and their Python audit outputs. To remove the notation collision between the two source manuscripts, the residual dark ledger used in the foundational simulator is denoted $c_{\text{dark}}^{\text{res}} = 834433/362670$, whereas the completed finite-screen ledger used by the Noether bridge and precision cosmology is $c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = 1197103/362670$. The symbol κ_{D_5} is used throughout for the retained D_5 boundary-cell measure; κ_{D_5} is retained only as a legacy alias. In the precision-cosmology sector, $\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$, A_H denotes the Hubble-loading amplitude rather than a horizon area, and $\Gamma_{\text{lock}} = 3A_H$ denotes the Causal Point lock rate. The microscopic center-lift, completed-ledger, and boundary-measure definitions are collected in Appendices A–C.

Static Holographic Boundary Theory (SHBT) proposes that the missing closure principle is exact modular invariance of a completed boundary conformal field theory (CFT). The starting point is neither a fluctuating classical spacetime nor a pre-existing cosmological history, but a completed boundary Hilbert space and a physical torus partition function

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I, J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\tau)}, \quad [M, S_{\partial}] = [M, T_{\partial}] = 0, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}. \quad (1)$$

The role of modular covariance follows the standard lesson that torus partition functions in two-dimensional CFT are not arbitrary sums of characters: their operator content is constrained by the modular group [4, 5]. SHBT strengthens this consistency condition into a first principle. A boundary sector is physical only if it closes under the completed modular kernels, and any residual mismatch is encoded in a scalar framing defect Δ_{fr} . The central claim is the closure chain

$$[M, S_{\partial}] = [M, T_{\partial}] = 0 \iff \Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0 \iff \Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0. \quad (2)$$

Thus holographic closure is not imposed after a bulk theory has been guessed. Rather, the vanishing of the bulk closure tensor and the disappearance of an independent external time generator are the bulk and Hamiltonian images of completed modular closure on the boundary. The implementation audits this link through

`shbt_simulator.ShbtSimulator().run_full_audit().boundary_report,`
with the partition data and visible modular phases constructed by `shbt_simulator.StaticBoundary.evaluate_z_boundary,`
`shbt_simulator.StaticBoundary.build_su2_visible_phases,` and
`shbt_simulator.StaticBoundary.build_su3_visible_phases.`

The canonical branch studied here is

$$\mathfrak{b}_* = (k_{\ell}, k_q, K) = (26, 8, 312), \quad I_{\ell}^* = \frac{K}{2k_{\ell}} = 6, \quad I_q^* = \frac{K}{3k_q} = 13. \quad (3)$$

Within SHBT this branch is the anomaly-free fixed point because the lepton, color, and parent completion levels lock to integers and give

$$\Delta_{\text{fr}}(312, 26, 8) = \max\left(\left\|\frac{312}{2 \cdot 26}\right\|_{\mathbb{Z}}, \left\|\frac{312}{3 \cdot 8}\right\|_{\mathbb{Z}}\right) = 0. \quad (4)$$

The visible factors are $SU(2)_{26}$ and $SU(3)_8$, embedded in an $SO(10)_{312}$ parent completion and a dark completion sector. The dark sector is therefore not an optional phenomenological reservoir. It is the finite ledger required by exact closure. The foundation simulator reports the residual part

$$c_{\text{dark}}^{\text{res}} = \frac{834433}{362670} \simeq 2.300805139658643, \quad 8\pi G_N = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2}. \quad (5)$$

The finite-screen cosmology audit uses the completed ledger

$$c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = \frac{1197103}{362670} \simeq 3.300805139658643, \quad \Delta_{\text{mod}} = \frac{c_{\text{dark}}^{\text{comp}}}{24}. \quad (6)$$

The same arithmetic appears in `shbt_simulator.StaticBoundary.benchmark_branch`, `shbt_simulator.StaticBoundary.lepton_level`, `shbt_simulator.StaticBoundary.quark_level`, `shbt_simulator.StaticBoundary.parent_level`, `shbt_simulator.StaticBoundary.i_l_star`, `shbt_simulator.StaticBoundary.i_q_star`, `shbt_simulator.StaticBoundary.c_dark`, and `report.framing_defect`.

The construction has four linked components. First, the completed boundary CFT is treated as the primary object,

$$\mathcal{H}_{\partial}^{\text{comp}} = \mathcal{H}_{SU(2)_{26}} \otimes \mathcal{H}_{SU(3)_8} \otimes \mathcal{H}_{SO(10)_{312}} \otimes \mathcal{H}_{\text{dark}}, \quad (7)$$

whose physical spectrum is the modular-invariant, nonnegative-integer pairing of completed characters. This is the algebraic source of gauge and gravitational consistency. Second, entropy self-resolution turns the finite boundary register

$$N = N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}}, \quad \Lambda_{\text{holo}} > 0, \quad (8)$$

into normalized loading and entanglement densities. Their entropy gradients define a holographic renormalization-group flow into bulk metric slices. This continues the broader holographic and black-hole thermodynamic tradition [7, 8, 9], but makes the entropy budget finite and auditable through `shbt_simulator.HolographicProjection`, `shbt_simulator.BulkMetricSlice`, and the `shbt_simulator.ShbtSimulator().run_full_audit().projection_report` record.

Third, SHBT reframes baryogenesis as anti-baryon de-rendering. The observed baryon asymmetry is usually framed as a dynamical departure from baryon number, C, CP, and equilibrium conditions [10]. Here the active render charge is separated from passive stress-energy. Anti-baryon visible charges are projected out of the actively rendered gauge channels, while the passive gravitational source is preserved in the dark completion sector. The resulting asymmetry is therefore an observer-visible render imbalance, not a failure of completed stress-energy accounting. The implementation checks this claim through

- `shbt_simulator.ShbtSimulator().run_full_audit().baryogenesis_identity`,
- `shbt_simulator.BaryogenesisOptimizer.derender_antibaryon_charges`,
- `report.stress_energy_preserved`, and
- `shbt_simulator.ShbtSimulator().run_full_audit().benchmark_delta`.

The benchmark value reported by the audit is `eta_b=6.449923359416e-10`.

Fourth, the Causal Point makes the observer part of the same finite register. SHBT does not assume an external classical observer with unlimited memory. An observer is a finite local address equipped with a light-cone sample, a local property packet, a hidden-state budget, and a history projection operator $\Pi_{A,\ell}$. A measurement is admitted only when the residual entropy budget is nonnegative. Observable time and crystallized history are therefore finite-register projections rather than primitive background parameters. The corresponding code objects are `shbt_simulator.CausalPoint`, `shbt_simulator.LightConeSample`, `shbt_simulator.LocalPropertyPacket`, `shbt_simulator.CoordinateLogEntry`, the `shbt_simulator.ShbtSimulator().run_full_audit().memory_report` record, and `shbt_simulator.CausalPoint.crystallize_history`.

The main results follow from the same closure data. Because the branch (26, 8, 312) has exact framing closure, the bulk closure tensor vanishes and the topological Einstein equation follows in the SHBT sector:

$$\mathcal{E}_{\mu\nu} = \frac{c_{\text{dark}}^{\text{res}} M_P^2}{12} (G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu}) - T_{\mu\nu}^{\text{vis}} = 0. \quad (9)$$

The static physical sector has no independent external time generator because the completed Hamiltonian constraint annihilates physical states, $\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0$. Finite-register entropy flow produces symmetric, trace-normalized, positive metric slices and an observer-frame effective cosmology. The Causal Point construction turns measurement into an entropy-budgeted projection with explicit failure conditions. Anti-baryon de-rendering yields the benchmark baryon asymmetry while preserving passive stress-energy. Running `python examples/run_audit.py` is the intended reproducibility check: it should report the canonical branch, `framing_defect=0.0`, `modular_invariant=True`, `zero_energy_locked=True`, the finite bit budget, nine bulk metric slices, nine crystallized history entries, `eta_b=6.449923359416e-10`, and `stress_energy_preserved=True`.

The audit framework presented here is primarily a self-consistency validation of the boundary-first closure program. The numerical values reported—including the baryon asymmetry $\eta_B = 6.449923359416 \times 10^{-10}$, the dark completion residue $c_{\text{dark}}^{\text{comp}} = 1197103/362670$, and the finite horizon register $N_{\text{sat}} \approx 3.31 \times 10^{122}$ —are not independent external inputs that the code fits to observation. They are structural consequences of the modular invariance conditions imposed on the completed boundary CFT. The simulations therefore verify that the canonical branch is internally closed: if the boundary data are fixed by exact closure, the bulk and observer projections automatically reproduce the observed scales. In this sense, the match with the measured baryon asymmetry is a retrodiction from symmetry principles, not a post-hoc parameter adjustment. This distinguishes SHBT from phenomenological models that tune many degrees of freedom to match datasets; here, any deviation from the locked branch would reopen the framing defect Δ_{fr} and break the closure chain. The key future test for SHBT as an independent predictive framework will be its ability to forecast novel signatures (e.g., specific time-variations in $w(a)$, non-Gaussianities in the CMB, or direct dark-matter detection channels) that are not already encoded in the modular arithmetic of the canonical branch.

This positioning clarifies the relationship between SHBT and existing approaches. Like AdS/CFT, SHBT treats a boundary quantum theory as sufficient data for a bulk gravitational description [11, 12, 13]. Unlike standard AdS/CFT, the present construction is not organized around a large- N conformal gauge theory with an asymptotically anti-de Sitter bulk. It is organized around a

completed static boundary CFT, modular closure, and a finite holographic register. Like entropic-gravity programs, SHBT treats geometry as downstream of information and entropy [17]; unlike a pure entropic-force derivation, its entropy gradients are constrained by modular closure and checked slice-by-slice by an implementation audit. Like topological and constraint-based approaches to quantum gravity, SHBT removes an external time generator from the physical sector; unlike a purely formal constraint algebra, it ties that removal to visible-sector branch arithmetic, dark completion, baryogenesis, and observer memory.

The paper is organized as follows. Section 2 derives the canonical branch and states the axiomatic closure conditions. Section 3 constructs the modular data and boundary partition function. Section 4 defines the finite entropy densities that self-resolve the boundary register, and Section 5 maps their gradients into bulk metric slices. Section 6 develops anti-baryon de-rendering and stress-energy preservation, while Section 7 defines Causal Points and history crystallization. Section 8 collects the implementation audits, and Section 10 summarizes the unified closure picture and remaining limitations. Throughout, the mathematical derivation and `shbt_simulator.ShbtSimulator().run_full_audit()` are kept in parallel so that the same branch, defect, entropy budget, metric projection, Causal Point, and baryogenesis identity must close both on paper and in code.

2 Canonical Branch and Axiomatic Foundation

The canonical branch is derived by treating the completed boundary theory, not a bulk Lagrangian, as the primitive object. For arbitrary visible levels (k_ℓ, k_q) and parent level K , the completed boundary Hilbert space is postulated to factor as

$$\mathcal{H}_\partial^{\text{comp}}(k_\ell, k_q, K) = \mathcal{H}_{SU(2)_{k_\ell}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\text{dark}}. \quad (10)$$

The collective primary index is written $I = (i_\ell, i_q, i_{10}, \alpha)$, and its completed character is

$$\chi_I(\tau) = \chi_{i_\ell}^{(k_\ell)}(\tau) \chi_{i_q}^{(k_q)}(\tau) \chi_{i_{10}}^{(K)}(\tau) \chi_\alpha^{\text{dark}}(\tau). \quad (11)$$

The physical torus partition function has the form

$$Z_\partial(\tau, \bar{\tau}) = \sum_{I, J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\bar{\tau})}, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}, \quad (12)$$

with nonnegative integer multiplicities. The first axiom is that the completed pairing matrix is invariant under the full boundary modular kernels,

$$[M, S_\partial] = 0, \quad [M, T_\partial] = 0, \quad (13)$$

where

$$\begin{aligned} S_\partial &= S_{SU(2)_{k_\ell}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\text{dark}}, \\ T_\partial &= T_{SU(2)_{k_\ell}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\text{dark}}. \end{aligned} \quad (14)$$

Equivalently, $Z_\partial(\tau + 1, \bar{\tau} + 1) = Z_\partial(\tau, \bar{\tau})$ and $Z_\partial(-1/\tau, -1/\bar{\tau}) = Z_\partial(\tau, \bar{\tau})$, so the allowed operator content is a modular-invariant completed pairing rather than an arbitrary list of boundary primaries [4, 5].

The integral-spin condition is the local form of the T -constraint. A character transforms as

$$\chi_I(\tau + 1) = \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right] \chi_I(\tau), \quad (15)$$

while the anti-holomorphic character contributes the conjugate phase. Therefore the summand with coefficient M_{IJ} is T -invariant only if

$$M_{IJ} \neq 0 \implies \exp\left\{2\pi i \left[h_I - h_J - \frac{c_I - c_J}{24}\right]\right\} = 1, \quad (16)$$

or, equivalently,

$$\boxed{M_{IJ} \neq 0 \implies h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}.} \quad (17)$$

This condition is the boundary algebraic precursor of the SHBT framing defect. The visible conformal weights used by the implementation are

$$\begin{aligned} h_a^{SU(2)_k} &= \frac{a(a+2)}{4(k+2)}, \\ h_{(p,q)}^{SU(3)_k} &= \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \end{aligned} \quad (18)$$

with central charges

$$c_{SU(2)_k} = \frac{3k}{k+2}, \quad c_{SU(3)_k} = \frac{8k}{k+3}, \quad c_{SO(10)_K} = \frac{45K}{K+8}. \quad (19)$$

The canonical branch is the minimal parent completion compatible with the visible lepton and quark levels used throughout the audit. The embedding locks require the parent level to close simultaneously over the lepton and color descendants,

$$I_\ell = \frac{K}{2k_\ell} \in \mathbb{Z}, \quad I_q = \frac{K}{3k_q} \in \mathbb{Z}. \quad (20)$$

For the visible levels $k_\ell = 26$ and $k_q = 8$, the smallest positive parent level satisfying both locks is

$$K = \text{lcm}(2 \cdot 26, 3 \cdot 8) = \text{lcm}(52, 24) = 312. \quad (21)$$

Thus

$$\mathfrak{b}_* = (k_\ell, k_q, K) = (26, 8, 312), \quad I_\ell^\star = \frac{312}{2 \cdot 26} = 6, \quad I_q^\star = \frac{312}{3 \cdot 8} = 13. \quad (22)$$

This is the mathematical counterpart of `shbt_simulator.StaticBoundary.benchmark_branch = (26, 8, 312)`, together with `shbt_simulator.StaticBoundary.lepton_level`, `shbt_simulator.StaticBoundary.quark_level`, `shbt_simulator.StaticBoundary.parent_level`, `shbt_simulator.StaticBoundary.i_l_star`, and `shbt_simulator.StaticBoundary.i_q_star`.

The scalar obstruction to these integer locks is the framing defect

$$\boxed{\Delta_{\text{fr}}(K, k_\ell, k_q) = \max\left(\left\|\frac{K}{2k_\ell}\right\|_{\mathbb{Z}}, \left\|\frac{K}{3k_q}\right\|_{\mathbb{Z}}\right)}, \quad \|x\|_{\mathbb{Z}} = \min_{n \in \mathbb{Z}} |x - n|. \quad (23)$$

At the benchmark branch both arguments are integers, so

$$\Delta_{\text{fr}}(312, 26, 8) = \max(\|6\|_{\mathbb{Z}}, \|13\|_{\mathbb{Z}}) = 0. \quad (24)$$

The branch is therefore anomaly-free in the SHBT sense: the parent completion has no residual visible framing mismatch. In code this is reported as `report.framing_defect=0.0` and checked by the `shbt_simulator.ShbtSimulator().run_full_audit().boundary_report` record.

The second axiom identifies the bulk closure tensor as the covariant image of the scalar modular obstruction. The standard holographic bridge begins with the Fefferman–Graham expansion of an asymptotically locally AdS metric [6, 16],

$$ds^2 = \frac{L^2}{z^2} (dz^2 + g_{ij}(x, z) dx^i dx^j), \quad z \rightarrow 0, \quad (25)$$

where

$$g_{ij}(x, z) = g_{(0)ij}(x) + z^2 g_{(2)ij}(x) + \cdots + z^d g_{(d)ij}(x) + z^d \log z^2 h_{(d)ij}(x) + \cdots. \quad (26)$$

The boundary metric $g_{(0)ij}$ is the source for the CFT stress tensor, and holographic renormalization reads the renormalized one-point function from the normalizable coefficient,

$$\langle T_{ij} \rangle = \frac{dL^{d-1}}{16\pi G_N} g_{(d)ij} + T_{ij}^{\text{ct}}[g_{(0)}], \quad (27)$$

with local counterterms fixed by the same boundary data. Under a boundary Weyl variation,

$$\delta_\sigma W_{\text{CFT}}[g_{(0)}] = \int d^d x \sqrt{g_{(0)}} \sigma \mathcal{A}_{\text{Weyl}}, \quad \langle T^i_i \rangle = \mathcal{A}_{\text{Weyl}}. \quad (28)$$

In SHBT the framing part of this anomaly is not an independent local curvature functional. It is the scalar obstruction left by the completed modular pairing:

$$\mathcal{A}_{\text{Weyl}}^{\text{fr}} = d \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}}, \quad \mathcal{E}_{ij}^{\partial} = \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{(0)ij}. \quad (29)$$

The unique covariant bulk continuation of this purely scalar boundary obstruction on a closed branch is therefore metric-proportional,

$$\boxed{\mathcal{E}_{\mu\nu} = \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{\mu\nu}}, \quad \mathcal{Q}_{\text{fr}} \neq 0. \quad (30)$$

Equivalently, the framing defect is a topological Weyl-anomaly source: if the integer center lifts close, the source vanishes; if they do not, the only allowed covariant obstruction is proportional to the metric.

The same tensor is defined by the completed SHBT stress accounting,

$$\mathcal{E}_{\mu\nu} \equiv \frac{c_{\text{dark}}^{\text{res}} M_P^2}{12} (G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu}) - T_{\mu\nu}^{\text{vis}}, \quad 8\pi G_N = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2}. \quad (31)$$

Combining Eqs. (30) and (31) gives the off-branch topological Einstein equation with a framing source,

$$G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2} (T_{\mu\nu}^{\text{vis}} + \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{\mu\nu}). \quad (32)$$

On the canonical branch, Eq. (24) gives $\Delta_{\text{fr}} = 0$, hence

$$\Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0. \quad (33)$$

Using Eq. (31), the vanishing condition is precisely

$$G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2} T_{\mu\nu}^{\text{vis}} = 8\pi G_N T_{\mu\nu}^{\text{vis}}. \quad (34)$$

Thus the complete boundary-to-bulk derivation chain is

$$\begin{aligned} [M, S_{\partial}] = [M, T_{\partial}] = 0 &\implies h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z} \implies \frac{K}{2k_{\ell}}, \frac{K}{3k_q} \in \mathbb{Z} \implies \Delta_{\text{fr}} = 0, \\ \Delta_{\text{fr}} = 0 &\implies \mathcal{A}_{\text{Weyl}}^{\text{fr}} = 0 \implies \mathcal{E}_{\mu\nu} = 0 \implies G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = 8\pi G_N T_{\mu\nu}^{\text{vis}}. \end{aligned} \quad (35)$$

The tensor equation is therefore not an independent tuning condition; it is the bulk representative of exact modular closure.

The final axiom is static global Hamiltonian cancellation. Let

$$H_{\text{total}} = H_{\partial} + H_{\text{bulk}} + H_{\text{stabilizer}}, \quad (36)$$

where H_{∂} acts on the completed boundary characters, H_{bulk} is the bulk image of the closed boundary data, and $H_{\text{stabilizer}}$ enforces the closure syndromes. The physical projector is defined by the simultaneous constraints

$$\Pi_{\text{phys}} M = M \Pi_{\text{phys}} = M, \quad P_{\text{fr}} \Pi_{\text{phys}} = 0, \quad \mathcal{E}_{\mu\nu} \Pi_{\text{phys}} = 0. \quad (37)$$

On this sector the stabilizer generator is fixed, not freely chosen:

$$H_{\text{stabilizer}} \Pi_{\text{phys}} = -(H_{\partial} + H_{\text{bulk}}) \Pi_{\text{phys}}. \quad (38)$$

Multiplying Eq. (36) by Π_{phys} on both sides gives

$$\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0. \quad (39)$$

Therefore, for every physical completed-boundary state satisfying $|\Psi_{\text{phys}}\rangle = \Pi_{\text{phys}} |\Psi_{\text{phys}}\rangle$,

$$\boxed{H_{\text{total}} |\Psi_{\text{phys}}\rangle = 0.} \quad (40)$$

This is the static-block statement that the completed physical sector has no independent external time generator.

The Rust/Python simulator mirrors these axioms through the `shbt_simulator.StaticBoundary` class. The constructor fixes the branch data; the visible modular phases are assembled by `shbt_simulator.StaticBoundary.build_su2_visible_phases` and `shbt_simulator.StaticBoundary.build_su3_visible_phases`; the finite partition check is performed by `shbt_simulator.StaticBoundary.evaluate_z_boundary`; the scalar obstruction is computed by `report.framing_defect`; and the closure chain, including $\Delta_{\text{fr}} = 0$, the Einstein lock, and global Hamiltonian cancellation, is audited by `shbt_simulator.ShbtSimulator().run_full_audit().boundary_report`.

3 Modular Data and Boundary Partition Function

This section makes explicit the modular data used by the completed boundary partition function. The general Wess–Zumino–Witten rule is that an affine factor G_k has integrable highest weights λ , modular character vector χ^{G_k} , dual Coxeter number $g^\vee$, central charge

$$c_{G_k} = \frac{k \dim G}{k + g^\vee}, \quad (41)$$

and diagonal modular- T entries

$$T_{\lambda\mu}^{G_k} = \delta_{\lambda\mu} \exp \left[2\pi i \left(h_\lambda^{G_k} - \frac{c_{G_k}}{24} \right) \right], \quad h_\lambda^{G_k} = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^\vee)}. \quad (42)$$

Here ρ is the Weyl vector and the inner product is normalized so long roots have length squared 2. The Rust/Python simulator stores the reduced visible phase $\exp(2\pi i h_\lambda)$ in `t_boundary`; the omitted scalar factor $\exp(-2\pi i c/24)$ is common to a fixed factor and cancels between the holomorphic and anti-holomorphic sectors, and it also leaves the commutator norms unchanged.

For $SU(2)_k$, the integrable labels are $a = 0, 1, \dots, k$. With spin $j = a/2$ and $C_2(j) = j(j+1) = a(a+2)/4$, Eq. (42) gives

$$h_a^{SU(2)_k} = \frac{a(a+2)}{4(k+2)}, \quad c_{SU(2)_k} = \frac{3k}{k+2}. \quad (43)$$

The A_1 Weyl sum has two terms and reduces to the sine kernel

$$S_{ab}^{SU(2)_k} = \sqrt{\frac{2}{k+2}} \sin \left(\frac{\pi(a+1)(b+1)}{k+2} \right), \quad T_{ab}^{SU(2)_k} = \delta_{ab} \exp \left[2\pi i \left(h_a^{SU(2)_k} - \frac{c_{SU(2)_k}}{24} \right) \right]. \quad (44)$$

For $SU(3)_k$, the integrable labels are Dynkin pairs (p, q) with $p, q \geq 0$ and $p+q \leq k$. The quadratic Casimir is

$$C_2(p, q) = \frac{p^2 + q^2 + pq + 3p + 3q}{3}, \quad (45)$$

so that

$$h_{(p,q)}^{SU(3)_k} = \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \quad c_{SU(3)_k} = \frac{8k}{k+3}. \quad (46)$$

It is useful to embed a Dynkin label in the traceless three-dimensional weight space by

$$v(p, q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, -\frac{p+2q}{3} \right), \quad \rho = v(1, 1) = (1, 0, -1). \quad (47)$$

Then the A_2 Weyl-sum formula used by the implementation is

$$S_{(p,q),(r,s)}^{SU(3)_k} = \frac{i^3}{\sqrt{3}(k+3)} \sum_{\sigma \in S_3} \text{sgn}(\sigma) \exp \left[-\frac{2\pi i}{k+3} (\sigma(v(p, q) + \rho), v(r, s) + \rho) \right], \quad (48)$$

with

$$T_{(p,q),(r,s)}^{SU(3)_k} = \delta_{(p,q),(r,s)} \exp \left[2\pi i \left(h_{(p,q)}^{SU(3)_k} - \frac{c_{SU(3)_k}}{24} \right) \right]. \quad (49)$$

For the parent completion $SO(10)_K$, the Lie algebra is D_5 , with $g^\vee = 8$, rank 5, $\dim SO(10) = 45$, and 20 positive roots. Thus

$$c_{SO(10)_K} = \frac{45K}{K+8}. \quad (50)$$

If Λ and M are integrable D_5 highest weights, the corresponding formal kernels are

$$\begin{aligned} S_{\Lambda M}^{SO(10)_K} &= \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \text{sgn}(w) \exp \left[-\frac{2\pi i}{K+8} (w(\Lambda + \rho), M + \rho) \right], \\ T_{\Lambda M}^{SO(10)_K} &= \delta_{\Lambda M} \exp \left[2\pi i \left(\frac{(\Lambda, \Lambda + 2\rho)}{2(K+8)} - \frac{45K}{24(K+8)} \right) \right]. \end{aligned} \quad (51)$$

The finite audit does not enumerate the full $SO(10)_{312}$ character table; instead, this parent factor enters through the branch locks, the framing-defect test, and the central charge in Eq. (50).

The affine data on the completed benchmark branch are summarized in Table 1. The visible central charge is the sum of the first two visible factors; the completed ledger is the finite-screen dark contribution used by the Noether bridge and cosmology.

Affine factor	Level	Central charge	Center data
$SU(2)_{k_\ell}$	$k_\ell = 26$	$\frac{39}{14} = 2.785714285714$	$\mathbb{Z}_2$ spin parity
$SU(3)_{k_q}$	$k_q = 8$	$\frac{64}{11} = 5.818181818182$	$\mathbb{Z}_3$ color triality
$SO(10)_K$	$K = 312$	$\frac{351}{8} = 43.875000000000$	$\mathbb{Z}_4$ spinor center
$\mathcal{H}_{\text{dark}}$	—	$c_{\text{dark}}^{\text{comp}} = \frac{1197103}{362670} = 3.300805139659$	—

Table 1: Affine algebraic parameters and central charges of the completed boundary CFT factors on the canonical branch.

At the benchmark branch, the visible block evaluated in code is the 3×3 slice

$$\text{charge_embedding} = (k_\ell - 4, k_\ell - 3, k_\ell) = (22, 23, 26), \quad (52)$$

$$\text{LOW_SU3_WEIGHTS} = ((0, 0), (1, 0), (0, 1)). \quad (53)$$

This slice is not the full completed Hilbert space. It is the visible finite-dimensional block on which `shbt_simulator.StaticBoundary.s_boundary`, `shbt_simulator.StaticBoundary.t_boundary`, and `shbt_simulator.StaticBoundary.evaluate_z_boundary` perform the numerical audit. The combined visible index is

$$A = (i, j), \quad a_i \in (22, 23, 26), \quad w_j \in ((0, 0), (1, 0), (0, 1)), \quad (54)$$

so the visible audit space has 9 basis entries. In that ordering,

$$S_{\partial}^{\text{code}} = \text{Re } S_{SU(2)_{26}}^{\text{vis}} \otimes \text{Re } S_{SU(3)_8}^{\text{vis}}, \quad T_{\partial}^{\text{code}} = \text{diag} \left(T_{SU(2)_{26}}^{\text{vis}} \otimes T_{SU(3)_8}^{\text{vis}} \right), \quad (55)$$

which is exactly the construction of the `s_boundary` and `t_boundary` properties.

The benchmark central charges and conformal weights are

$$\begin{aligned}
c_{SU(2)_{26}} &= \frac{39}{14} = 2.785714285714, & c_{SU(3)_8} &= \frac{64}{11} = 5.818181818182, \\
c_{SO(10)_{312}} &= \frac{351}{8} = 43.875, & c_{\text{vis}} &= \frac{1325}{154} = 8.603896103896, \\
h_{22}^{SU(2)_{26}} &= \frac{33}{7} = 4.714285714286, & h_{23}^{SU(2)_{26}} &= \frac{575}{112} = 5.133928571429, \\
h_{26}^{SU(2)_{26}} &= \frac{13}{2} = 6.5, & h_{(0,0)}^{SU(3)_8} &= 0, \\
h_{(1,0)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212, & h_{(0,1)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212.
\end{aligned} \tag{56}$$

The visible S blocks produced by `shbt_simulator.StaticBoundary.build_su2_visible_block` and `shbt_simulator.StaticBoundary.build_su3_visible_block` are

$$\begin{aligned}
S_{SU(2)_{26}}^{\text{vis}} &\simeq \begin{pmatrix} 0.088270792276 & -0.208953252971 & 0.142191553507 \\ -0.208953252971 & 0.260560444585 & -0.115960306962 \\ 0.142191553507 & -0.115960306962 & 0.029923764933 \end{pmatrix}, \\
\text{Re } S_{SU(3)_8}^{\text{vis}} &\simeq \begin{pmatrix} 0.018018539201 & 0.048334858719 & 0.048334858719 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \end{pmatrix}.
\end{aligned} \tag{58}$$

The imaginary parts of the displayed $SU(3)$ entries are at roundoff scale for the first row and are conjugate in the fundamental/anti-fundamental subblock; the implemented `s_boundary` uses the real part shown above. The reduced visible T phases used by the code are

$$\begin{aligned}
T_{SU(2)_{26}}^{\text{vis}} &\simeq \begin{pmatrix} -0.222520933956 - 0.974927912182i \\ 0.666346577952 + 0.745642164883i \\ -1.000000000000 \end{pmatrix}, \\
T_{SU(3)_8}^{\text{vis}} &\simeq \begin{pmatrix} 1 \\ 0.723734038105 + 0.690079011482i \\ 0.723734038105 + 0.690079011482i \end{pmatrix}.
\end{aligned} \tag{59}$$

The completed boundary partition function is the torus pairing of completed characters,

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\tau)}, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}, \tag{60}$$

where $I = (a, (p, q), \Lambda, \alpha)$ includes the $SU(2)$ label, $SU(3)$ label, $SO(10)$ parent label, and dark-sector label. In vector notation this is $Z_{\partial} = \chi^{\dagger} M \chi$, up to the conventional placement of the holomorphic index. Since the character vector transforms as

$$\chi(\tau + 1) = T_{\partial} \chi(\tau), \quad \chi(-1/\tau) = S_{\partial} \chi(\tau), \tag{61}$$

modular invariance follows from

$$[M, T_{\partial}] = 0, \quad [M, S_{\partial}] = 0. \tag{62}$$

Indeed,

$$Z_{\partial}(\tau + 1, \bar{\tau} + 1) = \chi(\tau)^{\dagger} T_{\partial}^{\dagger} M T_{\partial} \chi(\tau) = \chi(\tau)^{\dagger} M \chi(\tau), \tag{63}$$

$$Z_{\partial}(-1/\tau, -1/\bar{\tau}) = \chi(\tau)^{\dagger} S_{\partial}^{\dagger} M S_{\partial} \chi(\tau) = \chi(\tau)^{\dagger} M \chi(\tau), \tag{64}$$

using unitarity of the modular kernels. Therefore

$$\boxed{Z_{\partial}(\tau + 1, \bar{\tau} + 1) = Z_{\partial}(\tau, \bar{\tau}), \quad Z_{\partial}(-1/\tau, -1/\bar{\tau}) = Z_{\partial}(\tau, \bar{\tau}).} \quad (65)$$

The completed pairing matrices are therefore classified by the finite commutant of the two modular generators,

$$\mathfrak{M}_{\partial} = \{M \in \text{Mat}_N(\mathbb{Z}_{\geq 0}) \mid M_{00} = 1, [M, S_{\partial}] = 0, [M, T_{\partial}] = 0\}. \quad (66)$$

Writing

$$\theta_I = \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right], \quad (T_{\partial})_{IJ} = \theta_I \delta_{IJ}, \quad (67)$$

the T commutator is

$$([M, T_{\partial}])_{IJ} = M_{IJ}(\theta_J - \theta_I). \quad (68)$$

Hence every nonzero pairing lies inside a common modular-spin sector,

$$M_{IJ} \neq 0 \implies \theta_I = \theta_J \iff h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \quad (69)$$

This is the integral-spin closure condition of the completed CFT. The canonical locks in Eq. (22) refine this condition by resolving the visible center lifts: an admissible visible pairing must preserve both the $SU(2)$ spin-parity lift and the $SU(3)$ color-triality lift into the $SO(10)_{312}$ parent. Pairings that mix different lifted center classes would change one of the integer quotients $K/(2k_{\ell})$ or $K/(3k_q)$ and therefore contribute directly to Δ_{fr} .

On the 9-state visible block this leaves only pointwise pairings after the center-lift classes have been resolved. Thus the visible restriction has the form

$$M_{\text{vis}} = \text{diag}(m_1, \dots, m_9), \quad m_A \in \mathbb{Z}_{\geq 0}. \quad (70)$$

The remaining S constraint forces all diagonal multiplicities to coincide:

$$([M_{\text{vis}}, S_{\partial}^{\text{code}}])_{AB} = (m_A - m_B)(S_{\partial}^{\text{code}})_{AB} = 0. \quad (71)$$

Every entry of the displayed benchmark $S_{\partial}^{\text{code}}$ is nonzero because the visible $SU(2)_{26}$ and real $SU(3)_8$ subblocks in Eq. (58) have no zero entries. Therefore $m_A = m_B$ for all A, B , and the vacuum normalization $M_{00} = 1$ fixes the common value to one:

$$\boxed{M_{\text{vis}} = M_{\text{code}} = \mathbb{1}_9.} \quad (72)$$

This is the unique center-resolved visible-block representative in $\mathfrak{M}_{\partial}$. It keeps the visible sector pointwise paired at the self-dual point $\tau = i$ and avoids adding a spin-parity or color-triality framing defect.

The finite Rust/Python partition audit takes $M_{\text{code}} = \mathbb{1}_9$ and the leading visible character monomials

$$\chi_{(i,j)}^{\text{code}}(\tau) = q^{h_{a_i}^{SU(2)_{26}} + h_{w_j}^{SU(3)_8} - c_{\text{vis}}/24}, \quad q = e^{2\pi i \tau}, \quad c_{\text{vis}} = \frac{1325}{154}. \quad (73)$$

Thus `shbt_simulator.StaticBoundary.evaluate_z_boundary` evaluates

$$Z_{\partial}^{\text{code}}(\tau) = \overline{\chi^{\text{code}}(\tau)}^T M_{\text{code}} \chi^{\text{code}}(\tau). \quad (74)$$

For $\tau = i$ the code gives

$$\begin{aligned} Z_{\partial}^{\text{code}}(i) &= 2.441381789163 \times 10^{-24}, \\ Z_{\partial}^{\text{code}}(i+1) &= 2.441381789163 \times 10^{-24}, \end{aligned} \tag{75}$$

$$Z_{\partial}^{\text{code}}(-1/i) = 2.441381789163 \times 10^{-24}. \tag{76}$$

The equality under the S move in this numerical line is pointwise because i is the self-dual point of $\tau \mapsto -1/\tau$; the nontrivial finite-block audit of the two modular generators is the matrix commutator test. The `shbt_simulator.ShbtSimulator().run_full_audit().boundary_report` record reports

Audit quantity	Value
<code>s_boundary.shape</code>	(9, 9)
<code>t_boundary.shape</code>	(9, 9)
<code>z_boundary_matrix</code>	$\not\sim_9$
<code>modular_S_commutator_norm</code>	0.0
<code>modular_T_commutator_norm</code>	0.0
<code>modular_invariant</code>	True
<code>integral_spin_closure</code>	True
<code>all_checks_passed</code>	True

Thus the implemented boundary block satisfies

$$\|M_{\text{code}} S_{\partial}^{\text{code}} - S_{\partial}^{\text{code}} M_{\text{code}}\|_F = 0, \quad \|M_{\text{code}} T_{\partial}^{\text{code}} - T_{\partial}^{\text{code}} M_{\text{code}}\|_F = 0, \tag{77}$$

which is the finite visible-block realization of the modular closure condition used in the full SHBT boundary partition function.

4 Boundary Entropy Densities and Self-Resolution

The modular data of Section 3 also determine how the finite visible register is loaded. The relevant coordinate lattice is the 3×3 visible block

$$\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}, \quad c = (i, j) \in \mathcal{C}. \tag{78}$$

The two axes of this lattice are the displayed $SU(2)_{26}$ charge slice and the displayed $SU(3)_8$ low-weight slice. We write

$$q_i, \ell_j \in (22, 23, 26), \quad w_i, w_j \in ((0, 0), (1, 0), (0, 1)), \tag{79}$$

where q_i and ℓ_j are the row and column labels of the same visible $SU(2)$ block. The raw boundary loading assigned to $c = (i, j)$ is the product of the modular overlap probabilities in the two visible factors,

$$\tilde{\rho}_B(i, j) = \left| S_{q_i, \ell_j}^{SU(2)_{26}} \right|^2 \left| S_{w_i, w_j}^{SU(3)_8} \right|^2. \tag{80}$$

This is positive on the benchmark visible block, so the finite normalization constant

$$Z_B^{\text{load}} = \sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a,b) \quad (81)$$

is nonzero. The normalized loading density is therefore

$$\rho_B(i,j) = \frac{\tilde{\rho}_B(i,j)}{\sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a,b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_B(i,j) = 1. \quad (82)$$

The associated entanglement density is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i,j) = \frac{-\rho_B(i,j) \ln \rho_B(i,j)}{\sum_{(a,b) \in \mathcal{C}} -\rho_B(a,b) \ln \rho_B(a,b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_E(i,j) = 1. \quad (83)$$

The convention $0 \ln 0 = 0$ applies, although no zero entry occurs in the benchmark block.

The dominant loading sequence is the ordering of coordinates by decreasing normalized loading density,

$$\sigma = (c_1, \dots, c_9), \quad \rho_B(c_1) \geq \rho_B(c_2) \geq \dots \geq \rho_B(c_9). \quad (84)$$

Equal-density ties are physically degenerate and are resolved in the stable implementation order. This ordering is the audit field `dominant_loading_sequence`; it must agree with the coordinate order used by `sequence_bit_loading`. Equations (80)–(84) are implemented by the following audit helpers:

- `shbt_simulator.StaticBoundary.build_loading_density`,
- `shbt_simulator.StaticBoundary.build_entanglement_density`, and
- `shbt_simulator.StaticBoundary.build_dominant_sequence`.

Entropy self-resolution promotes the static loading order into an iterative finite-register update. Let $U_n \subset \mathcal{C}$ be the unresolved set at step n , with $c_n \in U_n$ the next coordinate selected by σ . Before the update, the cumulative resolved entropy state is

$$\Omega_{n-1}(i,j) = \sum_{m < n} \delta_{c_m, (i,j)} \frac{\Delta S_m}{N}. \quad (85)$$

The resolved state feeds back into each unresolved coordinate through the Manhattan-distance kernel

$$F_{n-1}(c) = \sum_{(i,j) \in \mathcal{C}} \frac{\Omega_{n-1}(i,j)}{1 + |i - c_i| + |j - c_j|}, \quad c = (c_i, c_j). \quad (86)$$

For feedback coupling $\lambda \geq 0$, the loading weight is dressed as

$$w_B(c) = \rho_B(c) (1 + \lambda F_{n-1}(c)). \quad (87)$$

The entropy channel is dressed by the same feedback kernel,

$$w_E(c) = \rho_E(c) (1 + \lambda F_{n-1}(c)). \quad (88)$$

With remaining bit and entropy budgets

$$R_{n-1}^B = N - \sum_{m < n} \Delta B_m, \quad R_{n-1}^E = N - \sum_{m < n} \Delta S_m, \quad (89)$$

the self-resolution update is

$$\Delta B_n = R_{n-1}^B \frac{w_B(c_n)}{\sum_{c \in U_n} w_B(c)},$$

$$\Delta S_n = R_{n-1}^E \frac{w_E(c_n)}{\sum_{c \in U_n} w_E(c)}, \quad (90)$$

$$\Delta T_n = \frac{\Delta S_n}{N}. \quad (91)$$

Thus ΔT_n is not an independent background clock tick. It is the dimensionless entropy residue left by resolving the next boundary coordinate. This update is the mathematical contract of `shbt_simulator.StaticBoundary.entropy_self_resolution()`, whose audit record stores `feedback_signal`, `delta_B`, `delta_S`, `delta_T`, and the updated unresolved set.

The dimensional projection is driven by the cumulative temporal residue. At step n the effective projection dimension is

$$D_n = 26 - 22 \sum_{m \leq n} \Delta T_m, \quad \Delta D_n = D_{n-1} - D_n = 22 \Delta T_n. \quad (92)$$

Since U_9 contains only the final unresolved coordinate, the last update spends the full remaining entropy budget. By induction on Eq. (90),

$$\sum_{n=1}^9 \Delta S_n = N, \quad \sum_{n=1}^9 \Delta T_n = 1. \quad (93)$$

Consequently the benchmark projection starts at $D_0 = 26$ and terminates at

$$D_9 = 26 - 22 = 4. \quad (94)$$

This proves the audit predicate `projection_dimension_26_to_4`: it holds precisely when the self-resolution run exhausts the normalized entropy budget and reports terminal dimension 4.

The same normalization gives the temporal kernel used for observer-frame cosmology. For a sampled expansion rate

$$H(t) = \frac{\dot{a}(t)}{a(t)}, \quad (95)$$

the total entropy-production rate is postulated to saturate the finite register as

$$\dot{S}_{\text{total}} = N H(t). \quad (96)$$

The local entanglement contribution is distributed by the normalized density,

$$\dot{S}_{ij} = \rho_E(i, j) \dot{S}_{\text{total}}. \quad (97)$$

Therefore

$$\dot{T} = \frac{1}{N} \sum_{(i,j) \in \mathcal{C}} \dot{S}_{ij} = \frac{\dot{S}_{\text{total}}}{N} \sum_{(i,j) \in \mathcal{C}} \rho_E(i, j) = H(t). \quad (98)$$

This is the perception identity: apparent temporal flow is the entropy flow per finite boundary bit. In the implementation it is audited by `shbt_simulator.StaticBoundary.derive_temporal_increment()`, with fields `local_entanglement_entropy_gradient`, `dot_S_total`, `dot_T`, and `perception_identity_holds`.

The full Section 4 implementation contract is therefore

Mathematical object	Rust/Python simulator object
ρ_B	<code>loading_density</code>
ρ_E	<code>entanglement_density</code>
σ	<code>dominant_loading_sequence</code>
F_{n-1}	<code>feedback_signal</code>
$(\Delta B_n, \Delta S_n, \Delta T_n)$	<code>entropy_self_resolution()</code>
$D_9 = 4$	<code>projection_dimension_26_to_4</code>
$\dot{T} = H(t)$	<code>derive_temporal_increment()</code>

The boundary entropy densities are therefore not phenomenological weights added after modular closure. They are the normalized modular-overlap probabilities of the finite visible block, and their self-resolution produces both the $26 \rightarrow 4$ dimensional projection and the local temporal kernel.

5 Holographic RG Flow: Boundary Entropy to Bulk Metric

The standard holographic starting point is the geometric dictionary for boundary entanglement. For a boundary subregion A , the Ryu–Takayanagi relation identifies its entanglement entropy with the area of the bulk minimal codimension-two surface γ_A homologous to A ,

$$S(A) = \frac{\text{Area}(\gamma_A)}{4G_{d+1}}. \quad (99)$$

This formula supplies the role of entanglement in the RG flow: changing the boundary resolution changes which extremal surfaces are sampled, and therefore which radial portion of the bulk geometry is probed [14, 11, 12, 13].

In an asymptotically anti-de Sitter geometry, the radial coordinate is the holographic renormalization scale. With the Fefferman–Graham form [6]

$$ds^2 = d\rho^2 + e^{2\rho/L} g_{ij}(x, \rho) dx^i dx^j = \frac{L^2}{z^2} (dz^2 + g_{ij}(x, z) dx^i dx^j), \quad (100)$$

the two common radial variables obey

$$z = L e^{-\rho/L}, \quad \frac{\mu(\rho)}{\mu_0} = e^{\rho/L} = \frac{L}{z}. \quad (101)$$

Thus moving outward toward the boundary corresponds to the ultraviolet of the dual CFT, while integrating out high-energy boundary modes corresponds to translating inward along the bulk radial direction. In the standard Hamilton–Jacobi formulation, the on-shell bulk action generates the boundary functional and radial evolution becomes a Callan–Symanzik flow,

$$\frac{\partial S_{\text{os}}}{\partial \rho} + \mathcal{H}_{\text{grav}} \left(\gamma_{ij}, \phi^I, \frac{\delta S_{\text{os}}}{\delta \gamma_{ij}}, \frac{\delta S_{\text{os}}}{\delta \phi^I} \right) = 0, \quad \mu \frac{dJ^I}{d\mu} = \beta^I(J). \quad (102)$$

This is the standard radial/Hamilton–Jacobi dictionary used in holographic renormalization [15, 16]. SHBT keeps the entanglement-to-geometry lesson of Eq. (99), but replaces the continuous large- N radial dynamics of Eq. (102) with a finite boundary-first projection. The input is the entropy self-resolution cascade of Section 4; the output is a trace-normalized static metric slice. In this section $\tau \in [0, 1]$ denotes the normalized entropy-resolution parameter, not an independent external time coordinate. At the discrete audit steps,

$$\tau_n = \sum_{m \leq n} \Delta T_m = \frac{1}{N} \sum_{m \leq n} \Delta S_m, \quad \tau_0 = 0, \quad \tau_9 = 1. \quad (103)$$

Thus the RG parameter is the same finite-register entropy residue that drives the $26 \rightarrow 4$ projection.

The finite SHBT replacement for the continuum boundary subregion is the visible coordinate lattice

$$\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}, \quad c = (i, j) \in \mathcal{C}. \quad (104)$$

The coordinates label the displayed $SU(2)_{26}$ charge slice and the $SU(3)_8$ low-weight slice. The raw boundary loading at $c = (i, j)$ is the product of the modular overlap probabilities in those two affine factors,

$$\tilde{\rho}_B(i, j) = \left| S_{q_i, \ell_j}^{SU(2)_{26}} \right|^2 \left| S_{w_i, w_j}^{SU(3)_8} \right|^2. \quad (105)$$

For the benchmark branch, the explicit kernels used in this loading are

$$S_{ab}^{SU(2)_{26}} = \sqrt{\frac{2}{28}} \sin \left(\frac{\pi(a+1)(b+1)}{28} \right), \quad S_{(p,q),(r,s)}^{SU(3)_8} = \frac{i^3}{\sqrt{3} 11} \sum_{\sigma \in S_3} \text{sgn}(\sigma) \exp \left[-\frac{2\pi i}{11} (\sigma(\mathbf{v}(p, q) + \varrho), \mathbf{v}(r, s) + \varrho) \right], \quad (106)$$

where $\mathbf{v}(p, q)$ is the weight vector with Dynkin labels (p, q) and ϱ is the Weyl vector. Normalizing the raw loading gives

$$\rho_B(i, j) = \frac{\tilde{\rho}_B(i, j)}{\sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a, b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_B(i, j) = 1. \quad (107)$$

The entropy density used by the RG projector is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i, j) = \frac{-\rho_B(i, j) \ln \rho_B(i, j)}{\sum_{(a,b) \in \mathcal{C}} -\rho_B(a, b) \ln \rho_B(a, b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_E(i, j) = 1. \quad (108)$$

Equations (104)–(108) are the Section 4 density definitions repeated here to make the RG derivation self-contained.

The prime lattice basis used by the metric projector is the five-point arithmetic skeleton

$$(p_0, p_1, p_2, p_3, p_4) = (2, 3, 5, 7, 11). \quad (109)$$

The choice of the prime basis $(2, 3, 5, 7, 11)$ is not a numerical coincidence or a compression trick. In information-theoretic terms, primes are the irreducible generators of the positive integers under multiplication; they represent the minimal set of atomic information channels required to encode the full arithmetic capacity of the finite register. The modulo-5 folding of the 9-component ordered entropy sequence onto this prime basis ensures that each residue class receives contributions from linearly independent arithmetic factors of the boundary load. This prevents spectral aliasing in the subsequent Gram projection: if the folding were performed onto a composite basis, the Euler flux Φ_s would suffer from redundant factor cancellations that could wash out entropy shear. The prime basis therefore provides a maximal-entropy encoding of the ordered boundary state into a five-component load vector $\ell_r(\tau)$, where the logarithmic radial RG coordinates $x_r = \ln p_r$ are naturally ordered and have distinct prime-power scaling. This construction can be seen as a finite-register analogue of the continuous radial coordinate ρ in standard holographic RG, with the prime exponents playing the role of discrete energy scales. The specific set $(2, 3, 5, 7, 11)$ is the smallest set of five primes whose mutual logarithmic differences span the typical entropy-gradient range of the visible block; replacing it with any other set of five primes changes the numerical values of the metric slices but preserves the closure structure, as long as the selected primes are distinct and ordered.

Its logarithms are the radial RG coordinates,

$$x_r = (\ln p_r), \quad r = 0, \dots, 4, \quad (110)$$

so nearest-neighbor finite differences on this lattice are naturally divided by $\ln(p_{s+1}/p_s) = x_{s+1} - x_s$. The role of the prime basis is not to introduce a new spectrum of particles. It is a fixed, finite coordinate frame in which the ordered boundary entropy state can be compressed into four Euler intervals and then lifted to a symmetric metric tensor.

Let

$$\sigma = (c_1, \dots, c_9) \quad (111)$$

be the dominant loading sequence of Eq. (84). The partially resolved entanglement state at RG parameter τ is the interpolation of the discrete self-resolution records,

$$\Omega_\tau(c) = \sum_{n=1}^9 \theta_n(\tau) \delta_{c_n, c} \Delta T_n, \quad (112)$$

where

$$\theta_n(\tau) = \begin{cases} 0, & \tau < \tau_{n-1}, \\ \frac{\tau - \tau_{n-1}}{\tau_n - \tau_{n-1}}, & \tau_{n-1} \leq \tau \leq \tau_n, \\ 1, & \tau > \tau_n. \end{cases} \quad (113)$$

Consequently $\sum_{c \in \mathcal{C}} \Omega_\tau(c) = \tau$. The metric projector uses both the static entanglement density and the resolved entropy state. Along the ordered sequence define

$$E_m(\tau) = \rho_E(c_m) + \Omega_\tau(c_m), \quad m = 1, \dots, 9. \quad (114)$$

Since $\sum_{m=1}^9 \rho_E(c_m) = 1$ and $\sum_{m=1}^9 \Omega_\tau(c_m) = \tau$, the total resolved load before folding is

$$\sum_{m=1}^9 E_m(\tau) = 1 + \tau. \quad (115)$$

The five-component load vector is obtained by folding this ordered sequence onto the prime lattice by residue class,

$$\begin{aligned} \widehat{\ell}_r(\tau) &= \sum_{m=1}^9 E_m(\tau) \mathbf{1}_{\{m-1 \equiv r \pmod{5}\}}, \quad r = 0, \dots, 4, \\ \ell_r(\tau) &= \frac{\widehat{\ell}_r(\tau)}{\sum_{q=0}^4 \widehat{\ell}_q(\tau)} = \frac{\widehat{\ell}_r(\tau)}{1 + \tau}. \end{aligned} \quad (116)$$

This gives

$$\ell_r(\tau) > 0, \quad \sum_{r=0}^4 \ell_r(\tau) = 1. \quad (117)$$

The dependence on σ is essential: the same multiset of boundary entropies would give a different prime-lattice load only if the modular loading order were changed. Equal-density ties remain the stable implementation ties already fixed in Section 4.

The Euler flux is the logarithmic finite difference of the load vector across adjacent prime sites. For $s = 0, 1, 2, 3$,

$$\Phi_s(\tau) = \frac{\ell_{s+1}(\tau) - \ell_s(\tau)}{\ln(p_{s+1}/p_s)}. \quad (118)$$

Thus Φ_s is the discrete RG gradient $\Delta \ell / \Delta \ln p$ on the finite prime lattice. The local mean loading and arithmetic hardness factor are

$$\bar{\ell}_s(\tau) = \frac{\ell_s(\tau) + \ell_{s+1}(\tau)}{2}, \quad h_s = \frac{1}{p_s} + \frac{1}{p_{s+1}}. \quad (119)$$

The closure-weighted contribution of the s th Euler interval is

$$W_s(\tau) = \lambda_{\text{closure}} (|\Phi_s(\tau)| + \bar{\ell}_s(\tau) + h_s), \quad \lambda_{\text{closure}} > 0. \quad (120)$$

Every term in Eq. (120) is nonnegative, so the outer-product part of the metric projection is positive semidefinite. The absolute flux measures local entropy shear, the mean load measures local occupancy, and h_s prevents a completely flat load vector from erasing the arithmetic interval.

The four execution vectors embed the four Euler intervals into the static metric block. Let $e_s^\mu = \delta_s^\mu$ for $\mu, s = 0, 1, 2, 3$, and read all load-vector indices modulo 5. The implemented execution vector is

$$v_s^\mu(\tau) = e_s^\mu + (\ell_s + \delta_{s0}, \ell_{s+1} + \delta_{s1}, \ell_{(s+2) \pmod{5}} + \delta_{s2}, |\Phi_s| + \ell_{(s+3) \pmod{5}} + \delta_{s3})^\mu. \quad (121)$$

The Kronecker terms mark the execution channel, while the cyclic load entries couple each interval to the neighboring prime sites. The fourth component also contains $|\Phi_s|$, so entropy shear enters the block as a metric direction rather than only as a scalar weight.

The unstabilized static metric is the weighted Gram matrix of these execution vectors, plus a positive diagonal closure floor,

$$\tilde{g}_{\mu\nu}(\tau) = \sum_{s=0}^3 W_s(\tau) v_{s\mu}(\tau) v_{s\nu}(\tau) + \text{diag}(d_0, d_1, d_2, d_3)_{\mu\nu}. \quad (122)$$

The diagonal entries used by the audit are

$$d_s(\tau) = \lambda_{\text{closure}} + W_s(\tau) + \ell_s(\tau) + \ell_{s+1}(\tau), \quad s = 0, 1, 2, 3. \quad (123)$$

Since $\lambda_{\text{closure}} > 0$ and the remaining terms are nonnegative, Eq. (123) gives a strictly positive diagonal regularizer. The tensor in Eq. (122) is therefore already a finite, positive candidate for the static block; the final step enforces exact symmetry and trace normalization in the same way for every slice.

Define

$$A_{\mu\nu}(\tau) = \text{Sym}(\tilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}} \delta_{\mu\nu}), \quad \text{Sym}(B)_{\mu\nu} = \frac{B_{\mu\nu} + B_{\nu\mu}}{2}. \quad (124)$$

The eigenvalue lift is chosen as

$$\epsilon_{\text{eig}} = \max(0, \epsilon_{\text{pos}} - \lambda_{\min}(\text{Sym} \tilde{g})), \quad \epsilon_{\text{pos}} > 0. \quad (125)$$

In exact arithmetic the positive diagonal floor usually makes the lift unnecessary, but the audit keeps Eq. (125) so that numerical roundoff cannot produce a nonpositive metric slice. The physical static metric is

$$g_{\mu\nu}(\tau) = \frac{A_{\mu\nu}(\tau)}{\text{Tr } A(\tau)} = \frac{\text{Sym}(\tilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}} \delta_{\mu\nu})}{\text{Tr}[\text{Sym}(\tilde{g}(\tau) + \epsilon_{\text{eig}} \mathbf{1}_4)]}. \quad (126)$$

It obeys the metric-slice checks

$$g_{\mu\nu} = g_{\nu\mu}, \quad \text{Tr } g = 1, \quad \lambda_{\min}(g) > 0. \quad (127)$$

The trace condition fixes the finite-register scale of the slice; it does not remove physical anisotropy, because anisotropy is carried by the normalized eigenvalue ratios and off-diagonal components inherited from the execution vectors.

The spatial bulk metric is the projection of the static 4×4 block onto the three rendered spatial directions. With

$$P_a{}^\mu = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a = 1, 2, 3, \quad \mu = 0, 1, 2, 3, \quad (128)$$

the induced spatial metric is

$$g_{ab}^{\text{bulk}}(\tau) = P_a{}^\mu g_{\mu\nu}^{\text{static}}(\tau) P_b{}^\nu. \quad (129)$$

Equivalently, the spatial block is the lower-right 3×3 principal block of the stabilized static metric in this benchmark projector. Other projectors could rotate the rendered spatial frame, but the audit uses the fixed matrix in Eq. (128) so that the benchmark is deterministic.

Combining the construction gives the complete holographic RG map

$$\boxed{\nabla_C S_\partial \mapsto \rho_E(i, j) \mapsto \ell_r(\tau) \mapsto \Phi_s(\tau), W_s(\tau), v_s^\mu(\tau) \mapsto g_{\mu\nu}(\tau) \mapsto g_{ab}^{\text{bulk}}(\tau)}. \quad (130)$$

The shortened form is

$$\nabla_C S_{\partial} \longrightarrow \rho_E(i, j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau). \quad (131)$$

Every arrow is fixed by the previous sections: the boundary entropy gradient comes from modular-overlap loading, ρ_E is the normalized entanglement density, ℓ_r is the prime-lattice compression of the ordered resolved state, and $g_{\mu\nu}$ is the stabilized trace-one Gram projection. No independent bulk equation of motion is inserted between the boundary entropy state and the static metric slice.

This is the precise deviation from standard holographic RG. The usual AdS/CFT construction treats the bulk metric as a dynamical saddle governed by the Einstein–Hilbert action; the boundary scale dependence is recovered by solving radial Hamilton–Jacobi flow equations. SHBT instead uses the closed finite boundary register as the primitive object. The replacement is

$$(S(A), \rho, S_{os})_{\text{standard}} \longrightarrow (\rho_B, \rho_E, \Omega_\tau, x_r = \ln p_r)_{\text{SHBT}}, \quad \partial_\rho \longrightarrow \Delta_{\ln p}. \quad (132)$$

Thus Eq. (102) is not solved as an independent bulk field equation. Its SHBT analogue is the finite algebraic pipeline

$$(\rho_B, \rho_E, \Omega_\tau) \longmapsto \ell_r(\tau) \longmapsto \Phi_s(\tau) \longmapsto \tilde{g}_{\mu\nu}(\tau) \longmapsto g_{\mu\nu}(\tau), \quad (133)$$

with closure, positivity, and trace normalization checked slice by slice.

Feature	Standard holographic RG	SHBT RG flow
Primitive data	Continuum boundary QFT and classical bulk saddle	Completed finite boundary CFT and modular-overlap densities
Radial scale	Continuous ρ or z , with $\mu \propto e^{\rho/L} \propto z^{-1}$	Discrete prime sites $x_r = \ln p_r$ and Euler intervals $\Delta_{\ln p}$
Entanglement role	RT surfaces measure entropy from a given bulk metric	Normalized entropy densities synthesize the metric slice
Dynamics	Einstein–Hilbert equations recast as Hamilton–Jacobi flow	Deterministic entropy self-resolution and weighted Gram projection
Deviation	Large- N semiclassical bulk is assumed	No independent bulk dynamics is assumed; geometry is an audited projection

The implementation contract for this section is

Mathematical object	Rust/Python simulator object
RG projection controller	<code>shbt_simulator.HolographicProjection</code>
$\rho_E, \sigma, \Omega_\tau \mapsto \ell_r$	<code>derive_load_vector()</code>
$\ell_r \mapsto \Phi_s, W_s, v_s^\mu, g_{\mu\nu}$	<code>metric_from_load_vector()</code>
$P_a^\mu g_{\mu\nu} P_b^\nu$	<code>project_static_block_to_bulk()</code>
Symmetry, trace, positivity, spatial block	<code>run_full_audit().projection_report</code>
Metric slice record	<code>shbt_simulator.BulkMetricSlice</code>

The class `shbt_simulator.HolographicProjection` receives the finite entropy cascade from `shbt_simulator.StaticBoundary.entropy_self_resolution()`. Its

`derive_load_vector()` method constructs Eq. (116); `metric_from_load_vector()` constructs Eqs. (118)–(126); `project_static_block_to_bulk()` applies Eq. (129); and `shbt_simulator.ShbtSimulator().run_full_audit().projection_report` checks Eq. (127), the 3×3 shape of `spatial_metric`, agreement with the dominant loading sequence, and the absence of negative eigenvalues after stabilization. The benchmark therefore exports a list of `shbt_simulator.BulkMetricSlice` records whose fields include `load_vector`, `euler_flux`, `execution_vectors`, `unstabilized_metric`, `metric_components`, `spatial_metric`, and `eigenvalues`.

6 Topological Baryogenesis and Anti-Baryon De-Rendering

The baryogenesis sector uses the completed boundary branch to separate two questions that are usually entangled. The first is the size of the visible baryon excess. The second is the fate of the anti-baryon stress-energy that would ordinarily accompany visible anti-baryon charge. SHBT answers the first question with a topological CP invariant and a modular restoration scale, and it answers the second with an active de-rendering map that strips visible color and electromagnetic charge while preserving the passive gravitational source. The mechanism is therefore not a dynamical erasure of stress-energy. It is a projection of active render channels into the dark completion sector, constrained by the same framing closure that fixes the branch.

The visible electroweak conversion factor is the standard sphaleron coefficient for the rendered Standard Model channel,

$$C_{\text{sph}} = \frac{28}{79}. \quad (134)$$

This coefficient converts the topological lepton-sector seed into the baryon number observed in the active visible register. In SHBT the CP-odd seed is not introduced as a free phase. It is the level-locked topological Jarlskog invariant

$$J_{CP}^{\text{topo}} = \frac{1}{K} \left(\frac{k_q}{k_\ell + 2} \right) \sqrt{1 - \kappa_{D_5}^2} \sin \left(\frac{2\pi k_q}{k_\ell} \right). \quad (135)$$

The factors have distinct origins. The parent completion contributes the dilution $1/K$; the visible level mismatch contributes $k_q/(k_\ell + 2)$, with the $+2$ the $SU(2)$ dual-Coxeter shift; the dark completion contributes the admissible overlap factor $\sqrt{1 - \kappa_{D_5}^2}$; and the modular phase difference between the quark and lepton clocks contributes $\sin(2\pi k_q/k_\ell)$. Thus CP violation is a residue of completed modular transport rather than an independently tunable angle.

The heavy restoration scale is fixed by the same branch arithmetic. Define the rank projection factor by

$$\Pi_{\text{rank}} = \sqrt{\frac{(h_{SO(10)}^\vee \dim SO(10))/12}{(h_{SU(3)}^\vee \dim SU(3))/12}} \sqrt{\frac{k_q + h_{SU(3)}^\vee}{K + h_{SO(10)}^\vee}}. \quad (136)$$

The residual matching of the 126 completion channel is written $\delta \Pi_{126}^{\text{match}}$. The modular restoration mass is then

$$M_N = M_{\text{GUT}} \exp \left[- \left(I_\ell^\star \Pi_{\text{rank}} + I_q^\star \delta \Pi_{126}^{\text{match}} \right) \right]. \quad (137)$$

For the canonical branch $I_\ell^\star = 6$ and $I_q^\star = 13$, so the exponent is a structural suppression determined by the completed embedding. It is not adjusted after the baryon asymmetry is computed.

The topological baryogenesis identity derived below is therefore a consistency check of the completed modular transport: the same algebra that fixes the visible levels and the dark completion residue also fixes the CP-odd seed and the restoration scale. The numerical outcome $\eta_B \approx 6.45 \times 10^{-10}$ is not a free parameter introduced to match observations; it is the residue of the modular phase mismatch between the quark and lepton clocks after the center lifts are locked. Thus the benchmark audit reported in Section 8 is a verification that the branch arithmetic reproduces the observed visible asymmetry, but it does not claim to be an independent, unconstrained prediction of the baryon abundance without the branch data.

The SHBT baryogenesis identity is therefore

$$\boxed{\eta_B = C_{\text{sph}} J_{CP}^{\text{topo}} \frac{M_N}{M_P}.} \quad (138)$$

The four ingredients in Eq. (138) are fixed by the electroweak conversion coefficient, the modular CP residue, the restoration scale, and the Planck normalization. The benchmark implementation evaluates this identity as `eta_b=6.449923359416e-10`.

The visible asymmetry becomes a complete accounting statement only after the anti-baryon channel is specified. Let $\mathcal{H}_B^{\text{vis}}$ denote the active visible anti-baryon Hilbert space. For a visible anti-baryon density matrix ρ , define the active render cost

$$\boxed{\mathcal{C}_{\bar{B}}(\rho) = \gamma_3 \text{Tr}(\rho Q_{SU(3)}^2) + \gamma_{\text{EM}} \text{Tr}(\rho Q_{\text{EM}}^2) + \gamma_{\text{fr}} \Delta_{\text{fr}}^2, \quad \gamma_3, \gamma_{\text{EM}}, \gamma_{\text{fr}} > 0.} \quad (139)$$

This is an active rendering functional, not a passive mass functional. It counts the cost of keeping anti-baryons visible as colored and electrically charged excitations, plus any residual framing mismatch. On the canonical branch $\Delta_{\text{fr}} = 0$, so the only nonzero active cost comes from visible gauge charge.

The de-rendering operator is the completion-sector projection

$$\mathcal{D}_{\bar{B}} : \mathcal{H}_B^{\text{vis}} \longrightarrow \mathcal{H}_{\text{dark}}^{B-L}, \quad \rho_B^{(n+1)} = \frac{\mathcal{D}_{\bar{B}} \rho_B^{(n)} \mathcal{D}_{\bar{B}}^\dagger}{\text{Tr}(\mathcal{D}_{\bar{B}} \rho_B^{(n)} \mathcal{D}_{\bar{B}}^\dagger)}. \quad (140)$$

The de-rendering map $\mathcal{D}_{\bar{B}}$ acts at the modular restoration scale M_N given in Eq. (137). This scale is determined by the same branch arithmetic that fixes the parent completion and the framing defect. In the boundary theory, the transition is a continuous, unitary projection in the completed Hilbert space: as the energy scale descends from the GUT scale toward M_N , the visible anti-baryon gauge currents lose their support on the active $SU(3)$ and $U(1)_{\text{EM}}$ channels. Equivalently, the projection is a constant phase-transition boundary condition in the register: the active render cost $\mathcal{C}_{\bar{B}}$ drops to zero exactly at the scale where the modular restoration exponent vanishes. For any finite Causal Point, however, this de-rendering is only observed as a local horizon-conditioned effect: if the observer's horizon fraction f_H (Eq. (145)) is too small, the active visible anti-baryon channels are already below the local entropy resolution limit, so the de-rendering appears to the local observer as a pre-existing completion asymmetry rather than a dynamical phase transition.

Thus the mechanism has two complementary descriptions: (i) a bulk phase transition at M_N in the completed boundary data, and (ii) an observer-conditional projection for any Causal Point with f_H below the threshold required to resolve active anti-baryon gauge charges. The benchmark branch fixes M_N once; the observer dependence only determines whether the local history record includes the transition, not whether the dark completion sector stores the passive stress-energy.

It is defined by the active charge-stripping constraints

$$\boxed{\mathcal{D}_{\bar{B}}^\dagger Q_{SU(3)} \mathcal{D}_{\bar{B}} = 0, \quad \mathcal{D}_{\bar{B}}^\dagger Q_{EM} \mathcal{D}_{\bar{B}} = 0.} \quad (141)$$

The dark target stores the conjugate $B - L$ residue, so the completed boundary state remains neutral even though the active visible anti-baryon gauge channels are no longer rendered.

Stress-energy preservation is imposed on the passive source, not on the active charge cost. The defining equality is

$$\boxed{\text{Tr}\left(\rho^{(n)} T_{\mu\nu}^{\bar{B}, \text{passive}}\right) = \text{Tr}\left(\rho^{(n+1)} T_{\mu\nu}^{\text{dark, passive}}\right).} \quad (142)$$

Thus the visible register can lose active anti-baryon color and electromagnetic support without violating the gravitational accounting used by the topological Einstein equation. Equivalently, the de-rendered anti-baryon contributes as a dark passive source rather than as an actively rendered visible particle.

The anti-baryon loop closes at the fixed point

$$\boxed{\rho_{\bar{B}}^* = \frac{\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^* \mathcal{D}_{\bar{B}}^\dagger}{\text{Tr}\left(\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^* \mathcal{D}_{\bar{B}}^\dagger\right)}, \quad \mathcal{C}_{\bar{B}}(\rho_{\bar{B}}^*) = 0, \quad \delta T_{\mu\nu}^{\text{passive}} = 0.} \quad (143)$$

At this point the anti-baryon is absent from the active visible gauge-rendered channels, while its passive stress-energy has been moved into the dark completion sector. The observed baryon asymmetry is therefore a render-channel asymmetry, not an unbalanced disappearance of energy-momentum.

The implementation contract for this section is

Mathematical object	Rust/Python simulator object
Baryogenesis identity record	<code>shbt_simulator.BaryogenesisIdentity</code>
$C_{\text{sph}}, J_{CP}^{\text{topo}}, M_N/M_P, \eta_B$	<code>run_full_audit().baryogenesis_identity</code>
$\mathcal{C}_{\bar{B}}$	<code>cpu_cycle_weight()</code>
$\mathcal{D}_{\bar{B}}$ charge stripping	<code>derender_antibaryon_charges()</code>
Field A versus Field B comparison	<code>run_full_audit().benchmark_delta</code>
Passive source equality	<code>report.stress_energy_preserved</code>

The simulator record `shbt_simulator.BaryogenesisIdentity` stores `sphaleron_coefficient`, `jarlskog_topological`, `Pi_rank`, `deltaPi_126_match`, `structural_exponent`, `modular_restoration_scale_gev`, `heavy_neutrino_to_planck_ratio`, and `eta_b`.

The `shbt_simulator.ShbtSimulator().run_full_audit().baryogenesis_identity` record stores the numerical fields appearing in Eqs. (134)–(138). The method `shbt_simulator.BaryogenesisOptimizer.cpu_cycle_weight()` evaluates the active cost in Eq. (139); `shbt_simulator.BaryogenesisOptimizer.derender_antibaryon_charges()` implements the stripping constraints in Eq. (141); and `report.stress_energy_preserved` reports the passive equality check in Eq. (142). Finally, the `shbt_simulator.ShbtSimulator().run_full_audit().benchmark_delta` record compares the actively rendered Field A calculation with the SHBT-optimized Field B calculation, reporting the changes in `cpu_cycle_weight`, `operation_count`, `memory_bytes`, `elapsed_s`, and the Boolean `stress_energy_preserved`.

7 Causal Point and History Crystallization

The Causal Point is the observer interface of SHBT. It is not an external classical system appended to the boundary theory, but a finite local address inside the completed register. Its job is to translate global holographic closure into the quantities an observer can actually sample: a local memory budget, a local valuation of hidden information, a past-light-cone loading flow, and a deterministic projection that crystallizes one history from a finite outcome ensemble. Thus observation is treated as an entropy-budgeted operation on the boundary register rather than as a primitive background process.

The global horizon follows directly from the saturated bit budget introduced in the Introduction. Since $N = 3\pi/(L_P^2 \Lambda_{\text{holo}})$, the horizon radius is

$$R_H = \sqrt{\frac{NL_P^2}{\pi}} = \sqrt{\frac{3}{\Lambda_{\text{holo}}}}. \quad (144)$$

For a Causal Point whose local sampling radius is R_{local} , define the horizon fraction and local area by

$$f_H = \frac{R_{\text{local}}}{R_H}, \quad A_{\text{local}} = 4\pi R_{\text{local}}^2, \quad 0 \leq f_H \leq 1. \quad (145)$$

The observer cannot address the full horizon register unless $f_H = 1$. The available local memory, hidden complement, and operational entropy limit are therefore

$$N_{\text{local}} = N f_H^2, \quad N_{\text{hidden}} = N - N_{\text{local}}, \quad N_{\text{limit}} = \min\left(N_{\text{local}}, \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right). \quad (146)$$

The first entry counts addressable register bits. The second counts degrees of freedom outside the local addressable patch. The third enforces the stricter of the local register count and the local area-law bound, with the factor $\ln 2$ converting the area entropy into bits.

The Causal Point's self-valuation is the scalar response of the local observer to residual framing and hidden-sector loading. Writing $\Delta_{\text{frame}} \equiv \Delta_{\text{fr}}$ and

$$f_{\text{hidden}} = \frac{N_{\text{hidden}}}{N} = 1 - \frac{N_{\text{local}}}{N}, \quad w(\xi) = \frac{\xi_\ell + \xi_q + \xi_s}{3}, \quad (147)$$

the valuation field is

$$\Sigma = (1 + \Delta_{\text{frame}})(1 + w(\xi)f_{\text{hidden}}). \quad (148)$$

On the canonical branch $\Delta_{\text{frame}} = 0$, but the formula is written in a branch-independent form so that any residual framing mismatch would be visible as a multiplicative observer cost. The localized entropy gradient is the valuation density across the local sampling radius,

$$\nabla_{\text{obs}}\Sigma = \frac{\Sigma f_{\text{hidden}}}{R_{\text{local}}}. \quad (149)$$

The apparent observer acceleration is then

$$a_{\text{obs}} = c^2 \nabla_{\text{obs}}\Sigma. \quad (150)$$

This acceleration is not inserted as an external force law. It is the local inertial response to the hidden entropy gradient seen by an entropy-limited observer.

The same local data define the observer-frame Jacobian. The anisotropy coordinates (ξ_ℓ, ξ_q, ξ_s) are the three directional entries stored in the local property packet, and the Causal Point projection matrix is

$$J^\mu{}_\nu = \begin{pmatrix} 1 + \Sigma f_{\text{hidden}} & -\Sigma \xi_\ell & -\Sigma \xi_q & -\Sigma \xi_s \\ 0 & 1 + \Sigma \xi_\ell & 0 & 0 \\ 0 & 0 & 1 + \Sigma \xi_q & 0 \\ 0 & 0 & 0 & 1 + \Sigma \xi_s \end{pmatrix}. \quad (151)$$

Applied to a bulk metric slice from Section 5, it gives the observer-frame metric

$$g_{\mu\nu}^{\text{obs}} = J_\mu{}^\alpha g_{\alpha\beta} J_\nu{}^\beta. \quad (152)$$

The off-diagonal entries in the first row encode the finite-register mixing between temporal sampling and the three anisotropic property channels, while the diagonal spatial entries encode direction-dependent local rescaling.

Past-light-cone flow is the observer-time image of a static boundary loading process. Let $N_{\text{load}}(t)$ be the number of loaded bits along the sampled history. The loading fraction and effective expansion rate are

$$f_{\text{load}}(t) = \frac{N_{\text{load}}(t)}{N}, \quad H_{\text{eff}}(t) = H_\Lambda + \frac{1}{3} \frac{df_{\text{load}}}{dt}. \quad (153)$$

Here H_Λ is the horizon rate associated with the static Λ_{holo} sector. The second term is not a new external clock; it is the one-dimensional residue of finite boundary loading as reconstructed by the Causal Point. Along the observer's past light cone, redshift parameterizes the same process through

$$\frac{dt}{dz} = -\frac{1}{(1+z)H_{\text{eff}}(z)}, \quad \frac{d\chi}{dz} = \frac{1}{H_{\text{eff}}(z)}. \quad (154)$$

The first equation gives the lookback-time increment, and the second gives the comoving radial increment in the same light-cone units. A sampled redshift therefore selects a boundary coordinate by the ordered loading sequence,

$$n(z) = 1 + \lfloor (|\sigma| - 1) f_{\text{load}}(z) \rfloor, \quad c(z) = \sigma_{n(z)} \in \mathcal{C}. \quad (155)$$

This gives the bridge between the apparent cosmological flow and the finite boundary address on which a measurement can be attempted.

History crystallization begins only after such a boundary address has been selected. Let $\mathcal{R}$ be the finite register visible to the Causal Point, let $A \in \mathcal{R}$ be the selected address, and let

$$\Omega_A = \{\omega_0, \omega_1, \dots, \omega_{m-1}\}, \quad \sum_{i=0}^{m-1} p_i = 1, \quad p_i \geq 0, \quad (156)$$

be the finite outcome ensemble and its normalized local loading weights. The entropy costs of a retrieval event are

$$\begin{aligned} C_{\text{addr}} &= \log_2 |\mathcal{R}|, & C_{\text{ens}} &= \log_2 |\Omega_A|, \\ C_{\text{get}} &= \max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}}), & R_{\text{entropy}} &= N_{\text{limit}} - C_{\text{get}}. \end{aligned} \quad (157)$$

The GET operation is admissible exactly when

$$R_{\text{entropy}} \geq 0. \quad (158)$$

If the residual is negative, the Causal Point has insufficient local entropy budget to crystallize the requested classical record.

When the budget condition holds, the deterministic collapse index is

$$\boxed{\iota = \text{Hash}(\text{observable_name}, A, f_H, N_{\text{local}}, m) \bmod m.} \quad (159)$$

The corresponding rank-one projection operator is

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \quad (160)$$

For a sequence of admissible Causal Point retrievals, the history density matrix updates by

$$\rho_{\text{hist}}^{(n+1)} = \frac{\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)} \Pi_{A_n, \iota_n}}{\text{Tr}(\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)})}. \quad (161)$$

The full crystallized history along a sampled past light cone is therefore the ordered product

$$\Pi_{\text{history}}(z_1, \dots, z_m) = \Pi_{A_m, \iota_m} \cdots \Pi_{A_2, \iota_2} \Pi_{A_1, \iota_1}, \quad A_n = A(c(z_n)). \quad (162)$$

Thus a classical past is not retrieved from an unlimited archive. It is the finite ordered record that survives all local entropy-budget checks.

The pointer wavefunction associated with the selected outcome is the visible and dark completion packet

$$\boxed{\Psi_\iota = (\text{amplitude} = p_\iota, \quad \text{c_vis} = -p_\iota, \quad \text{c_dark} = p_\iota).} \quad (163)$$

The opposite signs of the visible and dark coefficients encode the same completion accounting used elsewhere in SHBT: the visible pointer is rendered as a local classical record while the complementary dark coefficient preserves the finite-register balance.

The implementation contract for this section is

Mathematical object	Rust/Python simulator object
Causal observer interface	<code>shbt_simulator.CausalPoint</code>
Past-light-cone samples	<code>build_past_light_cone()</code>
Local anisotropy and packet data	<code>compute_property_packets()</code>
Entropy-budget admissibility	<code>run_full_audit().memory_report</code>
History projection and pointer packet	<code>crystallize_history()</code>

The class `shbt_simulator.CausalPoint` stores the local horizon fraction, hidden-bit fraction, self-valuation, observer Jacobian, and past-light-cone samples. Its `build_past_light_cone()` method evaluates Eqs. (153)–(155); `compute_property_packets()` stores the anisotropy data entering Eqs. (148) and (151); `shbt_simulator.ShbtSimulator().run_full_audit().memory_report` checks Eqs. (146) and (158); and `crystallize_history()` constructs Eqs. (159)–(163). The benchmark fields reported by this interface include `local_available_bits`, `hidden_bits`, `entropy_limit_bits`, `sigma`, `localized_entropy_gradient_per_m`, `gravitational_acceleration_m_per_s2`, `collapse_index`, `retrieval_cost_bits`, `entropy_budget_residual`, and `pointer_wavefunction`.

8 Numerical Verification and Performance

All audits in this section are verification checks, not phenomenological fits. The branch constants, central charges, and finite register size are fixed by the modular closure axioms of Sections 2–3; the numerical layer confirms that these axioms close consistently across the entropy, metric, baryogenesis, and observer sectors. Where a simulated value agrees with a measured cosmological quantity (e.g., η_B), that agreement is read as evidence that the completed boundary data are physically relevant, not as a validation of a free parameter.

The preceding sections define SHBT by algebraic closure, entropy self-resolution, holographic projection, Causal Point admissibility, and anti-baryon de-rendering. This section records the corresponding finite audit outputs from the canonical Rust/Python simulator. The simulator entry point is `shbt_simulator.ShbtSimulator().run_full_audit()`, and the reproducibility command is `python examples/run_audit.py`. The older standalone `shbt_core.py` audit is deprecated and removed from the paper-level specification; where legacy values differ, the new simulator output is taken as canonical because the audit uses 512-bit precision and parallel computation. The verification is not a separate phenomenological fit: every entry is evaluated on the same canonical branch

$$\boxed{\text{shbt_simulator.StaticBoundary.benchmark_branch} = (26, 8, 312), \quad \Delta_{\text{fr}} = 0.} \quad (164)$$

The purpose of the numerical layer is therefore to check that the branch data, normalization conventions, projector algebra, observer memory accounting, and baryogenesis optimizer all close simultaneously.

8.1 Boundary closure audit

The `shbt_simulator.StaticBoundary` component tests the canonical branch before any bulk metric or observer-frame reconstruction is used. The first requirement is exact framing closure. The audit evaluates

$$\Delta_{\text{fr}}(312, 26, 8) = \max\left(\left\|\frac{312}{2 \cdot 26}\right\|_{\mathbb{Z}}, \left\|\frac{312}{3 \cdot 8}\right\|_{\mathbb{Z}}\right) = 0, \quad (165)$$

and then checks that the loading and entanglement densities used by the entropy flow are normalized. The modular tests verify that the visible and completed partition data commute with the constructed S_{∂} and T_{∂} kernels to simulator precision:

$$\|[M, S_{\partial}]\| = 0.0, \quad \|[M, T_{\partial}]\| = 0.0. \quad (166)$$

The full boundary audit is summarized in Table 2.

Audit quantity	Simulator accessor	Rust/Python output
Branch levels	<code>report.branch</code>	(26, 8, 312)
Framing defect	<code>report.framing_defect</code>	0.0
$Z_{\partial}^{\text{code}}(i)$	<code>evaluate_z_boundary()</code>	2.441381789163 $\times 10^{-24}$
Loading density sum	<code>build_loading_density()</code>	1.0
Entanglement density sum	<code>build_entanglement_density()</code>	1.0
Dominant sequence length	<code>build_dominant_sequence()</code>	9
Modular S commutator	<code>boundary_report.modular_S_commutator</code>	0.0
Modular T commutator	<code>boundary_report.modular_T_commutator</code>	0.0
Modular invariant	<code>report.modular_invariant</code>	true
Zero-energy lock	<code>report.zero_energy_locked</code>	true
Visible projection	<code>report.projection_dimension_26_to_4</code>	true

Table 2: Static boundary verification results from the canonical Rust/Python simulator.

Together these checks establish the implementation form of the first closure equivalence:

$$[M, S_{\partial}] = [M, T_{\partial}] = 0 \iff \Delta_{\text{fr}} = 0 \implies \Pi_{\text{phys}} H_{\partial} \Pi_{\text{phys}} = 0. \quad (167)$$

The final two entries in Table 2 are especially important for reproducibility: `report.zero_energy_locked=True` confirms that the completed static boundary has no residual external time generator, while `report.projection_dimension_26_to_4=True` confirms that the visible lepton-sector projection used by the benchmark branch has the expected finite dimension.

8.2 Bulk metric projection audit

The `shbt_simulator.HolographicProjection` component starts from the normalized loading and entanglement densities and constructs the entropy cascade of metric slices. The benchmark run returns the full nine-slice cascade,

$$\text{projection_report.slice_count} = 9, \quad (168)$$

with a rank-three spatial projector at every sampled step. Each metric slice is then checked for the three algebraic properties required by the projection construction:

$$g_{ij} = g_{ji}, \quad \text{Tr}(g) = 1, \quad v^i g_{ij} v^j > 0 \quad \text{for } v \neq 0. \quad (169)$$

The resulting audit table is Table 3.

Audit quantity	Simulator accessor	Rust/Python output
Entropy cascade length	<code>projection_report.slice_count</code>	9
Spatial projector rank	<code>projection_report.projector_rank</code>	3
Metric symmetry	<code>projection_report.symmetric</code>	true
Trace normalization	<code>projection_report.trace_normalized</code>	true
Positive definiteness	<code>projection_report.positive_definiteness</code>	true

Table 3: Holographic projection verification results from the canonical Rust/Python simulator.

This confirms that the entropy flow does not merely produce arrays of numbers. It produces symmetric, trace-normalized, positive metric data with the intended rank-three spatial image, so the bulk slices satisfy the numerical version of the projection conditions stated in Section 5.

8.3 Causal Point audit

The `shbt_simulator.CausalPoint` component verifies that observer access is limited by a finite entropy budget. For a horizon fraction f_H , the local bit budget is checked against

$$N_{\text{local}}^{\text{audit}} \simeq N f_H^2, \quad N_{\text{hidden}}^{\text{audit}} = N - N_{\text{local}}^{\text{audit}}. \quad (170)$$

Here $N_{\text{local}}^{\text{audit}}$ and $N_{\text{hidden}}^{\text{audit}}$ are the `memory_report.local_available_bits` and `memory_report.hidden_bits` fields. The retrieval interface is admissible only when the entropy limit is positive, and the past-light-cone builder is checked against the requested redshift grid:

$$N_{\text{limit}}^{\text{audit}} > 0, \quad n_{\text{lightcone}}^{\text{audit}} = n_{\text{redshift}}. \quad (171)$$

These are the `memory_report.entropy_limit_bits`, `memory_report.past_light_cone_samples`, and `redshift_samples` checks in the report. The audit is summarized in Table 4.

These outputs check the observer side of the closure chain. The Causal Point does not access an unlimited record; it constructs exactly as many past-light cone entries as the redshift sampler asks for, constructs nine local property packets, and filters each attempted retrieval through the finite-memory inequality.

8.4 Baryogenesis and optimized field simulation

The baryogenesis benchmark compares a standard visible field simulation with the optimized SHBT render in which anti-baryon visible charge channels are de-rendered while their passive stress-energy

Audit quantity	Simulator accessor	Rust/Python output
Local available bits	<code>memory_report.local_available_bits</code>	$2.535748254483999 \times 10^{122}$
Hidden bits	<code>memory_report.hidden_bits</code>	$7.76249465658367 \times 10^{121}$
Entropy limit	<code>memory_report.entropy_limit_bits</code>	$2.535748254483999 \times 10^{122}$
Light-cone samples	<code>memory_report.past_light_cone_samples</code>	9
Property packets	<code>memory_report.property_packets</code>	9
Memory admissibility	<code>report.memory_all_passed</code>	true

Table 4: Causal Point verification results from the canonical Rust/Python simulator.

contribution remains in the dark completion sector. The simulator reports the visible baryon asymmetry as

$$\eta_B = 6.449923359416 \times 10^{-10}, \quad (172)$$

with `report.stress_energy_preserved=True`. In the benchmark implementation the optimization is also a performance statement: fewer actively rendered field channels are evolved, while the passive gravitational accounting is unchanged. The standard and optimized simulations are compared in Table 5.

Simulation mode	Active visible channels	CPU cycles	Operations	Memory
Standard field simulation	baryon and anti-baryon	1.0	1.0	1.0
Optimized SHBT simulation	baryon rendered, anti-baryon de-rendered	0.5	0.001533314747700058214747736093	
Reduction fraction	removed active anti-baryon render	0.5	0.998466685252309941785252263907	

Table 5: Normalized standard-versus-optimized field simulation cost from `benchmark_delta`.

The physical comparison is shown in Table 6. The optimized simulation is not allowed to discard energy: it removes anti-baryon charge from the actively rendered visible channel but preserves the passive source that enters the topological Einstein equation.

Quantity	Standard simulation	Optimized SHBT simulation
Visible baryon channel	actively rendered	actively rendered
Visible anti-baryon channel	actively rendered	de-rendered
Passive stress-energy	explicitly evolved	preserved in dark completion
Baryon asymmetry output	dynamical baseline target	$\eta_B = 6.449923359416 \times 10^{-10}$
Stress-energy check	baseline conservation test	true

Table 6: Physical accounting in the standard and optimized simulations.

The combined verification suite therefore checks both correctness and cost. On the cor-

rectness side, `shbt_simulator.ShbtSimulator().run_full_audit()` returns the same branch data with `report.modular_invariant=True`, `report.zero_energy_locked=True`, `report.memory_all_passed=True`, and `report.stress_energy_preserved=True`. On the performance side, the optimized render reduces CPU cycles by 50%, operation count by approximately 99.8467%, and memory use by approximately 99.4179% relative to the normalized standard field simulation, without violating the stress-energy preservation identity.

9 Precision Cosmological Implications

The foundation sections define a static closed boundary theory. The cosmology sector asks how a finite observer inside that closed register infers an expanding history. No new branch coordinate is introduced. The same center-locked parent level, dark completion, finite entropy register, and Causal Point light-cone rule are reinterpreted as a precision-cosmology effective field theory. The Python audit layer for this section is `noether_bridge.py` and `precision_cosmology_engine.py`.

9.1 Completion Notation and Entropy Debt

The source cosmology draft used an unsuperscripted dark-ledger symbol for the completed finite-screen ledger, while the foundational simulator used the same printed name for the residual part. In this unified manuscript the distinction is fixed by Eq. (6):

$$c_{\text{dark}}^{\text{res}} = \frac{834433}{362670}, \quad c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = \frac{1197103}{362670}, \quad \Delta_{\text{mod}} = \frac{c_{\text{dark}}^{\text{comp}}}{24} = \frac{1197103}{8704080}. \quad (173)$$

The residual quantity $c_{\text{dark}}^{\text{res}}$ is the foundational closure residue used by `shbt_simulator.StaticBoundary.c_dark`. The completed quantity $c_{\text{dark}}^{\text{comp}}$ is the finite-vacuum-subtracted ledger used by `noether_bridge.load_c_dark_completion_fraction` and by the clock-skew cosmology model. Appendix B, Eq. (251), gives the completed ledger from the coset subtraction, while Appendix A fixes the parent-level arithmetic used by both ledgers. This is the only dark-sector convention used below.

The finite horizon register is the saturated SHBT form of the holographic entropy bound. For a spatial region bounded by a cosmological horizon of area A , the Bekenstein–Hawking counting bound is

$$S \leq \frac{A}{4L_P^2 \ln 2}. \quad (174)$$

On the canonical branch the cosmological screen is fixed by Λ_{holo} , so the saturated bit capacity is

$$N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}}, \quad \Lambda_{\text{holo}} = 1.08913883 \times 10^{-52} \text{ m}^{-2}, \quad N_{\text{sat}} = 3.312593327986 \times 10^{122}. \quad (175)$$

This finite number is not a density parameter. It is the total horizon register against which the observer-frame loading process is charged.

Let $N_{\text{load}}(z)$ be the number of horizon bits actively loaded by the past-light-cone sample at redshift z . The dimensionless loading fraction and entropy debt are

$$f_{\text{load}}(z) = \frac{N_{\text{load}}(z)}{N_{\text{sat}}}, \quad S_{\text{debt}}(z) = N_{\text{load}}(z) = N_{\text{sat}} f_{\text{load}}(z), \quad 0 \leq f_{\text{load}}(z) \leq 1. \quad (176)$$

The horizon contribution is saturated on this branch, so the only active light-cone entropy increment in the cosmology audit is the debt term. If t_{lc} denotes the observer's past-light-cone acquisition parameter, then

$$\frac{dz}{dt_{\text{lc}}} = (1+z)H_{\text{SHBT}}(z). \quad (177)$$

The report imposes a constant Causal Point loading rate Γ_{lock} in the clock-normalized channel:

$$\frac{1}{N_{\text{sat}}} \frac{dS_{\text{total}}}{dt_{\text{lc}}} = \frac{\Gamma_{\text{lock}}}{1+z}. \quad (178)$$

Combining Eqs. (176)–(178) gives

$$\begin{aligned} \frac{dS_{\text{total}}}{dt_{\text{lc}}} &= \frac{dS_{\text{H}}}{dt_{\text{lc}}} + \frac{dS_{\text{debt}}}{dt_{\text{lc}}} = 0 + N_{\text{sat}} \frac{df_{\text{load}}}{dz} \frac{dz}{dt_{\text{lc}}} \\ &= N_{\text{sat}} \frac{df_{\text{load}}}{dz} (1+z)H_{\text{SHBT}}(z) = N_{\text{sat}} \frac{\Gamma_{\text{lock}}}{1+z} > 0. \end{aligned} \quad (179)$$

Solving the last equality for the loading fraction gives the redshift ODE

$$\boxed{\frac{df_{\text{load}}}{dz} = \frac{\Gamma_{\text{lock}}}{(1+z)^2 H_{\text{SHBT}}(z)}}. \quad (180)$$

Thus the generalized second law is not an independent late-time fluid assumption; it is the monotone light-cone image of finite-register loading.

The lock rate is fixed by the Causal Point clock relation

$$d\tau_{\text{lock}} = \frac{dt}{1+z}. \quad (181)$$

In the observer frame, the loaded clock channel contributes one third of its rate to the locally inferred Hubble intercept:

$$H_0(z) = H_{\Lambda} + \frac{1}{3(1+z)} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}. \quad (182)$$

The static horizon term is anchored to the CMB intercept, $H_{\Lambda} = H_0^{\text{CMB}}$. Evaluating Eq. (182) at $z = 0$ gives

$$H_0^{\text{loc}} = H_0^{\text{CMB}} + \frac{1}{3} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}, \quad \Gamma_{\text{lock}} \equiv \frac{df_{\text{load}}}{d\tau_{\text{lock}}} = 3(H_0^{\text{loc}} - H_0^{\text{CMB}}). \quad (183)$$

This identifies the entropy-debt loading rate with the measured separation between the local and CMB Hubble intercepts.

9.2 Noether Bridge and Newton Lock

The Noether bridge audit uses the completed ledger to evaluate

$$8\pi G_{\text{eff}} = \frac{12}{c_{\text{dark}}^{\text{comp}} M_P^2}, \quad \frac{d \ln G_{\text{eff}}}{dt} = -\frac{d \ln c_{\text{dark}}^{\text{comp}}}{dt} - 2 \frac{d \ln M_P}{dt}. \quad (184)$$

On the fixed branch both logarithmic derivatives vanish, so the Bianchi stationarity check gives $\dot{G}/G = 0$. The same report verifies the unity-of-scale identity. The retained D_5 cell factor κ_{D_5} entering this identity is defined in Appendix C, Eq. (259), and the neutrino eigenvalues used for the normal hierarchy are listed in Eqs. (262)–(264):

$$m_{\nu,1} = \kappa_{D_5} M_P N^{-1/4}, \quad \Lambda_{\text{holo}} = \frac{3\pi}{\kappa_{D_5}^4} G_{\text{top}} m_{\nu,1}^4, \quad \epsilon_\Lambda \leq \frac{1}{N}. \quad (185)$$

The key values returned by `python noether_bridge.py --precision 80` are collected in Table 7.

Audit quantity	Value	Status
$c_{\text{dark}}^{\text{comp}}$	1197103/362670 3.300805139659	= fixed input
κ_{D_5}	0.988769793998	fixed input
Λ_{holo}	$1.08913883 \times 10^{-52} \text{ m}^{-2}$	fixed input
M_P	$1.220890146939 \times 10^{28} \text{ eV}$	derived
$8\pi G_{\text{eff}}$	$2.438979677015 \times 10^{-56} \text{ eV}^{-2}$	locked
$G_{\text{top}}/G_{\text{eff}}$	6.913190118455	derived
Newton-lock residual	0.0 eV^{-2}	PASS
$\dot{G}/G$	0.0 yr^{-1}	PASS
$N(\Lambda_{\text{holo}})$	$3.312593327986 \times 10^{122}$	derived
$1/N$ register floor	$3.018782871871 \times 10^{-123}$	derived
$m_{\nu,1}$	$2.829630635353 \times 10^{-3} \text{ eV}$	derived
ϵ_Λ	$1.658583028709 \times 10^{-249}$	PASS
Normal-hierarchy sum	61.8 meV	1.68σ
Inverted-hierarchy sum	103 meV	2.80σ
Overall Noether bridge	all checks closed	PASS

Table 7: Key `noether_bridge.py` outputs for the completed finite-screen ledger.

9.3 Neutrino Hierarchy Lock

The same Noether bridge report fixes the neutrino floor before the Hubble loading law is applied. The lightest normal eigenstate is not a fit parameter: it is the finite-register quarter-root scale weighted by the retained D_5 boundary-cell measure,

$$m_{\nu,1} = \kappa_{D_5} M_P N_{\text{sat}}^{-1/4}. \quad (186)$$

Substituting the benchmark values used in the audit gives

$$\begin{aligned} m_{\nu,1} &= 0.988769793998 (1.220890146939 \times 10^{28} \text{ eV}) (3.312593327986 \times 10^{122})^{-1/4} \\ &= 2.829630635353 \times 10^{-3} \text{ eV}. \end{aligned} \quad (187)$$

The oscillation splittings then convert the boundary floor into absolute propagation eigenstates. The audit uses

$$\Delta m_{21}^2 \simeq 7.4 \times 10^{-5} \text{ eV}^2, \quad \Delta m_{31}^2 \simeq 2.5 \times 10^{-3} \text{ eV}^2. \quad (188)$$

For the normal hierarchy,

$$m_{\nu,2}^{\text{NH}} = \sqrt{m_{\nu,1}^2 + \Delta m_{21}^2} \simeq 8.92 \times 10^{-3} \text{ eV}, \quad m_{\nu,3}^{\text{NH}} = \sqrt{m_{\nu,1}^2 + \Delta m_{31}^2} \simeq 5.01 \times 10^{-2} \text{ eV}. \quad (189)$$

The normal-ordering sum is therefore

$$\Sigma_{\nu}^{\text{NH}} = m_{\nu,1}^{\text{NH}} + m_{\nu,2}^{\text{NH}} + m_{\nu,3}^{\text{NH}} = 61.8 \text{ meV}. \quad (190)$$

For the inverted hierarchy, the finite-register floor is assigned to the lightest third eigenstate,

$$m_{\nu,3}^{\text{IH}} = 2.829630635353 \times 10^{-3} \text{ eV}, \quad m_{\nu,1}^{\text{IH}} \simeq 4.87 \times 10^{-2} \text{ eV}, \quad m_{\nu,2}^{\text{IH}} \simeq 4.95 \times 10^{-2} \text{ eV}, \quad (191)$$

so that

$$\Sigma_{\nu}^{\text{IH}} = m_{\nu,3}^{\text{IH}} + \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{32}^2|} + \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{32}^2| + \Delta m_{21}^2} = 103 \text{ meV}. \quad (192)$$

The report's hierarchy statistic is the audit-normalized pull of each derived sum against the cosmological mass-sum likelihood used by `noether_bridge.py`:

$$\sigma_{\text{tens}}^{\text{NH}} = \frac{\Delta \Sigma_{\nu}^{\text{NH}}}{\sigma_{\Sigma, \text{audit}}^{\text{NH}}} = 1.68, \quad \sigma_{\text{tens}}^{\text{IH}} = \frac{\Delta \Sigma_{\nu}^{\text{IH}}}{\sigma_{\Sigma, \text{audit}}^{\text{IH}}} = 2.80. \quad (193)$$

Thus the normal ordering is the lower-tension finite-register completion. The explicit eigenstate ledger is shown in Table 8.

Quantity	Normal hierarchy	Inverted hierarchy
Floor eigenstate	$m_1 = 2.83 \times 10^{-3} \text{ eV}$	$m_3 = 2.83 \times 10^{-3} \text{ eV}$
Second eigenstate	$m_2 = 8.92 \times 10^{-3} \text{ eV}$	$m_1 = 4.87 \times 10^{-2} \text{ eV}$
Third eigenstate	$m_3 = 5.01 \times 10^{-2} \text{ eV}$	$m_2 = 4.95 \times 10^{-2} \text{ eV}$
Derived sum Σ_{ν}	61.8 meV	103 meV
Audit tension	1.68 σ	2.80 σ

Table 8: Neutrino propagation eigenstates, mass sums, and hierarchy pulls from the finite-register floor.

The hierarchy check is therefore tied to the same κ_{D_5} , N_{sat} , and completed-ledger inputs as the Newton lock.

9.4 Cosmology Code-to-Math Map

The cosmology audit uses a separate Python layer from the foundation simulator, but every displayed quantity below has a direct code counterpart:

Table 9 is also the cross-reference index for the new precision derivations: the completed ledger and finite screen are Eqs. (173)–(175); the light-cone loading and GSL identities are Eqs. (176)–(180); the Noether bridge and neutrino hierarchy are Eqs. (184)–(193); the observer-frame Hubble law is Eqs. (194)–(199); growth and cluster checks are tied to Eqs. (200)–(206) and Tables 11–13; ISW, BBN, and chronometer checks are Eqs. (207)–(221); and the GET measurement rule is Eqs. (222)–(231).

Mathematical quantity	Code object or audit entry	Role
$c_{\text{dark}}^{\text{comp}}$	<code>load_c_dark_completion_fraction</code>	completed ledger
κ_{D_5}	<code>PhysicsConstantConfig.kappa_d5</code>	retained D_5 measure
N_{sat}	<code>holographic_bits_from_lambda</code>	finite horizon register
$m_{\nu,1}$	<code>report.unity.lightest_mass_ev</code>	neutrino floor
$\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$	<code>benchmark_entropy_debt_fraction</code> <code>ClockSkewCosmology.delta_mod</code>	/ entropy-debt uplift
H_0^{loc}	<code>late_time_hubble_constant</code> <code>audit.h0_local_km_s_mpc</code>	/ local Hubble intercept
$A_H, \Gamma_{\text{lock}}$	<code>hubble_gradient</code> <code>loading_fraction_derivative</code>	/ loading amplitude and lock rate
$H_0(z)$	<code>redshift_dependent_hubble_constant</code> <code>audit.samples</code>	/ redshift loading law
$f_{\text{load}}(z)$ and $S_{\text{debt}}(z)$	<code>compute_lightcone_loaded_fraction</code> <code>compute_lightcone_entropy_debt</code>	/ finite-register debt
$f\sigma_8(z)$	<code>build_structure_growth_audit</code>	growth suppression
δ_c, σ_M , halo tail	<code>build_cluster_collapse_audit</code>	cluster mirage
$\Delta(\dot{H}/H^2)$	<code>audit_isw_stability</code>	ISW residual
BBN shift	<code>build_early_universe_stability_audit</code>	radiation-era stability
χ_ν^2	<code>build_cosmic_chronometer_validation</code>	CC covariance check
$dS_{\text{total}}/dt_{\text{lc}}$	<code>build_thermodynamic_arrow_audit</code>	light-cone entropy arrow
$C_{\text{get}}, R_{\text{entropy}}, \Pi_{A,\ell}$	<code>measurement_operator.py</code>	GET measurement cost

Table 9: Code-to-math map for the precision-cosmology audit layer.

9.5 Observer-Frame Hubble Loading

The clock-skew cosmology is built from the entropy-debt uplift

$$\frac{H_0^{\text{loc}}}{H_0^{\text{CMB}}} = \exp\left(\frac{\Delta_{\text{mod}}}{2}\right), \quad H_0^{\text{CMB}} = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (194)$$

For the benchmark branch this gives

$$\Delta_{\text{mod}} = 0.137533547486, \quad \exp(\Delta_{\text{mod}}/2) = 1.071186351229, \quad (195)$$

$$H_0^{\text{loc}} = 72.197960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (196)$$

Define the local loading amplitude

$$A_H = H_0^{\text{loc}} - H_0^{\text{CMB}} = H_0^{\text{CMB}} \left[\exp\left(\frac{\Delta_{\text{mod}}}{2}\right) - 1 \right] = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (197)$$

The Causal Point clock relation in Eq. (181) converts the local loading derivative into the redshift-dependent intercept sampled by a finite observer:

$$H_0(z) = H_\Lambda + \frac{1}{3(1+z)} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}, \quad H_\Lambda = H_0^{\text{CMB}}, \quad \frac{df_{\text{load}}}{d\tau_{\text{lock}}} = 3(H_0^{\text{loc}} - H_0^{\text{CMB}}). \quad (198)$$

Equivalently, Eq. (183) gives $\Gamma_{\text{lock}} = df_{\text{load}}/d\tau_{\text{lock}} = 3A_H$. Substituting this value gives the closed form

$$H_0(z) = H_0^{\text{CMB}} + \frac{A_H}{1+z} = H_0^{\text{CMB}} \left[1 + \frac{\exp(\Delta_{\text{mod}}/2) - 1}{1+z} \right]. \quad (199)$$

Thus the local uplift is active at $z = 0$ but decays as $(1+z)^{-1}$, so the early universe remains anchored to the CMB intercept. The audit values from `precision_cosmology_engine.py` are shown in Table 10.

z	Loading term [$\text{km s}^{-1} \text{ Mpc}^{-1}$]	$H_0(z)$ [$\text{km s}^{-1} \text{ Mpc}^{-1}$]
0.0	4.797960072861	72.197960072861
0.5	3.198640048574	70.598640048574
1.0	2.398980036431	69.798980036431
2.0	1.599320024287	68.999320024287
10.0	0.436178188442	67.836178188442
1100.0	0.004357820230	67.404357820230

Table 10: Redshift-ladder tracking from the finite-register loading law in Eq. (198).

9.6 Growth, Mirage, and Cluster Checks

The precision engine compares the clock-skew branch against a rigid Λ CDM intercept in standard observational channels. The structure-growth audit computes $f\sigma_8$ from the same background loading law while keeping G_{eff} fixed. The starting point is the standard linear matter perturbation equation

$$\ddot{\delta}_m + 2H\dot{\delta}_m - 4\pi G_{\text{eff}}\rho_m\delta_m = 0. \quad (200)$$

Writing $\delta_m = D(x)$, with $x = \ln a = -\ln(1+z)$, converts the time derivatives to

$$\dot{D} = H \frac{dD}{dx}, \quad \ddot{D} = H^2 \frac{d^2 D}{dx^2} + \dot{H} \frac{dD}{dx}. \quad (201)$$

Substitution into Eq. (200) gives

$$H^2 \frac{d^2 D}{dx^2} + \left(\dot{H} + 2H^2 \right) \frac{dD}{dx} - 4\pi G_{\text{eff}} \rho_m D = 0. \quad (202)$$

After division by H^2 , using

$$\frac{d \ln H}{dx} = \frac{\dot{H}}{H^2}, \quad \Omega_m(x) = \frac{8\pi G_{\text{eff}} \rho_m}{3H^2},$$

the linear growth ODE used by the audit is

$$\boxed{\frac{d^2 D}{dx^2} + \left[2 + \frac{d \ln H_{\text{SHBT}}}{dx} \right] \frac{dD}{dx} - \frac{3}{2} \Omega_m(x) D = 0, \quad f\sigma_8(z) = \sigma_{8,0} \frac{dD}{dx}.} \quad (203)$$

The loading law modifies this equation only through the background friction term. Writing

$$H_{\text{SHBT}}(z) = H_0(z) \left[\Omega_{m0}(1+z)^3 + \Omega_{r0}(1+z)^4 + \Omega_{\Lambda 0} \right]^{1/2}, \quad (204)$$

Eq. (199) gives

$$\begin{aligned} \frac{d \ln H_{\text{SHBT}}}{dx} &= \left(\frac{d \ln E}{dx} \right)_{\Lambda\text{CDM}} + \frac{d \ln H_0(z)}{dx}, \\ \frac{d \ln H_0(z)}{dx} &= \frac{1}{H_0(z)} \frac{dH_0}{dz} \frac{dz}{dx} = \frac{1}{H_0(z)} \left[-\frac{A_H}{(1+z)^2} \right] [-(1+z)] = \frac{A_H}{(1+z)H_0(z)}. \end{aligned} \quad (205)$$

The positive last term increases the late-time friction while Eq. (184) keeps G_{eff} fixed. The result is a kinematic suppression of growth rather than a modified Newton constant:

$$\frac{(f\sigma_8)_{\text{SHBT}}}{(f\sigma_8)_{\Lambda\text{CDM}}} - 1 = \{-9.8\%, -8.3\%\} \quad \text{at} \quad z = \{0.5, 0.8\}. \quad (206)$$

The corresponding audit table is Table 11.

z	$(f\sigma_8)_{\Lambda\text{CDM}}$	$(f\sigma_8)_{\text{SHBT}}$	Kinematic suppression
0.5	0.475	0.428	-9.8%
0.8	0.453	0.415	-8.3%

Table 11: Structure-growth suppression audit from `build_structure_growth_audit`.

When the same branch is interpreted as a conserved-fluid template, the inferred dark-energy density is only an effective image. The Python report labels this a high-redshift mirage: the template has a zero crossing at $z_{\text{cross}} = 1.865$, after which the conserved-fluid interpretation fails even though the underlying boundary-loading law remains well defined.

The cluster-collapse audit separates the collapse barrier from the fluctuation amplitude. The barrier changes by less than 0.3%, while the suppressed σ_M is exponentiated in the halo tail. The benchmark reference mass is $\log_{10}(M/M_{\odot}) = 14.5$ with $\sigma_M(z=0) = 0.6000$.

Feature	H_0^{SHBT}	ΔH_0	σ_H	Significance	$R_{\text{DE}}^{\text{eff}}$
$z = 1.5$	69.32	-2.88	0.75	-3.8σ	0.360
$z_{\text{cross}} = 1.865$	—	—	—	—	0.000
$z = 2.0$	69.00	-3.20	0.70	-4.6σ	-0.162
$z = 2.5$	68.77	-3.43	0.67	-5.1σ	-0.920

Table 12: High-redshift conserved-fluid mirage forecast.

z	$\delta_c^{\Lambda\text{CDM}}$	δ_c^{SHBT}	$\sigma_M^{\Lambda\text{CDM}}$	σ_M^{SHBT}	ν_{SHBT}	Abundance ratio
0.0	1.6760	1.6733	0.6000	0.5601	2.9873	6.10×10^{-1}
0.5	1.6821	1.6799	0.4614	0.4378	3.8370	5.15×10^{-1}
1.0	1.6844	1.6826	0.3641	0.3492	4.8186	4.21×10^{-1}

Table 13: Cluster-collapse mirage audit from `build_cluster_collapse_audit`.

9.7 Light-Cone, Stability, and Forecast Ledgers

The finite light-cone ledger integrates Eqs. (176) and (180). At recombination, the integral consumes about 10.75% of the finite register, leaving about 89.25% headroom. The sampled debt values are shown in Table 14.

z	$H_0(z)$	$H_{\text{SHBT}}(z)$	f_{load}	S_{debt} [bits]
0.0	72.1980	72.1980	0.000000	0.000000×10^{127}
0.5	70.5986	93.3432	0.060046	1.981525×10^{121}
1.0	69.7990	124.9577	0.082644	2.727249×10^{121}
2.0	68.9993	209.1715	0.098020	3.234674×10^{121}
10.0	67.8362	1390.1452	0.107076	3.533498×10^{121}
1100.0	67.4044	1.3820515142×10^6	0.107453	3.545951×10^{121}

Table 14: Information-theoretic past-light-cone ledger.

The stability checks verify that the late-time loading term is negligible in the early universe and that the integrated Sachs–Wolfe (ISW) residual stays small in the matter era. The ISW temperature source is

$$\left(\frac{\Delta T}{T}\right)_{\text{ISW}} = \int (\dot{\Phi} + \dot{\Psi}) dt. \quad (207)$$

In a pure matter era with unmodified Newton coupling, the linear gravitational potentials are static, so $\dot{\Phi} = \dot{\Psi} = 0$. In the SHBT observer frame the only matter-era source left in the audit is the logarithmic drift of the squared clock normalization $H_0(z)^2$. The residual is therefore

$$\Delta\left(\frac{\dot{H}}{H^2}\right) = \left(\frac{\dot{H}}{H^2}\right)_{\text{SHBT}} - \left(\frac{\dot{H}}{H^2}\right)_{\text{matter}}, \quad \left(\frac{\dot{H}}{H^2}\right)_{\text{matter}} = -\frac{3}{2}. \quad (208)$$

With $a = (1+z)^{-1}$,

$$d \ln a = -\frac{dz}{1+z}, \quad \frac{d \ln H_0}{d \ln a} = \frac{1}{H_0(z)} \frac{dH_0}{dz} [-(1+z)]. \quad (209)$$

Using $H_0(z) = H_0^{\text{CMB}} + A_H/(1+z)$ gives

$$\frac{d \ln H_0}{d \ln a} = \frac{1}{H_0(z)} \left[-\frac{A_H}{(1+z)^2} \right] [-(1+z)] = \frac{A_H}{(1+z)H_0(z)}. \quad (210)$$

Because the source is the drift of H_0^2 , the matter-era residual used in Table 15 is

$$\Delta_{\text{ISW}}(z) = -2 \frac{d \ln H_0}{d \ln a} = -\frac{2A_H}{(1+z)H_0(z)}. \quad (211)$$

For $z = 1100$, this gives $-2A_H/[(1101)H_0(1100)] = -1.29 \times 10^{-4}$, matching the recombination row of the ISW audit.

z	$\Omega_m(z)$	$\Delta(\dot{H}/H^2)$	SHBT $\dot{H}/H^2$
10.0	0.998369	-0.012762	-1.512762
50.0	0.999984	-0.002785	-1.502785
100.0	0.999998	-0.001408	-1.501408
1100.0	1.000000	-0.000129	-1.500129

Table 15: ISW stability audit from `audit_isw_stability`.

The BBN check follows from the radiation-era scaling. During nucleosynthesis,

$$\rho_{\text{rad}}(z) = \rho_{r0}(1+z)^4, \quad H_{\text{rad}}(z) = H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^2. \quad (212)$$

The SHBT clock load adds only

$$\delta H_{\text{load}}(z) = \frac{A_H}{1+z}, \quad H_{\text{SHBT}}(z) = H_{\text{rad}}(z) + \delta H_{\text{load}}(z) \quad \text{in the radiation audit.} \quad (213)$$

The stability requirement is

$$\frac{\delta H_{\text{load}}(z_{\text{BBN}})}{H_{\text{rad}}(z_{\text{BBN}})} \ll 1, \quad z_{\text{BBN}} \simeq 10^9. \quad (214)$$

For the branch value $A_H = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}$,

$$\delta H_{\text{load}}(10^9) = \frac{4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}}{1 + 10^9} = 4.797960068064 \times 10^{-9} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (215)$$

$$H_{\text{rad}}(10^9) \simeq 6.46 \times 10^{17} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (216)$$

$$\frac{\delta H_{\text{load}}}{H_{\text{rad}}(10^9)} = 7.421690135465 \times 10^{-27}. \quad (217)$$

The active loading term is therefore decoupled from the early thermal expansion rate by more than twenty-six orders of magnitude.

9.8 Cosmic Chronometer Covariance

The cosmic chronometer (CC) validation is a covariance test of the background expansion history. The method uses differential galaxy ages to infer

$$H(z) = -\frac{1}{1+z} \frac{dz}{dt}. \quad (218)$$

Let $\mathbf{H}_{\text{obs}} = (H_{\text{obs}}(z_1), \dots, H_{\text{obs}}(z_{32}))^T$ be the 32-point CC vector, and let $\mathbf{C}$ be the corresponding 32×32 statistical-plus-systematic covariance matrix. The validation statistic is

$$\chi^2 = \sum_{i,j=1}^{32} [H_{\text{obs}}(z_i) - H_{\text{SHBT}}(z_i)] [\mathbf{C}^{-1}]_{ij} [H_{\text{obs}}(z_j) - H_{\text{SHBT}}(z_j)]. \quad (219)$$

The audit treats H_0^{CMB} , Δ_{mod} , and Ω_{m0} as the three background parameters entering this check, so

$$\nu = N_{\text{data}} - N_{\text{params}} = 32 - 3 = 29. \quad (220)$$

The reported total chi-square is $\chi^2 = 30.16$, hence

$$\chi_\nu^2 = \frac{\chi^2}{\nu} = \frac{30.16}{29} = 1.04. \quad (221)$$

Metric	Cosmic chronometer audit value
Observational data points N_{data}	32
Model parameters N_{params}	3
Degrees of freedom ν	29
Total chi-square χ^2	30.16
Reduced chi-square χ_ν^2	1.04

Table 16: Cosmic chronometer covariance validation from `precision_cosmology_engine.py`.

The remaining precision-cosmology checks are summarized in Table 17. The forecasted model-selection rows are sensitivity forecasts rather than current empirical detections. The light-cone entropy row is the sampled form of Eq. (179); because $\Gamma_{\text{lock}} > 0$ and $N_{\text{sat}} > 0$, every sampled $z \geq 0$ has $dS_{\text{total}}/dt_{\text{lc}} > 0$.

Check	Python output	Status
Local uplift	72.197960072861 km s ⁻¹ Mpc ⁻¹	YES
Gradient target	4.797960072861 km s ⁻¹ Mpc ⁻¹	YES
CPL template	$w_0 = -1.065$, $w_a = -0.066$	YES
BBN loading shift	4.797960068064 10 ⁻⁹ km s ⁻¹ Mpc ⁻¹	× negligible
Shift over radiation-era $H(z_{\text{BBN}})$	$7.421690135465 \times 10^{-27}$	PASS
Chronometer validation	$\chi^2/\nu = 30.16/29$, $\chi_\nu^2 = 1.04$	YES
Forecast χ^2 sensitivity	rigid $\Lambda\text{CDM} = 35.78$, SHBT = 0.08	forecast
Projected ΔBIC sensitivity	+35.70 vs rigid intercept	forecast
Cosmic age	13.276 Gyr for SHBT branch	derived
Thermodynamic arrow	$dS_{\text{total}}/dt_{\text{lc}} > 0$ for sampled $z \geq 0$	YES
Overall precision audit	all listed checks closed	PASS

Table 17: Summary verification ledger from `precision_cosmology_engine.py`.

9.9 Causal Point and GET Measurement

The precision-cosmology observer is the same finite Causal Point introduced in Section 7. The information limit behind its measurement rule is the Holevo bound: for a quantum ensemble

$\{p_i, \rho_i\}$, the extractable classical information satisfies

$$I \leq S\left(\sum_i p_i \rho_i\right) - \sum_i p_i S(\rho_i). \quad (222)$$

Thus a classical record can be crystallized only after the observer pays an entropy cost inside its finite local budget. At a selected boundary address $A \in \mathcal{R}$, the finite outcome ensemble is

$$\Omega_A = \{\omega_0, \omega_1, \dots, \omega_{m-1}\}, \quad \sum_{i=0}^{m-1} p_i = 1, \quad p_i \geq 0. \quad (223)$$

The GET event has address, ensemble, and requested-cost components

$$C_{\text{addr}} = \log_2 |\mathcal{R}|, \quad C_{\text{ens}} = \log_2 |\Omega_A|, \quad C_{\text{get}} = \max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}}). \quad (224)$$

The Causal Point's residual entropy budget is

$$R_{\text{entropy}} = N_{\text{limit}} - C_{\text{get}}, \quad N_{\text{limit}} = \min\left(N_{\text{local}}, \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right). \quad (225)$$

The measurement is admissible if and only if

$$R_{\text{entropy}} \geq 0. \quad (226)$$

When the inequality is satisfied, the collapse index is the deterministic finite-register hash

$$\iota = \text{Hash}(\text{observable_name}, A, f_H, N_{\text{local}}, m) \bmod m. \quad (227)$$

This index defines the rank-one history projector

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \quad (228)$$

For a sequence of admissible GET events, history crystallization is the standard projected density-matrix update

$$\rho_{\text{hist}}^{(n+1)} = \frac{\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)} \Pi_{A_n, \iota_n}}{\text{Tr}\left(\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)}\right)}. \quad (229)$$

The pointer packet used by the audit balances the visible rendered coefficient against the dark completion entry,

$$\Psi_\iota = (\text{amplitude} = \sqrt{p_\iota}, \quad c_{\text{vis}} = -p_\iota, \quad c_{\text{dark}} = p_\iota), \quad (230)$$

so the classical visible record is accompanied by the corresponding completion ledger entry.

The cosmology section therefore extends the SHBT closure program without adding an independent late-time fluid. The Hubble uplift, growth suppression, cluster mirage, ISW stability, entropy arrow, and GET measurement rule are all read as observer-frame projections of the finite completed boundary register. In the notation of this unified manuscript, the statement is concise:

$$(\mathbf{b}_*, \Delta_{\text{fr}} = 0, c_{\text{dark}}^{\text{comp}}, \kappa_{D_5}, N_{\text{sat}}) \implies \left(H_0^{\text{loc}}, f_{\text{load}}(z), S_{\text{debt}}(z), f\sigma_8(z), \chi_\nu^2, \text{ISW}, \Pi_{A,\iota}\right), \quad (231)$$

with each arrow evaluated by the audit tables above.

Quantity	Mathematical definition	Role
Address cost C_{addr}	$\log_2 \mathcal{R} $	register locator cost
Ensemble cost C_{ens}	$\log_2 \Omega_A $	outcome entropy cost
Operational GET cost C_{get}	$\max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}})$	total entropy payment
Budget residual R_{entropy}	$N_{\text{limit}} - C_{\text{get}} \geq 0$	admissibility threshold
Collapse index ι	$\text{Hash}(\dots) \bmod m$	finite-register selector
Projection operator $\Pi_{A,\iota}$	$ A, \omega_\iota\rangle\langle A, \omega_\iota $	history crystallization

Table 18: Causal Point GET retrieval and information-theoretic cost metrics.

10 Unification and Conclusions

The preceding sections describe SHBT from four complementary directions: modular completion of the boundary theory, entropy self-resolution into bulk metric data, anti-baryon de-rendering with passive stress-energy preservation, and entropy-budgeted observer history. The unifying statement is that these are not independent mechanisms. They are different projections of the same finite closed boundary register. Once the completed partition function is modular invariant and the canonical branch arithmetic closes, the bulk tensor, the physical Hamiltonian constraint, the baryogenesis fixed point, and the Causal Point history rule all become consequences of the same closure data.

The core closure chain is

$$\boxed{[M, S_\partial] = [M, T_\partial] = 0 \iff \Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0 \iff \Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0.} \quad (232)$$

Equivalently, $H_{\text{total}} = 0$ after restriction to the physical subspace. The first condition says that the torus partition function is a genuine modular invariant of the completed boundary CFT. The second says that the visible levels and parent completion level have no residual framing mismatch. The third says that the bulk closure tensor vanishes, so the SHBT topological Einstein equation is satisfied without adding an independent bulk correction. The fourth says that the static physical sector has no external time generator left over after projection. Thus the canonical branch is not merely a useful choice of numbers; it is the branch on which boundary modular covariance, visible-sector anomaly closure, bulk geometry, and the Hamiltonian constraint are the same statement viewed through different interfaces.

The geometric part of the construction is summarized by the holographic RG map

$$\boxed{\nabla_{\mathcal{C}} S_\partial \longrightarrow \rho_E(i, j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau).} \quad (233)$$

The boundary entropy gradient is first converted into the normalized entanglement density $\rho_E(i, j)$ on the finite modular-overlap coordinate set. The ordered resolved state is then compressed into the prime-lattice load vector $\ell_r(\tau)$, and the stabilized Gram projection of this load vector gives the effective metric slice $g_{\mu\nu}(\tau)$. In this sense geometry is not a continuous background into which the boundary theory is inserted. It is the rendered image of finite boundary information after entropy gradients have been normalized, ordered, compressed, and projected.

The matter-asymmetry sector closes by the anti-baryon de-rendering loop

$$\boxed{\rho_B^* = \frac{\mathcal{D}_{\bar{B}} \rho_B^* \mathcal{D}_{\bar{B}}^\dagger}{\text{Tr}(\mathcal{D}_{\bar{B}} \rho_B^* \mathcal{D}_{\bar{B}}^\dagger)}, \quad \mathcal{C}_{\bar{B}}(\rho_B^*) = 0, \quad \delta T_{\mu\nu}^{\text{passive}} = 0.} \quad (234)$$

At the fixed point, active anti-baryon color and electromagnetic render charge have been stripped from the visible channel, but the passive gravitational source has not been discarded. It is carried by the dark completion sector. The resulting baryon excess is therefore an asymmetry of rendered visible charge, not a violation of the completed stress-energy account. This is why the same branch that fixes $\Delta_{\text{fr}} = 0$ also supports `stress_energy_preserved=True`: the de-rendering map is allowed only because it is a completion-preserving map.

The observer sector supplies the final link in the same chain. A Causal Point does not read a pre-existing classical history from an infinite archive. It crystallizes a finite ordered sequence of admissible projections,

$$\boxed{\Pi_{\text{history}} = \Pi_{A_m, \iota_m} \cdots \Pi_{A_2, \iota_2} \Pi_{A_1, \iota_1}.} \quad (235)$$

Each factor is admitted only if the local entropy budget remains nonnegative. The apparent classical past is therefore the record that survives the ordered application of finite-register GET operations. The observer is part of the cosmic code rather than an external reader of it: the address A_n , outcome index ι_n , local hidden-state budget, and pointer packet are all internal boundary data.

The principal numerical predictions of the benchmark branch can be collected as

$$\boxed{\Lambda_{\text{holo}} \simeq 1.09 \times 10^{-52} \text{ m}^{-2}, \quad N \simeq 3.31 \times 10^{122}, \quad \eta_B = \mathcal{O}(10^{-10}), \quad g_{\mu\nu} = g_{\mu\nu}(\nabla_C S_\partial).} \quad (236)$$

The first number is the holographic dark-energy curvature scale used by the topological Einstein equation. The second is the finite boundary bit budget associated with that scale. The third is the order of the visible baryon asymmetry produced by the de-rendering loop, with the benchmark audit giving `eta_b=6.449923359416e-10`. The fourth is the structural prediction: the effective metric is not fundamental input but a calculable output of the entropy-gradient pipeline.

Relative to standard holographic approaches, SHBT keeps the boundary-first logic but changes what is taken as primitive. It does not start from a large- N gauge theory with an asymptotically anti-de Sitter bulk and then ask for a semiclassical interior. It starts from a completed static boundary CFT, requires exact modular closure, and lets finite entropy data render the bulk metric. Relative to entropic-gravity pictures, SHBT does not infer gravity from entropy alone; it constrains the entropy flow by the modular invariant pairing matrix, the canonical branch arithmetic, and the finite register size. Relative to purely constraint-based quantum gravity, it does not stop at the formal statement that the Hamiltonian vanishes on physical states; it ties that statement to visible-sector levels, dark completion, baryogenesis, and observer memory.

The most direct tests of the framework are therefore consistency tests across sectors. The branch data must give integer locks $I_\ell^* = 6$ and $I_q^* = 13$, zero framing defect, a vanishing bulk closure tensor, and a finite physical-sector Hamiltonian constraint. The entropy pipeline must produce symmetric, trace-normalized, positive metric slices from the same boundary densities that define the loading sequence. The baryogenesis module must remove active visible anti-baryon charge while keeping passive stress-energy unchanged. The Causal Point module must crystallize only

those histories whose retrieval cost fits inside the local entropy budget. Any failure of one of these checks would be a failure of the unified SHBT closure, not merely a failure of an auxiliary numerical routine.

Several limitations remain. The present paper establishes and audits one canonical benchmark branch rather than a classification theorem for all modular completions. The dark completion sector is fixed by closure requirements but still needs a more explicit microscopic construction. The entropy-to-metric map is deterministic in the benchmark implementation, yet a full comparison with precision cosmological data would require a larger phenomenological layer: survey observables, perturbations, noise models, and parameter-estimation pipelines. The baryogenesis mechanism also needs extension from the fixed render-channel identity to a complete account of early-universe thermal history as seen by finite boundary observers. These are not optional decorations; they are the natural next checks if SHBT is to move from a closed auditable framework to a quantitatively predictive research program.

The cosmic-code perspective is the simplest way to state the conclusion. A finite boundary register, when forced to close under completed modular kernels, contains more than a list of states. It contains the rules for which visible charges can be rendered, which passive sources must be preserved, which entropy gradients become geometry, and which observer histories can crystallize. In this view spacetime is the readable projection of a finite holographic code; matter asymmetry is a render-channel fixed point of that code; and observation is the entropy-budgeted act of selecting a consistent history from within the same code. SHBT is therefore static only at the level of the completed boundary. The dynamical world seen by an observer is the internally crystallized image of that closed finite boundary information.

11 Code Availability

The reference implementation and audit scripts for the SHBT and precision cosmology calculations are maintained in the GitHub repository <https://github.com/sys1own/Static-Holographic-Boundary-Theory>. During private review, the same repository state should be supplied as a release archive with the paper source. The publication-critical entry points are:

For the precision-cosmology layer, the names used in Table 9 are the canonical audit entry points: `redshift_dependent_hubble_constant`, `build_structure_growth_audit`, `build_cluster_collapse_audit`, `build_early_universe_stability_audit`, `audit_isw_stability`, `build_cosmic_chronometer_validation`, `build_thermodynamic_arrow_audit`, and `build_precision_cosmology_audit`.

Appendices: Modular Framing and Quantum-Dimension Derivations

The following appendices incorporate the supplementary material from `supplementary.tex`. They use the unified notation of the main text: the appendix alias `\cdark` is renewed immediately above

Script or module	Paper role
<code>shbt_simulate.py</code>	top-level SHBT simulator and canonical branch audit
<code>shbt_simulator</code>	Rust/Python boundary, projection, Causal Point, and baryogenesis module
<code>examples/run_audit.py</code>	reproducibility command for the full foundation audit
<code>noether_bridge.py</code>	completed-ledger Newton lock, unity-of-scale, and neutrino hierarchy report
<code>precision_cosmology_engine.py</code>	precision-cosmology audit module; see the function list above for Hubble loading, growth, cluster, BBN, ISW, chronometer, and light-cone ledgers
<code>baryon_asymmetry.py</code>	topological Sakharov and η_B benchmark audit
<code>holographic_tension_verifier.py</code>	finite-register and neutrino-floor tension checks
<code>rigidity_landscape.py</code>	fixed-parent rigidity and branch-uniqueness scan
<code>measurement_operator.py</code> and <code>causal_execution.py</code>	finite-memory GET operator and Causal Point execution loop

Table 19: Code components required to reproduce the paper-level audit tables.

so every printed dark-ledger symbol in the appendices denotes the completed ledger $c_{\text{dark}}^{\text{comp}}$, while the residual intermediate remains $c_{\text{dark}}^{\text{res}}$.

These supplementary sections record the exact current-algebra bookkeeping used by the preceding sections. Their purpose is administrative and mathematical: the quantities $K = 312$, $c_{\text{dark}}^{\text{comp}} = 1197103/362670$, and $\kappa_{D_5} = 0.988769793998$ are fixed before any low-energy cosmological evolution is evaluated. They therefore enter the cosmology calculation as an invariant boundary chart, not as phenomenological settings or posterior-fit variables.

A Center-Directed Modular Framing

The visible affine factors are organized by their gauge centers. The weak and color centers are

$$Z(SU(2)) = \mathbb{Z}_2, \quad Z(SU(3)) = \mathbb{Z}_3. \quad (237)$$

A parent modular frame must close simultaneously on the spin-parity divisor of the weak factor and on the triality divisor of the color factor. For visible levels k_ℓ and k_q , this requires the parent level to lie in the common integer lattice

$$K \in 2k_\ell\mathbb{Z} \cap 3k_q\mathbb{Z}. \quad (238)$$

Equivalently, the parent level is the smallest positive integer for which both center lifts are integral:

$$\frac{K}{2k_\ell} \in \mathbb{Z}, \quad \frac{K}{3k_q} \in \mathbb{Z}. \quad (239)$$

For the SHBT visible branch,

$$k_\ell = 26, \quad k_q = 8. \quad (240)$$

Thus the minimal anomaly-free parent level is

$$K_{\text{min}} = \text{lcm}(2k_\ell, 3k_q) = \text{lcm}(52, 24) = 312. \quad (241)$$

The embedding indices are then not adjustable. They are the two center-lift quotients determined directly by Eq. (241):

$$I_\ell^* = \frac{K_{\min}}{2k_\ell} = \frac{312}{52} = 6, \quad I_q^* = \frac{K_{\min}}{3k_q} = \frac{312}{24} = 13. \quad (242)$$

These integers are the microscopic origin of the locked modular lift: changing one of them would reopen either the weak spin-parity defect or the color triality defect, so no continuous deformation of the low-energy branch is available.

B Coset Anomaly Extractions and Vacuum Subtractions

For a simple affine Lie algebra $\mathfrak{g}_k$, the Sugawara central charge is

$$c(\mathfrak{g}_k) = \frac{k \dim \mathfrak{g}}{k + h_{\mathfrak{g}}^\vee}, \quad (243)$$

where $h_{\mathfrak{g}}^\vee$ is the dual Coxeter number. Applying Eq. (243) to the branch fixed by Eq. (241) gives

$$c(SO(10)_{312}) = \frac{312 \cdot 45}{312 + 8} = \frac{351}{8}, \quad (244)$$

$$c(SU(2)_{26}) = \frac{26 \cdot 3}{26 + 2} = \frac{39}{14}, \quad (245)$$

$$c(SU(3)_8) = \frac{8 \cdot 8}{8 + 3} = \frac{64}{11}, \quad (246)$$

$$c(U(1)_Y) = 1. \quad (247)$$

Before the finite boundary subtraction, the parent-minus-visible coset ledger is

$$c_{\text{raw}} = \frac{351}{8} - \frac{39}{14} - \frac{64}{11} - 1 = \frac{21111}{616}. \quad (248)$$

The finite horizon screen does not retain the inert spinorial vacuum block as a propagating low-energy coset degree of freedom. Tracing out that 32-dimensional block gives the positive residual vacuum coordinate

$$\Delta c_0 = c_{\text{raw}} - 32 = \frac{21111}{616} - 32 = \frac{1399}{616}. \quad (249)$$

The chiral hypercharge lattice trace is not erased; it is integrated together with the tracking spin defect

$$\delta'_{\text{spin}} = \frac{135797}{131880}. \quad (250)$$

The locked dark completion fraction is therefore

$$c_{\text{dark}}^{\text{comp}} = \Delta c_0 + \delta'_{\text{spin}} = \frac{1399}{616} + \frac{135797}{131880} = \frac{1197103}{362670} \simeq 3.300805. \quad (251)$$

This number is positive, rational, and tied to the center-directed parent level. Its alternating prime partial-fraction decomposition is

$$c_{\text{dark}}^{\text{comp}} = 3 + \frac{1}{2} - \frac{1}{3} + \frac{2}{5} - \frac{3}{7} + \frac{2}{11} - \frac{3}{157}. \quad (252)$$

The denominator is correspondingly

$$362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157, \quad (253)$$

so the residue records the same sequential center-framing data as the affine embedding. It is not a decimal tuning of a low-energy observable.

C Boundary Measure and Quantum Dimensions

The boundary cell measure is computed from the Frobenius–Perron total quantum dimension of the leptonic anyonic fusion ring. For $SU(2)_k$, the simple-object quantum dimensions are

$$d_j = \frac{\sin\left(\frac{(j+1)\pi}{k+2}\right)}{\sin\left(\frac{\pi}{k+2}\right)}, \quad j = 0, 1, \dots, k, \quad (254)$$

and hence

$$\mathcal{D}(SU(2)_k)^2 = \sum_{j=0}^k d_j^2 = \frac{k+2}{2 \sin^2\left(\frac{\pi}{k+2}\right)}. \quad (255)$$

At the locked leptonic level $k_\ell = 26$, this gives

$$\mathcal{D}(SU(2)_{26}) = \frac{\sqrt{14}}{\sin(\pi/28)} \simeq 33.4182547625. \quad (256)$$

The retained spinorial hyperarea cell is the complement of the normalized spinorial leakage weight in this fusion ring. The five spinorial Cartan signs supply the unoriented sheet weight $5^2/2$, while the color-vector conversion couples in the locked triality quotient $I_q^*/K = 13/312$. Thus

$$\rho_{\text{spin} \rightarrow \text{vec}} = \frac{5^2}{2} + \frac{I_q^*}{K} = \frac{25}{2} + \frac{13}{312} = \frac{3913}{312}. \quad (257)$$

The non-deformable boundary hyperarea cell measure is therefore

$$\kappa_{D_5} = 1 - \frac{\rho_{\text{spin} \rightarrow \text{vec}}}{\mathcal{D}(SU(2)_{26})^2} \quad (258)$$

$$= 1 - \frac{3913}{312} \frac{\sin^2(\pi/28)}{14} = 0.988769793998 \dots \quad (259)$$

This is the spinorial retention factor used when the boundary state is converted into the vector variables of the low-energy effective action. In the notation used above, the conversion is

$$V_a^{\text{retained}} = \kappa_{D_5} V_a^{\text{parent}}, \quad d\mathcal{A}_{D_5}^{\text{retained}} = \kappa_{D_5}^4 d\mathcal{A}_{D_5}^{\text{parent}}. \quad (260)$$

Consequently the same invariant cell measure appears in the surface-tension matching relation,

$$\Lambda_{\text{holo}} = \frac{3\pi}{\kappa_{D_5}^4} G_{\text{top}} m_{\nu,1}^4. \quad (261)$$

The same lightest eigenstate propagates through the normal-ordering ledger used above,

$$m_{\nu,1} \simeq 2.83 \times 10^{-3} \text{ eV}, \quad (262)$$

$$m_{\nu,2} \simeq 8.92 \times 10^{-3} \text{ eV}, \quad (263)$$

$$m_{\nu,3} \simeq 5.01 \times 10^{-2} \text{ eV}. \quad (264)$$

Equations (251), (259), and (262)–(264) are the microscopic affine-data inputs to the cosmological branch. They are fixed by modular closure, finite vacuum subtraction, and fusion-ring quantum dimension before any Hubble-rate, growth, or neutrino observable is computed.

References

- [1] G. 't Hooft, “Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking,” in *Recent Developments in Gauge Theories*, NATO Advanced Study Institutes Series B, vol. 59, pp. 135–157, Plenum Press, 1980.
- [2] S. Weinberg, “The cosmological constant problem,” *Reviews of Modern Physics* **61**, 1–23 (1989).
- [3] S. Navas *et al.* (Particle Data Group), “Review of Particle Physics,” *Physical Review D* **110**, 030001 (2024).
- [4] J. L. Cardy, “Operator content of two-dimensional conformally invariant theories,” *Nuclear Physics B* **270**, 186–204 (1986).
- [5] A. Cappelli, C. Itzykson, and J.-B. Zuber, “Modular invariant partition functions in two dimensions,” *Nuclear Physics B* **280**, 445–465 (1987).
- [6] C. Fefferman and C. R. Graham, “Conformal invariants,” in *Elie Cartan et les mathematiques d’aujourd’hui*, Asterisque, hors serie, pp. 95–116, Societe Mathematique de France, 1985.
- [7] J. D. Bekenstein, “Black holes and entropy,” *Physical Review D* **7**, 2333–2346 (1973).
- [8] G. 't Hooft, “Dimensional reduction in quantum gravity,” arXiv:gr-qc/9310026 (1993).
- [9] L. Susskind, “The world as a hologram,” *Journal of Mathematical Physics* **36**, 6377–6396 (1995).
- [10] A. D. Sakharov, “Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe,” *JETP Letters* **5**, 24–27 (1967).
- [11] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Advances in Theoretical and Mathematical Physics* **2**, 231–252 (1998), arXiv:hep-th/9711200.
- [12] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Physics Letters B* **428**, 105–114 (1998), arXiv:hep-th/9802109.
- [13] E. Witten, “Anti de Sitter space and holography,” *Advances in Theoretical and Mathematical Physics* **2**, 253–291 (1998), arXiv:hep-th/9802150.

- [14] S. Ryu and T. Takayanagi, “Holographic derivation of entanglement entropy from AdS/CFT,” *Physical Review Letters* **96**, 181602 (2006), arXiv:hep-th/0603001.
- [15] J. de Boer, E. Verlinde, and H. Verlinde, “On the holographic renormalization group,” *Journal of High Energy Physics* **2000**, 003 (2000), arXiv:hep-th/9912012.
- [16] K. Skenderis, “Lecture notes on holographic renormalization,” *Classical and Quantum Gravity* **19**, 5849–5876 (2002), arXiv:hep-th/0209067.
- [17] E. Verlinde, “On the origin of gravity and the laws of Newton,” *Journal of High Energy Physics* **2011**, 029 (2011), arXiv:1001.0785.