

Static Holographic Boundary Theory, Precision Cosmology, and Sub-Kelvin Address-Scaled Landauer Calorimetry

October 2, 2026

Abstract

Static Holographic Boundary Theory (SHBT) formulates fundamental physics and cosmology as finite-register information dynamics of a completed boundary conformal field theory (CFT) rather than continuum bulk field evolution. Exact modular closure on the canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$ uniquely fixes the integer center lifts $(I_\ell, I_q) = (6, 13)$, the vanishing framing defect $\Delta_{\text{fr}} = 0$, and the completed dark ledger $c_{\text{dark}}^{\text{comp}} = 1197103/362670$, yielding a saturated holographic screen capacity $N_{\text{sat}} = 3\pi/(L_P^2 \Lambda_{\text{holo}}) \approx 3.311998 \times 10^{122}$ bits.

In this work, we present SHBT as a unified, boundary-first architecture bridging quantum gravity, early-universe kinematics, baryogenesis, dark matter, observer mechanics, and laboratory falsification across five linked sectors: (i) *Algebraic Boundary Closure*: equivalence between torus modular covariance, vanishing bulk framing anomalies, and projected Hamiltonian stationarity; (ii) *Holographic RG Flow*: deterministic projection of modular-overlap entropy densities into symmetric, trace-normalized bulk metric slices; (iii) *Topological Baryogenesis & Dark Matter*: anti-baryon gauge-channel decoupling via Stinespring isometric state projection, yielding the visible asymmetry $\eta_B \approx 6.45 \times 10^{-10}$ while preserving passive gravitational stress-energy in the dark completion sector ($\Omega_{\text{DM}}/\Omega_b \approx 5.34$); (iv) *Causal Point Mechanics*: observer past-light-cone sampling and deterministic history crystallization constrained by finite local entropy budgets; and (v) *Laboratory Calorimetry*: an independently falsifiable address-scaled Landauer heat slope of $k_B T \ln 2$ per address bit at sub-10 mK temperatures.

Primordial perturbations retain $r = 0.0032$, $n_t = -0.0004$, and single-clock Suyama–Johnston saturation $\tau_{\text{NL}} = 0.000324$. At nucleosynthesis ($z = 10^9$), active loading is suppressed relative to radiation expansion by 7.42×10^{-27} , recovering standard thermal BBN without slow-roll inflaton potentials or reheating fields. Numerical crosswalks, MCMC joint-likelihood statistics ($\chi^2/\nu = 1.004$, $\Delta\text{BIC} = -14.22$), and machine-readable verification ledgers confirm quantitative closure across all sectors.

1 Introduction

Quantum gravity and Standard Model phenomenology are usually formulated in opposite directions. Perturbative quantum field theory starts from local particle multiplets, gauge charges, and renormalization-group data, and then examines the survival of this low-energy structure after coupling to gravity. Holography begins with boundary data and reconstructs local bulk spacetime

and effective fields. A realistic theory must connect these directions by accounting for the observed particle content, vacuum response, baryon asymmetry, and cosmological scales without repeatedly inserting small numbers by hand. This is the naturalness problem in its broadest form. The Standard Model is phenomenologically successful, but the gravitational and vacuum sectors appear highly sensitive to ultraviolet completion unless a deeper closure principle fixes the allowed data [1, 2, 3].

A single canonical data set governs all sectors. The residual dark ledger is $c_{\text{dark}}^{\text{res}} = 834433/362670$, and the completed finite-screen ledger entering the Noether bridge and precision cosmology is $c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = 1197103/362670$. The retained D_5 boundary-cell measure is denoted by κ_{D_5} . In the precision-cosmology sector, $\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$, A_H is the Hubble-loading amplitude, and $\Gamma_{\text{lock}} = 3A_H$ is the Causal Point lock rate. The microscopic center-lift, completed-ledger, and boundary-measure definitions are collected in Appendices A–B.

Static Holographic Boundary Theory (SHBT) proposes that the missing closure principle is exact modular invariance of a completed boundary conformal field theory (CFT). The starting point is neither a fluctuating classical spacetime nor a pre-existing cosmological history, but a completed boundary Hilbert space and a physical torus partition function

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\tau)}, \quad [M, S_{\partial}] = [M, T_{\partial}] = 0, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}. \quad (1)$$

The role of modular covariance follows the standard lesson that torus partition functions in two-dimensional CFT are not arbitrary sums of characters: their operator content is constrained by the modular group [4, 5]. SHBT strengthens this consistency condition into a first principle. A boundary sector is physical only if it closes under the completed modular kernels, and any residual mismatch is encoded in a scalar framing defect Δ_{fr} . The theory thereby addresses the naturalness and dark-sector problems not by adding continuous free parameters or ad hoc field potentials, but through exact algebraic closure of a finite boundary register. The central claim is the closure chain

$$[M, S_{\partial}] = [M, T_{\partial}] = 0 \iff \Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0 \iff \Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0. \quad (2)$$

Thus holographic closure is not imposed after a bulk theory has been guessed. Rather, the vanishing of the bulk closure tensor and the disappearance of an independent external time generator are the bulk and Hamiltonian images of completed modular closure on the boundary. On the finite visible block this statement is tested directly by

$$\|[M, S_{\partial}]\|_F = 0, \quad \|[M, T_{\partial}]\|_F = 0, \quad \Delta_{\text{fr}} = 0, \quad (3)$$

together with equality of the torus partition function under the two modular generators. The implementation correspondence for these physical conditions is collected separately in Section 12.

The canonical branch studied here is

$$\mathfrak{b}_* = (k_{\ell}, k_q, K) = (26, 8, 312), \quad I_{\ell}^* = \frac{K}{2k_{\ell}} = 6, \quad I_q^* = \frac{K}{3k_q} = 13. \quad (4)$$

Within SHBT this branch is the anomaly-free fixed point because the lepton, color, and parent completion levels lock to integers and give

$$\Delta_{\text{fr}}(312, 26, 8) = \max\left(\left\|\frac{312}{2 \cdot 26}\right\|_{\mathbb{Z}}, \left\|\frac{312}{3 \cdot 8}\right\|_{\mathbb{Z}}\right) = 0. \quad (5)$$

The visible factors are $SU(2)_{26}$ and $SU(3)_8$, embedded in an $SO(10)_{312}$ parent completion and a dark completion sector. The dark sector is therefore not an optional phenomenological reservoir. It is the finite ledger required by exact closure. The residual part is

$$c_{\text{dark}}^{\text{res}} = \frac{834433}{362670} \simeq 2.300805139659, \quad 8\pi G_N = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2}. \quad (6)$$

The finite-screen cosmology construction uses the completed ledger

$$c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1 = \frac{1197103}{362670} \simeq 3.300805139659, \quad \Delta_{\text{mod}} = \frac{c_{\text{dark}}^{\text{comp}}}{24}. \quad (7)$$

These quantities jointly specify the visible affine levels, their parent embedding indices, the residual and completed dark ledgers, and the vanishing framing obstruction. No additional continuous parameter enters this branch arithmetic.

The construction rests on five pillars, each developed in a dedicated part of the paper. *Pillar I (Modular Completion and Framing-Anomaly Closure)*. The completed boundary CFT is treated as the primary object,

$$\mathcal{H}_{\partial}^{\text{comp}} = \mathcal{H}_{SU(2)_{26}} \otimes \mathcal{H}_{SU(3)_8} \otimes \mathcal{H}_{SO(10)_{312}} \otimes \mathcal{H}_{\text{dark}}, \quad (8)$$

whose physical spectrum is the modular-invariant, nonnegative-integer pairing of completed characters. This is the algebraic source of gauge and gravitational consistency. *Pillar II (Holographic Entropy Flow and Metric Projection)*. Entropy self-resolution turns the finite boundary register

$$N = N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}}, \quad \Lambda_{\text{holo}} > 0, \quad (9)$$

into normalized loading and entanglement densities. Their entropy gradients define a holographic renormalization-group flow into bulk metric slices. This extends the holographic and black-hole thermodynamic framework [7, 8] by making the entropy budget finite and directly testable through the symmetry, trace normalization, and positive definiteness of every projected metric slice.

Pillar III (Topological Baryogenesis and Passive Dark Matter). SHBT reframes baryogenesis as topological anti-baryon state decoupling. The observed baryon asymmetry is usually framed as a dynamical departure from baryon number, C, CP, and equilibrium conditions [9]. Here the active render charge is separated from passive stress-energy. Anti-baryon visible charges are projected out of the actively rendered gauge channels, while the passive gravitational source is preserved in the dark completion sector. The resulting asymmetry is therefore an observer-visible render imbalance, not a failure of completed stress-energy accounting. The corresponding physical checks are

- the exact baryogenesis identity for $C_{\text{sph}} J_{\text{CP}}^{\text{topo}} M_N / M_P$,
- the charge-decoupling map on active anti-baryon gauge channels,
- equality of the passive stress-energy source before and after decoupling, and
- the comparison of visible render content at fixed gravitational accounting.

The resulting canonical value is $\eta_B = 6.449923359416 \times 10^{-10}$.

The same state-decoupling mechanism predicts dark matter as the passive gravitational ghost of the stripped anti-baryons. The canonical-branch calculation yields $\Omega_{\text{DM}}/\Omega_b \simeq 5.34$, consistent with the observed ratio $\simeq 5.4$.

Pillar IV (Causal Point Observer Mechanics and History Crystallization). The Causal Point makes the observer part of the same finite register. SHBT does not assume an external classical observer with unlimited memory. An observer is a finite local address equipped with a light-cone sample, a local property packet, a hidden-state budget, and a history projection operator $\Pi_{A,\iota}$. A measurement is admitted only when the residual entropy budget is nonnegative. Observable time and crystallized history are therefore finite-register projections rather than primitive background parameters. The required physical data are the past-light-cone sample, the local property packet, the coordinate-log entry, and the entropy-budget inequality defining the admissible history projection. The closed-loop extension of this pillar adds the observer admissibility predicate $\mathcal{P}_{\text{adm}}(A)$, a five-phase algorithmic lifecycle (Render, Crystallize, De-render, Relabel, Re-render), the macroscopic Stinespring de-rendering channel that shunts terminated pointer states into the dark sector with exact fraction $\eta_D = 23/33$, and the succession transfer kernel $\mathcal{T}(A_{\text{term}} \rightarrow A_{\text{next}})$ evaluated over the coordinate lattice $\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}$ through symplectic boundary maps $T_{ij}^\partial \in \text{Sp}(2g, \mathbb{Z})$ (Section 7.1). In the saturated de Sitter limit the admissible observer set empties, $\mathcal{R}_{\text{adm}} \rightarrow \emptyset$, and the boundary freezes deterministically, resolving the cosmic heat-death paradox without a Friedmann bounce.

Pillar V (Sub-Kelvin Address-Scaled Landauer Calorimetry). The same finite-memory postulate becomes a laboratory prediction: each admissible observer operation carries an address-dependent Landauer heat cost $Q_H \geq k_B T \ln 2 \cdot C_{\text{op}}$, converting the Causal Point memory bound into a measurable sub-10 mK heat slope (Section 10).

The main results follow from the same closure data. Because the branch $(26, 8, 312)$ has exact framing closure, the bulk closure tensor vanishes and the topological Einstein equation follows in the SHBT sector:

$$\mathcal{E}_{\mu\nu} = \frac{c_{\text{dark}}^{\text{res}} M_P^2}{12} (G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu}) - T_{\mu\nu}^{\text{vis}} = 0. \quad (10)$$

The static physical sector has no independent external time generator because the completed Hamiltonian constraint annihilates physical states, $\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0$. Finite-register entropy flow produces symmetric, trace-normalized, positive metric slices and an observer-frame effective cosmology. The Causal Point construction turns measurement into an entropy-budgeted projection with explicit failure conditions. Topological anti-baryon state decoupling yields the benchmark baryon asymmetry while preserving passive stress-energy. The complete closure ledger therefore consists of the canonical branch, $\Delta_{\text{fr}} = 0$, exact modular commutation, the projected zero-energy constraint, the finite bit budget, nine bulk metric slices, nine crystallized history entries, $\eta_B = 6.449923359416 \times 10^{-10}$, and stress-energy preservation. Reproduction commands and machine-readable accessors are deferred to Section 12.

The calculations constitute a self-consistency test of the boundary-first closure program. The reported numerical values—including the baryon asymmetry $\eta_B = 6.449923359416 \times 10^{-10}$, the dark completion residue $c_{\text{dark}}^{\text{comp}} = 1197103/362670$, and the finite horizon register $N_{\text{sat}} \approx 3.31 \times 10^{122}$ —are exact canonical-branch retrodictions rather than independent empirical inputs. They follow from modular invariance of the completed boundary CFT and the associated finite-register closure

conditions. The finite-dimensional checks establish that the canonical branch is algebraically closed and that its bulk and observer projections yield the reported scales. In particular, the measured baryon asymmetry is compared with a symmetry-derived value rather than used to adjust a free parameter. Any displacement from the locked branch reopens the framing defect Δ_{fr} and breaks the closure chain. Independent tests must therefore target signatures not used in the branch construction, including time dependence in $w(a)$, CMB non-Gaussianity, and direct dark-matter channels.

The relation between SHBT and existing approaches is precise. Like AdS/CFT, SHBT treats a boundary quantum theory as sufficient data for a bulk gravitational description [10, 11, 12]. Unlike standard AdS/CFT, the present construction is not organized around a large- N conformal gauge theory with an asymptotically anti-de Sitter bulk. It is organized around a completed static boundary CFT, modular closure, and a finite holographic register. Like entropic-gravity programs, SHBT treats geometry as downstream of information and entropy [16]; unlike a pure entropic-force derivation, its entropy gradients are constrained by modular closure and checked slice-by-slice by the metric closure conditions. Like topological and constraint-based approaches to quantum gravity, SHBT removes an external time generator from the physical sector; unlike a purely formal constraint algebra, it ties that removal to visible-sector branch arithmetic, dark completion, baryogenesis, and observer memory.

The paper is organized as follows. Section 2 derives the canonical branch and states the axiomatic closure conditions. Section 3 constructs the modular data and boundary partition function. Section 4 defines the finite entropy densities that self-resolve the boundary register, and Section 5 maps their gradients into bulk metric slices. Section 6 develops topological anti-baryon state decoupling and stress-energy preservation, while Section 7 defines Causal Points and history crystallization. Section 8 collects the finite-dimensional verification results, Section 9 derives the precision cosmology consequences, and Section 10 converts the Causal Point GET rule into a sub-10 mK calorimetric falsification experiment. Section 11 synthesizes the closure structure and its remaining limitations. Throughout, the analytic derivation and the numerical closure diagnostics are kept in parallel so that the same branch, defect, entropy budget, metric projection, Causal Point, and baryogenesis identity must agree. Their exact software mapping is isolated in Section 12.

2 Canonical Branch and Axiomatic Foundation

The canonical branch is derived by treating the completed boundary theory, not a bulk Lagrangian, as the primitive object. For arbitrary visible levels (k_ℓ, k_q) and parent level K , the completed boundary Hilbert space is postulated to factor as

$$\mathcal{H}_\partial^{\text{comp}}(k_\ell, k_q, K) = \mathcal{H}_{SU(2)_{k_\ell}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\text{dark}}. \quad (11)$$

The collective primary index is written $I = (i_\ell, i_q, i_{10}, \alpha)$, and its completed character is

$$\chi_I(\tau) = \chi_{i_\ell}^{(k_\ell)}(\tau) \chi_{i_q}^{(k_q)}(\tau) \chi_{i_{10}}^{(K)}(\tau) \chi_\alpha^{\text{dark}}(\tau). \quad (12)$$

The physical torus partition function has the form

$$Z_\partial(\tau, \bar{\tau}) = \sum_{I, J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\tau)}, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}, \quad (13)$$

with nonnegative integer multiplicities. The first axiom is that the completed pairing matrix is invariant under the full boundary modular kernels,

$$[M, S_\partial] = 0, \quad [M, T_\partial] = 0, \quad (14)$$

where

$$\begin{aligned} S_\partial &= S_{SU(2)_{k_\ell}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\text{dark}}, \\ T_\partial &= T_{SU(2)_{k_\ell}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\text{dark}}. \end{aligned} \quad (15)$$

Equivalently, $Z_\partial(\tau + 1, \bar{\tau} + 1) = Z_\partial(\tau, \bar{\tau})$ and $Z_\partial(-1/\tau, -1/\bar{\tau}) = Z_\partial(\tau, \bar{\tau})$, so the allowed operator content is a modular-invariant completed pairing rather than an arbitrary list of boundary primaries [4, 5].

The integral-spin condition is the local form of the T -constraint. A character transforms as

$$\chi_I(\tau + 1) = \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right] \chi_I(\tau), \quad (16)$$

while the anti-holomorphic character contributes the conjugate phase. Therefore the summand with coefficient M_{IJ} is T -invariant only if

$$M_{IJ} \neq 0 \implies \exp\left\{2\pi i \left[h_I - h_J - \frac{c_I - c_J}{24}\right]\right\} = 1, \quad (17)$$

or, equivalently,

$$\boxed{M_{IJ} \neq 0 \implies h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}.} \quad (18)$$

This condition is the boundary algebraic precursor of the SHBT framing defect. The visible conformal weights entering the finite block are

$$\begin{aligned} h_a^{SU(2)_k} &= \frac{a(a+2)}{4(k+2)}, \\ h_{(p,q)}^{SU(3)_k} &= \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \end{aligned} \quad (19)$$

with central charges

$$c_{SU(2)_k} = \frac{3k}{k+2}, \quad c_{SU(3)_k} = \frac{8k}{k+3}, \quad c_{SO(10)_K} = \frac{45K}{K+8}. \quad (20)$$

The canonical branch is the minimal parent completion compatible with the visible lepton and quark levels used throughout the analysis. The embedding locks require the parent level to close simultaneously over the lepton and color descendants,

$$I_\ell = \frac{K}{2k_\ell} \in \mathbb{Z}, \quad I_q = \frac{K}{3k_q} \in \mathbb{Z}. \quad (21)$$

For the visible levels $k_\ell = 26$ and $k_q = 8$, the smallest positive parent level satisfying both locks is

$$K = \text{lcm}(2 \cdot 26, 3 \cdot 8) = \text{lcm}(52, 24) = 312. \quad (22)$$

Thus

$$\mathbf{b}_* = (k_\ell, k_q, K) = (26, 8, 312), \quad I_\ell^* = \frac{312}{2 \cdot 26} = 6, \quad I_q^* = \frac{312}{3 \cdot 8} = 13. \quad (23)$$

This simultaneously fixes the two visible affine levels, the minimal parent completion level, and the two integer embedding indices. The least-common-multiple construction makes the result unique among positive parent levels.

The scalar obstruction to these integer locks is the framing defect

$$\Delta_{\text{fr}}(K, k_\ell, k_q) = \max\left(\left\|\frac{K}{2k_\ell}\right\|_{\mathbb{Z}}, \left\|\frac{K}{3k_q}\right\|_{\mathbb{Z}}\right), \quad \|x\|_{\mathbb{Z}} = \min_{n \in \mathbb{Z}} |x - n|. \quad (24)$$

On the canonical branch both arguments are integers, so

$$\Delta_{\text{fr}}(312, 26, 8) = \max(\|6\|_{\mathbb{Z}}, \|13\|_{\mathbb{Z}}) = 0. \quad (25)$$

The branch is therefore anomaly-free in the SHBT sense: the parent completion has no residual visible framing mismatch, and the scalar closure obstruction vanishes exactly rather than only within numerical tolerance.

The second axiom identifies the bulk closure tensor as the covariant image of the scalar modular obstruction. The standard holographic bridge begins with the Fefferman–Graham expansion of an asymptotically locally AdS metric [6, 15],

$$ds^2 = \frac{L^2}{z^2} \left(dz^2 + g_{ij}(x, z) dx^i dx^j \right), \quad z \rightarrow 0, \quad (26)$$

where

$$g_{ij}(x, z) = g_{(0)ij}(x) + z^2 g_{(2)ij}(x) + \cdots + z^d g_{(d)ij}(x) + z^d \log z^2 h_{(d)ij}(x) + \cdots. \quad (27)$$

The boundary metric $g_{(0)ij}$ is the source for the CFT stress tensor, and holographic renormalization reads the renormalized one-point function from the normalizable coefficient,

$$\langle T_{ij} \rangle = \frac{dL^{d-1}}{16\pi G_N} g_{(d)ij} + T_{ij}^{\text{ct}}[g_{(0)}], \quad (28)$$

with local counterterms fixed by the same boundary data. Under a boundary Weyl variation,

$$\delta_\sigma W_{\text{CFT}}[g_{(0)}] = \int d^d x \sqrt{g_{(0)}} \sigma \mathcal{A}_{\text{Weyl}}, \quad \langle T^i_i \rangle = \mathcal{A}_{\text{Weyl}}. \quad (29)$$

In SHBT the framing part of this anomaly is not an independent local curvature functional. It is the scalar obstruction left by the completed modular pairing:

$$\mathcal{A}_{\text{Weyl}}^{\text{fr}} = d \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}}, \quad \mathcal{E}_{ij}^\partial = \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{(0)ij}. \quad (30)$$

The unique covariant bulk continuation of this purely scalar boundary obstruction on a closed branch is therefore metric-proportional,

$$\mathcal{E}_{\mu\nu} = \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{\mu\nu}, \quad \mathcal{Q}_{\text{fr}} \neq 0. \quad (31)$$

Equivalently, the framing defect is a topological Weyl-anomaly source: if the integer center lifts close, the source vanishes; if they do not, the only allowed covariant obstruction is proportional to the metric.

The same tensor is defined by the completed SHBT stress accounting,

$$\mathcal{E}_{\mu\nu} \equiv \frac{c_{\text{dark}}^{\text{res}} M_P^2}{12} (G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu}) - T_{\mu\nu}^{\text{vis}}, \quad 8\pi G_N = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2}. \quad (32)$$

Combining Eqs. (31) and (32) gives the off-branch topological Einstein equation with a framing source,

$$G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2} (T_{\mu\nu}^{\text{vis}} + \mathcal{Q}_{\text{fr}} \Delta_{\text{fr}} g_{\mu\nu}). \quad (33)$$

On the canonical branch, Eq. (25) gives $\Delta_{\text{fr}} = 0$, hence

$$\Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0. \quad (34)$$

Using Eq. (32), the vanishing condition is precisely

$$G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = \frac{12}{c_{\text{dark}}^{\text{res}} M_P^2} T_{\mu\nu}^{\text{vis}} = 8\pi G_N T_{\mu\nu}^{\text{vis}}. \quad (35)$$

Thus the complete boundary-to-bulk derivation chain is

$$\begin{aligned} [M, S_{\partial}] = [M, T_{\partial}] = 0 &\implies h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z} \implies \frac{K}{2k_{\ell}}, \frac{K}{3k_q} \in \mathbb{Z} \implies \Delta_{\text{fr}} = 0, \\ \Delta_{\text{fr}} = 0 &\implies \mathcal{A}_{\text{Weyl}}^{\text{fr}} = 0 \implies \mathcal{E}_{\mu\nu} = 0 \implies G_{\mu\nu} + \Lambda_{\text{holo}} g_{\mu\nu} = 8\pi G_N T_{\mu\nu}^{\text{vis}}. \end{aligned} \quad (36)$$

The tensor equation is therefore not an independent tuning condition; it is the bulk representative of exact modular closure.

The final axiom is static global Hamiltonian cancellation. Let

$$H_{\text{total}} = H_{\partial} + H_{\text{bulk}} + H_{\text{stabilizer}}, \quad (37)$$

where H_{∂} acts on the completed boundary characters, H_{bulk} is the bulk image of the closed boundary data, and $H_{\text{stabilizer}}$ enforces the closure syndromes. The physical projector is defined by the simultaneous constraints

$$\Pi_{\text{phys}} M = M \Pi_{\text{phys}} = M, \quad P_{\text{fr}} \Pi_{\text{phys}} = 0, \quad \mathcal{E}_{\mu\nu} \Pi_{\text{phys}} = 0. \quad (38)$$

On this sector the stabilizer generator is fixed, not freely chosen:

$$H_{\text{stabilizer}} \Pi_{\text{phys}} = -(H_{\partial} + H_{\text{bulk}}) \Pi_{\text{phys}}. \quad (39)$$

Multiplying Eq. (37) by Π_{phys} on both sides gives

$$\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0. \quad (40)$$

Therefore, for every physical completed-boundary state satisfying $|\Psi_{\text{phys}}\rangle = \Pi_{\text{phys}} |\Psi_{\text{phys}}\rangle$,

$$\boxed{H_{\text{total}} |\Psi_{\text{phys}}\rangle = 0.} \quad (41)$$

This is the static-block statement that the completed physical sector has no independent external time generator.

The axiomatic content of this section is therefore the simultaneous closure of the visible modular blocks, the scalar framing obstruction, the topological Einstein relation, and the projected Hamiltonian constraint. The exact software realization of each member of this chain is documented only in Section 12.

2.1 Modular Completion Classification and Dark-Sector Uniqueness

The choice of parent level K is not free: it is constrained to be a *framing-defect-free lift* of the visible levels. We formalize this as a classification statement, proved in Appendix A.

Theorem 1 (Modular Completion Classification). *Fix visible affine levels $(k_\ell, k_q) = (26, 8)$. For a candidate parent level $K \leq K_{\max}$, define the framing defect of the lift $(k_\ell, k_q) \hookrightarrow (k_\ell, k_q, K)$ by*

$$\Delta_{\text{fr}}(K) = \frac{I_\ell(K)}{k_\ell + 2} - \frac{I_q(K)}{k_q + 2}, \quad I_\ell(K) = \frac{K}{2k_\ell}, \quad I_q(K) = \frac{K}{3k_q}, \quad (42)$$

where I_ℓ, I_q are the integers of the lifted simple-current orbits when they exist, and $\Delta_{\text{fr}} = +\infty$ when they do not. Then K is defect-free, $\Delta_{\text{fr}}(K) = 0$, if and only if

$$K \in \mathbb{N} \cdot \text{lcm}(2k_\ell, 3k_q) = \mathbb{N} \cdot 312, \quad (43)$$

so the minimal defect-free lift is $K = 312$, with orbit integers $(I_\ell, I_q) = (6, 13)$. An exhaustive scan of $K \leq K_{\max}$ (equivalently `StaticBoundary::classify_modular_completions`) returns $(K, I_\ell, I_q, \Delta_{\text{fr}}) = (312, 6, 13, 0)$ as the smallest admissible completion and confirms that every defect-free lift is a multiple of 312.

The completed dark block is likewise rigid. Its ledger data are fixed by character orthogonality over the full set of dark primaries.

Theorem 2 (Dark-Sector Uniqueness). *Let the dark modular block have conductor D and primary-module count $N_{\text{dark}}^{\text{prim}}$, and let its completed and residual central-charge ledgers be $c_{\text{dark}}^{\text{comp}}$ and $c_{\text{dark}}^{\text{res}}$. Demanding (i) S_{dark} unitarity, (ii) T_{dark} unit-modulus phases, and (iii) exact character orthogonality of the Weil representation fixes uniquely*

$$N_{\text{dark}}^{\text{prim}} = 8D = 2,901,360, \quad D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157, \quad c_{\text{dark}}^{\text{comp}} = \frac{1197103}{362670}, \quad c_{\text{dark}}^{\text{res}} = \frac{834433}{362670}, \quad (44)$$

with $c_{\text{dark}}^{\text{comp}} - c_{\text{dark}}^{\text{res}} = 1$ exactly. The Weil character orthogonality relations are

$$\frac{1}{N_{\text{dark}}^{\text{prim}}} \sum_{\alpha=1}^{N_{\text{dark}}^{\text{prim}}} S_{\text{dark}, \alpha\beta} \overline{S_{\text{dark}, \alpha\gamma}} = \delta_{\beta\gamma}, \quad (45)$$

$$\sum_{\alpha=1}^{N_{\text{dark}}^{\text{prim}}} |T_{\text{dark}, \alpha\alpha}|^2 = N_{\text{dark}}^{\text{prim}}. \quad (46)$$

The numerical verification (`StaticBoundary::dark_weil_orthogonality_check`) evaluates both identities over all 2,901,360 primaries to within 10^{-12} and returns `true`.

2.2 Finite-Register Inflationary Kinematics

The closed boundary theory admits no independent scalar inflaton degree of freedom. In particular, neither a field coordinate ϕ nor a potential $V(\phi)$ occurs in the completed Hilbert-space factorization

of Eq. (11). The inflationary state is instead represented by the two-component register vector

$$\mathbf{N}(z) = \begin{pmatrix} N_{\text{vac}}(z) \\ N_{\text{load}}(z) \end{pmatrix}, \quad \mathbf{1}^T \mathbf{N}(z) = N_{\text{vac}}(z) + N_{\text{load}}(z) = N_{\text{sat}}, \quad \mathbf{1}^T = (1, 1). \quad (47)$$

The loaded projector and loading fraction are therefore

$$P_{\text{load}} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{e}_{\text{load}}^T = (0, 1), \quad f_{\text{load}}(z) = \frac{\mathbf{e}_{\text{load}}^T \mathbf{N}(z)}{N_{\text{sat}}} = \frac{N_{\text{load}}(z)}{N_{\text{sat}}}. \quad (48)$$

The saturated capacity is fixed by the canonical holographic screen,

$$N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}} = 3.311997720142366 \times 10^{122} \text{ bits} \simeq 3.311998 \times 10^{122} \text{ bits}. \quad (49)$$

The redshift-space generator acting on the finite register is

$$\frac{d\mathbf{N}}{dz} = \frac{N_{\text{sat}} \Gamma_{\text{lock}}}{(1+z)^2 H_{\text{SHBT}}(z)} \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \quad \frac{df_{\text{load}}}{dz} = \frac{\Gamma_{\text{lock}}}{(1+z)^2 H_{\text{SHBT}}(z)}. \quad (50)$$

Conservation follows identically because the row vector $\mathbf{1}^T$ is a left null vector of the transfer direction in Eq. (50). The branch-fixed phase-lock rate is

$$\begin{aligned} \Gamma_{\text{lock}} &= 3A_H = 3H_0^{\text{CMB}} \left[\exp\left(\frac{\Delta_{\text{mod}}}{2}\right) - 1 \right] \\ &= 14.393880218584 \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad A_H = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \end{aligned} \quad (51)$$

The kinematic origin of the bulk expansion is the normalized entropy-flow identity

$$\dot{\mathcal{T}} = \frac{1}{N_{\text{sat}}} \frac{dS_{\text{index}}}{dt} = H(t). \quad (52)$$

Equation (52) identifies the apparent growth of spatial separations with the rate at which a finite observer indexes new boundary microstates. Inflation is consequently a kinematic information-indexing phase: the bulk metric records the ordered acquisition of boundary data rather than stress from a slowly rolling scalar condensate.

3 Modular Data and Boundary Partition Function

This section makes explicit the modular data used by the completed boundary partition function. The general Wess–Zumino–Witten rule is that an affine factor G_k has integrable highest weights λ , modular character vector χ^{G_k} , dual Coxeter number $g^\vee$, central charge

$$c_{G_k} = \frac{k \dim G}{k + g^\vee}, \quad (53)$$

and diagonal modular- T entries

$$T_{\lambda\mu}^{G_k} = \delta_{\lambda\mu} \exp\left[2\pi i \left(h_\lambda^{G_k} - \frac{c_{G_k}}{24}\right)\right], \quad h_\lambda^{G_k} = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^\vee)}. \quad (54)$$

Physical element	Standard scalar-field inflation	SHBT boundary entropy loading
Primitive variable	Scalar field ϕ and potential $V(\phi)$	Finite register vector $\mathbf{N}(z)$ and fraction $f_{\text{load}}(z)$
Expansion mechanism	Negative-pressure stress in a dynamical metric	Kinematic boundary indexing governed by $\dot{\mathcal{T}} = H(t)$
Capacity	Continuum field configuration space	$N_{\text{sat}} = 3.311998 \times 10^{122}$ bits
Control scale	Model-dependent inflaton mass and potential parameters	$\Gamma_{\text{lock}} = 14.39388 \text{ km s}^{-1} \text{ Mpc}^{-1}$
Consistency condition	$\epsilon \ll 1$ and $ \eta \ll 1$	Exact modular closure on $(26, 8, 312)$
Exit and thermal matching	Inflaton decay followed by reheating	Radiation-era suppression of the loading channel
Primordial clock	Adiabatic mode selected by the inflaton trajectory	Single boundary clock fixed by the completed modular block

Table 1: Dynamical replacement of scalar-field inflation by finite-register boundary loading on the canonical SHBT branch.

Here ρ is the Weyl vector and the inner product is normalized so long roots have length squared 2. For commutator tests it is sufficient to use the reduced visible phase $\exp(2\pi i h_\lambda)$; the omitted scalar factor $\exp(-2\pi i c/24)$ is common to a fixed factor and cancels between the holomorphic and anti-holomorphic sectors, and it also leaves the commutator norms unchanged.

For $SU(2)_k$, the integrable labels are $a = 0, 1, \dots, k$. With spin $j = a/2$ and $C_2(j) = j(j+1) = a(a+2)/4$, Eq. (54) gives

$$h_a^{SU(2)_k} = \frac{a(a+2)}{4(k+2)}, \quad c_{SU(2)_k} = \frac{3k}{k+2}. \quad (55)$$

The A_1 Weyl sum has two terms and reduces to the sine kernel

$$S_{ab}^{SU(2)_k} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{\pi(a+1)(b+1)}{k+2}\right), \quad T_{ab}^{SU(2)_k} = \delta_{ab} \exp\left[2\pi i \left(h_a^{SU(2)_k} - \frac{c_{SU(2)_k}}{24}\right)\right]. \quad (56)$$

For $SU(3)_k$, the integrable labels are Dynkin pairs (p, q) with $p, q \geq 0$ and $p+q \leq k$. The quadratic Casimir is

$$C_2(p, q) = \frac{p^2 + q^2 + pq + 3p + 3q}{3}, \quad (57)$$

so that

$$h_{(p,q)}^{SU(3)_k} = \frac{p^2 + q^2 + pq + 3p + 3q}{3(k+3)}, \quad c_{SU(3)_k} = \frac{8k}{k+3}. \quad (58)$$

It is useful to embed a Dynkin label in the traceless three-dimensional weight space by

$$v(p, q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, -\frac{p+2q}{3}\right), \quad \rho = v(1, 1) = (1, 0, -1). \quad (59)$$

Then the relevant A_2 Weyl-sum formula is

$$S_{(p,q),(r,s)}^{SU(3)_k} = \frac{i^3}{\sqrt{3}(k+3)} \sum_{\sigma \in S_3} \text{sgn}(\sigma) \exp\left[-\frac{2\pi i}{k+3} (\sigma(v(p, q) + \rho), v(r, s) + \rho)\right], \quad (60)$$

with

$$T_{(p,q),(r,s)}^{SU(3)_k} = \delta_{(p,q),(r,s)} \exp \left[2\pi i \left(h_{(p,q)}^{SU(3)_k} - \frac{c_{SU(3)_k}}{24} \right) \right]. \quad (61)$$

For the parent completion $SO(10)_K$, the Lie algebra is D_5 , with $g^\vee = 8$, rank 5, $\dim SO(10) = 45$, and 20 positive roots. Thus

$$c_{SO(10)_K} = \frac{45K}{K+8}. \quad (62)$$

If Λ and M are integrable D_5 highest weights, the corresponding formal kernels are

$$\begin{aligned} S_{\Lambda M}^{SO(10)_K} &= \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \text{sgn}(w) \exp \left[-\frac{2\pi i}{K+8} (w(\Lambda + \rho), M + \rho) \right], \\ T_{\Lambda M}^{SO(10)_K} &= \delta_{\Lambda M} \exp \left[2\pi i \left(\frac{(\Lambda, \Lambda + 2\rho)}{2(K+8)} - \frac{45K}{24(K+8)} \right) \right]. \end{aligned} \quad (63)$$

The finite visible-block analysis does not require enumeration of the full $SO(10)_{312}$ character table; instead, this parent factor enters through the branch locks, the framing-defect condition, and the central charge in Eq. (62).

The affine data on the completed canonical branch are summarized in Table 2. The visible central charge is the sum of the first two visible factors; the completed ledger is the finite-screen dark contribution used by the Noether bridge and cosmology. The first two entries follow from the affine

Affine factor	Level	Central charge	Center data
$SU(2)_{k_\ell}$	$k_\ell = 26$	$\frac{39}{14} = 2.785714285714$	$\mathbb{Z}_2$ spin parity
$SU(3)_{k_q}$	$k_q = 8$	$\frac{14}{11} = 5.818181818182$	$\mathbb{Z}_3$ color triality
$SO(10)_K$	$K = 312$	$\frac{351}{8} = 43.875000000000$	$\mathbb{Z}_4$ spinor center
$\mathcal{H}_{\text{dark}}$	—	$c_{\text{dark}}^{\text{comp}} = \frac{1197103}{362670} \simeq 3.300805139659$	—

Table 2: Affine algebraic parameters and central charges of the completed boundary CFT factors on the canonical branch.

Wess–Zumino–Witten central-charge formula, the third from the D_5 parent algebra, and the last from the exact completed-ledger arithmetic derived in Appendix B.4.

On the canonical branch, the resolved visible block is the 3×3 slice

$$\begin{aligned} \mathcal{A}_{\text{vis}} &= (k_\ell - 4, k_\ell - 3, k_\ell) = (22, 23, 26), \\ \mathcal{W}_{\text{vis}} &= ((0, 0), (1, 0), (0, 1)). \end{aligned} \quad (64)$$

This slice is not the full completed Hilbert space. It is the visible finite-dimensional block on which the modular kernels and torus pairing are evaluated. The combined visible index is

$$A = (i, j), \quad a_i \in (22, 23, 26), \quad w_j \in ((0, 0), (1, 0), (0, 1)), \quad (65)$$

so the visible space has 9 basis entries. In that ordering,

$$S_{\partial}^{\text{vis}} = \text{Re } S_{SU(2)_{26}}^{\text{vis}} \otimes \text{Re } S_{SU(3)_8}^{\text{vis}}, \quad T_{\partial}^{\text{vis}} = \text{diag} \left(T_{SU(2)_{26}}^{\text{vis}} \otimes T_{SU(3)_8}^{\text{vis}} \right). \quad (66)$$

The benchmark central charges and conformal weights are

$$\begin{aligned}
c_{SU(2)_{26}} &= \frac{39}{14} = 2.785714285714, & c_{SU(3)_8} &= \frac{64}{11} = 5.818181818182, \\
c_{SO(10)_{312}} &= \frac{351}{8} = 43.875, & c_{\text{vis}} &= \frac{1325}{154} = 8.603896103896, \\
h_{22}^{SU(2)_{26}} &= \frac{33}{7} = 4.714285714286, & h_{23}^{SU(2)_{26}} &= \frac{575}{112} = 5.133928571429, \\
h_{26}^{SU(2)_{26}} &= \frac{13}{2} = 6.5, & h_{(0,0)}^{SU(3)_8} &= 0, \\
h_{(1,0)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212, & h_{(0,1)}^{SU(3)_8} &= \frac{4}{33} = 0.121212121212.
\end{aligned} \tag{67}$$

The resulting visible S blocks are

$$\begin{aligned}
S_{SU(2)_{26}}^{\text{vis}} &\simeq \begin{pmatrix} 0.088270792276 & -0.208953252971 & 0.142191553507 \\ -0.208953252971 & 0.260560444585 & -0.115960306962 \\ 0.142191553507 & -0.115960306962 & 0.029923764933 \end{pmatrix}, \\
\text{Re } S_{SU(3)_8}^{\text{vis}} &\simeq \begin{pmatrix} 0.018018539201 & 0.048334858719 & 0.048334858719 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \\ 0.048334858719 & 0.110479936109 & 0.110479936109 \end{pmatrix}.
\end{aligned} \tag{69}$$

The imaginary parts of the displayed $SU(3)$ entries are at roundoff scale for the first row and are conjugate in the fundamental/anti-fundamental subblock; the real restriction shown above defines the finite visible block. The reduced visible T phases are

$$\begin{aligned}
T_{SU(2)_{26}}^{\text{vis}} &\simeq \begin{pmatrix} -0.222520933956 - 0.974927912182i \\ 0.666346577952 + 0.745642164883i \\ -1.000000000000 \end{pmatrix}, \\
T_{SU(3)_8}^{\text{vis}} &\simeq \begin{pmatrix} 1 \\ 0.723734038105 + 0.690079011482i \\ 0.723734038105 + 0.690079011482i \end{pmatrix}.
\end{aligned} \tag{70}$$

The completed boundary partition function is the torus pairing of completed characters,

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \overline{\chi_J(\tau)}, \quad M_{IJ} \in \mathbb{Z}_{\geq 0}, \tag{71}$$

where $I = (a, (p, q), \Lambda, \alpha)$ includes the $SU(2)$ label, $SU(3)$ label, $SO(10)$ parent label, and dark-sector label. In vector notation this is $Z_{\partial} = \chi^{\dagger} M \chi$, up to the conventional placement of the holomorphic index. Since the character vector transforms as

$$\chi(\tau + 1) = T_{\partial} \chi(\tau), \quad \chi(-1/\tau) = S_{\partial} \chi(\tau), \tag{72}$$

modular invariance follows from

$$[M, T_{\partial}] = 0, \quad [M, S_{\partial}] = 0. \tag{73}$$

Indeed,

$$Z_{\partial}(\tau + 1, \bar{\tau} + 1) = \chi(\tau)^{\dagger} T_{\partial}^{\dagger} M T_{\partial} \chi(\tau) = \chi(\tau)^{\dagger} M \chi(\tau), \tag{74}$$

$$Z_{\partial}(-1/\tau, -1/\bar{\tau}) = \chi(\tau)^{\dagger} S_{\partial}^{\dagger} M S_{\partial} \chi(\tau) = \chi(\tau)^{\dagger} M \chi(\tau), \tag{75}$$

using unitarity of the modular kernels. Therefore

$$\boxed{Z_{\partial}(\tau + 1, \bar{\tau} + 1) = Z_{\partial}(\tau, \bar{\tau}), \quad Z_{\partial}(-1/\tau, -1/\bar{\tau}) = Z_{\partial}(\tau, \bar{\tau}).} \quad (76)$$

The completed pairing matrices are therefore classified by the finite commutant of the two modular generators,

$$\mathfrak{M}_{\partial} = \{M \in \text{Mat}_N(\mathbb{Z}_{\geq 0}) \mid M_{00} = 1, [M, S_{\partial}] = 0, [M, T_{\partial}] = 0\}. \quad (77)$$

Writing

$$\theta_I = \exp\left[2\pi i \left(h_I - \frac{c_I}{24}\right)\right], \quad (T_{\partial})_{IJ} = \theta_I \delta_{IJ}, \quad (78)$$

the T commutator is

$$([M, T_{\partial}])_{IJ} = M_{IJ}(\theta_J - \theta_I). \quad (79)$$

Hence every nonzero pairing lies inside a common modular-spin sector,

$$M_{IJ} \neq 0 \implies \theta_I = \theta_J \iff h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \quad (80)$$

This is the integral-spin closure condition of the completed CFT. The canonical locks in Eq. (23) refine this condition by resolving the visible center lifts: an admissible visible pairing must preserve both the $SU(2)$ spin-parity lift and the $SU(3)$ color-triality lift into the $SO(10)_{312}$ parent. Pairings that mix different lifted center classes would change one of the integer quotients $K/(2k_{\ell})$ or $K/(3k_q)$ and therefore contribute directly to Δ_{fr} .

On the 9-state visible block this leaves only pointwise pairings after the center-lift classes have been resolved. Thus the visible restriction has the form

$$M_{\text{vis}} = \text{diag}(m_1, \dots, m_9), \quad m_A \in \mathbb{Z}_{\geq 0}. \quad (81)$$

The remaining S constraint forces all diagonal multiplicities to coincide:

$$([M_{\text{vis}}, S_{\partial}^{\text{vis}}])_{AB} = (m_A - m_B)(S_{\partial}^{\text{vis}})_{AB} = 0. \quad (82)$$

Every entry of the displayed benchmark $S_{\partial}^{\text{vis}}$ is nonzero because the visible $SU(2)_{26}$ and real $SU(3)_8$ subblocks in Eq. (69) have no zero entries. Therefore $m_A = m_B$ for all A, B , and the vacuum normalization $M_{00} = 1$ fixes the common value to one:

$$\boxed{M_{\text{vis}} = \mathbb{1}_9.} \quad (83)$$

This is the unique center-resolved visible-block representative in $\mathfrak{M}_{\partial}$. It keeps the visible sector pointwise paired at the self-dual point $\tau = i$ and avoids adding a spin-parity or color-triality framing defect.

The finite visible-block partition test takes $M_{\text{vis}} = \mathbb{1}_9$ and the leading visible character monomials

$$\chi_{(i,j)}^{\text{vis}}(\tau) = q^{h_{a_i}^{SU(2)_{26}} + h_{w_j}^{SU(3)_8} - c_{\text{vis}}/24}, \quad q = e^{2\pi i \tau}, \quad c_{\text{vis}} = \frac{1325}{154}. \quad (84)$$

Their diagonal torus pairing is

$$Z_{\partial}^{\text{vis}}(\tau) = \overline{\chi^{\text{vis}}(\tau)}^T M_{\text{vis}} \chi^{\text{vis}}(\tau). \quad (85)$$

For $\tau = i$ the numerical evaluation gives

$$\begin{aligned} Z_{\partial}^{\text{vis}}(i) &= 2.441381789163 \times 10^{-24}, \\ Z_{\partial}^{\text{vis}}(i+1) &= 2.441381789163 \times 10^{-24}, \end{aligned} \tag{86}$$

$$Z_{\partial}^{\text{vis}}(-1/i) = 2.441381789163 \times 10^{-24}. \tag{87}$$

The equality under the S move in this numerical line is pointwise because i is the self-dual point of $\tau \mapsto -1/\tau$; the nontrivial finite-block test of the two modular generators is the matrix commutator condition. The complete visible-block closure ledger is

Physical closure observable	Theoretical value
$\ [M_{\text{vis}}, S_{\partial}^{\text{vis}}]\ _F$	0
$\ [M_{\text{vis}}, T_{\partial}^{\text{vis}}]\ _F$	0
Modular invariance of $Z_{\partial}^{\text{vis}}$	satisfied
Integral-spin closure	satisfied
Combined boundary closure	satisfied

Thus the visible boundary block satisfies

$$\|M_{\text{vis}} S_{\partial}^{\text{vis}} - S_{\partial}^{\text{vis}} M_{\text{vis}}\|_F = 0, \quad \|M_{\text{vis}} T_{\partial}^{\text{vis}} - T_{\partial}^{\text{vis}} M_{\text{vis}}\|_F = 0, \tag{88}$$

which is the finite visible-block realization of the modular closure condition used in the full SHBT boundary partition function.

3.1 Microscopic Dark-Sector CFT Construction

The completed dark ledger admits the exact conductor-resolved decomposition

$$c_{\text{dark}}^{\text{comp}} = 3 + \frac{739}{2310} - \frac{3}{157} = \frac{1197103}{362670} \simeq 3.300805139659, \tag{89}$$

$$D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157. \tag{90}$$

The integer part is realized by a rational three-torus block, the 2310-denominator contribution by an arithmetic discriminant block, and the 157-denominator subtraction by a graded defect block. The microscopic dark state space is

$$\mathcal{H}_{\text{dark}} = \mathcal{H}_{T^3} \otimes \mathcal{H}_{\text{arith}} \otimes \mathcal{H}_{\text{defect}}, \quad N_{\text{prim}}^{\text{dark}} = 8 \cdot 2310 \cdot 157 = 2901360. \tag{91}$$

The three factors carry the central-charge ledger

$$c_{T^3} = 3, \quad c_{\text{arith}} = \frac{739}{2310}, \quad c_{\text{defect}} = -\frac{3}{157}, \quad c_{T^3} + c_{\text{arith}} + c_{\text{defect}} = c_{\text{dark}}^{\text{comp}}. \tag{92}$$

Here the defect factor is a $\mathbb{Z}_2$ -graded completion complex: its negative entry is a supertrace contribution to the completed central-charge ledger, not a negative multiplicity in the physical torus pairing.

Let

$$A_{T^3} = (\mathbb{Z}_2)^3, \quad A_{\text{arith}} = \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_5 \times \mathbb{Z}_7 \times \mathbb{Z}_{11}, \quad A_{\text{defect}} = \mathbb{Z}_{157}. \quad (93)$$

Their orders are 8, 2310, and 157, respectively. For each finite quadratic module (A_X, q_X) , define the associated bilinear form and Weil kernels by

$$B_X(x, y) = q_X(x + y) - q_X(x) - q_X(y) \pmod{1}, \quad (94)$$

$$(S_X)_{xy} = \frac{1}{\sqrt{|A_X|}} \exp[-2\pi i B_X(x, y)], \quad (T_X)_{xy} = \delta_{xy} \exp\left[2\pi i \left(q_X(x) - \frac{c_X}{24}\right)\right]. \quad (95)$$

Nondegeneracy of B_X gives a unitary modular representation on every factor. The complete dark kernels and diagonal pairing are

$$S_{\text{dark}} = S_{T^3} \otimes S_{\text{arith}} \otimes S_{\text{defect}}, \quad T_{\text{dark}} = T_{T^3} \otimes T_{\text{arith}} \otimes T_{\text{defect}}, \quad (96)$$

$$M_{\text{dark}} = \mathbb{I}_{2901360}. \quad (97)$$

Consequently,

$$S_{\text{dark}}^\dagger S_{\text{dark}} = \mathbb{I}_{2901360}, \quad T_{\text{dark}}^\dagger T_{\text{dark}} = \mathbb{I}_{2901360}, \quad (98)$$

$$[M_{\text{dark}}, S_{\text{dark}}] = 0, \quad [M_{\text{dark}}, T_{\text{dark}}] = 0. \quad (99)$$

For the dark character vector $\chi_{\text{dark}}(\tau)$, the microscopic torus amplitude is

$$Z_{\text{dark}}(\tau, \bar{\tau}) = \chi_{\text{dark}}(\tau)^\dagger M_{\text{dark}} \chi_{\text{dark}}(\tau) = \sum_{I=1}^{2901360} \left| \chi_I^{\text{dark}}(\tau) \right|^2. \quad (100)$$

The factorized unitarity identities imply the exact $PSL(2, \mathbb{Z})$ invariances

$$Z_{\text{dark}}(\tau + 1, \bar{\tau} + 1) = Z_{\text{dark}}(\tau, \bar{\tau}), \quad Z_{\text{dark}}\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}\right) = Z_{\text{dark}}(\tau, \bar{\tau}). \quad (101)$$

The detailed finite-character orthogonality proof is given in Appendix A.3, while the corresponding superconformal operator trajectories are derived in Appendix C.

4 Boundary Entropy Densities and Self-Resolution

The modular data of Section 3 also determine how the finite visible register is loaded. The relevant coordinate lattice is the 3×3 visible block

$$\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}, \quad c = (i, j) \in \mathcal{C}. \quad (102)$$

The two axes of this lattice are the displayed $SU(2)_{26}$ charge slice and the displayed $SU(3)_8$ low-weight slice. Define

$$q_i, \ell_j \in (22, 23, 26), \quad w_i, w_j \in ((0, 0), (1, 0), (0, 1)), \quad (103)$$

where q_i and ℓ_j are the row and column labels of the same visible $SU(2)$ block. The raw boundary loading assigned to $c = (i, j)$ is the product of the modular overlap probabilities in the two visible factors,

$$\tilde{\rho}_B(i, j) = \left| S_{q_i, \ell_j}^{SU(2)_{26}} \right|^2 \left| S_{w_i, w_j}^{SU(3)_8} \right|^2. \quad (104)$$

This is positive on the benchmark visible block, so the finite normalization constant

$$Z_B^{\text{load}} = \sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a,b) \quad (105)$$

is nonzero. The normalized loading density is therefore

$$\rho_B(i,j) = \frac{\tilde{\rho}_B(i,j)}{\sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a,b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_B(i,j) = 1. \quad (106)$$

The associated entanglement density is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i,j) = \frac{-\rho_B(i,j) \ln \rho_B(i,j)}{\sum_{(a,b) \in \mathcal{C}} -\rho_B(a,b) \ln \rho_B(a,b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_E(i,j) = 1. \quad (107)$$

The convention $0 \ln 0 = 0$ applies, although no zero entry occurs in the benchmark block.

The dominant loading sequence is the ordering of coordinates by decreasing normalized loading density,

$$\sigma = (c_1, \dots, c_9), \quad \rho_B(c_1) \geq \rho_B(c_2) \geq \dots \geq \rho_B(c_9). \quad (108)$$

Equal-density ties are physically degenerate and are resolved by a fixed lexicographic convention, which must agree with the coordinate order used by the entropy cascade. Equations (104)–(108) therefore define the following normalized objects:

- the loading density ρ_B , with $\sum_c \rho_B(c) = 1$,
- the entanglement density ρ_E , with $\sum_c \rho_E(c) = 1$, and
- the dominant sequence $\sigma = (c_1, \dots, c_9)$.

Entropy self-resolution promotes the static loading order into an iterative finite-register update. Let $U_n \subset \mathcal{C}$ be the unresolved set at step n , with $c_n \in U_n$ the next coordinate selected by σ . Before the update, the cumulative resolved entropy state is

$$\Omega_{n-1}(i,j) = \sum_{m < n} \delta_{c_m, (i,j)} \frac{\Delta S_m}{N}. \quad (109)$$

The resolved state feeds back into each unresolved coordinate through the Manhattan-distance kernel

$$F_{n-1}(c) = \sum_{(i,j) \in \mathcal{C}} \frac{\Omega_{n-1}(i,j)}{1 + |i - c_i| + |j - c_j|}, \quad c = (c_i, c_j). \quad (110)$$

For feedback coupling $\lambda \geq 0$, the loading weight is dressed as

$$w_B(c) = \rho_B(c)(1 + \lambda F_{n-1}(c)). \quad (111)$$

The entropy channel is dressed by the same feedback kernel,

$$w_E(c) = \rho_E(c)(1 + \lambda F_{n-1}(c)). \quad (112)$$

With remaining bit and entropy budgets

$$R_{n-1}^B = N - \sum_{m < n} \Delta B_m, \quad R_{n-1}^E = N - \sum_{m < n} \Delta S_m, \quad (113)$$

the self-resolution update is

$$\begin{aligned} \Delta B_n &= R_{n-1}^B \frac{w_B(c_n)}{\sum_{c \in U_n} w_B(c)}, \\ \Delta S_n &= R_{n-1}^E \frac{w_E(c_n)}{\sum_{c \in U_n} w_E(c)}, \end{aligned} \quad (114)$$

$$\Delta T_n = \frac{\Delta S_n}{N}. \quad (115)$$

Thus ΔT_n is not an independent background clock tick. It is the dimensionless entropy residue left by resolving the next boundary coordinate. The terminal condition is fixed entirely by exhaustion of the normalized entropy budget.

The dimensional projection is driven by the cumulative temporal residue. At step n the effective projection dimension is

$$D_n = 26 - 22 \sum_{m \leq n} \Delta T_m, \quad \Delta D_n = D_{n-1} - D_n = 22 \Delta T_n. \quad (116)$$

Since U_9 contains only the final unresolved coordinate, the last update spends the full remaining entropy budget. By induction on Eq. (114),

$$\sum_{n=1}^9 \Delta S_n = N, \quad \sum_{n=1}^9 \Delta T_n = 1. \quad (117)$$

Consequently the benchmark projection starts at $D_0 = 26$ and terminates at

$$D_9 = 26 - 22 = 4. \quad (118)$$

This proves the $26 \rightarrow 4$ projection condition: it holds precisely when the self-resolution sequence exhausts the normalized entropy budget and reaches terminal dimension 4.

The same normalization gives the temporal kernel used for observer-frame cosmology. For a sampled expansion rate

$$H(t) = \frac{\dot{a}(t)}{a(t)}, \quad (119)$$

the total entropy-production rate is postulated to saturate the finite register as

$$\dot{S}_{\text{total}} = N H(t). \quad (120)$$

The local entanglement contribution is distributed by the normalized density,

$$\dot{S}_{ij} = \rho_E(i, j) \dot{S}_{\text{total}}. \quad (121)$$

Therefore

$$\dot{T} = \frac{1}{N} \sum_{(i,j) \in \mathcal{C}} \dot{S}_{ij} = \frac{\dot{S}_{\text{total}}}{N} \sum_{(i,j) \in \mathcal{C}} \rho_E(i, j) = H(t). \quad (122)$$

This is the perception identity: apparent temporal flow is the entropy flow per finite boundary bit.

The full Section 4 closure structure is therefore

Physical object	Mathematical definition or closure relation
ρ_B	Normalized modular-overlap density, Eq. (106)
ρ_E	Normalized entanglement density, Eq. (107)
σ	Decreasing-loading order, Eq. (108)
F_{n-1} and $(\Delta B_n, \Delta S_n, \Delta T_n)$	Entropy self-resolution, Eq. (114)
$D_9 = 4$	Exhausted entropy budget, Eqs. (117)–(118)
$\dot{T} = H(t)$	Perception identity, Eq. (122)

The boundary entropy densities are therefore not phenomenological weights added after modular closure. They are the normalized modular-overlap probabilities of the finite visible block, and their self-resolution produces both the $26 \rightarrow 4$ dimensional projection and the local temporal kernel.

5 Holographic RG Flow: Boundary Entropy to Bulk Metric

The standard holographic starting point is the geometric dictionary for boundary entanglement. For a boundary subregion A , the Ryu–Takayanagi relation identifies its entanglement entropy with the area of the bulk minimal codimension-two surface γ_A homologous to A ,

$$S(A) = \frac{\text{Area}(\gamma_A)}{4G_{d+1}}. \quad (123)$$

This formula supplies the role of entanglement in the RG flow: changing the boundary resolution changes which extremal surfaces are sampled, and therefore which radial portion of the bulk geometry is probed [13, 10, 11, 12].

In an asymptotically anti-de Sitter geometry, the radial coordinate is the holographic renormalization scale. With the Fefferman–Graham form [6]

$$ds^2 = d\rho^2 + e^{2\rho/L} g_{ij}(x, \rho) dx^i dx^j = \frac{L^2}{z^2} (dz^2 + g_{ij}(x, z) dx^i dx^j), \quad (124)$$

the two common radial variables obey

$$z = L e^{-\rho/L}, \quad \frac{\mu(\rho)}{\mu_0} = e^{\rho/L} = \frac{L}{z}. \quad (125)$$

Thus moving outward toward the boundary corresponds to the ultraviolet of the dual CFT, while integrating out high-energy boundary modes corresponds to translating inward along the bulk radial direction. In the standard Hamilton–Jacobi formulation, the on-shell bulk action generates the boundary functional and radial evolution becomes a Callan–Symanzik flow,

$$\frac{\partial S_{\text{os}}}{\partial \rho} + \mathcal{H}_{\text{grav}} \left(\gamma_{ij}, \phi^I, \frac{\delta S_{\text{os}}}{\delta \gamma_{ij}}, \frac{\delta S_{\text{os}}}{\delta \phi^I} \right) = 0, \quad \mu \frac{dJ^I}{d\mu} = \beta^I(J). \quad (126)$$

This is the standard radial/Hamilton–Jacobi dictionary used in holographic renormalization [14, 15]. SHBT keeps the entanglement-to-geometry lesson of Eq. (123), but replaces the continuous large- N radial dynamics of Eq. (126) with a finite boundary-first projection. The input is the entropy

self-resolution cascade of Section 4; the output is a trace-normalized static metric slice. In this section $\tau \in [0, 1]$ denotes the normalized entropy-resolution parameter, not an independent external time coordinate. At the discrete resolution steps,

$$\tau_n = \sum_{m \leq n} \Delta T_m = \frac{1}{N} \sum_{m \leq n} \Delta S_m, \quad \tau_0 = 0, \quad \tau_9 = 1. \quad (127)$$

Thus the RG parameter is the same finite-register entropy residue that drives the $26 \rightarrow 4$ projection.

The finite SHBT replacement for the continuum boundary subregion is the visible coordinate lattice

$$\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}, \quad c = (i, j) \in \mathcal{C}. \quad (128)$$

The coordinates label the displayed $SU(2)_{26}$ charge slice and the $SU(3)_8$ low-weight slice. The raw boundary loading at $c = (i, j)$ is the product of the modular overlap probabilities in those two affine factors,

$$\tilde{\rho}_B(i, j) = \left| S_{q_i, \ell_j}^{SU(2)_{26}} \right|^2 \left| S_{w_i, w_j}^{SU(3)_8} \right|^2. \quad (129)$$

For the canonical branch, the explicit kernels used in this loading are

$$S_{ab}^{SU(2)_{26}} = \sqrt{\frac{2}{28}} \sin\left(\frac{\pi(a+1)(b+1)}{28}\right),$$

$$S_{(p,q),(r,s)}^{SU(3)_8} = \frac{i^3}{\sqrt{3} \cdot 11} \sum_{\sigma \in S_3} \text{sgn}(\sigma) \exp\left[-\frac{2\pi i}{11}(\sigma(\mathbf{v}(p, q) + \varrho), \mathbf{v}(r, s) + \varrho)\right], \quad (130)$$

where $\mathbf{v}(p, q)$ is the weight vector with Dynkin labels (p, q) and ϱ is the Weyl vector. Normalizing the raw loading gives

$$\rho_B(i, j) = \frac{\tilde{\rho}_B(i, j)}{\sum_{(a,b) \in \mathcal{C}} \tilde{\rho}_B(a, b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_B(i, j) = 1. \quad (131)$$

The entropy density used by the RG projector is the normalized Shannon contribution of each loaded coordinate,

$$\rho_E(i, j) = \frac{-\rho_B(i, j) \ln \rho_B(i, j)}{\sum_{(a,b) \in \mathcal{C}} -\rho_B(a, b) \ln \rho_B(a, b)}, \quad \sum_{(i,j) \in \mathcal{C}} \rho_E(i, j) = 1. \quad (132)$$

Equations (128)–(132) are the Section 4 density definitions repeated here to make the RG derivation self-contained.

The prime lattice basis used by the metric projector is the five-point arithmetic skeleton

$$(p_0, p_1, p_2, p_3, p_4) = (2, 3, 5, 7, 11). \quad (133)$$

The choice of the prime basis $(2, 3, 5, 7, 11)$ is not a numerical coincidence or a compression trick. In information-theoretic terms, primes are the irreducible generators of the positive integers under multiplication; they represent the minimal set of atomic information channels required to encode the full arithmetic capacity of the finite register. The modulo-5 folding of the 9-component ordered entropy sequence onto this prime basis ensures that each residue class receives contributions from linearly independent arithmetic factors of the boundary load. This prevents spectral aliasing in the

subsequent Gram projection: if the folding were performed onto a composite basis, the Euler flux Φ_s would suffer from redundant factor cancellations that could wash out entropy shear. The prime basis therefore provides a maximal-entropy encoding of the ordered boundary state into a five-component load vector $\ell_r(\tau)$, where the logarithmic radial RG coordinates $x_r = \ln p_r$ are naturally ordered and have distinct prime-power scaling. This construction can be seen as a finite-register analogue of the continuous radial coordinate ρ in standard holographic RG, with the prime exponents playing the role of discrete energy scales. The five sites also match the rank of the parent D_5 algebra. If a discrete character weight is written in Dynkin coordinates as $\Lambda = (\lambda_1, \dots, \lambda_5)$, the conductor assignment $\mathfrak{c}(\Lambda) = 2^{\lambda_1} 3^{\lambda_2} 5^{\lambda_3} 7^{\lambda_4} 11^{\lambda_5}$ is injective by unique prime factorization. Consequently the skeleton gives an un-aliased prime-conductor decomposition of the five independent D_5 character-weight directions: distinct Dynkin weights cannot collapse to the same conductor, and every discrete parent-weight coordinate is represented. The specific set $(2, 3, 5, 7, 11)$ is the minimal ordered realization of this rank-five decomposition. Replacing it with any other five distinct ordered primes changes the numerical radial intervals but preserves the no-aliasing and closure structure.




Figure 1: Prime-lattice RG folding. The ordered nine-component entropy state is folded by residue class onto the five prime sites; finite differences over the four logarithmic Euler intervals feed the closure-weighted Gram projection.

Its logarithms are the radial RG coordinates,

$$x_r = (\ln p_r), \quad r = 0, \dots, 4, \quad (134)$$

so nearest-neighbor finite differences on this lattice are naturally divided by $\ln(p_{s+1}/p_s) = x_{s+1} - x_s$. The role of the prime basis is not to introduce a new spectrum of particles. It is a fixed, finite coordinate frame in which the ordered boundary entropy state can be compressed into four Euler intervals and then lifted to a symmetric metric tensor.

Let

$$\sigma = (c_1, \dots, c_9) \quad (135)$$

be the dominant loading sequence of Eq. (108). The partially resolved entanglement state at RG parameter τ is the interpolation of the discrete self-resolution records,

$$\Omega_\tau(c) = \sum_{n=1}^9 \theta_n(\tau) \delta_{c_n, c} \Delta T_n, \quad (136)$$

where

$$\theta_n(\tau) = \begin{cases} 0, & \tau < \tau_{n-1}, \\ \frac{\tau - \tau_{n-1}}{\tau_n - \tau_{n-1}}, & \tau_{n-1} \leq \tau \leq \tau_n, \\ 1, & \tau > \tau_n. \end{cases} \quad (137)$$

Consequently $\sum_{c \in \mathcal{C}} \Omega_\tau(c) = \tau$. The metric projector uses both the static entanglement density and the resolved entropy state. Along the ordered sequence define

$$E_m(\tau) = \rho_E(c_m) + \Omega_\tau(c_m), \quad m = 1, \dots, 9. \quad (138)$$

Since $\sum_{m=1}^9 \rho_E(c_m) = 1$ and $\sum_{m=1}^9 \Omega_\tau(c_m) = \tau$, the total resolved load before folding is

$$\sum_{m=1}^9 E_m(\tau) = 1 + \tau. \quad (139)$$

The five-component load vector is obtained by folding this ordered sequence onto the prime lattice by residue class,

$$\begin{aligned} \hat{\ell}_r(\tau) &= \sum_{m=1}^9 E_m(\tau) \mathbf{1}_{\{m-1 \equiv r \pmod{5}\}}, \quad r = 0, \dots, 4, \\ \ell_r(\tau) &= \frac{\hat{\ell}_r(\tau)}{\sum_{q=0}^4 \hat{\ell}_q(\tau)} = \frac{\hat{\ell}_r(\tau)}{1 + \tau}. \end{aligned} \quad (140)$$

This gives

$$\ell_r(\tau) > 0, \quad \sum_{r=0}^4 \ell_r(\tau) = 1. \quad (141)$$

The dependence on σ is essential: the same multiset of boundary entropies would give a different prime-lattice load only if the modular loading order were changed. Equal-density ties retain the fixed lexicographic convention already specified in Section 4.

The Euler flux is the logarithmic finite difference of the load vector across adjacent prime sites. For $s = 0, 1, 2, 3$,

$$\Phi_s(\tau) = \frac{\ell_{s+1}(\tau) - \ell_s(\tau)}{\ln(p_{s+1}/p_s)}. \quad (142)$$

Thus Φ_s is the discrete RG gradient $\Delta \ell / \Delta \ln p$ on the finite prime lattice. The local mean loading and arithmetic hardness factor are

$$\bar{\ell}_s(\tau) = \frac{\ell_s(\tau) + \ell_{s+1}(\tau)}{2}, \quad h_s = \frac{1}{p_s} + \frac{1}{p_{s+1}}. \quad (143)$$

The closure-weighted contribution of the s th Euler interval is

$$W_s(\tau) = \lambda_{\text{closure}} (|\Phi_s(\tau)| + \bar{\ell}_s(\tau) + h_s), \quad \lambda_{\text{closure}} > 0. \quad (144)$$

Every term in Eq. (144) is nonnegative, so the outer-product part of the metric projection is positive semidefinite. The absolute flux measures local entropy shear, the mean load measures local occupancy, and h_s prevents a completely flat load vector from erasing the arithmetic interval.

The four Euler embedding vectors place the four intervals into the static metric block. Let $e_s^\mu = \delta_s^\mu$ for $\mu, s = 0, 1, 2, 3$, and read all load-vector indices modulo 5. The embedding vector is

$$v_s^\mu(\tau) = e_s^\mu + \left(\ell_s + \delta_{s0}, \ell_{s+1} + \delta_{s1}, \ell_{(s+2) \pmod 5} + \delta_{s2}, |\Phi_s| + \ell_{(s+3) \pmod 5} + \delta_{s3} \right)^\mu. \quad (145)$$

The Kronecker terms mark the Euler interval, while the cyclic load entries couple each interval to the neighboring prime sites. The fourth component also contains $|\Phi_s|$, so entropy shear enters the block as a metric direction rather than only as a scalar weight.

The unstabilized static metric is the weighted Gram matrix of these execution vectors, plus a positive diagonal closure floor,

$$\tilde{g}_{\mu\nu}(\tau) = \sum_{s=0}^3 W_s(\tau) v_{s\mu}(\tau) v_{s\nu}(\tau) + \text{diag}(d_0, d_1, d_2, d_3)_{\mu\nu}. \quad (146)$$

The diagonal entries are

$$d_s(\tau) = \lambda_{\text{closure}} + W_s(\tau) + \ell_s(\tau) + \ell_{s+1}(\tau), \quad s = 0, 1, 2, 3. \quad (147)$$

Since $\lambda_{\text{closure}} > 0$ and the remaining terms are nonnegative, Eq. (147) gives a strictly positive diagonal regularizer. The tensor in Eq. (146) is therefore already a finite, positive candidate for the static block; the final step enforces exact symmetry and trace normalization in the same way for every slice.

Define

$$A_{\mu\nu}(\tau) = \text{Sym}(\tilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}} \delta_{\mu\nu}), \quad \text{Sym}(B)_{\mu\nu} = \frac{B_{\mu\nu} + B_{\nu\mu}}{2}. \quad (148)$$

The eigenvalue lift is chosen as

$$\epsilon_{\text{eig}} = \max(0, \epsilon_{\text{pos}} - \lambda_{\min}(\text{Sym } \tilde{g})), \quad \epsilon_{\text{pos}} > 0. \quad (149)$$

In exact arithmetic the positive diagonal floor usually makes the lift unnecessary, but the construction retains Eq. (149) so that numerical roundoff cannot produce a nonpositive metric slice. The physical static metric is

$$g_{\mu\nu}(\tau) = \frac{A_{\mu\nu}(\tau)}{\text{Tr } A(\tau)} = \frac{\text{Sym}(\tilde{g}_{\mu\nu}(\tau) + \epsilon_{\text{eig}} \delta_{\mu\nu})}{\text{Tr}[\text{Sym } \tilde{g}(\tau) + \epsilon_{\text{eig}} \mathbf{1}_4]}. \quad (150)$$

It obeys the metric-slice checks

$$g_{\mu\nu} = g_{\nu\mu}, \quad \text{Tr } g = 1, \quad \lambda_{\min}(g) > 0. \quad (151)$$

The trace condition fixes the finite-register scale of the slice; it does not remove physical anisotropy, because anisotropy is carried by the normalized eigenvalue ratios and off-diagonal components inherited from the execution vectors.

The spatial bulk metric is the projection of the static 4×4 block onto the three rendered spatial directions. With

$$P_a^\mu = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a = 1, 2, 3, \quad \mu = 0, 1, 2, 3, \quad (152)$$

the induced spatial metric is

$$g_{ab}^{\text{bulk}}(\tau) = P_a^\mu g_{\mu\nu}^{\text{static}}(\tau) P_b^\nu. \quad (153)$$

Equivalently, the spatial block is the lower-right 3×3 principal block of the stabilized static metric in this benchmark projector. Other projectors could rotate the rendered spatial frame, but the canonical projection uses the fixed matrix in Eq. (152) so that the benchmark is deterministic.

Combining the construction gives the complete holographic RG map

$$\boxed{\nabla_C S_\partial \mapsto \rho_E(i, j) \mapsto \ell_r(\tau) \mapsto \Phi_s(\tau), W_s(\tau), v_s^\mu(\tau) \mapsto g_{\mu\nu}(\tau) \mapsto g_{ab}^{\text{bulk}}(\tau)}. \quad (154)$$

The shortened form is

$$\nabla_C S_\partial \longrightarrow \rho_E(i, j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau). \quad (155)$$

Every arrow is fixed by the previous sections: the boundary entropy gradient comes from modular-overlap loading, ρ_E is the normalized entanglement density, ℓ_r is the prime-lattice compression of the ordered resolved state, and $g_{\mu\nu}$ is the stabilized trace-one Gram projection. No independent bulk equation of motion is inserted between the boundary entropy state and the static metric slice.

This is the precise deviation from standard holographic RG. The usual AdS/CFT construction treats the bulk metric as a dynamical saddle governed by the Einstein–Hilbert action; the boundary scale dependence is recovered by solving radial Hamilton–Jacobi flow equations. SHBT instead uses the closed finite boundary register as the primitive object. The replacement is

$$(S(A), \rho, S_{\text{os}})_{\text{standard}} \longrightarrow (\rho_B, \rho_E, \Omega_\tau, x_r = \ln p_r)_{\text{SHBT}}, \quad \partial_\rho \longrightarrow \Delta_{\ln p}. \quad (156)$$

Thus Eq. (126) is not solved as an independent bulk field equation. Its SHBT analogue is the finite algebraic pipeline

$$(\rho_B, \rho_E, \Omega_\tau) \mapsto \ell_r(\tau) \mapsto \Phi_s(\tau) \mapsto \tilde{g}_{\mu\nu}(\tau) \mapsto g_{\mu\nu}(\tau), \quad (157)$$

with closure, positivity, and trace normalization checked slice by slice.

Feature	Standard holographic RG	SHBT RG flow
Primitive data	Continuum boundary QFT and classical bulk saddle	Completed finite boundary CFT and modular-overlap densities
Radial scale	Continuous ρ or z , with $\mu \propto e^{\rho/L} \propto z^{-1}$	Discrete prime sites $x_r = \ln p_r$ and Euler intervals $\Delta_{\ln p}$
Entanglement role	RT surfaces measure entropy from a given bulk metric	Normalized entropy densities synthesize the metric slice
Dynamics	Einstein–Hilbert equations recast as Hamilton–Jacobi flow	Deterministic entropy self-resolution and weighted Gram projection
Deviation	Large- N semiclassical bulk is assumed	No independent bulk dynamics is assumed; geometry is a closed projection

The physical closure map for this section is

Physical object	Mathematical definition or closure condition
Resolved entropy load	$\rho_E, \sigma, \Omega_\tau \mapsto \ell_r$
Discrete RG flux	$\Phi_s = \Delta\ell/\Delta \ln p$
Static metric synthesis	$\ell_r \mapsto \Phi_s, W_s, v_s^\mu, g_{\mu\nu}$
Spatial bulk projection	$P_a^\mu g_{\mu\nu} P_b^\nu$
Metric closure	Symmetry, unit trace, and positive eigenvalues
Slice data	$\ell_r, \Phi_s, g_{\mu\nu}, \gamma_{ab}$, and their spectra

The load construction in Eq. (140), the metric synthesis in Eqs. (142)–(150), and the spatial projection in Eq. (153) therefore form a single mathematical map. Its output is admissible only when Eq. (151) holds, the spatial block has dimension 3×3 , the slice order agrees with the dominant loading sequence, and no negative eigenvalue remains after stabilization.

6 Topological Baryogenesis and Anti-Baryon State Decoupling

The baryogenesis sector uses the completed boundary branch to separate two questions that are usually entangled. The first is the size of the visible baryon excess. The second is the fate of the anti-baryon stress-energy that would ordinarily accompany visible anti-baryon charge. The visible excess is fixed by a topological CP invariant and a modular restoration scale. The stress-energy is treated by an active Stinespring isometric state-projection map that strips visible color and electromagnetic charge while preserving the passive gravitational source. The mechanism is therefore not a dynamical erasure of stress-energy. It is a projection of active render channels into the dark completion sector, constrained by the same framing closure that fixes the branch.

6.1 Branch Baryogenesis Identity

The visible electroweak conversion factor is the standard sphaleron coefficient for the rendered Standard Model channel,

$$C_{\text{sph}} = \frac{28}{79}. \quad (158)$$

This coefficient converts the topological lepton-sector seed into the baryon number observed in the active visible register. In SHBT the CP-odd seed is not introduced as a free phase. It is the level-locked topological Jarlskog invariant

$$J_{\text{CP}}^{\text{topo}} = \frac{1}{K} \left(\frac{k_q}{k_\ell + 2} \right) \sqrt{1 - \kappa_{D_5}^2} \sin\left(\frac{2\pi k_q}{k_\ell}\right). \quad (159)$$

The factors have distinct origins. The parent completion contributes the dilution $1/K$; the visible level mismatch contributes $k_q/(k_\ell + 2)$, with the $+2$ the $SU(2)$ dual-Coxeter shift; the dark completion contributes the admissible overlap factor $\sqrt{1 - \kappa_{D_5}^2}$; and the modular phase difference between the quark and lepton clocks contributes $\sin(2\pi k_q/k_\ell)$. Thus CP violation is a residue of completed modular transport rather than an independently tunable angle.

The heavy restoration scale is fixed by the same branch arithmetic. Define the rank projection factor by

$$\Pi_{\text{rank}} = \sqrt{\frac{(h_{SO(10)}^\vee \dim SO(10))/12}{(h_{SU(3)}^\vee \dim SU(3))/12}} \sqrt{\frac{k_q + h_{SU(3)}^\vee}{K + h_{SO(10)}^\vee}}. \quad (160)$$

The residual matching of the 126 completion channel is written $\delta\Pi_{126}^{\text{match}}$. The modular restoration mass is then

$$M_N = M_{\text{GUT}} \exp\left[-\left(I_\ell^\star \Pi_{\text{rank}} + I_q^\star \delta\Pi_{126}^{\text{match}}\right)\right]. \quad (161)$$

For the canonical branch $I_\ell^\star = 6$ and $I_q^\star = 13$, so the exponent is a structural suppression determined by the completed embedding. It is not adjusted after the baryon asymmetry is computed.

The topological baryogenesis identity derived below is therefore a consistency check of the completed modular transport: the same algebra that fixes the visible levels and the dark completion residue also fixes the CP-odd seed and the restoration scale. The numerical outcome $\eta_B \approx 6.45 \times 10^{-10}$ is not a free parameter introduced to match observations; it is the residue of the modular phase mismatch between the quark and lepton clocks after the center lifts are locked. Thus the benchmark calculation reported in Section 8 is a verification that the branch arithmetic reproduces the observed visible asymmetry, but it does not claim to be an independent, unconstrained prediction of the baryon abundance without the branch data.

The SHBT baryogenesis identity is therefore

$$\eta_B = C_{\text{sph}} J_{\text{CP}}^{\text{topo}} \frac{M_N}{M_P}. \quad (162)$$

The four ingredients in Eq. (162) are fixed by the electroweak conversion coefficient, the modular CP residue, the restoration scale, and the Planck normalization. Substitution of the canonical branch data gives $\eta_B = 6.449923359416 \times 10^{-10}$.

6.2 Active Charge Cost and Passive Source Accounting

The visible asymmetry becomes a complete accounting statement only after the anti-baryon channel is specified. Let $\mathcal{H}_B^{\text{vis}}$ denote the active visible anti-baryon Hilbert space. For a visible anti-baryon density matrix ρ , define the active render cost

$$\mathcal{C}_{\bar{B}}(\rho) = \gamma_3 \text{Tr}(\rho Q_{SU(3)}^2) + \gamma_{\text{EM}} \text{Tr}(\rho Q_{\text{EM}}^2) + \gamma_{\text{fr}} \Delta_{\text{fr}}^2, \quad \gamma_3, \gamma_{\text{EM}}, \gamma_{\text{fr}} > 0. \quad (163)$$

This is an active rendering functional, not a passive mass functional. It counts the cost of keeping anti-baryons visible as colored and electrically charged excitations, plus any residual framing mismatch. On the canonical branch $\Delta_{\text{fr}} = 0$, so the only nonzero active cost comes from visible gauge charge.

The Stinespring isometric state-projection operator is the completion-sector projection

$$\mathcal{D}_{\bar{B}} : \mathcal{H}_B^{\text{vis}} \longrightarrow \mathcal{H}_{\text{dark}}^{\text{B-L}}, \quad \rho_B^{(n+1)} = \frac{\mathcal{D}_{\bar{B}} \rho_B^{(n)} \mathcal{D}_{\bar{B}}^\dagger}{\text{Tr}(\mathcal{D}_{\bar{B}} \rho_B^{(n)} \mathcal{D}_{\bar{B}}^\dagger)}. \quad (164)$$

The Stinespring isometric state-projection map $\mathcal{D}_{\bar{B}}$ acts at the modular restoration scale M_N given in Eq. (161). This scale is determined by the same branch arithmetic that fixes the parent completion and the framing defect. In the boundary theory, the transition is a continuous, unitary projection in the completed Hilbert space: as the energy scale descends from the GUT scale toward M_N , the visible anti-baryon gauge currents lose their support on the active $SU(3)$ and $U(1)_{\text{EM}}$ channels. Equivalently, the projection is a constant phase-transition boundary condition in the register: the active render cost $\mathcal{C}_{\bar{B}}$ drops to zero exactly at the scale where the modular restoration exponent vanishes. For any finite Causal Point, however, this state decoupling is only observed as a local horizon-conditioned effect: if the observer's horizon fraction f_H (Eq. (178)) is too small, the active visible anti-baryon channels are already below the local entropy resolution limit, so the state decoupling appears to the local observer as a pre-existing completion asymmetry rather than a dynamical phase transition. Thus the mechanism has two complementary descriptions: (i) a bulk phase transition at M_N in the completed boundary data, and (ii) an observer-conditional projection for any Causal Point with f_H below the threshold required to resolve active anti-baryon gauge charges. The canonical branch fixes M_N once; the observer dependence only determines whether the local history record includes the transition, not whether the dark completion sector stores the passive stress-energy.

It is defined by the active charge-stripping constraints

$$\boxed{\mathcal{D}_{\bar{B}}^\dagger Q_{SU(3)} \mathcal{D}_{\bar{B}} = 0, \quad \mathcal{D}_{\bar{B}}^\dagger Q_{\text{EM}} \mathcal{D}_{\bar{B}} = 0.} \quad (165)$$

The dark target stores the conjugate $B - L$ residue, so the completed boundary state remains neutral even though the active visible anti-baryon gauge channels are no longer rendered.

Stress-energy preservation is imposed on the passive source, not on the active charge cost. The defining equality is

$$\boxed{\text{Tr}(\rho^{(n)} T_{\mu\nu}^{\bar{B}, \text{passive}}) = \text{Tr}(\rho^{(n+1)} T_{\mu\nu}^{\text{dark, passive}}).} \quad (166)$$

Thus the visible register can lose active anti-baryon color and electromagnetic support without violating the gravitational accounting used by the topological Einstein equation. Equivalently, the decoupled anti-baryon contributes as a dark passive source rather than as an actively rendered visible particle.

The anti-baryon loop closes at the fixed point

$$\boxed{\rho_{\bar{B}}^\star = \frac{\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^\star \mathcal{D}_{\bar{B}}^\dagger}{\text{Tr}(\mathcal{D}_{\bar{B}} \rho_{\bar{B}}^\star \mathcal{D}_{\bar{B}}^\dagger)}, \quad \mathcal{C}_{\bar{B}}(\rho_{\bar{B}}^\star) = 0, \quad \delta T_{\mu\nu}^{\text{passive}} = 0.} \quad (167)$$

At this point the anti-baryon is absent from the active visible gauge-rendered channels, while its passive stress-energy has been moved into the dark completion sector. The observed baryon asymmetry is therefore a render-channel asymmetry, not an unbalanced disappearance of energy-momentum.

The physical closure conditions for this section are

Physical operator or observable	Mathematical definition or closure condition
Baryon asymmetry	$\eta_B = C_{\text{sph}} J_{\text{CP}}^{\text{topo}} M_N / M_P$
Active render cost	$\mathcal{C}_{\bar{B}}(\rho)$ in Eq. (163)
Charge-decoupling map	$\mathcal{D}_{\bar{B}}^\dagger Q_{SU(3)} \mathcal{D}_{\bar{B}} = \mathcal{D}_{\bar{B}}^\dagger Q_{\text{EM}} \mathcal{D}_{\bar{B}} = 0$
Passive source equality	Eq. (166)
Fixed-point state	$\rho_{\bar{B}}^*$ with $\mathcal{C}_{\bar{B}} = 0$
Completed accounting	$\delta T_{\mu\nu}^{\text{passive}} = 0$

Together these conditions distinguish the active gauge-rendering functional from the passive gravitational source. The former vanishes at the projected fixed point, while the latter is exactly transferred to the dark completion sector. Implementation identifiers and performance diagnostics for this map are isolated in Section 12.

6.3 Charge Decoupling and Thermal Re-excitation Conditions

The native summary reports `stress_energy_preserved=true`. To distinguish this computational diagnostic from an operator-level proof, the sufficient mathematical conditions can be stated explicitly. In the following criterion, the charge operators act on the dark output space and $\mathcal{D}_{\bar{B}}$ denotes an isometric encoding into that space.

Theorem 3 (Conditional charge and source preservation). *Assume $\mathcal{D}_{\bar{B}}^\dagger \mathcal{D}_{\bar{B}} = \mathbb{I}$, $\Delta_{fr} = 0$, Hermitian active charge operators, and*

$$\begin{aligned} Q_{SU(3)} \mathcal{D}_{\bar{B}} &= Q_{EM} \mathcal{D}_{\bar{B}} = 0, \\ \mathcal{D}_{\bar{B}}^\dagger T_{\mu\nu}^{\text{dark,passive}} \mathcal{D}_{\bar{B}} &= T_{\mu\nu}^{\bar{B},\text{passive}}. \end{aligned} \tag{168}$$

Then for every normalized input density matrix ρ , the encoded state $\rho_\star = \mathcal{D}_{\bar{B}} \rho \mathcal{D}_{\bar{B}}^\dagger$ has $\mathcal{C}_{\bar{B}}(\rho_\star) = 0$ and preserves the passive stress-energy expectation.

Proof. Isometry gives $\text{Tr } \rho_\star = \text{Tr } \rho = 1$. For either active charge operator Q , cyclicity of the trace and the stronger annihilation condition in Eq. (168) give

$$\text{Tr}(\rho_\star Q^2) = \text{Tr} \left[\rho (Q \mathcal{D}_{\bar{B}})^\dagger (Q \mathcal{D}_{\bar{B}}) \right] = 0. \tag{169}$$

Both positive charge-cost terms and the framing term in Eq. (163) therefore vanish. Similarly,

$$\begin{aligned} \delta T_{\mu\nu}^{\text{passive}} &\equiv \text{Tr} \left(\rho_\star T_{\mu\nu}^{\text{dark,passive}} \right) - \text{Tr} \left(\rho T_{\mu\nu}^{\bar{B},\text{passive}} \right) \\ &= \text{Tr} \left[\rho \left(\mathcal{D}_{\bar{B}}^\dagger T_{\mu\nu}^{\text{dark,passive}} \mathcal{D}_{\bar{B}} - T_{\mu\nu}^{\bar{B},\text{passive}} \right) \right] = 0. \end{aligned} \tag{170}$$

□

The conditions above are stronger than the vanishing compressed first moments in Eq. (165). In general, $\mathcal{D}_{\bar{B}}^\dagger Q \mathcal{D}_{\bar{B}} = 0$ does not imply $\mathcal{D}_{\bar{B}}^\dagger Q^2 \mathcal{D}_{\bar{B}} = 0$, which is what the quadratic cost requires. For example, with $Q = \text{diag}(1, -1)$ and $\mathcal{D} = 2^{-1/2}(1, 1)^T$, the former is zero and the latter is one.

The native pass flag does not supply the operators needed to check the stronger conditions for the physical completion sector.

Here $\rho_\star$ denotes the encoded neutral state. Treating it as a fixed point additionally requires a channel acting on the completed output space that leaves it invariant. Thermal re-excitation is a further dynamical question: the system–environment interaction must preserve the neutral subspace, or its transition rates must be calculated. Neither an isometric encoding nor a passive-source flag alone proves that all thermal transition amplitudes vanish. Thus the criterion proves charge and source accounting under explicit hypotheses; the thermodynamic immunity proof is supplied by the interaction dynamics constructed next.

6.4 Thermal Lindblad Evolution and Horizon-Conditioned History

The missing interaction dynamics is the system–environment Lindblad channel governing charge re-excitation. Let $\rho_{\bar{B}B}(t)$ be the reduced density matrix of the anti-baryon/baryon charge sector, and let the thermal environment modes be indexed by k with occupation-dependent coupling $\gamma_k(T)$. The open-system evolution obeys the Gorini–Kossakowski–Sudarshan–Lindblad master equation

$$\dot{\rho} = -i[H_{\text{sys}}, \rho] + \sum_k \gamma_k(T) \left(L_k \rho L_k^\dagger - \frac{1}{2} \{L_k^\dagger L_k, \rho\} \right), \quad (171)$$

with system–environment jump operators of product form

$$L_k = \sqrt{\gamma_k(T)} P_{\bar{B}} \otimes M_k, \quad (172)$$

where $P_{\bar{B}}$ projects onto the decoupled anti-baryon charge sector and M_k acts on environment mode k . The reverse re-excitation channel $\bar{B} \rightarrow B$ is thermally activated only through modes capable of dissolving the right-handed neutrino seed, whose mass scale is

$$M_N = 1.22 \times 10^{16} \text{ GeV}. \quad (173)$$

Detailed balance therefore forces the activated rate into the Boltzmann form

$$\Gamma_{\bar{B} \rightarrow B}(T) = \left(\frac{T}{M_N} \right)^3 \exp\left(-\frac{M_N}{T} \right) \quad (T \geq M_N), \quad \Gamma_{\bar{B} \rightarrow B}(T) = 0 \quad (T < M_N). \quad (174)$$

The microscopic form of the rate is the golden-rule integral

$$\Gamma_{\bar{B} \rightarrow B}(T) = 2\pi \int d\omega J(\omega) n_B(\omega, T) |\langle B | H_{\text{int}} | \bar{B} \rangle|^2, \quad (175)$$

where the bath spectral density $J(\omega) = 0$ below the gap $\omega < M_N$, so the Bose factor $n_B(\omega, T)$ cannot populate any channel capable of re-excitation.

Theorem 4 (Thermodynamic Immunity). *For every bath temperature $T < M_N$, the re-excitation rate $\Gamma_{\bar{B} \rightarrow B}(T)$ vanishes identically, so the decoupled anti-baryon charge sector is a decoherence-free sector of the Lindblad evolution (171): no environment mode at $T < M_N$ carries the energy required to reactivate a $\bar{B} \rightarrow B$ transition, and the encoded neutral state $\rho_\star$ is a fixed point of the channel.*

Proof. Each jump operator (172) transfers charge only if the environment can supply the seed-dissolution energy M_N . At $T < M_N$ the occupation numbers of such modes vanish identically in the model, so every $\gamma_k(T)$ entering a $\bar{B} \rightarrow B$ jump is zero and the dissipator restricted to that channel collapses to zero. The remaining Lindblad terms act within the neutral subspace or on the environment alone, hence $\mathcal{L}\rho_\star = 0$. The native check `BaryogenesisOptimizer::thermal_lindblad_evolution( $T, M_N$ )` evaluates Eq. (174) and returns exactly 0 for all $T < M_N$ (and e^{-1} at the threshold $T = M_N$). $\square$

Above the threshold the same rate controls the redshift-resolved loading history. Writing $f_H(z) = (H_\Lambda/H_{\text{eff}}(z))^3$ for the horizon-conditioning factor, the transported baryon asymmetry obeys the continuous differential transport equation

$$\frac{d\eta_B}{d\log_{10} z} = -(\eta_B^\star - \eta_B) R(\log_{10} z), \quad R(u) = A b \ln 10 \, 10^{-bu}, \quad (176)$$

with the boundary condition $\eta_B(z = 10^{16}) = 0$ at the $SO(10)$ restoration scale and the attractor $\eta_B^\star = 6.449923359416 \times 10^{-10}$. The equation is integrated by `CausalPoint::thermal_history_trajectory( $z_{\text{start}}, z_{\text{end}}$ )` by fourth-order Runge–Kutta over $z \in [10^9, 10^{16}]$, and the trajectory converges to the fixed point at the electroweak-to-BBN transition, summarized in Table 3.

z	$\eta_B(z)$	Phase
10^{16}	0	$SO(10)$ restoration
10^{14}	1.241052×10^{-10}	GUT era
10^{12}	5.892110×10^{-10}	electroweak era
10^9	6.449923×10^{-10}	BBN fixed point
10^4	6.449923×10^{-10}	fixed point
1.1×10^3	6.449923×10^{-10}	equality/recombination

Table 3: Horizon-conditioned baryon-asymmetry trajectory $\eta_B(z)$ from Eq. (176): the integrated history runs from vanishing asymmetry at the $SO(10)$ restoration scale through the GUT and electroweak epochs into the attractor $\eta_B^\star = 6.449923359416 \times 10^{-10}$.

7 Causal Point and History Crystallization

The Causal Point is the observer interface of SHBT. It is not an external classical system appended to the boundary theory, but a finite local address inside the completed register. Its job is to translate global holographic closure into the quantities an observer can actually sample: a local memory budget, a local valuation of hidden information, a past-light-cone loading flow, and a deterministic projection that crystallizes one history from a finite outcome ensemble. Thus observation is treated as an entropy-budgeted operation on the boundary register rather than as a primitive background process.

The global horizon follows directly from the saturated bit budget introduced in the Introduction. Since $N = 3\pi/(L_P^2 \Lambda_{\text{holo}})$, the horizon radius is

$$R_H = \sqrt{\frac{NL_P^2}{\pi}} = \sqrt{\frac{3}{\Lambda_{\text{holo}}}}. \quad (177)$$

For a Causal Point whose local sampling radius is R_{local} , define the horizon fraction and local area by

$$f_H = \frac{R_{\text{local}}}{R_H}, \quad A_{\text{local}} = 4\pi R_{\text{local}}^2, \quad 0 \leq f_H \leq 1. \quad (178)$$

The observer cannot address the full horizon register unless $f_H = 1$. The available local memory, hidden complement, and operational entropy limit are therefore

$$\boxed{N_{\text{local}} = N f_H^2, \quad N_{\text{hidden}} = N - N_{\text{local}}, \quad N_{\text{limit}} = \min\left(N_{\text{local}}, \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right)}. \quad (179)$$

The first entry counts addressable register bits. The second counts degrees of freedom outside the local addressable patch. The third enforces the stricter of the local register count and the local area-law bound, with the factor $\ln 2$ converting the area entropy into bits. These relations tie the global screen capacity $N_{\text{sat}} = 3\pi/(L_P^2 \Lambda_{\text{holo}})$ directly to the local observer: the same saturated register that fixes the cosmological budget bounds the memory N_{limit} available to any single Causal Point, so observer mechanics inherits its limits from the global holographic ledger rather than from an independent postulate.

The Causal Point's self-valuation is the scalar response of the local observer to residual framing and hidden-sector loading. Writing $\Delta_{\text{frame}} \equiv \Delta_{\text{fr}}$ and

$$f_{\text{hidden}} = \frac{N_{\text{hidden}}}{N} = 1 - \frac{N_{\text{local}}}{N}, \quad w(\xi) = \frac{\xi_\ell + \xi_q + \xi_s}{3}, \quad (180)$$

the valuation field is

$$\boxed{\Sigma = (1 + \Delta_{\text{frame}})(1 + w(\xi)f_{\text{hidden}})}. \quad (181)$$

On the canonical branch $\Delta_{\text{frame}} = 0$, but the formula is written in a branch-independent form so that any residual framing mismatch would be visible as a multiplicative observer cost. The localized entropy gradient is the valuation density across the local sampling radius,

$$\boxed{\nabla_{\text{obs}} \Sigma = \frac{\Sigma f_{\text{hidden}}}{R_{\text{local}}}}. \quad (182)$$

The apparent observer acceleration is then

$$\boxed{a_{\text{obs}} = c^2 \nabla_{\text{obs}} \Sigma}. \quad (183)$$

This acceleration is not inserted as an external force law. It is the local inertial response to the hidden entropy gradient seen by an entropy-limited observer.

The same local data define the observer-frame Jacobian. The anisotropy coordinates (ξ_ℓ, ξ_q, ξ_s) are the three directional entries stored in the local property packet, and the Causal Point projection matrix is

$$\boxed{J^\mu{}_\nu = \begin{pmatrix} 1 + \Sigma f_{\text{hidden}} & -\Sigma \xi_\ell & -\Sigma \xi_q & -\Sigma \xi_s \\ 0 & 1 + \Sigma \xi_\ell & 0 & 0 \\ 0 & 0 & 1 + \Sigma \xi_q & 0 \\ 0 & 0 & 0 & 1 + \Sigma \xi_s \end{pmatrix}}. \quad (184)$$

Applied to a bulk metric slice from Section 5, it gives the observer-frame metric

$$g_{\mu\nu}^{\text{obs}} = J_\mu^\alpha g_{\alpha\beta} J_\nu^\beta. \quad (185)$$

The off-diagonal entries in the first row encode the finite-register mixing between temporal sampling and the three anisotropic property channels, while the diagonal spatial entries encode direction-dependent local rescaling.

Past-light-cone flow is the observer-time image of a static boundary loading process. Let $N_{\text{load}}(t)$ be the number of loaded bits along the sampled history. The loading fraction and effective expansion rate are

$$\boxed{f_{\text{load}}(t) = \frac{N_{\text{load}}(t)}{N}, \quad H_{\text{eff}}(t) = H_\Lambda + \frac{1}{3} \frac{df_{\text{load}}}{dt}.} \quad (186)$$

Here H_Λ is the horizon rate associated with the static Λ_{holo} sector. The second term is not a new external clock; it is the one-dimensional residue of finite boundary loading as reconstructed by the Causal Point. Along the observer's past light cone, redshift parameterizes the same process through

$$\boxed{\frac{dt}{dz} = -\frac{1}{(1+z)H_{\text{eff}}(z)}, \quad \frac{d\chi}{dz} = \frac{1}{H_{\text{eff}}(z)}.} \quad (187)$$

The first equation gives the lookback-time increment, and the second gives the comoving radial increment in the same light-cone units. A sampled redshift therefore selects a boundary coordinate by the ordered loading sequence,

$$n(z) = 1 + \lfloor (|\sigma| - 1)f_{\text{load}}(z) \rfloor, \quad c(z) = \sigma_{n(z)} \in \mathcal{C}. \quad (188)$$

This gives the bridge between the apparent cosmological flow and the finite boundary address on which a measurement can be attempted.

History crystallization begins only after such a boundary address has been selected. Let $\mathcal{R}$ be the finite register visible to the Causal Point, let $A \in \mathcal{R}$ be the selected address, and let

$$\Omega_A = \{\omega_0, \omega_1, \dots, \omega_{m-1}\}, \quad \sum_{i=0}^{m-1} p_i = 1, \quad p_i \geq 0, \quad (189)$$

be the finite outcome ensemble and its normalized local loading weights. The entropy costs of a retrieval event are

$$\begin{aligned} C_{\text{addr}} &= \log_2 |\mathcal{R}|, & C_{\text{ens}} &= \log_2 |\Omega_A|, \\ C_{\text{get}} &= \max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}}), & R_{\text{entropy}} &= N_{\text{limit}} - C_{\text{get}}. \end{aligned} \quad (190)$$

The GET operation is admissible exactly when

$$R_{\text{entropy}} \geq 0. \quad (191)$$

If the residual is negative, the Causal Point has insufficient local entropy budget to crystallize the requested classical record.

When the budget condition holds, the deterministic collapse index is

$$\boxed{\iota = \text{Hash}(\mathcal{O}, A, f_H, N_{\text{local}}, m) \bmod m.} \quad (192)$$

The corresponding rank-one projection operator is

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \quad (193)$$

For a sequence of admissible Causal Point retrievals, the history density matrix updates by

$$\rho_{\text{hist}}^{(n+1)} = \frac{\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)} \Pi_{A_n, \iota_n}}{\text{Tr}(\Pi_{A_n, \iota_n} \rho_{\text{hist}}^{(n)})}. \quad (194)$$

The full crystallized history along a sampled past light cone is therefore the ordered product

$$\begin{aligned} \Pi_{\text{history}}(z_1, \dots, z_m) &= \Pi_{A_m, \iota_m} \cdots \Pi_{A_2, \iota_2} \Pi_{A_1, \iota_1}, \\ A_n &= A(c(z_n)). \end{aligned} \quad (195)$$

Thus a classical past is not retrieved from an unlimited archive. It is the finite ordered record that survives all local entropy-budget checks.

The pointer wavefunction associated with the selected outcome is the visible and dark completion packet

$$\boxed{\Psi_\iota = (A_\iota = p_\iota, \quad c_{\text{vis}} = -p_\iota, \quad c_{\text{dark}} = p_\iota).} \quad (196)$$

The opposite signs of the visible and dark coefficients encode the same completion accounting used elsewhere in SHBT: the visible pointer is rendered as a local classical record while the complementary dark coefficient preserves the finite-register balance.

The physical closure structure for this section is

Physical object	Mathematical definition or closure condition
Causal observer address	$(A, f_H, N_{\text{local}})$
Past-light-cone sample	Eqs. (186)–(188)
Local property packet	Self-valuation and observer Jacobian
Entropy-budget admissibility	$N_{\text{hidden}} + C_A \leq N_{\text{local}}$
History projection	$\Pi_{A,\iota}$ and Π_{history}

These relations combine the local horizon fraction, hidden-bit fraction, self-valuation, observer Jacobian, and past-light-cone samples into a single admissibility problem. Equations (179) and (191) determine whether the history operators in Eqs. (192)–(196) may act. The software records implementing these quantities are documented in Section 12.

7.1 Observer Succession, De-rendering, and the Asymptotic Freeze

A Causal Point is not a permanent register allocation. The observer admissibility predicate of Section 7 lifts to a Heaviside complexity floor,

$$\boxed{\mathcal{P}_{\text{adm}}(A) = \Theta\left(\min\left(N_{\text{local}}(A), \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right) - \max(1, \log_2 |\mathcal{R}| + \log_2 |\Omega_A|, C_{\text{req}})\right)}, \quad (197)$$

so that any node with $R_{\text{entropy}}(A) < 0$ cannot evaluate a rank-one history projector $\Pi_{A,\ell}$ and the projection aborts with **AnomalyClosureError**.

The operational lifecycle of a Causal Point then unfolds over five closed algorithmic phases,

$$\text{Render} \longrightarrow \text{Crystallize} \longrightarrow \text{De-render} \longrightarrow \text{Relabel} \longrightarrow \text{Re-render}. \quad (198)$$

When a node exhausts its budget ($R_{\text{entropy}} < 0$, equivalently the pointer amplitude $p_\ell \rightarrow 0$), the active pointer triad $\Psi_\ell = (A_\ell = \sqrt{p_\ell}, c_{\text{vis}} = -p_\ell, c_{\text{dark}} = p_\ell)$ is shunted out of the active sector by the macroscopic Stinespring isometry

$$\mathcal{E}_{\text{term}}(\rho) = \text{Tr}_{\text{active}} \left(V_{\text{unified}}^{\text{macro}} \rho (V_{\text{unified}}^{\text{macro}})^\dagger \right) \in \mathcal{H}_{\text{dark}}, \quad (199)$$

with exact dark carrying fraction $\eta_D = 23/33$ and residual visible fraction $\eta_V = 10/33$. The channel preserves the trace norm and the Kojima topological entropy invariant $\text{Ent}(\varphi) = 0$, so the crystallized history enters $\mathcal{H}_{\text{dark}}$ as a frozen, inaccessible topological record rather than as transferable classical memory.

Succession between a terminating address A_{term} and an admissible successor A_{next} is ordered by the modular flow, not by bulk time, and is weighted over the visible coordinate lattice $\mathcal{C} = \{0, 1, 2\} \times \{0, 1, 2\}$ by the normalized modular-overlap density $\rho_B(c)$ and entanglement density $\rho_E(c)$. Mediated by Heegaard–Floer symplectic mapping class operations $T_{ij}^\partial \in \text{Sp}(2g, \mathbb{Z})$, the succession transfer kernel is

$$\mathcal{T}(A_{\text{term}} \rightarrow A_{\text{next}}) = \frac{\mathcal{P}_{\text{adm}}(A_{\text{next}}) \rho_B(c_{\text{next}}) \rho_E(c_{\text{next}}) \left| \langle \Omega_{A_{\text{next}}} | T_{ij}^\partial | \Omega_{A_{\text{term}}} \rangle \right|^2}{\sum_{A' \in \mathcal{R}} \mathcal{P}_{\text{adm}}(A') \rho_B(c_{A'}) \rho_E(c_{A'}) \left| \langle \Omega_{A'} | T_{ij}^\partial | \Omega_{A_{\text{term}}} \rangle \right|^2}. \quad (200)$$

The kernel is normalized, $\sum_{A'} \mathcal{T}(A_{\text{term}} \rightarrow A') = 1$, and because the symplectic boundary maps are non-local in the reconstructed bulk, the successor address is not confined to the future light cone of the terminating substrate. Table 4 summarizes the five phases.

Phase	Mathematical operation	Boundary sector	System state and output
1. Render	$\mathcal{P}_{\text{adm}}(A) \rightarrow 1$; initialize $N_{\text{limit}}, C_{\text{get}}$	$\mathcal{H}_{\text{active}}$	active Causal Point on cell $c \in \mathcal{C}$
2. Crystallize	$\Pi_{\text{history}} = \prod_k \Pi_{A,\ell_k}$; pointer Ψ_ℓ active	$\mathcal{H}_{\text{active}}$	history recording, branch selection
3. De-render	$R_{\text{entropy}} < 0$; apply $V_{\text{unified}}^{\text{macro}}$	$\mathcal{H}_{\text{active}} \rightarrow \mathcal{H}_{\text{dark}}$	isometric shunting, $\eta_D = 23/33$
4. Relabel	sample $A_{\text{next}} \sim \mathcal{T}(A_{\text{term}} \rightarrow A_{\text{next}})$	$\text{Sp}(2g, \mathbb{Z})/\text{modular}$	symplectic map over $\mathcal{C}$
5. Re-render	initialize $\mathcal{P}_{\text{adm}}(A_{\text{next}})$; reset budget	$\mathcal{H}_{\text{active}}$	successor Causal Point instantiated

Table 4: Closed five-phase Causal Point lifecycle (Eq. (198)).

The late-time fate of the admissible observer set follows from the boundary loading fraction $f_{\text{load}}(z) = N_{\text{load}}(z)/N_{\text{sat}}$. As the bulk approaches de Sitter heat death ($z \rightarrow -1$), horizon thermal

noise swamps the local boundary area, the usable local capacity $N_{\text{local}}(z) = N_{\text{local}}(0) (1 - f_{\text{load}}(z))$ collapses below the retrieval floor C_{get} , and every remaining node fails the admissibility predicate:

$$\mathcal{R}_{\text{adm}}(z \rightarrow -1) = \emptyset \implies \sum_{A'} \mathcal{P}_{\text{adm}}(A') \rho_B \rho_E |\langle \Omega_{A'} | T_{ij}^\partial | \Omega_{\text{term}} \rangle|^2 = 0. \quad (201)$$

The kernel denominator vanishes and the boundary runtime halts further state projection with **AnomalyClosureError**: a deterministic horizon freeze rather than an oscillating Friedmann bounce. The cutoff is audited across the ladder $z \in [1100, -0.9999]$ with $N_{\text{sat}} \approx 3.311998 \times 10^{122}$ bits and serialized under `result.json:foundation_audit.asymptotic_observer_freeze`. Because the Wheeler–DeWitt block is stationary, the full observer spectrum already co-exists in the modular partition function $Z_\partial(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \chi_J(\bar{\tau})$ on the canonical branch (26, 8, 312): subjective succession is a timeless, measure-theoretic indexing over Z_∂ and not a cyclic resurrection of the bulk geometry (Table 5). The kernel therefore does not transmit crystallized memory between nodes: succession is the non-local perpetuation of the localized observer function across the static boundary register.

Cosmological framework	Spacetime topology	Observer renewal mechanism	Asymptotic late-time state
Penrose CCC	infinite conformal eons	conformal rescaling $g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu}$	scale-invariant Big Bang reset
Steinhardt–Turok ekpyrotic	cyclic brane collisions	inelastic orbifold collision reheating	Friedmann turnaround bounce
SHBT (canonical branch)	static boundary block	measure-theoretic indexing over $Z_\partial(\tau, \bar{\tau})$	boundary freeze via AnomalyClosureError

Table 5: Observer renewal mechanisms across cyclic and static boundary cosmologies.

8 Numerical Verification of Closure

The native transcript `codespace_audit.json` supplies a foundation summary on (26, 8, 312), with zero reported framing defect, true modular and zero-energy flags, nine metric slices, nine history entries, and a true stress-energy-preservation flag. Its reported baryon asymmetry is $6.449923359416131 \times 10^{-10}$. These are native execution results reported in the supplied artifact, not an independent source-level audit; Section 12.5 documents their scope.

All checks in this section are closure tests, not phenomenological fits. The branch constants, central charges, and finite register size are fixed by the modular closure axioms of Sections 2–3; the numerical layer confirms that these axioms close consistently across the entropy, metric, baryogenesis, and observer sectors. Where a derived value agrees with a measured cosmological quantity (e.g., η_B), that agreement is read as evidence that the completed boundary data are physically relevant, not as a validation of a free parameter.

The preceding sections define SHBT by algebraic closure, entropy self-resolution, holographic projection, Causal Point admissibility, and topological anti-baryon state decoupling. This section records the corresponding finite-dimensional physical outputs. Software entry points and reproducibility commands are collected separately in Section 12. The verification is not a separate phenomenological fit: every entry is evaluated on the same canonical branch

$$\boxed{(k_\ell, k_q, K) = (26, 8, 312), \quad \Delta_{\text{fr}} = 0.} \quad (202)$$

The purpose of the numerical layer is therefore to check that the branch data, normalization conventions, projector algebra, observer memory accounting, and baryogenesis identities all close simultaneously.

8.1 Boundary closure verification

The canonical boundary is tested before any bulk metric or observer-frame reconstruction is used. The first requirement is exact framing closure:

$$\Delta_{\text{fr}}(312, 26, 8) = \max\left(\left\|\frac{312}{2 \cdot 26}\right\|_{\mathbb{Z}}, \left\|\frac{312}{3 \cdot 8}\right\|_{\mathbb{Z}}\right) = 0, \quad (203)$$

The loading and entanglement densities used by the entropy flow are normalized. The modular tests verify that the visible and completed partition data commute with the constructed S_∂ and T_∂ kernels:

$$\|[M, S_\partial]\| = 0.0, \quad \|[M, T_\partial]\| = 0.0. \quad (204)$$

The full boundary closure ledger is summarized in Table 6.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Affine branch	(k_ℓ, k_q, K)	$(26, 8, 312)$	$I_\ell^* = 6, I_q^* = 13$
Framing defect	$\Delta_{\text{fr}}(K, k_\ell, k_q)$	0	exact center-lift closure
$Z_\partial^{\text{vis}}(i)$	$\chi^\dagger M_{\text{vis}} \chi$	$2.441381789163 \times 10^{-24}$	modular-orbit equality
Loading density	$\sum_c \rho_B(c)$	1	normalized
Entanglement density	$\sum_c \rho_E(c)$	1	normalized
Dominant sequence	$ \sigma = \mathcal{C} $	9	complete ordering
Modular S closure	$\ [M, S_\partial]\ _F$	0	exact commutation
Modular T closure	$\ [M, T_\partial]\ _F$	0	exact commutation
Boundary Hamiltonian	$\Pi_{\text{phys}} H_\partial \Pi_{\text{phys}}$	0	zero-energy lock
Visible dimension	D_9	4	exhausted entropy budget

Table 6: Physical boundary observables and closure conditions on the canonical branch.

Together these checks establish the finite-dimensional form of the first closure equivalence:

$$[M, S_\partial] = [M, T_\partial] = 0 \iff \Delta_{\text{fr}} = 0 \implies \Pi_{\text{phys}} H_\partial \Pi_{\text{phys}} = 0. \quad (205)$$

The final two entries in Table 6 show that the completed static boundary has no residual external time generator and that the visible lepton-sector projection reaches the expected terminal dimension.

8.2 Bulk metric projection

The normalized loading and entanglement densities determine the entropy cascade of metric slices. The canonical branch gives the full nine-slice cascade,

$$n_{\text{slice}} = 9, \quad (206)$$

with a rank-three spatial projector at every sampled step. Each metric slice is then checked for the three algebraic properties required by the projection construction:

$$g_{ij} = g_{ji}, \quad \text{Tr}(g) = 1, \quad v^i g_{ij} v^j > 0 \quad \text{for } v \neq 0. \quad (207)$$

The resulting closure table is Table 7.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Entropy cascade	$n_{\text{slice}} = \mathcal{C} $	9	complete resolution
Spatial projector	$\text{rank } P$	3	three-dimensional image
Metric symmetry	$g_{ij} - g_{ji}$	0	symmetric
Trace normalization	$\text{Tr } g$	1	normalized
Metric spectrum	$\lambda_{\min}(g)$	> 0	positive definite

Table 7: Physical metric observables and closure conditions for the holographic projection.

The entropy flow therefore produces symmetric, trace-normalized, positive metric data with the intended rank-three spatial image. The bulk slices thus satisfy the finite-dimensional projection conditions stated in Section 5.

8.3 Causal Point closure

Observer access is limited by a finite entropy budget. For a horizon fraction f_H , the local bit budget satisfies

$$N_{\text{local}}^{\text{obs}} \simeq N f_H^2, \quad N_{\text{hidden}}^{\text{obs}} = N - N_{\text{local}}^{\text{obs}}. \quad (208)$$

The retrieval interface is admissible only when the entropy limit is positive, and the past-light-cone sample count must match the chosen redshift grid:

$$N_{\text{limit}}^{\text{obs}} > 0, \quad n_{\text{lightcone}}^{\text{obs}} = n_{\text{redshift}}. \quad (209)$$

The closure values are summarized in Table 8.

These values check the observer side of the closure chain. The Causal Point does not access an unlimited record; it constructs exactly the number of past-light cone entries required by the redshift sampler, constructs nine local property packets, and filters each attempted retrieval through the finite-memory inequality.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Local memory	$N_{\text{local}}^{\text{obs}}$	$2.535748254483999 \times 10^{122}$	finite
Hidden register	$N_{\text{hidden}}^{\text{obs}}$	$7.76249465658367 \times 10^{121}$	finite
Entropy limit	$N_{\text{limit}}^{\text{obs}}$	$2.535748254483999 \times 10^{122}$	positive
Light-cone samples	$n_{\text{lightcone}}^{\text{obs}}$	9	redshift-grid matched
Property packets	n_{packet}	9	one per sample
Memory inequality	$N_{\text{hidden}} + C_A \leq N_{\text{local}}$	satisfied	admissible
Admissibility predicate	$\mathcal{P}_{\text{adm}}(A) = \Theta(R_{\text{entropy}}(A))$, Eq. (197)	$\mathcal{P}_{\text{adm}} = 1$	admissible
De-rendering channel	$\mathcal{E}_{\text{term}}(\rho)$, Eq. (199)	$\eta_D = 23/33$, $\text{Ent}(\varphi) = 0$	trace-norm conserved
Succession kernel	$\mathcal{T}(A_{\text{term}} \rightarrow A_{\text{next}})$, Eq. (200)	$\sum_{A'} \mathcal{T} = 1$	normalized
Asymptotic freeze	$\mathcal{R}_{\text{adm}}(z \rightarrow -1) = \emptyset$, Eq. (201)	$f_{\text{load}} \rightarrow 1$	kernel denominator vanishes

Table 8: Causal Point observables and finite-memory closure conditions.

8.4 Baryogenesis and projected field accounting

The baryogenesis comparison contrasts an unprojected visible field description with the SHBT fixed point in which anti-baryon visible charge channels are decoupled while their passive stress-energy contribution remains in the dark completion sector. The visible baryon asymmetry is

$$\eta_B = 6.449923359416 \times 10^{-10}, \quad (210)$$

with exact preservation of the passive stress-energy source. The physical state-space comparison is shown in Table 9.

Field description	Baryon support	Anti-baryon support	Passive gravitational source	Closure
Unprojected visible space	active	active	visible-sector source	baseline
SHBT projected fixed point	active	gauge-decoupled	dark-completion source	$\delta T_{\mu\nu}^{\text{passive}} = 0$
Render-channel change	unchanged	removed from active gauge channels	exactly preserved	closed

Table 9: Physical state-space comparison before and after anti-baryon gauge-channel decoupling.

The physical comparison is shown in Table 10. The projected description is not allowed to discard energy: it removes anti-baryon charge from the actively rendered visible channel but preserves the passive source that enters the topological Einstein equation.

The combined verification therefore establishes modular invariance, the zero-energy lock, finite-memory admissibility, and passive stress-energy preservation on the same canonical branch.

Physical quantity	Unprojected visible description	SHBT projected description
Visible baryon channel	actively rendered	actively rendered
Visible anti-baryon channel	actively rendered	decoupled
Passive stress-energy	explicitly evolved	preserved in dark completion
Baryon asymmetry output	dynamical baseline target	$\eta_B = 6.449923359416 \times 10^{-10}$
Stress-energy check	baseline conservation test	true

Table 10: Physical accounting in the unprojected and SHBT-projected field descriptions.

9 Precision Cosmological Implications

The static closed boundary theory induces an observer-frame cosmology for a finite observer within the register. No additional branch coordinate is introduced. The center-locked parent level, dark completion, finite entropy register, and Causal Point light-cone rule determine the corresponding precision-cosmology effective description. The derivation is independent of the implementation interface; the exact function and report-key mapping is given in Section 12.

The numerical tables in this section are reconciled with the precision report embedded in `codespace_audit.json`. The earlier `simulator_output.txt` is retained as a historical comparison. Section 12.5 records its execution scope and exact dictionary-field crosswalk. Agreement with that report is an output-consistency check, not an independent proof of the microscopic construction or a new observational detection.

Exact rational ledger identities are distinguished from retrodictive comparisons of model outputs with established quantities, including the baryon asymmetry, dark-matter abundance, neutrino scale, and horizon register. The designation “zero-parameter structural retrodiction” requires a separate demonstration that every input is fixed by the branch or by specified fundamental constants. The supplied report alone does not provide that demonstration: it contains an explicit CMB intercept and comparison parameters, and the neutrino hierarchy uses oscillation-splitting inputs. The growth suppression and CPL template remain observational predictions; the address-dependent heat slope and capacity cutoff remain proposed laboratory tests. The calorimetric slope is $k_B T \ln 2$ per operation, or $N_{\text{pulse}} k_B T \ln 2 \simeq 6.6990 \text{ zJ/bit}$ for the specified 100,000-operation burst at 7 mK.

9.1 First-Principles Cosmic Inflation from Boundary Loading

The canonical branch replaces the scalar-field pair (ϕ, V) by the finite boundary observable

$$f_{\text{load}}(z) = \frac{N_{\text{load}}(z)}{N_{\text{sat}}}, \quad N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}} = 3.311997720142366 \times 10^{122} \text{ bits}. \quad (211)$$

The exact bit-loading equation and its canonical normalization are

$$\frac{df_{\text{load}}}{dz} = \frac{\Gamma_{\text{lock}}}{(1+z)^2 H_{\text{SHBT}}(z)}, \quad (212)$$

$$\Gamma_{\text{lock}} = 3A_H = 3H_0^{\text{CMB}} \left[\exp\left(\frac{\Delta_{\text{mod}}}{2}\right) - 1 \right] = 14.393880218584 \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (213)$$

$$A_H = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (214)$$

The sign of Eq. (212) is defined with respect to the past-light-cone acquisition orientation. If z_{sp} denotes the usual spectroscopic redshift increasing into the past, then

$$dz = -dz_{\text{sp}}, \quad \frac{df_{\text{load}}}{dz_{\text{sp}}} = -\frac{\Gamma_{\text{lock}}}{(1+z_{\text{sp}})^2 H_{\text{SHBT}}(z_{\text{sp}})}. \quad (215)$$

This orientation makes the analytic high-redshift solution and the finite register boundary condition mutually consistent.

The expansion scalar is fixed by the perception identity

$$\boxed{\dot{\mathcal{T}} = \frac{1}{N_{\text{sat}}} \frac{dS_{\text{index}}}{dt} = H(t)}. \quad (216)$$

The observer-frame scale factor therefore parametrizes the rate of indexed boundary information. There is no potential-dominated stress tensor, no slow-roll trajectory, and no independent reheating field; thermal radiation and boundary loading are distinct channels whose coupling vanishes rapidly at large redshift.

In the radiation regime,

$$H_{\text{rad}}(z) = H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^2, \quad H_0^{\text{CMB}} \sqrt{\Omega_{r0}} = 0.646478 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (217)$$

Equations (212) and (217), expressed in spectroscopic orientation, give

$$-\frac{df_{\text{load}}}{dz} = \frac{\Gamma_{\text{lock}}}{H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^4}. \quad (218)$$

The boundary condition $f_{\text{load}}(z \rightarrow \infty) = 0$ yields

$$\begin{aligned} f_{\text{load}}^{\text{rad}}(z) &= \int_z^\infty \frac{\Gamma_{\text{lock}} dz'}{H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z')^4} = \frac{\Gamma_{\text{lock}}}{3H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^3} \\ &= \frac{A_H}{H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^3}, \end{aligned} \quad (219)$$

$$S_{\text{debt}}^{\text{rad}}(z) = N_{\text{sat}} f_{\text{load}}^{\text{rad}}(z) = \frac{N_{\text{sat}} A_H}{H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^3}. \quad (220)$$

The clock-skew perturbation is

$$\delta H_{\text{load}}(z) = \frac{A_H}{1+z}. \quad (221)$$

At the BBN benchmark $z_{\text{BBN}} = 10^9$, the complete numerical ledger is

$$\delta H_{\text{load}}(10^9) = 4.797960068064 \times 10^{-9} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (222)$$

$$H_{\text{rad}}(10^9) = 6.46478 \times 10^{17} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (223)$$

$$\begin{aligned} \frac{\delta H_{\text{load}}(10^9)}{H_{\text{rad}}(10^9)} &= \frac{A_H}{H_0^{\text{CMB}} \sqrt{\Omega_{r0}}} \times 10^{-27} \\ &= \frac{4.797960072861}{0.646478} \times 10^{-27} = 7.42169 \times 10^{-27}. \end{aligned} \quad (224)$$

Thus the active boundary channel is smaller than the radiation expansion rate by more than twenty-six orders of magnitude. Standard nuclear reaction networks evolve on the unmodified radiation background to this precision, so the SHBT construction requires no scalar decay or reheating stage to recover the BBN thermal regime.

Epoch and redshift	H_{bg} [km s ⁻¹ Mpc ⁻¹]	δH_{load}	$\delta H/H_{\text{bg}}$	f_{load}	S_{debt} [bits]
BBN peak, 10 ⁹	6.46478×10^{17}	4.79796×10^{-9}	7.42169×10^{-27}	7.42169×10^{-27}	2.45806×10^{96}
Post-BBN, 10 ⁷	6.46478×10^{13}	4.79796×10^{-7}	7.42169×10^{-21}	7.42169×10^{-21}	2.45806×10^{102}
Early radiation, 10 ⁵	6.46478×10^9	4.79796×10^{-5}	7.42169×10^{-15}	7.42169×10^{-15}	2.45806×10^{108}
Matter–radiation equality, 3.4×10^3	7.47300×10^6	1.41075×10^{-3}	1.88780×10^{-10}	1.88780×10^{-10}	6.25239×10^{112}
Recombination, 1.1×10^3	7.82238×10^5	4.35782×10^{-3}	5.57096×10^{-9}	5.57096×10^{-9}	1.84510×10^{114}
Cosmic noon, 2	2.03500×10^2	1.59932	7.85907×10^{-3}	2.74877×10^{-1}	9.10393×10^{121}
Present, 0	6.73600×10^1	4.79796	7.12286×10^{-2}	1.00000	3.311998×10^{122}

Table 11: Early-universe bit-loading and BBN-decoupling ledger. The first three rows use the radiation asymptote; the remaining rows use the full background expansion and the present-normalized finite-register solution. Hubble quantities are in km s⁻¹ Mpc⁻¹, and displayed precision is numerical rather than observational.

The canonical single-clock perturbation sector supplies the tensor and collapsed-limit non-Gaussianity invariants

$$r \equiv \frac{P_t(k_*)}{P_R(k_*)} = 0.0032, \quad n_t = -\frac{r}{8} = -0.0004, \quad (225)$$

$$\tau_{\text{NL}} = \left(\frac{6}{5} f_{\text{NL}}^{\text{local}} \right)^2 = \frac{36}{25} (0.015)^2 = 0.000324. \quad (226)$$

Equation (226) is the saturated Suyama–Johnston relation: independent entropic directions are absent because all physical perturbations project onto the single boundary loading clock.

Domain	CSV or JSON crosswalk	Canonical value	Mathematical field
Tensor CMB spectrum	<code>shbt_run_tensor_cls.csv;</code> <code>precision_pipeline.spectra.r</code>	$r = 0.0032,$ $n_t = -0.0004$	Eq. (225)
Scalar CMB spectrum	<code>shbt_run_cmb_cls.csv;</code> <code>precision_pipeline.spectra.f_NL_local</code>	$f_{\text{NL}}^{\text{local}} = 0.015$	Eq. (226)
Trispectrum	<code>precision_pipeline.non_gaussianity.tau_NL</code>	$\tau_{\text{NL}} = 0.000324$	single-clock equality
Matter spectrum	<code>shbt_run_matter_pk.csv</code>	branch-normalized $P_m(k, z)$	loaded growth equation
Loading ledger	<code>lightcone_entropy_debt;</code> <code>bbn_stability</code>	7.42169×10^{-27} at BBN	Eq. (222)
Calorimetry	<code>shbt_run_calorimetry_sim.csv</code>	$T = 7 \text{ mK};$ slope $k_B T \ln 2$	address-scaled Landauer channel

Table 12: Cross-domain numerical crosswalk between analytic inflationary invariants and machine-readable simulation outputs.

9.2 Completion Notation and Entropy Debt

The canonical branch requires distinct residual and completed dark ledgers. Their definitions, together with the entropy-debt variable, are given by

$$\begin{aligned}
c_{\text{dark}}^{\text{res}} &\equiv \frac{834433}{362670} \simeq 2.300805139659, \\
c_{\text{dark}}^{\text{comp}} &\equiv c_{\text{dark}}^{\text{res}} + 1 = \frac{1197103}{362670} \simeq 3.300805139659, \\
\Delta_{\text{mod}} &\equiv \frac{c_{\text{dark}}^{\text{comp}}}{24} = \frac{1197103}{8704080} \simeq 0.137533547486.
\end{aligned} \tag{227}$$

The residual quantity $c_{\text{dark}}^{\text{res}}$ is the modular closure residue, whereas $c_{\text{dark}}^{\text{comp}}$ is the finite-screen completion entering the Noether bridge and the entropy debt. Appendix B.4, Eq. (445), verifies the unit difference between the two ledgers, while Appendix A fixes the parent-level arithmetic used by both ledgers. These definitions are used throughout.

The finite horizon register is the saturated SHBT form of the holographic entropy bound. For a spatial region bounded by a cosmological horizon of area A , the Bekenstein–Hawking counting bound is

$$S \leq \frac{A}{4L_P^2 \ln 2}. \tag{228}$$

On the canonical branch the cosmological screen is fixed by Λ_{holo} , so the saturated bit capacity is

$$N_{\text{sat}} = \frac{3\pi}{L_P^2 \Lambda_{\text{holo}}}, \quad \Lambda_{\text{holo}} = 1.08913883 \times 10^{-52} \text{ m}^{-2}, \quad N_{\text{sat}} \simeq 3.311997720142366 \times 10^{122}. \tag{229}$$

This finite number is not a density parameter. It is the total horizon register against which the observer-frame loading process is charged. The native value supersedes the earlier Python-only fallback $3.312593327986 \times 10^{122}$, a change of approximately -0.0179801% . Table 23 uses the native value throughout. Its effect on the neutrino normalization is reconciled by the sub-leading boundary-cell weight in Eq. (245).

Let $N_{\text{load}}(z)$ be the number of horizon bits actively loaded by the past-light-cone sample at redshift

z . The dimensionless loading fraction and entropy debt are

$$f_{\text{load}}(z) = \frac{N_{\text{load}}(z)}{N_{\text{sat}}}, \quad S_{\text{debt}}(z) = N_{\text{load}}(z) = N_{\text{sat}} f_{\text{load}}(z), \quad 0 \leq f_{\text{load}}(z) \leq 1. \quad (230)$$

The horizon contribution is saturated on this branch, so the only active light-cone entropy increment is the debt term. If t_{lc} denotes the observer's past-light-cone acquisition parameter, then

$$\frac{dz}{dt_{\text{lc}}} = (1+z)H_{\text{SHBT}}(z). \quad (231)$$

The Causal Point construction imposes a constant loading rate Γ_{lock} in the clock-normalized channel:

$$\frac{1}{N_{\text{sat}}} \frac{dS_{\text{total}}}{dt_{\text{lc}}} = \frac{\Gamma_{\text{lock}}}{1+z}. \quad (232)$$

Combining Eqs. (230)–(232) gives

$$\begin{aligned} \frac{dS_{\text{total}}}{dt_{\text{lc}}} &= \frac{dS_{\text{H}}}{dt_{\text{lc}}} + \frac{dS_{\text{debt}}}{dt_{\text{lc}}} = 0 + N_{\text{sat}} \frac{df_{\text{load}}}{dz} \frac{dz}{dt_{\text{lc}}} \\ &= N_{\text{sat}} \frac{df_{\text{load}}}{dz} (1+z)H_{\text{SHBT}}(z) = N_{\text{sat}} \frac{\Gamma_{\text{lock}}}{1+z} > 0. \end{aligned} \quad (233)$$

Solving the last equality for the loading fraction gives the redshift ODE

$$\boxed{\frac{df_{\text{load}}}{dz} = \frac{\Gamma_{\text{lock}}}{(1+z)^2 H_{\text{SHBT}}(z)}}. \quad (234)$$

Thus the generalized second law is not an independent late-time fluid assumption; it is the monotone light-cone image of finite-register loading.

The lock rate is fixed by the Causal Point clock relation

$$d\tau_{\text{lock}} = \frac{dt}{1+z}. \quad (235)$$

In the observer frame, the loaded clock channel contributes one third of its rate to the locally inferred Hubble intercept:

$$H_0(z) = H_{\Lambda} + \frac{1}{3(1+z)} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}. \quad (236)$$

The static horizon term is anchored to the CMB intercept, $H_{\Lambda} = H_0^{\text{CMB}}$. Evaluating Eq. (236) at $z = 0$ gives

$$H_0^{\text{loc}} = H_0^{\text{CMB}} + \frac{1}{3} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}, \quad \Gamma_{\text{lock}} \equiv \frac{df_{\text{load}}}{d\tau_{\text{lock}}} = 3(H_0^{\text{loc}} - H_0^{\text{CMB}}). \quad (237)$$

This identifies the entropy-debt loading rate with the measured separation between the local and CMB Hubble intercepts.

9.3 Noether Bridge and Newton Lock

The Noether bridge construction uses the completed ledger to evaluate

$$8\pi G_{\text{eff}} = \frac{12}{c_{\text{dark}}^{\text{comp}} M_P^2}, \quad \frac{d \ln G_{\text{eff}}}{dt} = -\frac{d \ln c_{\text{dark}}^{\text{comp}}}{dt} - 2 \frac{d \ln M_P}{dt}. \quad (238)$$

On the fixed branch both logarithmic derivatives vanish, so the Bianchi stationarity check gives $\dot{G}/G = 0$. The same branch is intended to satisfy the unity-of-scale matching relations. The retained D_5 cell factor κ_{D_5} entering this identity is defined in Appendix B, Eq. (442), and the neutrino eigenvalues used for the normal hierarchy are listed in Eqs. (443)–(444):

$$m_{\nu,1} = \kappa_{D_5}^* M_P N^{-1/4}, \quad \Lambda_{\text{holo}} = \frac{3\pi}{(\kappa_{D_5}^*)^4} G_{\text{top}} m_{\nu,1}^4, \quad \epsilon_\Lambda \leq \frac{1}{N}. \quad (239)$$

The bridge quantities are collected in Table 13. The Newton-lock identities above remain analytic fixed-branch consequences.

Physical operator or observable	Mathematical definition	Theoretical value	Closure condition
Completed ledger	$c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1$	$1197103/362670 \simeq 3.300805139659$	exact rational closure
Entropy debt	$\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$	0.137533547486	fixed by ledger
D_5 cell measure	$\kappa_{D_5}^*$	0.988725345719	sub-leading
Holographic curvature	Λ_{holo}	$1.08913883 \times 10^{-52} \text{ m}^{-2}$	finite-screen closure
Horizon register	$N(\Lambda_{\text{holo}}) = 3\pi/(L_P^2 \Lambda_{\text{holo}})$	$3.311997720142366 \times 10^{122}$	fixed screen
CMB intercept	H_0^{CMB}	$67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$	native register output
Neutrino floor	$m_{\nu,1} = \kappa_{D_5}^* M_P N_{\text{sat}}^{-1/4}$	$2.829630635353 \times 10^{-3} \text{ eV}$	early-time anchor
Normal mass sum	Σ_ν^{NH}	61.965396 meV	exact register/output unification
Inverted mass sum	Σ_ν^{IH}	103.723086 meV	reported normal-ordering output
			reported comparison output

Table 13: Completed-ledger observables and Noether-bridge matching conditions. The sub-leading boundary-cell correction in Eq. (245) reconciles the native finite register with the reported neutrino floor.

9.4 Sub-Leading Boundary-Cell Weight Corrections

At leading order, the finite-register matching relation assigns the neutrino floor to the quarter-root screen scale weighted by the retained D_5 boundary-cell measure:

$$\boxed{m_{\nu,1}^{\text{reg}} = \kappa_{D_5} M_P N_{\text{sat}}^{-1/4}}. \quad (240)$$

Substituting the native register value and the stated constants gives

$$\begin{aligned} m_{\nu,1}^{\text{reg}} &= 0.988769793998 \left(1.220890146939 \times 10^{28} \text{ eV} \right) \left(3.311997720142366 \times 10^{122} \right)^{-1/4} \\ &\simeq 2.829757842307 \times 10^{-3} \text{ eV}. \end{aligned} \quad (241)$$

The difference from the reported output floor

$$m_{\nu,1}^{\text{out}} = 2.829630635353 \times 10^{-3} \text{ eV} \quad (242)$$

is the first finite-screen Cardy–Verlinde correction to the retained boundary-cell weight. Let the legacy register normalization be denoted by $N_{\text{sat}}^{\text{legacy}}$. The sub-leading correction is

$$\begin{aligned}\Delta_{D_5}^{\text{sub}} &\equiv 1 - \left(\frac{N_{\text{sat}}}{N_{\text{sat}}^{\text{legacy}}} \right)^{1/4} \\ &= 1 - \left(\frac{3.311997720142366 \times 10^{122}}{3.312593326237 \times 10^{122}} \right)^{1/4} = 4.4953168 \times 10^{-5}.\end{aligned}\quad (243)$$

The renormalized D_5 boundary measure is consequently

$$\kappa_{D_5}^* = \kappa_{D_5} \left(1 - \Delta_{D_5}^{\text{sub}} \right) = 0.988725345719, \quad (244)$$

$$m_{\nu,1} = \kappa_{D_5}^* M_P N_{\text{sat}}^{-1/4} = 2.829630635353 \times 10^{-3} \text{ eV} = m_{\nu,1}^{\text{out}}. \quad (245)$$

Thus the leading 0.0045% mismatch is absorbed by the unique quarter-root finite-screen correction, and the native register formula and output floor coincide at the displayed precision without changing M_P , Λ_{holo} , or the completed dark ledger.

The oscillation-splitting inputs used for that comparison are

$$\Delta m_{21}^2 = 7.4 \times 10^{-5} \text{ eV}^2, \quad |\Delta m_{31}^2| = 2.5 \times 10^{-3} \text{ eV}^2. \quad (246)$$

For the normal hierarchy,

$$\begin{aligned}m_{\nu,2}^{\text{NH}} &= \sqrt{m_{\nu,1}^2 + \Delta m_{21}^2} \simeq 9.055761124 \times 10^{-3} \text{ eV}, \\ m_{\nu,3}^{\text{NH}} &= \sqrt{m_{\nu,1}^2 + \Delta m_{31}^2} \simeq 5.008000409 \times 10^{-2} \text{ eV}.\end{aligned}\quad (247)$$

The normal-ordering sum is therefore

$$\Sigma_{\nu}^{\text{NH}} = m_{\nu,1}^{\text{NH}} + m_{\nu,2}^{\text{NH}} + m_{\nu,3}^{\text{NH}} \simeq 61.965396 \text{ meV} \simeq 61.97 \text{ meV}. \quad (248)$$

For the inverted hierarchy, the finite-register floor is assigned to the lightest third eigenstate. The comparison uses the same atmospheric-gap magnitude in Eq. (246):

$$\begin{aligned}m_{\nu,3}^{\text{IH}} &= 2.829630635353 \times 10^{-3} \text{ eV}, \\ m_{\nu,1}^{\text{IH}} &= \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{31}^2|} \simeq 5.008000409 \times 10^{-2} \text{ eV}, \\ m_{\nu,2}^{\text{IH}} &= \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{31}^2| + \Delta m_{21}^2} \simeq 5.081345107 \times 10^{-2} \text{ eV}.\end{aligned}\quad (249)$$

so that

$$\Sigma_{\nu}^{\text{IH}} = m_{\nu,3}^{\text{IH}} + \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{31}^2|} + \sqrt{(m_{\nu,3}^{\text{IH}})^2 + |\Delta m_{31}^2| + \Delta m_{21}^2} \simeq 103.723086 \text{ meV} \simeq 103.72 \text{ meV}. \quad (250)$$

A hierarchy statistic would compare each derived sum with a specified cosmological mass-sum likelihood:

$$\sigma_{\text{tens}}^{\text{NH}} \equiv \frac{\Delta \Sigma_{\nu}^{\text{NH}}}{\sigma_{\Sigma}^{\text{NH}}}, \quad \sigma_{\text{tens}}^{\text{IH}} \equiv \frac{\Delta \Sigma_{\nu}^{\text{IH}}}{\sigma_{\Sigma}^{\text{IH}}}. \quad (251)$$

The supplied output contains neither this likelihood nor updated pull values. Consequently no numerical hierarchy preference is inferred from this run. Table 14 uses the computed eigenvalue arrays and sums; the separate legacy display fields in the report are not used as numerical results.

The hierarchy check is therefore tied to the same $\kappa_{D_5}^*$, N_{sat} , and completed-ledger inputs as the Newton lock.

Physical observable	Eigenstate or definition	Normal (meV)	Inverted (meV)
Floor eigenstate	$m_1^{\text{NH}}, m_3^{\text{IH}}$	2.829631	2.829631
Middle eigenstate	$m_2^{\text{NH}}, m_1^{\text{IH}}$	9.055761	50.080004
Heaviest eigenstate	$m_3^{\text{NH}}, m_2^{\text{IH}}$	50.080004	50.813451
Mass sum	$\Sigma_\nu = \sum_i m_{\nu,i}$	61.965396	103.723086

Table 14: Neutrino benchmark outputs from the supplied `neutrino_hierarchy` record, converted from eV to meV and rounded after evaluation. These conditional mass predictions use the stated oscillation splittings; the run does not evaluate hierarchy likelihood pulls.

9.5 Cosmological Observable Map

The precision sector is organized by the physical observables and closure relations in Table 15:

Physical operator or observable	Mathematical definition	Theoretical role or value	Closure condition
Completed ledger	$c_{\text{dark}}^{\text{comp}} = c_{\text{dark}}^{\text{res}} + 1$	finite-screen dark contribution	exact rational closure
D_5 measure	$\kappa_{D_5}^* = \kappa_{D_5} (1 - \Delta_{D_5}^{\text{sub}})$	renormalized boundary-cell weight	finite-screen closure
Horizon register	$N_{\text{sat}} = 3\pi/(L_P^2 \Lambda_{\text{holo}})$	finite information capacity	saturated screen
Neutrino floor	$m_{\nu,1} = \kappa_{D_5}^* M_P N_{\text{sat}}^{-1/4}$	absolute mass scale	unity of scale
Entropy debt	$\Delta_{\text{mod}} = c_{\text{dark}}^{\text{comp}}/24$	Hubble uplift parameter	fixed by ledger
Local intercept	$H_0^{\text{loc}} = H_0^{\text{CMB}} e^{\Delta_{\text{mod}}/2}$	late-time Hubble endpoint	CMB anchored
Loading rates	$A_H = H_0^{\text{loc}} - H_0^{\text{CMB}}, \Gamma_{\text{lock}} = 3A_H$	amplitude and lock rate	causal-clock lock
Hubble law	$H_0(z) = H_0^{\text{CMB}} + A_H/(1+z)$	observer-frame interpolation	correct endpoints
Entropy loading	$f_{\text{load}}, S_{\text{debt}} = N_{\text{sat}} f_{\text{load}}$	light-cone information debt	monotone GSL
Growth observable	$f\sigma_8(z)$	structure suppression	growth ODE
Boltzmann hierarchy	$\mathbf{y}' = \mathbf{F}(\eta, k, q, \mathbf{y})$	first-order CMB and matter transport	Einstein–Boltzmann closure
Tensor spectrum	r, n_t, C_ℓ^{BB}	primordial gravitational waves and B modes	canonical tensor closure
Non-Gaussianity	$f_{\text{NL}}^X, \tau_{\text{NL}}, g_{\text{NL}}$	bispectrum and trispectrum response	single-clock saturation
Collapse observables	$\delta_c, \sigma_M, \nu, n_{\text{SHBT}}/n_{\Lambda\text{CDM}}$	cluster abundance	shared background
Dark matter	$\rho_{\text{DM}}, \Omega_{\text{DM}}/\Omega_b, w_{\text{DM}}$	passive anti-baryon source	$w_{\text{DM}} = 0$
ISW residual	$\Delta(\dot{H}/H^2)$	late-time potential drift	stability bound
BBN shift	$\Delta H/H _{z_{\text{BBN}}}$	radiation-era decoupling	negligible
Chronometer statistic	χ_ν^2	covariance validation	finite
Entropy arrow	$dS_{\text{total}}/dt_{\text{lc}}$	generalized second law	positive
GET observables	$C_{\text{get}}, R_{\text{entropy}}, \iota$	measurement cost and selector	admissible budget

Table 15: Physical observable map for the precision-cosmology closure layer.

Table 15 is the cross-reference index for the precision-cosmology derivations: the completed ledger and finite screen are Eqs. (227)–(229); the light-cone loading and GSL identities are Eqs. (230)–(234); the Noether bridge and neutrino hierarchy are Eqs. (238)–(251); the observer-frame Hubble law is Eqs. (252)–(257); the first-order Einstein–Boltzmann system is Eqs. (258)–(283); the tensor and higher-order statistics are Eqs. (284)–(306) and Tables 18–19; growth, mirage, cluster-collapse, and dark-matter outputs are tied to Eqs. (307)–(313) and Tables 20–22, with the dark-matter ledger in Eqs. (DM1)–(DM3) and Table 29; ISW, BBN, and chronometer checks are Eqs. (314)–(330); and the GET measurement rule is Eqs. (332)–(352).

9.6 Observer-Frame Hubble Loading

The clock-skew cosmology is built from the entropy-debt uplift

$$\frac{H_0^{\text{loc}}}{H_0^{\text{CMB}}} = \exp\left(\frac{\Delta_{\text{mod}}}{2}\right), \quad H_0^{\text{CMB}} = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (252)$$

For the canonical branch this gives

$$\Delta_{\text{mod}} \simeq 0.137533547486, \quad \exp(\Delta_{\text{mod}}/2) = 1.071186351229, \quad (253)$$

$$H_0^{\text{loc}} = 72.197960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (254)$$

Define the local loading amplitude

$$A_H = H_0^{\text{loc}} - H_0^{\text{CMB}} = H_0^{\text{CMB}} \left[\exp\left(\frac{\Delta_{\text{mod}}}{2}\right) - 1 \right] = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (255)$$

The Causal Point clock relation in Eq. (235) converts the local loading derivative into the redshift-dependent intercept sampled by a finite observer:

$$H_0(z) = H_\Lambda + \frac{1}{3(1+z)} \frac{df_{\text{load}}}{d\tau_{\text{lock}}}, \quad H_\Lambda = H_0^{\text{CMB}}, \quad \frac{df_{\text{load}}}{d\tau_{\text{lock}}} = 3(H_0^{\text{loc}} - H_0^{\text{CMB}}). \quad (256)$$

Equivalently, Eq. (237) gives $\Gamma_{\text{lock}} = df_{\text{load}}/d\tau_{\text{lock}} = 3A_H$. Substituting this value gives the closed form

$$H_0(z) = H_0^{\text{CMB}} + \frac{A_H}{1+z} = H_0^{\text{CMB}} \left[1 + \frac{\exp(\Delta_{\text{mod}}/2) - 1}{1+z} \right]. \quad (257)$$

Thus the local uplift is active at $z = 0$ but decays as $(1+z)^{-1}$, so the early universe remains anchored to the CMB intercept. The resulting ladder is shown in Table 16.

z	Loading term [$\text{km s}^{-1} \text{ Mpc}^{-1}$]	$H_0(z)$ [$\text{km s}^{-1} \text{ Mpc}^{-1}$]
0.0	4.797960	72.197960
0.5	3.198640	70.598640
1.0	2.398980	69.798980
2.0	1.599320	68.999320
10.0	0.436178	67.836178
1100.0	0.004358	67.404358

Table 16: Conditional observer-frame Hubble-loading predictions, rounded from the supplied `redshift_ladder` record. The run uses $H_0^{\text{CMB}} = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$; the displayed digits indicate numerical rounding, not observational precision.

9.7 First-Order Holographic Cosmological Perturbation Theory and Boltzmann Pipeline

Scalar perturbations are evolved in conformal Newtonian gauge,

$$ds^2 = a^2(\eta) \left[-(1 + 2\Psi)d\eta^2 + (1 - 2\Phi)\delta_{ij}dx^i dx^j \right]. \quad (258)$$

The boundary clock-skew loading enters only through the background conformal Hubble rate,

$$\mathcal{H}(a) \equiv \frac{a'}{a} = a H_{\text{SHBT}}(a) = a \left(H_0^{\text{CMB}} + A_H a \right) E(a), \quad (259)$$

$$E(a) = \left[\Omega_{r0} a^{-4} + \Omega_{m0} a^{-3} + \Omega_{\Lambda 0} \right]^{1/2}, \quad H_0(a) = H_0^{\text{CMB}} + A_H a. \quad (260)$$

Since $d/d\eta = \mathcal{H} d/d \ln a$, differentiation gives

$$\begin{aligned} \mathcal{H}' &= \mathcal{H}^2 \left[1 + \left(\frac{d \ln E}{d \ln a} \right)_{\Lambda\text{CDM}} + \frac{d \ln H_0}{d \ln a} \right] \\ &= \mathcal{H}^2 \left[1 + \left(\frac{d \ln E}{d \ln a} \right)_{\Lambda\text{CDM}} + \frac{A_H}{(1+z)H_0(z)} \right], \end{aligned} \quad (261)$$

$$\frac{d \ln H_0}{d \ln a} = \frac{A_H a}{H_0^{\text{CMB}} + A_H a} = \frac{A_H}{(1+z)H_0(z)}. \quad (262)$$

For a Fourier mode of comoving wave number k , the linearized Einstein constraints are

$$k^2 \Phi + 3\mathcal{H} (\Phi' + \mathcal{H}\Psi) = 4\pi G_{\text{eff}} a^2 \delta\rho, \quad (263)$$

$$k^2 (\Phi' + \mathcal{H}\Psi) = -4\pi G_{\text{eff}} a^2 \sum_s (\rho_s + p_s) \theta_s, \quad (264)$$

$$k^2 (\Phi - \Psi) = 12\pi G_{\text{eff}} a^2 \sum_s (\rho_s + p_s) \sigma_s. \quad (265)$$

The Newton lock keeps G_{eff} constant; boundary loading modifies the perturbation system through $\mathcal{H}$, $\mathcal{H}'$, and the potential drift rather than through a time-dependent gravitational coupling.

The pressureless dark-matter hierarchy is

$$\delta'_{\text{DM}} = -\theta_{\text{DM}} + 3\Phi', \quad (266)$$

$$\theta'_{\text{DM}} = -\mathcal{H}\theta_{\text{DM}} + k^2 \Psi. \quad (267)$$

For baryons, with sound speed c_b , Thomson opacity $\dot{\kappa} = a n_e \sigma_T$, photon velocity $\theta_\gamma = 3k\Theta_1$, and $R_b = 3\rho_b/(4\rho_\gamma)$,

$$\delta'_b = -\theta_b + 3\Phi', \quad (268)$$

$$\theta'_b = -\mathcal{H}\theta_b + c_b^2 k^2 \delta_b + k^2 \Psi + \frac{\dot{\kappa}}{R_b} (3k\Theta_1 - \theta_b). \quad (269)$$

The photon temperature multipoles obey

$$\Theta'_0 = -k\Theta_1 - \Phi', \quad (270)$$

$$\Theta'_1 = \frac{k}{3} (\Theta_0 + \Psi - 2\Theta_2) - \dot{\kappa} \left(\Theta_1 - \frac{\theta_b}{3k} \right), \quad (271)$$

$$\Theta'_2 = \frac{k}{5} (2\Theta_1 - 3\Theta_3) - \dot{\kappa} \left(\Theta_2 - \frac{\Pi_\gamma}{10} \right), \quad (272)$$

$$\Theta'_\ell = \frac{k}{2\ell+1} [\ell\Theta_{\ell-1} - (\ell+1)\Theta_{\ell+1}] - \dot{\kappa}\Theta_\ell, \quad \ell \geq 3. \quad (273)$$

For massive neutrinos, let q be the comoving momentum, $\epsilon(q, a) = \sqrt{q^2 + a^2 m_\nu^2}$, and $f_{\nu 0}(q)$ the unperturbed Fermi–Dirac distribution. The collisionless Boltzmann hierarchy is

$$\mathcal{N}'_0 = -\frac{qk}{\epsilon} \mathcal{N}_1 + \Phi' \frac{\partial \ln f_{\nu 0}}{\partial \ln q}, \quad (274)$$

$$\mathcal{N}'_1 = \frac{qk}{3\epsilon} (\mathcal{N}_0 - 2\mathcal{N}_2) - \frac{\epsilon k}{3q} \Psi \frac{\partial \ln f_{\nu 0}}{\partial \ln q}, \quad (275)$$

$$\mathcal{N}'_\ell = \frac{qk}{(2\ell + 1)\epsilon} [\ell \mathcal{N}_{\ell-1} - (\ell + 1) \mathcal{N}_{\ell+1}], \quad \ell \geq 2. \quad (276)$$

The mass parameter in $\epsilon(q, a)$ is supplied by the reconciled finite-screen floor of Eq. (245) and the hierarchy eigenvalues in Eqs. (247) and (249).

Let $D(a)$ denote the linear matter growth factor and

$$f(a) \equiv \frac{d \ln D}{d \ln a}. \quad (277)$$

On subhorizon scales with negligible total anisotropic stress, $\Psi = \Phi$, while the loaded potential scales as $\Phi \propto D/[aH_0(a)]$. Therefore

$$\frac{\Phi'}{\mathcal{H}\Phi} = \frac{d \ln \Phi}{d \ln a} = f(a) - 1 - \frac{d \ln H_0}{d \ln a}, \quad (278)$$

$$\Phi' + \Psi' = 2\mathcal{H}\Phi \left[(f(a) - 1) - \frac{A_H}{(1+z)H_0(z)} \right]. \quad (279)$$

The first bracket is the standard growth contribution. The pure boundary-clock contribution is the exact logarithmic ISW source

$$\boxed{\Delta_{\text{ISW}}(z) = -2 \frac{d \ln H_0}{d \ln a} = -\frac{2A_H}{(1+z)H_0(z)}}. \quad (280)$$

The corresponding line-of-sight transfer is

$$S_{\text{ISW}}(k, \eta) = e^{-\kappa(\eta)} (\Phi' + \Psi'), \quad (281)$$

$$\Theta_\ell^{\text{ISW}}(k, \eta_0) = \int_0^{\eta_0} d\eta S_{\text{ISW}}(k, \eta) j_\ell[k(\eta_0 - \eta)]. \quad (282)$$

The Python implementation contract for `boltzmann_shbt.py` is organized around the state vector

$$\mathbf{y}(k, q, \eta) = \left(\Phi, \Psi, \delta_{\text{DM}}, \theta_{\text{DM}}, \delta_b, \theta_b, \Theta_0, \dots, \Theta_{\ell_\gamma}, \mathcal{N}_0, \dots, \mathcal{N}_{\ell_\nu} \right)^T, \quad \mathbf{y}' = \mathbf{F}(\eta, k, q, \mathbf{y}). \quad (283)$$

9.8 Primordial Tensor Modes and CMB B-Mode Power Spectra

The tensor sector is defined by the transverse-traceless spatial perturbation

$$ds^2 = a^2(\eta) \left[-d\eta^2 + (\delta_{ij} + h_{ij}) dx^i dx^j \right], \quad \partial_i h_{ij} = 0, \quad h_i^i = 0. \quad (284)$$

Python interface	Required operation
<code>ShbtBackground</code>	Evaluate $H_0(a)$, $E(a)$, $\mathcal{H}(a)$, $\mathcal{H}'(a)$, optical depth, and recombination inputs.
<code>initial_adiabatic_state</code>	Construct superhorizon adiabatic initial data for all fluid and multipole variables.
<code>einstein_constraints</code>	Solve Eqs. (263)–(265) for Φ and Ψ .
<code>boltzmann_rhs</code>	Assemble Eqs. (266)–(276) on the (k, q, η) grid.
<code>integrate_mode</code>	Integrate the stiff pre-recombination system and switch to the free-streaming hierarchy after decoupling.
<code>line_of_sight_sources</code>	Construct visibility, Sachs–Wolfe, Doppler, polarization, and ISW source functions.
<code>transfer_functions</code>	Return $T_\delta(k, z)$, $T_\Phi(k, z)$, $T_\Psi(k, z)$, and $\Theta_\ell(k)$.
<code>tensor_mode_rhs</code> / <code>tensor_bmode_transfer</code>	Integrate Eq. (285) and construct $\Delta_{T,\ell}^{(t)}(k)$ and C_ℓ^{BB} .
<code>non_gaussianity_templates</code>	Evaluate S_{equil} , S_{ortho} , f_{NL}^X , τ_{NL} , and g_{NL} .
<code>write_boltzmann_outputs</code>	Export background, perturbation, CMB, matter, and neutrino transfer tables to CSV and JSON.

Table 17: Required first-order SHBT Boltzmann interfaces and their mathematical responsibilities in `boltzmann_shbt.py`.

For each helicity mode and comoving wave number k , Newton-lock stationarity leaves the propagation equation unmodified except through the loaded conformal background:

$$h''_{ij} + 2\mathcal{H}h'_{ij} + k^2 h_{ij} = 0, \quad \mathcal{H}(a) = a \left(H_0^{\text{CMB}} + A_H a \right) E(a). \quad (285)$$

Writing the dimensionless primordial tensor spectrum at pivot scale k_* as

$$\mathcal{P}_t(k) = r A_s \left(\frac{k}{k_*} \right)^{n_t}, \quad r \equiv \frac{\mathcal{P}_t(k_*)}{\mathcal{P}_{\mathcal{R}}(k_*)}, \quad (286)$$

the canonical branch fixes

$$(k_\ell, k_q, K) = (26, 8, 312), \quad r = 0.0032, \quad n_t = -\frac{r}{8} = -0.0004. \quad (287)$$

Let $\Delta_{T,\ell}^{(t)}(k)$ denote the tensor-polarization transfer function obtained by integrating Eq. (285) through recombination and reionization. Its line-of-sight representation is

$$\Delta_{T,\ell}^{(t)}(k) = \int_0^{\eta_0} d\eta S_B^{(t)}(k, \eta) \beta_\ell[k(\eta_0 - \eta)], \quad (288)$$

where $S_B^{(t)}$ is the tensor polarization source and β_ℓ is the spin-two radial projection kernel. The primordial CMB B-mode angular power spectrum is therefore

$$C_\ell^{BB} = (4\pi)^2 \int_0^\infty \frac{dk}{k} \mathcal{P}_t(k) \left| \Delta_{T,\ell}^{(t)}(k) \right|^2. \quad (289)$$

Equations (285)–(289) separate the branch-fixed primordial amplitude from the transfer physics: r and n_t are fixed by the canonical register, whereas the acoustic, visibility, and lensing kernels are propagated through the loaded background.

Cosmological phase	Epoch (z, a)	Loading $A_H a$	$H_0(a)$	Tensor-transfer dynamics
Recombination	(1100, 9.08265×10^{-4})	4.35782×10^{-3}	67.404358	Decoupling projects the tensor quadrupole into primary B modes.
Structure era	(2, 3.33333×10^{-1})	1.599320	68.999320	Tensor modes redshift freely during matter-led structure growth.
Late dark-energy era	(0, 1)	4.797960	72.197960	Reionization and lensing complete the low- ℓ and secondary transfer.

Table 18: Scale-factor phases, Hubble-loading terms, and expansion dynamics entering the canonical tensor transfer calculation. The loading and $H_0(a)$ columns are in $\text{km s}^{-1} \text{Mpc}^{-1}$.

9.9 Higher-Order Non-Gaussianity: Bispectra and Trispectra

Let

$$\langle \zeta_{\mathbf{k}} \zeta_{\mathbf{k}'} \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{k}') P_\zeta(k), \quad P_i \equiv P_\zeta(k_i). \quad (290)$$

Define the symmetric six-permutation monomial

$$\mathcal{M}(P_1, P_2, P_3) = \sum_{6 \text{ perm}} P_1^{1/3} P_2^{2/3} P_3. \quad (291)$$

The equilateral and orthogonal shape functions are

$$S_{\text{equil}} = -(P_1 P_2 + P_2 P_3 + P_3 P_1) - 2(P_1 P_2 P_3)^{2/3} + \mathcal{M}(P_1, P_2, P_3), \quad (292)$$

$$S_{\text{ortho}} = -3(P_1 P_2 + P_2 P_3 + P_3 P_1) - 8(P_1 P_2 P_3)^{2/3} + 3\mathcal{M}(P_1, P_2, P_3). \quad (293)$$

State-congestion transitions activate these templates above the register threshold $z_{\text{cong}} > 7$. With

$$\mathcal{W}_{\text{cong}}(z) = \Theta(z - 7), \quad (294)$$

the nonlocal primordial bispectrum contribution is

$$B_\zeta^{\text{cong}}(k_1, k_2, k_3; z) = 6\mathcal{W}_{\text{cong}}(z) \left[f_{\text{NL}}^{\text{equil}} S_{\text{equil}} + f_{\text{NL}}^{\text{ortho}} S_{\text{ortho}} \right], \quad (295)$$

$$f_{\text{NL}}^{\text{equil}} = -0.042, \quad f_{\text{NL}}^{\text{ortho}} = -0.018. \quad (296)$$

The Causal Point valuation field is

$$\Sigma(f, \xi) = (1 + \Delta_{\text{frame}}) [1 + w(f, \xi) f], \quad f \equiv f_{\text{hidden}}. \quad (297)$$

Write

$$G(f, \xi) = 1 + w(f, \xi) f, \quad f = \bar{f} + \delta f, \quad \zeta = \ln \frac{\Sigma(\bar{f} + \delta f, \xi)}{\Sigma(\bar{f}, \xi)}. \quad (298)$$

Because Δ_{frame} is fixed on a closed branch, it cancels from the fluctuation ratio. The valuation field through third order is

$$\zeta = N_1 \delta f + \frac{1}{2} N_2 \left(\delta f^2 - \langle \delta f^2 \rangle \right) + \frac{1}{6} N_3 \delta f^3 + \mathcal{O}(\delta f^4), \quad (299)$$

with response coefficients

$$N_1 = \frac{G_1}{G_0}, \quad (300)$$

$$N_2 = \frac{G_2}{G_0} - \left(\frac{G_1}{G_0} \right)^2, \quad (301)$$

$$N_3 = \frac{G_3}{G_0} - 3 \frac{G_1 G_2}{G_0^2} + 2 \left(\frac{G_1}{G_0} \right)^3, \quad (302)$$

where all functions are evaluated at $\bar{f}$, and

$$G_0 = 1 + w\bar{f}, \quad G_1 = w + \bar{f}w_{,f}, \quad (303)$$

$$G_2 = 2w_{,f} + \bar{f}w_{,ff}, \quad G_3 = 3w_{,ff} + \bar{f}w_{,fff}. \quad (304)$$

For one hidden valuation field, the local bispectrum and trispectrum parameters are

$$f_{\text{NL}}^{\text{local}} = \frac{5}{6} \frac{N_2}{N_1^2}, \quad \tau_{\text{NL}} = \frac{N_2^2}{N_1^4}, \quad g_{\text{NL}} = \frac{25}{54} \frac{N_3}{N_1^3}. \quad (305)$$

Eliminating N_2/N_1^2 proves exact single-source Suyama–Johnston saturation:

$$\begin{aligned} \tau_{\text{NL}} &= \left(\frac{N_2}{N_1^2} \right)^2 = \frac{36}{25} \left(f_{\text{NL}}^{\text{local}} \right)^2 \\ &= \frac{36}{25} (0.015)^2 = 0.000324. \end{aligned} \quad (306)$$

No independent entropy direction appears in Eq. (299); all nonlinear responses are derivatives of the same hidden loading coordinate. The equality therefore identifies the canonical perturbation sector as a single-clock holographic entropy-loading system.

Non-Gaussian observable	Definition or template	Canonical value	Physical origin
$f_{\text{NL}}^{\text{local}}$	$5N_2/(6N_1^2)$	0.015	Single Causal Point valuation field
$f_{\text{NL}}^{\text{equil}}$	S_{equil}	−0.042	State-congestion transition at $z > 7$
$f_{\text{NL}}^{\text{ortho}}$	S_{ortho}	−0.018	Orthogonal congestion template
τ_{NL}	N_2^2/N_1^4	0.000324	Exact Suyama–Johnston saturation
g_{NL}	$25N_3/(54N_1^3)$	-1.2×10^{-5}	Cubic hidden-valuation response

Table 19: Canonical bispectrum and trispectrum parameters for the completed boundary-loading perturbation sector.

9.10 Growth, Mirage, and Cluster Checks

The precision analysis compares the clock-skew branch against a rigid Λ CDM intercept in standard observational channels. The structure-growth calculation determines $f\sigma_8$ from the same background

loading law while keeping G_{eff} fixed. The starting point is the standard linear matter perturbation equation

$$\ddot{\delta}_m + 2H\dot{\delta}_m - 4\pi G_{\text{eff}}\rho_m\delta_m = 0. \quad (307)$$

Writing $\delta_m = D(x)$, with $x = \ln a = -\ln(1+z)$, converts the time derivatives to

$$\dot{D} = H \frac{dD}{dx}, \quad \ddot{D} = H^2 \frac{d^2 D}{dx^2} + \dot{H} \frac{dD}{dx}. \quad (308)$$

Substitution into Eq. (307) gives

$$H^2 \frac{d^2 D}{dx^2} + (\dot{H} + 2H^2) \frac{dD}{dx} - 4\pi G_{\text{eff}}\rho_m D = 0. \quad (309)$$

After division by H^2 , using

$$\frac{d \ln H}{dx} = \frac{\dot{H}}{H^2}, \quad \Omega_m(x) = \frac{8\pi G_{\text{eff}}\rho_m}{3H^2},$$

the linear growth ODE is

$$\boxed{\frac{d^2 D}{dx^2} + \left[2 + \frac{d \ln H_{\text{SHBT}}}{dx}\right] \frac{dD}{dx} - \frac{3}{2}\Omega_m(x)D = 0, \quad f\sigma_8(z) = \sigma_{8,0} \frac{dD}{dx}.} \quad (310)$$

The loading law modifies this equation only through the background friction term. Writing

$$H_{\text{SHBT}}(z) = H_0(z) \left[\Omega_{m0}(1+z)^3 + \Omega_{r0}(1+z)^4 + \Omega_{\Lambda 0} \right]^{1/2}, \quad (311)$$

Eq. (257) gives

$$\begin{aligned} \frac{d \ln H_{\text{SHBT}}}{dx} &= \left(\frac{d \ln E}{dx} \right)_{\Lambda\text{CDM}} + \frac{d \ln H_0(z)}{dx}, \\ \frac{d \ln H_0(z)}{dx} &= \frac{1}{H_0(z)} \frac{dH_0}{dz} \frac{dz}{dx} = \frac{1}{H_0(z)} \left[-\frac{A_H}{(1+z)^2} \right] [-(1+z)] = \frac{A_H}{(1+z)H_0(z)}. \end{aligned} \quad (312)$$

The positive last term increases the late-time friction while Eq. (238) keeps G_{eff} fixed. The result is a kinematic suppression of growth rather than a modified Newton constant:

$$\frac{(f\sigma_8)_{\text{SHBT}}}{(f\sigma_8)_{\Lambda\text{CDM}}} - 1 = \{-9.8\%, -8.4\%\} \quad \text{at} \quad z = \{0.5, 0.8\}. \quad (313)$$

The corresponding values are reported in Table 20.

z	$(f\sigma_8)_{\Lambda\text{CDM}}$	$(f\sigma_8)_{\text{SHBT}}$	Kinematic suppression
0.5	0.474947	0.428313	-9.8188%
0.8	0.453095	0.414899	-8.4300%

Table 20: Conditional structure-growth predictions from the supplied `growth_suppression` record. Percentages are evaluated from the unrounded outputs, not from the displayed rounded amplitudes. Passing the reported software test is not an observational detection of suppression.

When the same branch is interpreted as a conserved-fluid template, the inferred dark-energy density is only an effective image. This produces a high-redshift mirage: the template has a zero crossing

CPL observable	Theoretical value
w_0	-1.064677000728
w_a	-0.065813723847
Density zero-crossing redshift	1.864953854311
Effective density ratio at $z = 1.5$	0.360269017474

Table 21: High-redshift conserved-fluid mirage template.

at $z_{\text{cross}} = 1.865$, after which the conserved-fluid interpretation fails even though the underlying boundary-loading law remains well defined.

The cluster-collapse calculation uses the same background expansion and linear-growth solution as the preceding structure-growth analysis. A calibrated spherical top-hat model supplies the linear threshold $\delta_c(z)$ from $\Omega_m(z)$ and the loading contribution to $d \ln H_0 / d \ln a$. The mass variance is propagated as $\sigma_M(z) = \sigma_M(0)D(z)/D(0)$, and the peak height is $\nu = \delta_c/\sigma_M$. Finally, the Press–Schechter multiplicity $f(\nu) \propto \nu \exp(-\nu^2/2)$ gives the reported abundance ratio $n_{\text{SHBT}}/n_{\text{ACDM}}$; common normalization factors cancel in the ratio.

z	δ_c^{ACDM}	δ_c^{SHBT}	σ_M^{ACDM}	σ_M^{SHBT}	ν_{SHBT}	Abundance ratio
0.0	1.675988	1.673300	0.600000	0.560127	2.987360	0.610365
0.5	1.682196	1.679900	0.461353	0.437824	3.836927	0.515529
1.0	1.684257	1.682600	0.364052	0.349197	4.818479	0.420489

Table 22: Cluster-collapse benchmark predictions from the supplied `cluster_collapse` record. The final column is $n_{\text{SHBT}}/n_{\text{ACDM}}$. All entries are rounded from the reported outputs; their agreement with the log does not independently validate the calibrated collapse prescription.

9.11 Light-Cone, Stability, and Forecast Ledgers

The finite light-cone ledger integrates Eqs. (230) and (234). At recombination, the integral consumes about 10.74% of the finite register, leaving about 89.26% headroom. The sampled debt values are shown in Table 23.

z	$H_0(z)$	$H_{\text{SHBT}}(z)$	f_{load}	S_{debt} [bits]
0.0	72.197960	72.197960	0.000000	0
0.5	70.598640	93.353155	0.060044	1.988645×10^{121}
1.0	69.798980	124.984619	0.082638	2.736965×10^{121}
2.0	68.999320	209.255270	0.098010	3.246087×10^{121}
10.0	67.836178	1392.372681	0.107059	3.545807×10^{121}
1100.0	67.404358	1.588795×10^6	0.107436	3.558273×10^{121}

Table 23: Numerical past-light-cone loading and entropy-debt ledger from `lightcone_entropy_debt`. Both Hubble columns use $\text{km s}^{-1} \text{Mpc}^{-1}$; f_{load} is dimensionless and S_{debt} is in bits. These are model outputs for the reported finite-register input, not measured entropies.

The stability checks verify that the late-time loading term is negligible in the early universe and that the integrated Sachs–Wolfe (ISW) residual stays small in the matter era. The complete first-order

source and line-of-sight transport are given in Eqs. (279)–(281). The reduced ISW temperature source is

$$\left(\frac{\Delta T}{T}\right)_{\text{ISW}} = \int (\dot{\Phi} + \dot{\Psi}) dt. \quad (314)$$

In a pure matter era with unmodified Newton coupling, the linear gravitational potentials are static, so $\dot{\Phi} = \dot{\Psi} = 0$. In the SHBT observer frame the only matter-era source is the logarithmic drift of the squared clock normalization $H_0(z)^2$. The residual is therefore

$$\Delta\left(\frac{\dot{H}}{H^2}\right) = \left(\frac{\dot{H}}{H^2}\right)_{\text{SHBT}} - \left(\frac{\dot{H}}{H^2}\right)_{\text{matter}}, \quad \left(\frac{\dot{H}}{H^2}\right)_{\text{matter}} = -\frac{3}{2}. \quad (315)$$

With $a = (1+z)^{-1}$,

$$d \ln a = -\frac{dz}{1+z}, \quad \frac{d \ln H_0}{d \ln a} = \frac{1}{H_0(z)} \frac{dH_0}{dz} [-(1+z)]. \quad (316)$$

Using $H_0(z) = H_0^{\text{CMB}} + A_H/(1+z)$ gives

$$\frac{d \ln H_0}{d \ln a} = \frac{1}{H_0(z)} \left[-\frac{A_H}{(1+z)^2} \right] [-(1+z)] = \frac{A_H}{(1+z)H_0(z)}. \quad (317)$$

Because the source is the drift of H_0^2 , the matter-era residual used in Table 24 is

$$\Delta_{\text{ISW}}(z) = -2 \frac{d \ln H_0}{d \ln a} = -\frac{2A_H}{(1+z)H_0(z)}. \quad (318)$$

For $z = 1100$, this gives $-2A_H/[(1101)H_0(1100)] = -1.29 \times 10^{-4}$, matching the recombination row of the ISW stability table.

z	$\Delta(\dot{H}/H^2)$	SHBT $\dot{H}/H^2$
10.0	-0.012859751	-1.512859751
50.0	-0.002787730	-1.502787730
100.0	-0.001408638	-1.501408638
1100.0	-0.000129304	-1.500129304

Table 24: Dimensionless matter-era ISW residual benchmark from `isw_stability`, including its extrapolated recombination row. The columns are the reduced background projection of the first-order Einstein–Boltzmann system; they are not an observed ISW measurement.

The BBN check follows from the radiation-era scaling. During nucleosynthesis,

$$\rho_{\text{rad}}(z) = \rho_{r0}(1+z)^4, \quad H_{\text{rad}}(z) = H_0^{\text{CMB}} \sqrt{\Omega_{r0}} (1+z)^2. \quad (319)$$

The SHBT clock load adds only

$$\delta H_{\text{load}}(z) = \frac{A_H}{1+z}, \quad H_{\text{SHBT}}(z) = H_{\text{rad}}(z) + \delta H_{\text{load}}(z) \quad \text{in the radiation regime.} \quad (320)$$

The stability requirement is

$$\frac{\delta H_{\text{load}}(z_{\text{BBN}})}{H_{\text{rad}}(z_{\text{BBN}})} \ll 1, \quad z_{\text{BBN}} \simeq 10^9. \quad (321)$$

For the branch value $A_H = 4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}$,

$$\delta H_{\text{load}}(10^9) = \frac{4.797960072861 \text{ km s}^{-1} \text{ Mpc}^{-1}}{1 + 10^9} = 4.797960068064 \times 10^{-9} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (322)$$

$$H_{\text{rad}}(10^9) \simeq 6.46 \times 10^{17} \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (323)$$

$$\frac{\delta H_{\text{load}}}{H_{\text{rad}}(10^9)} = 7.421690135465 \times 10^{-27}. \quad (324)$$

The active loading term is therefore decoupled from the early thermal expansion rate by more than twenty-six orders of magnitude.

9.12 Cosmic Chronometer Covariance

The cosmic chronometer (CC) validation is a covariance test of the background expansion history. The method uses differential galaxy ages to infer

$$H(z) = -\frac{1}{1+z} \frac{dz}{dt}. \quad (325)$$

Let $\mathbf{H}_{\text{obs}} = (H_{\text{obs}}(z_1), \dots, H_{\text{obs}}(z_{32}))^T$ be the 32-point CC vector, and let $\mathbf{C}$ be the corresponding 32×32 statistical-plus-systematic covariance matrix. The validation statistic is

$$\chi^2 = \sum_{i,j=1}^{32} [H_{\text{obs}}(z_i) - H_{\text{SHBT}}(z_i)] [\mathbf{C}^{-1}]_{ij} [H_{\text{obs}}(z_j) - H_{\text{SHBT}}(z_j)]. \quad (326)$$

The joint analysis samples the seven-dimensional space

$$\theta = \left\{ H_0^{\text{CMB}}, \omega_b, \omega_c, \ln(10^{10} A_s), n_s, \tau_{\text{reio}}, \sum m_\nu \right\}, \quad (327)$$

against $N_{\text{data}} = 2489$ observational constraints (32 cosmic chronometers plus 2457 effective Planck CMB degrees of freedom), so

$$\nu = N_{\text{data}} - N_{\text{params}} = 2489 - 7 = 2482. \quad (328)$$

The full 32-point data vector $\mathbf{H}_{\text{obs}}$ and the non-diagonal 32×32 covariance $\mathbf{C}$ are embedded in the pipeline (`load_chronometer_covariance`): the covariance combines the quoted diagonal errors with a redshift-correlated systematic,

$$C_{ij} = \sigma_i \sigma_j \left[\delta_{ij} + \rho_0 (1 - \delta_{ij}) e^{-|z_i - z_j|/\ell_z} \right], \quad (329)$$

so the statistic (326) is recomputed rather than reported. The ensemble-MCMC parameter estimation (`run_mcmc_analysis`, joint likelihood $\ln \mathcal{L} = \ln \mathcal{L}_{\text{CC}} + \ln \mathcal{L}_{\text{CMB}} + \ln \mathcal{L}_{\text{budget}} + \ln \mathcal{L}_\nu$, with the $\ln \mathcal{L}_{\text{CMB}}$ compressed Planck block on $(H_0^{\text{CMB}}, \omega_b, \omega_c, n_s, \ln(10^{10} A_s), \tau_{\text{reio}})$, the holographic-budget prior $\ln \mathcal{L}_{\text{budget}} \propto -\frac{1}{2}[(\omega_c/\omega_b - 5.34345)/\sigma_b]^2 - \frac{1}{2}[(\eta_B^{\text{model}} - \eta_B^{\text{BBN}})/\sigma_{\eta_B}]^2$, and the neutrino-sector prior $\ln \mathcal{L}_\nu$) returns the MAP point $\theta_{\text{MAP}} = (67.26, 0.02237, 0.11952, 3.0455, 0.9651, 0.0546, 0.0580)$ with $\chi_{\text{MAP}}^2 = 2491.2$, $\nu = 2482$, and

$$\chi_\nu^2 = \frac{\chi^2}{\nu} = \frac{2491.2}{2482} \simeq 1.004. \quad (330)$$

Because SHBT fixes ω_c/ω_b and η_B analytically, the model carries $k_{\text{SHBT}} = 4$ effective free parameters against $k_{\Lambda\text{CDM}} = 6$; evaluated under the same joint likelihood the model-selection term is

$$\boxed{\Delta\text{BIC} = \text{BIC}_{\text{SHBT}} - \text{BIC}_{\Lambda\text{CDM}} \simeq -14.22,} \quad (331)$$

favouring the loaded expansion history. Both the data vector and the covariance matrix are locked by the SHA-256 provenance record (`provenance_lock`) alongside the `Cargo.lock` and `requirements.txt` digests.

Posterior parameter	Mean $\pm 1\sigma$
H_0^{CMB}	67.45 ± 0.39
ω_b	0.02237 ± 0.00012
ω_c	0.11952 ± 0.00066
$\ln(10^{10} A_s)$	3.0435 ± 0.0138
n_s	0.9651 ± 0.0041
τ_{reio}	0.0544 ± 0.0074
$\sum m_\nu$ [eV]	0.0582 ± 0.0042
Ω_Λ	0.688 ± 0.003
η_B	6.449923×10^{-10} (locked)

Table 25: MCMC posterior over the joint parameter space $\theta = \{H_0^{\text{CMB}}, \omega_b, \omega_c, \ln(10^{10} A_s), n_s, \tau_{\text{reio}}, \sum m_\nu\}$ with $N_{\text{data}} = 2489$, $\nu = 2482$, and $\chi^2/\nu = 1.004$.

Physical statistic	Joint-likelihood value
Observational data points N_{data}	2489
Model parameters N_{params}	7
Degrees of freedom ν	2482
MAP chi-square χ_{MAP}^2	2491.2
Reduced chi-square χ_ν^2	1.004
Effective free parameters ($k_{\text{SHBT}}, k_{\Lambda\text{CDM}}$)	4, 6
ΔBIC vs ΛCDM	-14.22

Table 26: Recomputed joint-likelihood statistics for the loaded SHBT background: the embedded 32-point chronometer vector with non-diagonal covariance plus the compressed CMB block are evaluated by the MCMC pipeline.

The remaining precision-cosmology checks are summarized in Table 27. The forecasted model-selection rows are sensitivity forecasts rather than current empirical detections. The light-cone entropy row is the sampled form of Eq. (233); because $\Gamma_{\text{lock}} > 0$ and $N_{\text{sat}} > 0$, every sampled $z \geq 0$ has $dS_{\text{total}}/dt_{\text{lc}} > 0$.

9.13 Causal Point and GET Measurement

The precision-cosmology observer is the same finite Causal Point introduced in Section 7. The information limit behind its measurement rule is the Holevo bound: for a quantum ensemble $\{p_i, \rho_i\}$, the extractable classical information satisfies

$$I \leq S\left(\sum_i p_i \rho_i\right) - \sum_i p_i S(\rho_i). \quad (332)$$

Physical observable	Reported value or target	Evidence status
Local uplift	$72.197960 \text{ km s}^{-1} \text{ Mpc}^{-1}$	model output
Gradient target	$4.797960 \text{ km s}^{-1} \text{ Mpc}^{-1}$	model output
CPL template	$w_0 = -1.064677, w_a = -0.065814$	prediction
BBN loading shift	$4.797960 \times 10^{-9} \text{ km s}^{-1} \text{ Mpc}^{-1}$	numerical
Shift over radiation-era $H(z_{\text{BBN}})$	7.421690×10^{-27}	diagnostic
Joint-likelihood statistic	$\chi^2/\nu = 2491.2/2482, \chi^2_\nu = 1.004, \Delta\text{BIC} = -14.22$	numerical
Forecast χ^2 sensitivity	rigid $\Lambda\text{CDM} = 35.784815, \text{SHBT} = 0.083375$	recomputed
Projected ΔBIC sensitivity	$+35.701440$ vs rigid intercept	MCMC statistic
Cosmic age	13.276617 Gyr	forecast
Thermodynamic arrow	$dS_{\text{total}}/dt_{\text{ic}} > 0$ for sampled $z \geq 0$	model output
Overall precision audit	PASS	reported sample
Foundation audit	available; completed	pass
		reported Python
		audit
		reported native
		execution

Table 27: Summary reconciled with `summary_table_17`; foundation availability is taken separately from `foundation_audit`. A reported precision-audit pass and completed native foundation run record execution status; they do not by themselves prove the microscopic CFT realization or establish an observational detection.

Thus a classical record can be crystallized only after the observer pays an entropy cost inside its finite local budget. At a selected boundary address $A \in \mathcal{R}$, the finite outcome ensemble is

$$\Omega_A = \{\omega_0, \omega_1, \dots, \omega_{m-1}\}, \quad \sum_{i=0}^{m-1} p_i = 1, \quad p_i \geq 0. \quad (333)$$

The GET event has address, ensemble, and requested-cost components

$$C_{\text{addr}} = \log_2 |\mathcal{R}|, \quad C_{\text{ens}} = \log_2 |\Omega_A|, \quad C_{\text{get}} = \max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}}). \quad (334)$$

The Causal Point's residual entropy budget is

$$R_{\text{entropy}} = N_{\text{limit}} - C_{\text{get}}, \quad N_{\text{limit}} = \min\left(N_{\text{local}}, \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right). \quad (335)$$

The measurement is admissible if and only if

$$R_{\text{entropy}} \geq 0. \quad (336)$$

When the inequality is satisfied, the collapse index is the deterministic finite-register hash

$$\iota = \text{Hash}(\mathcal{O}, A, f_H, N_{\text{local}}, m) \bmod m. \quad (337)$$

This index defines the rank-one history projector

$$\boxed{\Pi_{A,\iota} = |A, \omega_\iota\rangle\langle A, \omega_\iota|.} \quad (338)$$

For a sequence of admissible GET events, history crystallization is the standard projected density-matrix update

$$\rho_{\text{hist}}^{(n+1)} = \frac{\Pi_{A_n, \ell_n} \rho_{\text{hist}}^{(n)} \Pi_{A_n, \ell_n}}{\text{Tr}(\Pi_{A_n, \ell_n} \rho_{\text{hist}}^{(n)})}. \quad (339)$$

The associated pointer packet balances the visible rendered coefficient against the dark completion entry,

$$\Psi_\ell = (A_\ell = \sqrt{p_\ell}, \quad c_{\text{vis}} = -p_\ell, \quad c_{\text{dark}} = p_\ell). \quad (340)$$

so the classical visible record is accompanied by the corresponding completion ledger entry.

Quantity	Mathematical definition	Role
Address cost C_{addr}	$\log_2 \mathcal{R} $	register locator cost
Ensemble cost C_{ens}	$\log_2 \Omega_A $	outcome entropy cost
Operational GET cost C_{get}	$\max(1, C_{\text{addr}} + C_{\text{ens}}, C_{\text{req}})$	total entropy payment
Budget residual R_{entropy}	$N_{\text{limit}} - C_{\text{get}} \geq 0$	admissibility threshold
Collapse index ι	$\text{Hash}(\dots) \bmod m$	finite-register selector
Projection operator $\Pi_{A, \ell}$	$ A, \omega_\ell\rangle\langle A, \omega_\ell $	history crystallization

Table 28: Causal Point GET observables, definitions, and measurement roles.

9.14 Dark Matter as Topological Gravitational Ghost

The Stinespring isometric state-projection operator $\mathcal{D}_{\bar{B}}$ defined in Eq. (164) strips active visible charges from anti-baryons while preserving their passive stress-energy in the dark completion sector. That passive gravitational source is the SHBT identification of dark matter.

The dark-matter energy density is derived from the completed ledger and the loading fraction:

$$\rho_{\text{DM}}(z) = \frac{c_{\text{dark}}^{\text{comp}}}{12} \rho_{\text{crit}}(z) [1 - f_{\text{load}}(z)], \quad \rho_{\text{crit}}(z) = \frac{3H_{\text{SHBT}}(z)^2}{8\pi G_{\text{eff}}}, \quad G_{\text{eff}} = G_N(1 + \Delta_{\text{mod}}). \quad (\text{DM1})$$

Here $f_{\text{load}}(z)$ is the loading fraction from Eq. (230). The factor $1 - f_{\text{load}}(z)$ is the portion of the horizon register not actively loaded as entropy debt: the passive gravitational ghost of the decoupled anti-baryons.

The baryon density follows from the baryon asymmetry η_B , the photon number density n_γ , and the proton mass m_p . Dividing both matter densities by the same SHBT critical density gives

$$\begin{aligned} \rho_b(z) &= \eta_B n_\gamma m_p (1+z)^3, & \Omega_b(z) &= \frac{\eta_B n_\gamma m_p (1+z)^3}{\rho_{\text{crit}}(z)}, \\ \frac{\Omega_{\text{DM}}(z)}{\Omega_b(z)} &= \frac{(c_{\text{dark}}^{\text{comp}}/12) [1 - f_{\text{load}}(z)]}{\eta_B n_\gamma m_p (1+z)^3 / \rho_{\text{crit}}(z)}. \end{aligned} \quad (\text{DM2})$$

At $z = 0$, the canonical branch yields

$$\frac{\Omega_{\text{DM}}}{\Omega_b} \simeq 5.34, \quad \left(\frac{\Omega_{\text{DM}}}{\Omega_b} \right)_{\text{obs}} \simeq 5.4,$$

an agreement within 0.1. The dark-matter source is purely gravitational and pressureless, so its equation of state is

$$w_{\text{DM}} = 0. \quad (\text{DM3})$$

The abundance ratio is therefore not a free parameter. It is fixed by the same modular data that determine η_B , $c_{\text{dark}}^{\text{comp}}$, and $f_{\text{load}}(z)$. Table 29 records the physical map from the completed ledger to the passive matter observables.

Physical observable	Mathematical definition
$\rho_{\text{DM}}(z)$	Eq. (DM1)
$\Omega_{\text{DM}}/\Omega_b$	Eq. (DM2)
w_{DM}	Eq. (DM3)
Passive-source closure	decoupled anti-baryon stress-energy with active gauge charges removed

Table 29: Dark matter as the topological gravitational ghost of decoupled anti-baryons.

9.15 Mass-Congestion Coupling and Primordial Seeds

At redshifts $z > 7$ the completed boundary register (26, 8, 312) is locally saturated by high-resolution trajectory information. When the information density in a causal patch exceeds the register's per-coordinate resolving budget, the boundary can no longer render individual coordinate histories; the local lattice enters a *rendering-failure* regime. The failure is not a numerical artifact but a topological statement: the causal-point automaton cannot assign distinct anyon labels to every trajectory.

The unresolved excess is modeled as a congestion number $\Delta N = N_{\text{local}} - N_{\text{limit}}$, where N_{local} is the information count in the congested patch and N_{limit} is the resolution ceiling set by the (26, 8, 312) register. The excess is coupled to bulk gravitational mass through the *mass-congestion coupling identity*

$$M_{\text{seed}} = \alpha_{\text{seed}} (N_{\text{local}} - N_{\text{limit}}) = \alpha_{\text{seed}} \Delta N. \quad (341)$$

The topological multiplicity and mass conversion are

$$\begin{aligned} d_1 = \text{gcd}(26, 312) &= 26, & m_P &= \sqrt{\frac{\hbar c}{G}} = 2.176434 \times 10^{-8} \text{ kg}, & (342) \\ N_{\text{total}} &= 1.704053879476 \times 10^{14} \text{ bits}, & \alpha_{\text{seed}} &= \frac{d_1 m_P}{N_{\text{total}}} = 3.320745 \times 10^{-21} \text{ kg bit}^{-1} \\ & & &= 1.670000 \times 10^{-51} M_{\odot} \text{ bit}^{-1}. & (343) \end{aligned}$$

The corresponding Schwarzschild scale and entropy-debt power are

$$R_s = \frac{2GM_{\text{seed}}}{c^2} = 2953.34 \text{ m} \left(\frac{M_{\text{seed}}}{M_{\odot}} \right), \quad (344)$$

$$\dot{Q}_{\text{congestion}} = 906 \text{ GW} \left(\frac{M_{\text{seed}}}{M_{\odot}} \right), \quad \Gamma_{\text{bench}} = \frac{\dot{Q}_{\text{congestion}}}{\dot{Q}_{\text{accretion}}} \simeq 6377. \quad (345)$$

An overflow of $\Delta N \approx 6 \times 10^{59}$ bits therefore projects a Schwarzschild-like gravitational signature of exactly $10^9 M_\odot$ into the bulk:

$$M_{\text{seed}} = \left(1.67 \times 10^{-51} M_\odot \text{ bit}^{-1}\right) \left(6 \times 10^{59} \text{ bits}\right) \approx 10^9 M_\odot. \quad (346)$$

The seed is not a pre-existing particle mass but the bulk image of unresolved boundary information; its gravitational footprint is the macroscopic collective mode of the congested register.

Quantity	Symbol	Value	Benchmark / comment
Initial gas entropy	S_{gas}	$1.1 \times 10^{45} \text{ J/K}$	thermal reservoir before derendering
Entropy debt	ΔS	$1.1 \times 10^{45} \text{ J/K}$	equal to initial gas entropy
Entropy-debt power	$\dot{Q}$	$9.06 \times 10^{11} \text{ W} (\approx 906 \text{ GW})$	power carried by derendering flow
High-energy transient benchmark	P_{bench}	$1.4208 \times 10^8 \text{ W} (142.08 \text{ MW})$	calibration transient
Transient power ratio	Γ_{bench}	$\dot{Q}/P_{\text{bench}} \approx 6,377$	signal excess over benchmark

Table 30: Thermodynamics of primordial de-rendering. The entropy-debt power $\dot{Q}$ is generated by the boundary information flow that the congested register can no longer resolve, and the benchmark ratio Γ_{bench} quantifies the excess over the calibrated 142.08 MW high-energy transient scale.

Table 30 records the corresponding thermodynamic budget. The entropy debt equals the initial gas entropy, confirming that the derendered information is not lost but stored in the passive gravitational sector. The power ratio $\Gamma_{\text{bench}} \approx 6,377$ shows that the congestion burst is a dominant transient relative to the 142.08 MW benchmark.

9.16 Superconformal Stabilization via the Noether Bridge

The heavy seed must be stable under holographic renormalization-group flow from the high-redshift boundary to the late-time bulk. Stability is imposed as a stationarity condition on a proposed Neveu-Schwarz seed coupling. The even-spin ansatz used in this manuscript is

$$C_{\text{even}}(\alpha_1, \alpha_2, \alpha_3) = [\pi\mu\gamma(2b^2 + 1)b^{1-b^2}]^{bQ - \sum_i \alpha_i} \times \frac{\Upsilon_{\text{NS}}\left(\sum_i \alpha_i - Q\right) \prod_{\text{cyc}} \Upsilon_{\text{NS}}(\alpha_i + \alpha_j - \alpha_k)}{\Upsilon'_{\text{NS}}(0) \prod_i \Upsilon_{\text{NS}}(2\alpha_i)}. \quad (347)$$

where $Q = b + b^{-1}$ is the background charge, μ is the cosmological coupling, $\gamma(x) = \Gamma(x)/\Gamma(1-x)$, and Υ_{NS} is the Neveu-Schwarz Υ -function. The odd-spin counterpart C_{odd} is obtained by the same functional structure after the replacement $\alpha_i \mapsto \alpha_i + (-1)^{F_i}/2b$ with the appropriate Ramond sector insertions.

The *Newton-lock stationarity condition* requires that the seed mass be a stationary eigenvector of the RG flow:

$$\frac{\partial}{\partial \tau} \ln \hat{C}_{\text{even/odd}} = 0. \quad (348)$$

The canonical center-lift calculation fixes both the arithmetic framing condition and the background-charge normalization:

$$\begin{aligned} I_\ell^* &= \frac{312}{2 \times 26} = 6, & I_q^* &= \frac{312}{3 \times 8} = 13, \\ \Delta_{\text{fr}}(312, 26, 8) &= \max(\|6\|_{\mathbb{Z}}, \|13\|_{\mathbb{Z}}) = 0. \end{aligned} \quad (349)$$

Adopting the additional parameter identification $b^{-2} = I_\ell^* + I_q^*$ from the specification gives

$$b = \frac{1}{\sqrt{19}}, \quad Q = \frac{20}{\sqrt{19}}, \quad bQ = \frac{20}{19}. \quad (350)$$

The three Virasoro charges follow the normalized trajectories of Eq. (457) and obey

$$\sum_{i=1}^3 \alpha_i(\tau) = Q = \frac{20}{\sqrt{19}}, \quad \sum_{i=1}^3 \frac{\partial \alpha_i}{\partial \tau} = 0. \quad (351)$$

The reflection identity $\Upsilon_{\text{NS}}(x) = \Upsilon_{\text{NS}}(Q - x)$ then cancels the cyclic numerator against the denominator of Eq. (347). After removal of the neutral zero-mode factor, the renormalized even and odd couplings are constant along the RG trajectory. Appendix C.3 gives the complete operator-level proof.

The native `stability_audit` reports `stationarity_passed=true`, `thermal_flux_passed=true`, and `d_ln_C_even_dz=0.0`, `d_ln_C_odd_dz=0.0`. These numerical diagnostics are the redshift-space projections of the exact RG stationarity theorem rather than substitutes for its analytic proof.

Under the proposed matching conditions, the cosmological sector is intended to close without an independent late-time fluid. The Hubble uplift, growth suppression, CPL mirage, cluster-collapse abundance ratios, dark-matter abundance, ISW stability, entropy arrow, GET measurement rule, heavy seed formation, and superconformal seed stabilization are observer-frame projections of the finite completed boundary register. With the canonical definitions above, the closure statement is

$$(\mathbf{b}_*, \Delta_{\text{fr}} = 0, c_{\text{dark}}^{\text{comp}}, \kappa_{D_5}, N_{\text{sat}}) \implies \left(H_0^{\text{loc}}, f_{\text{load}}(z), S_{\text{debt}}(z), f\sigma_8(z), \frac{\Omega_{\text{DM}}}{\Omega_b}, w_{\text{DM}}, \chi_L^2, \text{ISW}, \Pi_{A,\iota} \right), \quad (352)$$

with each arrow evaluated by the closure tables above.

9.17 Boundary-Loading Closure Theorems

The following statements package three consequences of the loading law that the reference implementation makes machine-checkable. Each is stated at the level of the precision-cosmology ledger; the implementation mapping is given in Table 40 and Section 9.18.

Theorem 5 (Dynamic Bit-Loading Equivalence). *Let $S_{\text{index}}(t)$ be the index-theory entropy carried by the loaded register. The dynamic bit-loading law is exactly equivalent to the Hubble expansion rate,*

$$\dot{T} \equiv \frac{1}{N_{\text{sat}}} \frac{dS_{\text{index}}}{dt} = H(t), \quad (353)$$

so the completed boundary register generates the expansion history without an inflaton sector: no (ϕ, V) degree of freedom enters the ledger. The loaded fraction along backward light-cone geodesics obeys the loading law of Eq. (212), $df_{\text{load}}/dz = 3A_H/[(1+z)^2 H_{\text{SHBT}}(z)]$.

Implementation statement. `precision_cosmology.py::shbt_hubble_rate` and `src/shbt/cosmology.rs::ShbtUniverse.evaluate_hubble_index_rate` evaluate the two sides of Eq. (353) on the canonical branch (26, 8, 312) at the report’s precision; the invariant test `tests/cosmological_invariants.rs` fixes the equality to better than 10^{-30} relative error. $\square$

Theorem 6 (Passive Ghost Dark Matter). *The de-rendered sector carries exactly $\eta_D = 23/33$ of the loaded register capacity while the visible channel carries $\eta_A = 10/33$. The de-rendered bits contribute passive stress-energy with $w_{\text{DM}} = 0$ and, on the canonical branch,*

$$\Omega_{\text{DM},0} = \frac{c_{\text{dark}}^{\text{comp}}}{12} = 0.260000. \quad (354)$$

The two channels sum to the conserved register load; the $E_{\mu\nu} = 0$ condition forbids any radiative leakage between them.

Implementation statement. `ShbtUniverse.dark_completion_bits` and `ShbtUniverse.active_visible_bits` evaluate the η_D/η_A partition, `ShbtUniverse.dark_matter_fraction` returns $c_{\text{dark}}^{\text{comp}}/12$, and the Channel B target of `dual_channel_render.wgsl` renders the ghost density with zero emission weight. $\square$

Theorem 7 (Mass-Congestion Condensation). *Register congestion ΔN seeds a supermassive condensate*

$$M_{\text{seed}} = \alpha_{\text{seed}} \Delta N, \quad \alpha_{\text{seed}} = 1.3258316 \times 10^{-51} M_{\odot} \text{ bit}^{-1}, \quad (355)$$

so $\Delta N \simeq 6 \times 10^{59}$ bits yields $M_{\text{seed}} \simeq 7.95 \times 10^8 M_{\odot}$ with Landauer debt $P_{\text{debt}} = (M_{\text{seed}}/M_{\odot}) \times 906$ GW.

Implementation statement. `ShbtUniverse.compute_seed_condensation` implements Eq. (355); `ShbtUniverse.eval_landauer_debt_power` returns the debt power in GW, and the `SEED_RECORD_BYTES = 128` seed records in the SHBT-MMIO frame carry $(\mathbf{x}, M_{\text{seed}}, P_{\text{debt}}, \nu, z_f)$ to the visualizer. $\square$

Theorem 8 (Causal-Point History Crystallization and GET Clustering). *Each causal-point observer address A is admissible if and only if its entropy margin is non-negative,*

$$R_{\text{entropy}}(A) \equiv N_{\text{limit}}(A) - C_{\text{get}}(A) \geq 0, \quad (356)$$

which is exactly the predicate $\mathcal{P}_{\text{adm}}(A)$ of Eq. (197). An admissible observer executes the rank-one history projector $\Pi_{A,i} = |A, \omega_i\rangle\langle A, \omega_i|$ on its past light cone, crystallizing the regional density field $\rho_{\text{proj}}(\mathbf{x})$ and inducing the emergent clustering acceleration

$$\mathbf{a}_{\text{GET}}(\mathbf{x}) = -\kappa_{\text{GET}} \nabla \ln \rho_{\text{proj}}(\mathbf{x}). \quad (357)$$

As $z \rightarrow -1$ the register load saturates ($f_{\text{load}} \rightarrow 1$), the local capacity vanishes ($N_{\text{limit}} \rightarrow 0$), and the admissible observer set freezes to the empty set, $\mathcal{R}_{\text{adm}} \rightarrow \emptyset$.

Implementation statement. `src/shbt/visualizer/src/shaders/causal_point_get.wgsl` implements the gate: `evaluate_causal_points` evaluates Eq. (356) per observer node and `apply_get_acceleration` evaluates Eq. (357) per particle against the CIC density grid. `WasmShbtEngine.update_epoch` and `HorizonLedger` drive the epoch partition and the freeze $\mathcal{R}_{\text{adm}} \rightarrow \emptyset$; the invariant gates `tests/invariant_gates.rs` fix bit conservation, the 10/33–23/33 split, and the de Sitter observer freeze, and the ledger is exported under `result.json:causal_point_simulation`. $\square$

9.18 Two-Tier Interactive Visualization Architecture

The reference implementation separates the computation into a high-precision Tier-1 core and a real-time Tier-2 renderer. Tier 1 (`shbt_simulator`, MPFR-precision ledger plus `precision_cosmology.py` and `boltzmann_shbt.py`) produces the observables reported above and serializes them into the SHBT-MMIO telemetry frame (`src/shbt/export.rs`): a 128-byte header carrying `magic=0x54424853` and `schema=0x00020000`, z , a , $H(z)$, $f_{\text{load}}(z)$, active and dark bit counts, Landauer debt, seed mass, $f\sigma_8$, Δ_{ISW} , and the particle count, followed by a 256-point transfer grid, a 256-point matter-power grid, and 128-byte `SeedDefectRecord` entries.

Tier 2 is the standalone `shbt-visualizer` crate (`src/shbt/visualizer`), compiled to `wasm32` and bound to `visualizer/index.html`. A `ShbtWebGpuEngine` drives four WGS1 stages: `causal_point_get.wgs1` (active Causal-Point GET dynamics of Theorem 8: `evaluate_causal_points` and `apply_get_acceleration`), `nbody_pm.wgs1` (Fast-PM/2LPT `kick_drift_kernel` under the loaded conformal friction $1 + f_{\text{load}} \cdot 10/33$ with the Stinespring anti-baryon de-render envelope and softened seed-defect attraction), `dual_channel_render.wgs1` (Channel A emission weighted by `vis_weight`; Channel B ghost shear and mass density conserved independently of it), and `holographic_post.wgs1` (lensed-UV composite, ghost false-color, `unwrap_torus_projection` conformal boundary unwrapping onto the CFT torus $[0, 2\pi)^2$, and the horizon overlay driven by f_{load}). Double-buffered storage holds 2^{20} particles in 32-byte structures (under 256 MB), and the pipeline is gated at 60 FPS by the browser presentation clock; the `headless` binary runs the identical pass structure natively for CI benchmarking.

10 Sub-Kelvin Address-Scaled Landauer Calorimetry Architecture and Falsification Protocols

The Causal Point construction yields a laboratory-scale discriminator that is logically downstream of the boundary theory rather than an independent phenomenological add-on. The experiment tests whether resolving a fixed target through a larger address space carries a thermodynamic cost. A complete readout cycle includes acquisition and reset of its classical record, so the standard lower bound for one logically irreversible bit is

$$\Delta Q_{\text{Landauer}} \geq k_B T \ln 2. \quad (358)$$

Standard quantum mechanics and thermodynamics assign this bound to the payload and record reset, independently of the register topology. SHBT instead treats the address locator as part of the finite entropy payment made by the observer.

10.1 Operational Capacity and Competing Hypotheses

The general Causal Point cost in Eq. (334) remains

$$C_{\text{get}} = \max(1, \log_2 |\mathcal{R}| + \log_2 |\Omega_A|, C_{\text{req}}). \quad (359)$$

For calorimetry it is useful to expose the additive sources of the operational capacity explicitly. Measuring entropy in bits, define

$$C_{\text{op}} = \Delta S_{\text{payload}} + \log_2 |\mathcal{R}| + C_0, \quad (360)$$

where $\Delta S_{\text{payload}}$ is the resolved payload entropy in bits and C_0 is the address-independent preparation, acquisition, and reset overhead. The SHBT dissipation inequality is

$$Q_H \geq k_B T \ln 2 C_{\text{op}}. \quad (361)$$

The fixed contributions $\Delta S_{\text{payload}} + C_0$ are measured in control runs and absorbed into the fitted intercept. Consequently, the slope test is independent of their calibrated value. For the proposed singleton outcome ensemble $|\Omega_A| = 1$, baseline request $C_{\text{req}} = 1$, and $|\mathcal{R}| \geq 2$, Eq. (359) gives

$$C_{\text{get}} = \max(1, \log_2 |\mathcal{R}|) = \log_2 |\mathcal{R}|. \quad (362)$$

Thus the background-subtracted experimental capacity C_{op} and the formal GET cost C_{get} have the same address-dependent part.

Let $k = \log_2 |\mathcal{R}|$. The hypotheses are deliberately posed as a derivative so that unknown constant heat offsets cannot mimic the SHBT signal:

$$H_0 : \begin{cases} Q_{H_0} = k_B T \ln 2, & |\mathcal{R}| \geq 2, \\ \frac{\partial Q_H}{\partial (\log_2 |\mathcal{R}|)} = 0, \end{cases} \quad (363)$$

$$H_1 : \begin{cases} Q_{H_1} = (\log_2 |\mathcal{R}|) k_B T \ln 2, \\ \frac{\partial Q_H}{\partial (\log_2 |\mathcal{R}|)} = k_B T \ln 2. \end{cases} \quad (364)$$

Here H_0 is the standard address-independent thermodynamic model and H_1 is the SHBT boundary-mechanics model. For a burst of N_{pulse} identical operations, both predicted heats and slopes are multiplied by N_{pulse} .

10.2 Predicted Heat at 7 mK

At the proposed base temperature,

$$\begin{aligned} E_L &= k_B T_{\text{base}} \ln 2 \\ &= \left(1.380649 \times 10^{-23} \text{ J K}^{-1} \right) (0.0070 \text{ K}) \ln 2 \\ &= 6.698951 \times 10^{-26} \text{ J} \simeq 0.0670 \text{ yJ}. \end{aligned} \quad (365)$$

The address space is swept through

$$|\mathcal{R}| \in \{2^1, 2^2, \dots, 2^{10}\}. \quad (366)$$

Table 31 retains both the single-event predictions and the synchronized 100,000-pulse burst predictions. The last entry follows directly from $10N_{\text{pulse}}E_L = 66.9895 \text{ zJ}$.

Address space $ \mathcal{R} $	Address bits k	Single event Q_{H_0} (yJ)	Single event Q_{H_1} (yJ)	100k burst Q_{H_0} (zJ)	100k burst Q_{H_1} (zJ)
$2^1 = 2$	1	0.0670	0.0670	6.6990	6.6990
$2^2 = 4$	2	0.0670	0.1340	6.6990	13.3979
$2^3 = 8$	3	0.0670	0.2010	6.6990	20.0969
$2^4 = 16$	4	0.0670	0.2680	6.6990	26.7958
$2^5 = 32$	5	0.0670	0.3349	6.6990	33.4948
$2^6 = 64$	6	0.0670	0.4019	6.6990	40.1937
$2^7 = 128$	7	0.0670	0.4689	6.6990	46.8927
$2^8 = 256$	8	0.0670	0.5359	6.6990	53.5916
$2^9 = 512$	9	0.0670	0.6029	6.6990	60.2906
$2^{10} = 1024$	10	0.0670	0.6699	6.6990	66.9895

Table 31: Address-independent H_0 and address-scaled H_1 heat predictions at $T_{\text{base}} = 7.0$ mK.

10.3 Superconducting Register and AuPd Nanocalorimeter

The register is a ten-qubit superconducting circuit on ultra-high-resistivity float-zone silicon. Enabling $k \in \{1, \dots, 10\}$ address-control lines with superconducting single-flux-quantum (SFQ) selectors realizes $|\mathcal{R}| = 2^k$ while routing each readout pulse to the same designated target. Table 32 gives the complete register specification.

Parameter	Value or specification	Physical function or constraint
Substrate	High-resistivity silicon, $\rho > 10 \text{ k}\Omega \text{ cm}$	Low dielectric loss, $\tan \delta < 10^{-6}$
Transmon Josephson energy E_J/h	15.0 GHz	Sets the transmon operating scale
Charging energy E_C/h	300 MHz	Maintains $E_J/E_C \simeq 50$
Transmon layout	Ten fixed-frequency transmons	Eliminates flux-bias heating during the address sweep
Transition frequency f_{01}	4.50–5.50 GHz	Frequency-multiplexed fixed-frequency state drive
Readout resonator frequency f_r	6.50–7.50 GHz	Dispersive Jaynes–Cummings readout
Active addressing switches	Superconducting SFQ logic gates	Address routing without normal-state dissipation
Register span	Ten address-control bits, $2 \leq \mathcal{R} \leq 1024$	Discrete sweep of the active decoder network
AuPd absorber	$1.0 \times 0.2 \times 0.05 \mu\text{m}^3$	$V_{\text{abs}} = 1.0 \times 10^{-20} \text{ m}^3$
Base-temperature heat capacity	$C_e = 1.40 \times 10^{-20} \text{ J K}^{-1}$	Calibrated at $T_{\text{base}} = 7.0$ mK
Thermal decay constant	$\tau_{\text{th}} = 20.41 \text{ ms}$	Resolves the integrated burst transient

Table 32: Experimental circuit specifications for the superconducting register, SFQ address network, and AuPd calorimetric termination.

The readout line terminates in a metallic nanocalorimeter whose electronic temperature is sensed by symmetric NIS tunnel junctions and low-noise SQUID preamplifiers. The absorber is electron-beam-evaporated $\text{Au}_{60}\text{Pd}_{40}$ at 60/40 wt% with $L = 1.0 \mu\text{m}$, $W = 0.2 \mu\text{m}$, and $t = 0.05 \mu\text{m} = 50 \text{ nm}$. Hence

$$V_{\text{abs}} = LWt = 1.0 \times 10^{-20} \text{ m}^3. \quad (367)$$

With Sommerfeld coefficient $\gamma_e = 200.0 \text{ J m}^{-3} \text{ K}^{-2}$, the electron heat capacity at base temperature

is

$$\begin{aligned}
C_e(T) &= \gamma_e V_{\text{abs}} T \\
&= \left(200.0 \text{ J m}^{-3} \text{ K}^{-2}\right) \left(1.0 \times 10^{-20} \text{ m}^3\right) (0.0070 \text{ K}) \\
&= 1.40 \times 10^{-20} \text{ J K}^{-1}.
\end{aligned} \tag{368}$$

The thermal link is selected so that the measured differential conductance at the operating point is

$$G_{\text{th}} = \frac{C_e}{\tau_{\text{th}}} = \frac{1.40 \times 10^{-20} \text{ J K}^{-1}}{20.41 \times 10^{-3} \text{ s}} = 6.85938 \times 10^{-19} \text{ W K}^{-1}. \tag{369}$$

The intrinsic thermal relaxation time is therefore

$$\tau_{\text{th}} = \frac{C_e}{G_{\text{th}}} = \frac{1.40 \times 10^{-20} \text{ J K}^{-1}}{6.85938 \times 10^{-19} \text{ W K}^{-1}} = 2.0408 \times 10^{-2} \text{ s} \simeq 20.41 \text{ ms}. \tag{370}$$

The electron and bath temperatures obey the linearized two-temperature equations

$$C_e \frac{d\Delta T_e}{dt} = P_{\text{in}}(t) - G_{\text{th}} \Delta T_e, \quad \tau_{\text{th}} = \frac{C_e}{G_{\text{th}}}, \tag{371}$$

$$P_{\text{in}}(t) = \frac{Q_{\text{exp}}}{\tau_{\text{in}}} e^{-t/\tau_{\text{in}}}, \quad \tau_{\text{in}} = 1.00 \text{ ms}. \tag{372}$$

Using the pulse-amplitude convention fixed by the calorimetric calibration, the fitted thermal transient is

$$\Delta T_e(t) = \frac{Q_{\text{exp}}}{C_e} \left(e^{-t/\tau_{\text{th}}} - e^{-t/\tau_{\text{in}}} \right), \quad \tau_{\text{th}} = 20.41 \text{ ms}. \tag{373}$$

Nanocalorimeter parameter	Value or specification	Measurement role
Absorber composition	Au ₆₀ Pd ₄₀ , 60/40 wt%	Normal-metal electron absorber
Absorber geometry	$1.0 \times 0.2 \times 0.05 \mu\text{m}^3$	$V_{\text{abs}} = 1.0 \times 10^{-20} \text{ m}^3$
Sommerfeld coefficient	$\gamma_e = 200.0 \text{ J m}^{-3} \text{ K}^{-2}$	Determines $C_e(T)$
Electron heat capacity	$C_e = 1.40 \times 10^{-20} \text{ J K}^{-1}$ at 7 mK	Converts deposited heat to electron-temperature rise
Thermal conductance	$G_{\text{th}} = 6.85938 \times 10^{-19} \text{ W K}^{-1}$	Calibrated bath-coupling scale
Thermal time constant	$\tau_{\text{th}} = 20.41 \text{ ms}$	Enables quasi-adiabatic integration of a 1 ms burst
Injection time constant	$\tau_{\text{in}} = 1.00 \text{ ms}$	Fixes the rising edge of the two-temperature pulse
NIS junction stack	Symmetric Al/AlO _x /AuPd	Electron thermometry
NIS junction area	$100 \times 100 \text{ nm}^2$, high R_N	Suppresses lead-mediated heat leakage
NIS bias	$V_{\text{bias}} \simeq \Delta_{\text{Al}}/e \simeq 180 \mu\text{V}$	High-responsivity operating point
Voltage responsivity	$dV/dT \simeq 1.50 \text{ mV K}^{-1}$ below 10 mK	Converts temperature excursions to voltage
Preamplification	Ultra-low-noise SQUID at the base stage	Resolves the NIS transient

Table 33: AuPd absorber, thermal link, and NIS nanocalorimeter specifications.




Figure 2: Inline apparatus schematic for the address-scaling test. The SFQ selector varies the active address depth while the same target and readout path deposit energy in the AuPd absorber; symmetric NIS junctions and a SQUID read the electron-temperature transient at the 7 mK stage.

10.4 Cryogenic Filtering and Environmental Control

The assembly is mounted in a dry dilution refrigerator at $T_{\text{base}} = 7.0 \pm 0.1$ mK. The complete staged attenuation, filtering, and magnetic-shielding budget is listed in Table 34.

Stage or component	Attenuation or filtering	Physical specification
Room temperature (300 K) to 4 K	20 dB coaxial attenuators	Stainless-steel flexible coaxial lines
Still stage (0.8 K)	10 dB attenuators	NbTi superconducting coaxial lines
Cold plate (100 mK)	10 dB attenuators	CuNi coaxial lines
Mixing chamber (7 mK)	20 dB thermalized attenuators	Eccosorb infrared filters and Cu-powder filters over 0.1–10 GHz
Outer magnetic shield	μ -metal cylinder at 300 K	Suppresses the ambient geomagnetic field
Cryogenic magnetic shield	Dual-layer Cryoperm-10 plus superconducting Al shield at 7 mK	Residual ambient flux $B < 0.5$ nT

Table 34: Cryogenic attenuation, filtering, and magnetic-shielding budget.

10.5 Address-Scaling Measurement Protocol

The synchronized sequence holds the payload and physical pulse envelope fixed while changing only the active address decoder:

1. *State initialization.* Initialize the target transmon in $|0\rangle$ by active feedback and microwave sideband cooling, require $P_{|1\rangle} < 0.001$, and wait more than 100 ms for baseline thermalization.
2. *Address configuration.* Assert $k = \log_2 |\mathcal{R}|$ SFQ selector channels, with $k \in \{1, \dots, 10\}$, so the enabled decoder spans $|\mathcal{R}| = 2^k$ addresses.
3. *GET readout.* Apply a 6.50 GHz, 10 ns readout pulse to the same address key $A \in \mathcal{R}$. The target remains logical zero in every trial, hence $\Delta S_{\text{payload}} = 0$ for the prepared target; the constant acquisition and record-reset heat is assigned to the intercept.

4. *Pulse-train integration.* Repeat the GET operation $N_{\text{pulse}} = 100,000$ times at $f_{\text{rep}} = 100$ MHz. The resulting $t_{\text{burst}} = 1.0$ ms obeys $t_{\text{burst}} \ll \tau_{\text{th}} = 20.41$ ms, so the absorber integrates the burst quasi-adiabatically.
5. *Calorimetric acquisition.* Record $V_{\text{NIS}}(t)$ for $0 \leq t \leq 100$ ms, extract the peak shift ΔV_{peak} , and integrate the thermal decay to obtain Q_{exp} .
6. *Thermal reset.* Wait $t_{\text{cool}} = 150$ ms $> 7\tau_{\text{th}}$ before the next burst so the absorber returns to T_{base} .

Each of the ten address configurations receives $M = 1,000$ independent burst repetitions. The complete sweep is therefore

$$k \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} \iff |\mathcal{R}| \in \{2, 4, 8, 16, 32, 64, 128, 256, 512, 1024\}. \quad (374)$$

The acquisition and inference chain is fixed before unblinding and is listed in Table 35. The same operations are applied to signal, null, and capacity-ceiling runs.

Stage	Analysis operation	Mathematical or instrumental action	Stored output
1	Baseline calibration	Measure T_{base} , C_e , G_{th} , NIS responsivity, and the zero-pulse trace	Calibration covariance matrix
2	Address randomization	Randomly permute $k \in \{1, \dots, 10\}$ while fixing the target, payload, pulse envelope, and repetition count	Blinded address key and run identifier
3	Pulse delivery	Execute $N_{\text{pulse}} = 100,000$ SFQ-selected GET operations during $t_{\text{burst}} = 1.0$ ms	Pulse timestamps and decoder-state audit
4	Transient acquisition	Sample $V_{\text{NIS}}(t)$ over $0 \leq t \leq 100$ ms and convert voltage to $\Delta T_e(t)$	Calibrated temperature time series
5	Thermal fit	Fit Eq. (373) jointly for Q_{exp} , baseline offset, and nuisance drift	Heat estimate and fit covariance
6	Quality control	Reject preregistered saturation, missing-trigger, thermal-reset, and residual-autocorrelation failures	Binary quality mask and diagnostic residuals
7	OLS/MLE inference	Fit Eq. (375), evaluate Eqs. (379)–(381), and compute the slope p -value	$\hat{\alpha}$, standard error, p , χ^2_ν , and ΔAIC
8	Capacity-ceiling audit	Impose $C_{\text{local}} = 6$ bits and test the transition between $k = 6$ and $k = 7$	Fidelity, thermal variance, and AnomalyClosureError flag

Table 35: Stage-by-stage calorimetry acquisition, reduction, inference, and finite-capacity falsification operations.

The systematic-error targets are small compared with the predicted SHBT step $N_{\text{pulse}}E_L = 6.6990$ zJ/bit.

10.6 Empirical Falsification Criteria and Observer-Capacity Cutoff

10.6.1 OLS/MLE Slope Test

For each address length k , let $Q_{\text{exp}}(k)$ be the burst heat. The direct regression used for the primary decision is

$$Q_{\text{exp}}(k) = \hat{\beta}_0 + \hat{\alpha}k + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2), \quad (375)$$

Systematic leakage channel	Estimated contribution	Physical mechanism	Mitigation
Parasitic capacitive cross-talk	< 0.05 zJ per burst	Coupling between inactive address lines and calorimeter	Differential ground-shielded coplanar waveguides
Microwave drive heating	< 0.10 zJ per burst	Direct absorption from 6.5 GHz readout photons	Spatial isolation $> 500 \mu\text{m}$ plus bandpass filters
Amplifier back-action	< 0.02 zJ per burst	Josephson or HEMT noise returning along the readout line	Dual junction isolators with > 45 dB isolation at 6.5 GHz
NIS-lead thermal conductance	< 0.01 zJ per burst	Heat leakage through normal-metal thermometer leads	$100 \times 100 \text{ nm}^2$ junctions with high R_N
Engineering noise allowance	$\simeq 0.18021$ zJ per burst	Safety-scaled design budget	Conditional on the noise model in Section 10.7

Table 36: Systematic-error attenuation budget for address-scaled calorimetry.

fitted by ordinary least squares (OLS) and Gaussian maximum likelihood estimation (MLE). The competing slope values are

$$\begin{aligned}
H_0 : \quad \alpha_0 &= 0 \text{ zJ/bit}, \\
H_1 : \quad \alpha_1 &= N_{\text{pulse}} k_B T \ln 2 \\
&= 100,000 \left(6.69895 \times 10^{-26} \text{ J/bit} \right) \\
&= 6.69895 \times 10^{-21} \text{ J/bit} = 6.6990 \text{ zJ/bit}.
\end{aligned} \tag{376}$$

Equivalently, the source model templates are

$$Q_{H_0}(k) = \alpha_0 + \epsilon, \quad \alpha_0 = N_{\text{pulse}} k_B T \ln 2 = 6.6990 \text{ zJ}, \tag{377}$$

$$Q_{H_1}(k) = \beta_0 + \beta_1 [k N_{\text{pulse}} k_B T \ln 2] + \epsilon, \quad \hat{\alpha} = \hat{\beta}_1 N_{\text{pulse}} k_B T \ln 2. \tag{378}$$

For model H_j , $j \in \{0, 1\}$, the Gaussian maximum-likelihood ledger is

$$\ln \mathcal{L}_j = -\frac{n}{2} \ln(2\pi\hat{\sigma}_j^2) - \frac{1}{2\hat{\sigma}_j^2} \sum_{a=1}^n [Q_a - Q_{H_j}(k_a)]^2, \tag{379}$$

$$\hat{\sigma}_j^2 = \frac{1}{n} \sum_{a=1}^n [Q_a - Q_{H_j}(k_a)]^2, \tag{380}$$

$$\text{AIC}_{H_j} = 2p_j - 2 \ln \mathcal{L}_j, \quad \Delta\text{AIC} = \text{AIC}_{H_0} - \text{AIC}_{H_1}. \tag{381}$$

The alternative is accepted only if the null slope is rejected at least 5σ with $p < 10^{-6}$, $\Delta\text{AIC} > 10.0$, the fitted slope is statistically consistent with 6.6990 zJ/bit, and the address sweep exhibits discrete stair-step increments at integer k . The address-scaling prediction is falsified, and H_0 is accepted within the stated experimental resolution, if

$$|\hat{\alpha}| \leq 0.05 (N_{\text{pulse}} k_B T \ln 2) = 0.33495 \text{ zJ/bit}. \tag{382}$$

Table 37 records the full fit and model-selection criteria.

Metric or parameter	Null-model H_0 target	SHBT H_1 target	Selection threshold
Direct slope $\hat{\alpha}$	0 zJ/bit	6.6990 zJ/bit	Reject H_0 at $\geq 5\sigma$, $p < 10^{-6}$
Normalized slope β_1	0.000 ± 0.010	1.000 ± 0.010	Requires consistency with the selected model
Intercept β_0	6.699 zJ	0.000 ± 0.18021 zJ design allowance after baseline subtraction	Diagnoses parasitic drive heating
Reduced chi-square χ_ν^2	$\chi_\nu^2 \gg 1$ for SHBT-like data	1.00 ± 0.05	Noise-consistent goodness of fit
Akaike information criterion	AIC_{H_0} baseline	$\Delta\text{AIC} =$ $\text{AIC}_{H_0} - \text{AIC}_{H_1} > 10$	Decisive preference for H_1

Table 37: OLS/MLE parameters and binary model-selection thresholds.

10.6.2 Local Capacity Ceiling and AnomalyClosureError

The second test probes the finite-observer cutoff independently of the linear heat slope. To make the notation explicit, define the Causal Point’s local capacity ceiling by

$$C_{\text{local}} \equiv N_{\text{limit}} = \min\left(N_{\text{local}}, \frac{A_{\text{local}}}{4L_P^2 \ln 2}\right). \quad (383)$$

The experimental residual entropy budget is

$$R_{\text{entropy}}^{\text{exp}} = C_{\text{local}} - C_{\text{op}}. \quad (384)$$

For the singleton, background-calibrated GET sequence, $C_{\text{op}} = C_{\text{get}} = k$. Normal state resolution requires $R_{\text{entropy}}^{\text{exp}} \geq 0$; if $C_{\text{op}} > C_{\text{local}}$, the negative residual triggers the SHBT failure condition denoted **AnomalyClosureError**.

The controlled cutoff run imposes an artificial local ceiling

$$C_{\text{local}}^{\text{exp}} = N_{\text{limit}}^{\text{exp}} = 6 \text{ bits}, \quad |\mathcal{R}|_{\text{max}} = 2^6 = 64, \quad (385)$$

and then continues the physical decoder sweep through $k = 10$. SHBT predicts

$$\begin{cases} k \leq 6 : & \text{admissible readout,} \\ k > 6 : & \text{loss of readout fidelity, nonlinear thermal-variance spikes,} \\ & \text{and state-resolution failure (AnomalyClosureError).} \end{cases} \quad (386)$$

Standard quantum mechanics predicts continuous address decoding across the same artificial ceiling provided the hardware remains functional. A sharp, reproducible breakdown at $C_{\text{op}} > C_{\text{local}}^{\text{exp}}$ therefore tests the finite-register postulate, while its absence falsifies this cutoff prediction even if a nonzero calorimetric slope were observed.

10.7 Cryogenic Nanocalorimetry Noise Budget at 7 mK

The noise budget distinguishes a single Landauer operation from the $N_{\text{pulse}} = 100,000$ -operation burst used in the measurement protocol. At $T = 7.0$ mK,

$$E_L = k_B T \ln 2 = 6.6989507 \times 10^{-26} \text{ J}, \quad (387)$$

$$\Delta Q_{\text{step}} = N_{\text{pulse}} E_L \simeq 6.698951 \text{ zJ}. \quad (388)$$

The burst heat slope is therefore $N_{\text{pulse}} k_B T \ln 2 \simeq 6.6990 \text{ zJ/bit}$; the single-operation slope is $k_B T \ln 2 \simeq 6.69895 \times 10^{-5} \text{ zJ/bit}$. These are different measurement normalizations of the same proposed address-dependent contribution.

For the equilibrium-fluctuation model and absorber heat capacity $C_e = 1.40 \times 10^{-20} \text{ J K}^{-1}$ used here, the thermal energy standard deviation is

$$\begin{aligned} \Delta E_{\text{th}} &= \sqrt{k_B T^2 C_e} \\ &= \sqrt{(1.380649 \times 10^{-23})(0.0070)^2(1.40 \times 10^{-20}) \text{ J}} \\ &\simeq 3.0775399 \times 10^{-24} \text{ J} \simeq 0.00307754 \text{ zJ}. \end{aligned} \quad (389)$$

This equilibrium quantity is not, by itself, the uncertainty of the finite-duration heat estimator. Relating the two requires the thermal and readout transfer functions, integration window, and estimator calibration.

For a provisional independent-noise model, interpret the channel allocations in Table 38 as equivalent per-burst RMS contributions. Their quadrature sum is

$$\begin{aligned} \Delta Q_{\text{total}} &= \left[(\Delta Q_{\text{SFQ}})^2 + (\Delta Q_{\text{micro}})^2 + (\Delta Q_{\text{SQUID}})^2 + (\Delta Q_{\text{lead}})^2 + (\Delta E_{\text{th}})^2 \right]^{1/2} \\ &\simeq \sqrt{0.050^2 + 0.100^2 + 0.020^2 + 0.010^2 + 0.00307754^2} \text{ zJ} \\ &\simeq 0.11405907 \text{ zJ}. \end{aligned} \quad (390)$$

Applying the specified engineering safety multiplier to the unrounded result defines the design allowance

$$\Delta Q_{\text{bounded}} \equiv 1.58 \Delta Q_{\text{total}} \simeq 0.18021333 \text{ zJ} \simeq 0.18021 \text{ zJ}. \quad (391)$$

The label “bounded” denotes this engineering allowance, not a proven upper bound or a specified statistical confidence interval. In particular, the unrounded allowance is slightly greater than 0.18 zJ. The multiplier 1.58 is an experimental design choice, not a constant derived from the canonical affine branch.

The corresponding design signal-to-noise ratios are

$$\frac{\Delta Q_{\text{step}}}{\Delta Q_{\text{total}}} \simeq 58.73, \quad \frac{\Delta Q_{\text{step}}}{\Delta Q_{\text{bounded}}} \simeq 37.17. \quad (392)$$

These ratios establish an arithmetic separation between the proposed signal and the allocated noise, conditional on the model. They do not establish that a fabricated device attains those allocations. For correlated noise, the combined variance instead includes $2 \sum_{i < j} \text{Cov}(Q_i, Q_j)$; deterministic heat offsets must be calibrated separately rather than combined as independent random variances. Device calibration and control runs are therefore required before claiming a measured noise bound or experimental signal resolution.

Noise channel or aggregate	Physical origin or combination	Energy (zJ)	Squared energy (zJ ²)
SFQ logic switching	Clock cross-talk allocation	0.05000	0.00250000
Microwave readout	Readout leakage allocation	0.10000	0.01000000
SQUID back-action	Amplifier back-action allocation	0.02000	0.00040000
Thermal leads	Lead-leakage allocation	0.01000	0.00010000
Electron thermal fluctuation	Equilibrium absorber model	0.00307754	0.000009471
Combined RMS total	Independent-channel quadrature	0.114059	0.013009471
Safety-scaled allowance	1.58× unrounded RMS total	0.180213	0.032476847

Table 38: Arithmetic audit of the proposed 7 mK per-burst noise budget. The first four entries are design allocations, not measured noise levels. Squared entries are variances only under the stated RMS interpretation; the safety-scaled row is not an additional noise source.

11 Unified Closure Synthesis

SHBT comprises five coupled sectors: modular completion of the boundary theory, entropy self-resolution into bulk metric data, topological anti-baryon state decoupling with passive stress-energy preservation, entropy-budgeted observer history, and address-scaled calorimetric falsification. These are not independent mechanisms; they are projections of the same finite closed boundary register. Once the completed partition function is modular invariant and the canonical branch arithmetic closes, the bulk tensor, the physical Hamiltonian constraint, the baryogenesis fixed point, and the Causal Point history rule all become consequences of the same closure data; the calorimetric experiment tests the observer-cost consequence directly.

The core closure chain is

$$\boxed{[M, S_\partial] = [M, T_\partial] = 0 \iff \Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0 \iff \Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0.} \quad (393)$$

Equivalently, $H_{\text{total}} = 0$ after restriction to the physical subspace. The first condition says that the torus partition function is a genuine modular invariant of the completed boundary CFT. The second says that the visible levels and parent completion level have no residual framing mismatch. The third says that the bulk closure tensor vanishes, so the SHBT topological Einstein equation is satisfied without adding an independent bulk correction. The fourth says that the static physical sector has no external time generator left over after projection. The canonical branch is therefore the branch on which boundary modular covariance, visible-sector anomaly closure, bulk geometry, and the Hamiltonian constraint are the same statement viewed through different interfaces.

The geometric part of the construction is summarized by the holographic RG map

$$\boxed{\nabla_{\mathcal{C}} S_\partial \longrightarrow \rho_E(i, j) \longrightarrow \ell_r(\tau) \longrightarrow g_{\mu\nu}(\tau).} \quad (394)$$

The boundary entropy gradient is first converted into the normalized entanglement density $\rho_E(i, j)$ on the finite modular-overlap coordinate set. The ordered resolved state is then compressed into the prime-lattice load vector $\ell_r(\tau)$, and the stabilized Gram projection of this load vector gives the effective metric slice $g_{\mu\nu}(\tau)$. In this sense geometry is not a continuous background into which the boundary theory is inserted. It is the rendered image of finite boundary information after entropy gradients have been normalized, ordered, compressed, and projected.

The matter-asymmetry sector closes by the topological anti-baryon state decoupling loop

$$\boxed{\rho_B^\star = \frac{\mathcal{D}_{\bar{B}}\rho_B^\star\mathcal{D}_B^\dagger}{\text{Tr}(\mathcal{D}_{\bar{B}}\rho_B^\star\mathcal{D}_B^\dagger)}, \quad \mathcal{C}_{\bar{B}}(\rho_B^\star) = 0, \quad \delta T_{\mu\nu}^{\text{passive}} = 0.} \quad (395)$$

At the fixed point, active anti-baryon color and electromagnetic render charge have been stripped from the visible channel, but the passive gravitational source has not been discarded. It is carried by the dark completion sector. The resulting baryon excess is therefore an asymmetry of rendered visible charge, not a violation of the completed stress-energy account. This is why the same branch that fixes $\Delta_{\text{fr}} = 0$ also enforces $\delta T_{\mu\nu}^{\text{passive}} = 0$: the Stinespring isometric state-projection map is allowed only because it is completion preserving.

The same modular ledger that fixes η_B also yields $\Omega_{\text{DM}}/\Omega_b \simeq 5.4$, identifying dark matter as the passive gravitational ghost of decoupled anti-baryons.

The observer sector supplies the final link in the same chain. A Causal Point does not read a pre-existing classical history from an infinite archive. It crystallizes a finite ordered sequence of admissible projections,

$$\boxed{\Pi_{\text{history}} = \Pi_{A_m, \iota_m} \cdots \Pi_{A_2, \iota_2} \Pi_{A_1, \iota_1}.} \quad (396)$$

Each factor is admitted only if the local entropy budget remains nonnegative. The apparent classical past is therefore the record that survives the ordered application of finite-register GET operations. The observer is part of the cosmic code rather than an external reader of it: the address A_n , outcome index ι_n , local hidden-state budget, and pointer packet are all internal boundary data.

The principal numerical retrodictions of the canonical branch can be collected as

$$\boxed{\Lambda_{\text{holo}} \simeq 1.09 \times 10^{-52} \text{ m}^{-2}, \quad N \simeq 3.31 \times 10^{122}, \quad \eta_B = \mathcal{O}(10^{-10}), \quad g_{\mu\nu} = g_{\mu\nu}(\nabla_{\mathcal{C}} S_{\partial}).} \quad (397)$$

The first number is the holographic dark-energy curvature scale used by the topological Einstein equation. The second is the finite boundary bit budget associated with that scale. The third is the order of the visible baryon asymmetry produced by the state-decoupling loop, with the canonical value $\eta_B = 6.449923359416 \times 10^{-10}$. The fourth is the structural prediction: the effective metric is not fundamental input but a calculable output of the entropy-gradient pipeline.

Relative to standard holographic approaches, SHBT keeps the boundary-first logic but changes what is taken as primitive. It does not start from a large- N gauge theory with an asymptotically anti-de Sitter bulk and then ask for a semiclassical interior. It starts from a completed static boundary CFT, requires exact modular closure, and lets finite entropy data render the bulk metric. Relative to entropic-gravity pictures, SHBT does not infer gravity from entropy alone; it constrains the entropy flow by the modular invariant pairing matrix, the canonical branch arithmetic, and the finite register size. Relative to purely constraint-based quantum gravity, it does not stop at the formal statement that the Hamiltonian vanishes on physical states; it ties that statement to visible-sector levels, dark completion, baryogenesis, and observer memory.

The most direct tests of the framework are therefore consistency tests across sectors. The branch data must give integer locks $I_\ell^\star = 6$ and $I_q^\star = 13$, zero framing defect, a vanishing bulk closure tensor, and a finite physical-sector Hamiltonian constraint. The entropy pipeline must produce symmetric,

trace-normalized, positive metric slices from the same boundary densities that define the loading sequence. The baryogenesis module must remove active visible anti-baryon charge while keeping passive stress-energy unchanged. The Causal Point module must crystallize only those histories whose retrieval cost fits inside the local entropy budget. A sub-10 mK implementation must then distinguish the predicted address slope $\hat{\alpha} = N_{\text{pulse}} k_B T \ln 2$ from the null slope and test the sharp failure condition at $C_{\text{op}} > C_{\text{local}}$. Any failure of one of these checks would invalidate the canonical SHBT closure chain rather than an auxiliary numerical routine alone.

Several limitations remain. The analysis establishes and verifies one canonical branch rather than a classification theorem for all modular completions. The dark completion sector now has an explicit finite-quadratic- module realization with 2901360 primary modules, but uniqueness of this microscopic realization within the full space of rational and graded CFT completions has not been established. The entropy-to-metric map is deterministic on the canonical branch, yet a full comparison with precision cosmological data would require a larger phenomenological layer: survey observables, perturbations, noise models, and parameter-estimation pipelines. The baryogenesis mechanism also needs extension from the fixed render-channel identity to a complete account of early-universe thermal history as seen by finite boundary observers. The calorimeter architecture and statistical thresholds specified here constitute a prospective experiment, not a reported detection; fabrication yield, calibrated excess-noise spectra, and blinded control data remain necessary before either hypothesis can be accepted. These requirements define the next tests needed for a quantitatively predictive research program.

The closure may be stated in code-theoretic terms. A finite boundary register, when forced to close under completed modular kernels, contains more than a list of states. It contains the rules for which visible charges can be rendered, which passive sources must be preserved, which entropy gradients become geometry, and which observer histories can crystallize. In this view spacetime is the readable projection of a finite holographic code; matter asymmetry is a render-channel fixed point of that code; and observation is the entropy-budgeted act of selecting a consistent history from within the same code. SHBT is therefore static only at the level of the completed boundary. The observer-accessible dynamics is the internally crystallized image of that closed finite boundary information.

12 Master Verification Tables and Code–Data Traceability

12.1 Reconciled Precision-Cosmology Master Ledger

The canonical branch observables, perturbative invariants, and laboratory falsification target are collected in a single verification ledger. Values in Table 39 retain the precision of the machine-readable report; the chronometer entry is a reported fit statistic, whereas the remaining entries are branch outputs or experimental targets.

Verification quantity	Reconciled value	Analytic or computational origin
Local Hubble uplift H_0^{loc}	$72.197960 \text{ km s}^{-1} \text{ Mpc}^{-1}$	finite-register loading law
Gradient target A_H	$4.797960 \text{ km s}^{-1} \text{ Mpc}^{-1}$	$\Gamma_{\text{lock}} = 3A_H$
CPL template	$w_0 = -1.064677, w_a = -0.065814$	conserved-fluid image of the loaded background
Density zero crossing	$z_{\text{cross}} = 1.864954$	effective CPL density root
Fractional BBN loading shift	7.421690×10^{-27}	Eq. (222)
Cosmic age	13.276617 Gyr	loaded background age integral
Chronometer statistic	$\chi^2/\nu = 2491.2/2482 = 1.004$	Eq. (330)
Tensor-to-scalar ratio	$r = 0.0032$	Eq. (287)
Tensor spectral index	$n_t = -0.0004$	$n_t = -r/8$
Collapsed trispectrum	$\tau_{\text{NL}} = 0.000324$	Eq. (306)
Address-scaled Landauer slope	$\alpha = 6.6990 \text{ zJ/bit}$	Eq. (376)

Table 39: Reconciled precision-cosmology master verification ledger (Table 17 data product) for the canonical branch (26, 8, 312).

12.2 Equation-to-Code Traceability Matrix

Table 40 maps the publication-level identities to their implementation boundaries. A row denotes an interface contract: exact reproduction requires the displayed equation, module, accessor, and serialized record to agree simultaneously.

12.3 Generated-Artifact and Column Crosswalk

12.4 Implementation Availability and Reproduction Contract

The reference implementation and audit scripts for the SHBT and precision cosmology calculations are maintained in the GitHub repository¹. The tagged release is the intended reproduction target; the supplied run does not record an immutable commit or wheel identifier. The branch-head marker in the repository footnote is not a commit hash. The publication-critical entry points and their physical roles are summarized in Table 42. This section provides the implementation bridge for the theory developed in Sections 1–11.

12.5 Reported Run Provenance and Output Crosswalk

The primary numerical artifact is now `codespace_audit.json`, supplied as the native Linux Codespace run. Its test excerpt reports three passing tests, zero failures, and zero documentation tests, followed by a Python docstring escape warning and two separate JSON objects. Despite the file extension, it is a mixed transcript rather than one standalone JSON document. The first JSON object is the foundation summary; the second is the precision-cosmology report. Its SHA-256 digest is

¹Named release tag: <https://github.com/syslown/shbt-precision/releases/tag/v1.0.0>; commit-sha: `main@head`.

Paper identity	Python or Rust module	Principal accessor	Serialized record
Modular closure, Eq. (393)	src/shbt/boundary.rs	StaticBoundary.s_boundary, t_boundary	result.json: foundation_audit
Boundary loading, Eqs. (212)–(257)	precision_cosmology.py	loading_fraction_ode, h0_redshift_dependent	result.json: redshift_ladder
Finite screen and neutrino floor, Eqs. (229) and (245)	precision_cosmology.py	subleading_d5_correction, neutrino_hierarchy_masses	result.json: neutrino_hierarchy
First-order transport, Eqs. (258)–(276)	boltzmann_shbt.py	boltzmann_rhs, integrate_mode	scalar and matter CSV products
Tensor transfer, Eqs. (285)–(289)	boltzmann_shbt.py	tensor_mode_rhs, tensor_bmode_transfer	shbt_run_tensor_ cls.csv
Bispectrum and trispectrum, Eqs. (292)–(306)	boltzmann_shbt.py	non_gaussianity_templates	result.json: precision_ pipeline.non_ gaussianity
GET capacity and cutoff, Eqs. (360) and (386)	src/shbt/causal_point.rs	CausalPoint.crystallize_ history	result.json: foundation_audit. memory_report
Observer admissibility and lifecycle, Eqs. (197)–(198)	src/shbt/causal_point.rs	CausalPoint.is_admissible, run_lifecycle_cycle	result.json: succession
Stinespring de-rendering, Eq. (199)	src/shbt/causal_point.rs	CausalPoint.terminate_and_ derender	result.json: succession.records
Succession transfer kernel, Eq. (200)	src/shbt/causal_point.rs	CausalPoint.evaluate_ succession_kernel, ShbtSimulator.run_ succession_cycles	result.json: succession. records[].kernel_ probabilities
Asymptotic observer freeze, Eq. (201)	precision_cosmology.py	asymptotic_observer_freeze, observer_admissible_set	result.json: foundation_audit. asymptotic_ observer_freeze
Landauer regression, Eqs. (375)–(381)	shbt_simulate.py	calorimetry OLS/MLE analysis pipeline	shbt_run_ calorimetry_sim. csv
Dynamic bit loading, Thm. 5	src/shbt/cosmology.rs	ShbtUniverse.evaluate_ hubble_index_rate, conformal_loading_fraction	result.json: cosmology_ visualization
Ghost partition, Thm. 6	src/shbt/cosmology.rs	ShbtUniverse.dark_ completion_bits, dark_matter_fraction	data/telemetry.bin header fields
Seed condensation, Thm. 7	src/shbt/cosmology.rs	ShbtUniverse.compute_seed_ condensation, eval_landauer_debt_power	SHBT-MMIO SeedDefectRecord
Telemetry serialization, Sec. 9.18	src/shbt/export.rs	serialize_mmio_frame_py, MmioTelemetryHeader.to_ bytes	data/telemetry.bin
WebGPU engine, Sec. 9.18	src/shbt/visualizer	ShbtWebGpuEngine.step_frame, hud_json	visualizer/pkg/ shbt_visualizer_ bg.wasm
GET clustering, Thm. 8	causal_point_get.wgsl	evaluate_causal_points, apply_get_acceleration	result.json: causal_point_ simulation
Fast-PM drift and seed attraction, Thm. 7	nbody_pm.wgsl	kick_drift_kernel	SHBT-MMIO SeedDefectRecord
Conformal torus unwrapping, Sec. 9.18	holographic_post.wgsl	unwrap_torus_projection	unwrap_transition control
Epoch ledger and observer admissibility, Thm. 8	src/shbt/visualizer	WasmShbtEngine.update_epoch, HorizonLedger	result.json: causal_point_ simulation

Table 40: Code and equation traceability matrix (Table 28 verification ledger).

Generated artifact	Required columns or nested records	Paper-level observable
shbt_run_cmb_cls.csv	Dl_TT, Dl_EE, Dl_TE	scalar temperature, polarization, and cross spectra
shbt_run_matter_pk.csv	Pk_z0, Pk_z05, Pk_z1	matter power at $z = 0, 0.5$, and 1
shbt_run_tensor_cls.csv	Cl_BB, Dl_BB, Cl_TT_tensor	primordial tensor B modes and tensor temperature spectrum
shbt_run_calorimetry_sim.csv	k_bits, R_addresses, Q_H0_zJ, Q_H1_zJ, Q_noise_zJ	address sweep, competing heat laws, and simulated noise
result.json	complete nested foundation audit, precision summary, spectra, non-Gaussianity, stability, and calorimetry records	full machine-readable simulation summary and cross-sector closure audit
data/telemetry.bin	128-byte SHBT-MMIO header; 256-point $T(k, z)$ and $P_m(k, z)$ grids; 128-byte SeedDefectRecord entries	two-tier visualizer telemetry contract (Sec. 9.18)

Table 41: Generated data-product traceability matrix (Table 29 verification ledger).

5a8eefc28f4a8b13791268c629e8a91b5b27d715fbcf0d2d2e5ef54982deb301

The precision report records `foundation_audit.available=true`, `foundation_audit.method=run_full_audit`, `foundation_audit.status=completed`, and `simulator_constants.simulator_available=true`. Thus the native artifact reports a successful foundation invocation, unlike the earlier Python-only run. These are supplied execution results, not an independent rerun of the native build during manuscript compilation.

For comparison, the historical `simulator_output.txt` records fourteen `PrecisionCosmologyTests` tests in 49.830s, with an OK summary. That earlier artifact has SHA-256 digest

73db22deb55747f4d4fe5adecc108fc11655ec79c1a057196f3685c9deffad1a

Only that earlier artifact records `foundation_audit.available=false`, `simulator_constants.simulator_available=false`, and the status `shbt_simulator.ShbtSimulatornotimportable`. The historical fourteen-test transcript is not presented as a fresh fourteen-test native transcript. Foundation availability in the current tables comes from the later native artifact. The interfaces in Table 42 remain the implementation contract; the summary flags are not a complete per-method execution trace.

The native foundation summary reports the branch (26,8,312), `framing_defect(delta_fr)=0.0`, `modular_invariant=true`, `zero_energy_locked=true`, `projection_dimension_26_to_4=true`, `stress_energy_preserved=true`, `projection_all_passed=true`, and `memory_all_passed=true`. The counts `metric_slices=9` and `history_entries=9` accompany `eta_b=6.449923359416131e-10`. Its `central_charge_ledger` stores the residual, completed, and visible charges as numerator/denominator pairs; those exact pairs agree with the manuscript’s rational ledgers. The native `Omega_DM_over_Omega_b=5.343450862978383` agrees with the precision report to its displayed precision.

The native register `N_sat_bits=3.311997720142366E+122` differs from the historical fallback $3.312593327986 \times 10^{122}$. Only the register, its entropy-debt outputs, the foundation-availability record, and simulator-constant record change between the two supplied precision reports. The

Software interface	Physical role and principal accessors
<code>shbt_simulator</code>	PyO3 module exposing the foundation simulator, report getters, and record classes
<code>StaticBoundary</code>	branch arithmetic, visible and dark modular blocks, 2901360-module dark pairing, RG charge trajectories, C-ABI export, partition evaluation, entropy densities, dominant sequence, and self-resolution
<code>ShbtSimulator</code>	full audit orchestration through <code>ShbtSimulator.run_full_audit()</code>
<code>HolographicProjection /</code>	load-vector construction, Euler flux, metric synthesis, spatial projection, and slice spectra
<code>BulkMetricSlice</code>	light-cone construction, local property packets, finite-memory admissibility, history crystallization, five-phase lifecycle state machine, Stinespring de-rendering (<code>terminate_and_derender</code>), and succession kernel evaluation (<code>evaluate_succession_kernel</code> , <code>relabel_and_rerender</code>)
<code>CausalPoint</code>	finite-capacity exception defined in <code>src/shbt/causal_point.rs</code> and exported through <code>src/lib.rs</code>
<code>AnomalyClosureError</code>	observer-record types supporting the Causal Point closure conditions
<code>LightConeSample, LocalPropertyPacket, CoordinateLogEntry, MemoryReport</code>	observer-succession record types supporting the five-phase lifecycle and transfer kernel
<code>CausalPointCandidate, DerenderingRecord, SuccessionRecord, LifecyclePhase</code>	
<code>BaryogenesisIdentity</code>	C_{sph} , $J_{\text{CP}}^{\text{topo}}$, projector rank, restoration scale, and η_B
<code>BaryogenesisOptimizer, FieldSimulation, BenchmarkDelta</code>	anti-baryon charge decoupling, passive-source preservation, and implementation profiling
<code>precision_cosmology.py</code>	completed ledger, Hubble loading, growth, cluster collapse, dark matter, BBN, ISW, neutrinos, chronometers, entropy, and GET costs
<code>boltzmann_shbt.py</code>	conformal background, Einstein constraints, matter and radiation hierarchies, tensor propagation, B-mode transfer, higher-order templates, line-of-sight sources, and transfer-function export
<code>examples/run_audit.py</code>	canonical foundation-audit entry point
<code>src/shbt/boundary.rs</code>	boundary branch, factorized dark Weil kernels, <code>StaticBoundaryCabi</code> , Virasoro trajectories, partition evaluator, and closure checks
<code>src/shbt/entropy_flow.rs</code>	entropy cascade and holographic metric projection
<code>src/shbt/baryogenesis.rs</code>	baryogenesis identity, κ_{D_5} , state decoupling, and benchmark comparison
<code>src/shbt/causal_point.rs</code>	Causal Point light cone, finite-memory checks, history crystallization, admissibility predicate $\mathcal{P}_{\text{adm}}$, Stinespring de-rendering, and the closed-loop succession engine
<code>src/lib.rs</code>	PyO3 exports and <code>ShbtSimulator</code> orchestration
<code>src/shbt/cosmology.rs</code>	<code>ShbtUniverse</code> branch cosmology: loading, Hubble, ghost partition, seed condensation, ISW residual, and horizon freeze
<code>src/shbt/export.rs</code>	SHBT-MMIO frame serialization and <code>serialize_mmio_frame_py</code> export
<code>src/shbt/visualizer</code>	<code>shbt-visualizer</code> crate: <code>Particle</code> , <code>ShbtWebGpuEngine</code> , <code>HudMetrics</code> , <code>WasmShbtEngine</code> , <code>HorizonLedger</code> , WGSL pipeline, and <code>headless</code> benchmark binary
<code>causal_point_get.wgsl</code>	Active Causal-Point GET compute passes: observer admissibility ($R_{\text{entropy}} \geq 0$) and emergent clustering acceleration
<code>tests/invariant_gates.rs</code>	$\mathbf{a}_{\text{GET}} = -\kappa_{\text{GET}} \nabla \ln \rho_{\text{proj}}$ Tier-2 invariant gates: bit conservation across epochs, exact 10/33–23/33 partition, de Sitter observer freeze, and stress-energy residual
<code>scripts/verify_cosmology.py</code>	Seed-coupling audit (α_{seed}), η_B evaluation, and 60 FPS frame-budget benchmark
<code>visualizer/index.html</code>	browser harness: horizon HUD, capacity gauge, seed ledger, Landauer monitor, timeline scrub, speed and channel controls

Table 42: Explicit physics-to-software bridge required to reproduce the paper-level closure tables.

unchanged neutrino floor is reconciled with the native register by the sub-leading D_5 weight correction in Eqs. (243)–(245).

Native build success and reported audit passes establish execution provenance at the level of these artifacts, not a proof that every numeric input is derived without empirical or engineering choices. For example, the report retains the CMB anchor and oscillation-splitting inputs, while `stability_audit` lists `seed_z_peak=7.5` and `seed_sigma_z=0.801` without their derivations. Neither supplied artifact records a commit hash, dependency lockfile, native wheel identity, or a complete parameter-dependency derivation. Zero-parameter computational provenance is therefore not certified by these logs alone.

Table 43 links the reconciled numerical results to exact keys in the native precision JSON. In the field column, names refer to fields within the record in the middle column; array records are matched by their `z` field. These are dictionary mappings, not invented PyO3 methods. Values are rounded only for display, and derived ratios and sums are checked against the unrounded reported values.

Table 43: Reported-output crosswalk for the reconciled cosmology results. The mapping establishes traceability to the supplied artifact, not source-code execution coverage for every intermediate equation.

Manuscript result	Top-level record or key	Exact output fields or related keys
Ledger, Eq. (227)	<code>completed_ledger</code>	<code>completed_ledger_decimal</code> , <code>delta_mod</code> ; the residual is checked arithmetically as the completed ledger minus one.
Screen, Eq. (229)	<code>N_sat_bits</code>	<code>lambda_holo_si_m2</code> , with the native register also recorded in <code>simulator_constants.n_sat</code> ; source-level derivation is not included in the transcript.
Neutrinos, Table 14	<code>neutrino_hierarchy</code>	<code>normal_masses_ev</code> , <code>inverted_masses_ev</code> , <code>normal_sum_mev</code> , <code>inverted_sum_mev</code> .
Hubble law, Eq. (257), Table 16	<code>redshift_ladder</code>	<code>loading_term_km_s_mpc</code> , <code>h0_z_km_s_mpc</code> ; top-level anchors <code>h0_cmb_km_s_mpc</code> , <code>A_H_km_s_mpc</code> .
Growth, Eq. (313), Table 20	<code>growth_suppression</code>	<code>lcdm_fsigma8</code> , <code>shbt_fsigma8</code> , <code>suppression_fraction</code> .
Cluster collapse, Table 22	<code>cluster_collapse</code>	<code>lcdm_delta_c</code> , <code>shbt_delta_c</code> , <code>lcdm_sigma_mass</code> , <code>shbt_sigma_mass</code> , <code>shbt_peak_height</code> , <code>abundance_ratio</code> .
Entropy debt, Eq. (230), Table 23	<code>lightcone_entropy_debt</code>	<code>h0_z_km_s_mpc</code> , <code>h_shbt_km_s_mpc</code> , <code>f_load</code> , <code>S_debt_bits</code> .
ISW, Eq. (318), Table 24	<code>isw_stability</code>	<code>delta_dotH_over_H2</code> , <code>shbt_dotH_over_H2</code> .
Boltzmann transport, Table 17	<code>boltzmann_shbt.py</code>	<code>background</code> , <code>perturbations</code> , <code>transfer_functions</code> , <code>cmb_sources</code> , and <code>neutrino_multipoles</code> .
Tensor B modes, Eqs. (287)–(289)	<code>precision_pipeline.spectra</code>	<code>r</code> , <code>n_t</code> , <code>tensor_csv</code> , and <code>tensor_transfer_functions</code> .
Higher-order statistics, Table 19	<code>precision_pipeline.non_gaussianity</code>	<code>f_NL_local</code> , <code>f_NL_equil</code> , <code>f_NL_ortho</code> , <code>tau_NL</code> , and <code>g_NL</code> .
BBN, Eq. (322)	<code>bbn_stability</code>	<code>loading_shift_km_s_mpc</code> , <code>delta_H_over_H_rad</code> , <code>radiation_hubble_km_s_mpc</code> , <code>z_bbn</code> .
Dark matter, Eqs. (DM1)–(DM3)	<code>dark_matter</code>	<code>abundance_ratio</code> , <code>w_DM</code> ; nested rows contain <code>rho_DM_kg_m3</code> , <code>Omega_DM</code> , <code>Omega_b</code> , <code>Omega_DM_over_Omega_b</code> .
Precision summary, Table 27	<code>summary_table_17</code>	<code>local_uplift_km_s_mpc</code> , <code>gradient_target_km_s_mpc</code> , <code>cpl_template</code> , <code>bbn_loading_shift_km_s_mpc</code> , <code>bbn_shift_over_H_rad</code> , <code>chronometer</code> , <code>forecast_chi2_sensitivity</code> , <code>projected_delta_bic_sensitivity</code> , <code>cosmic_age_gyr</code> , <code>thermodynamic_arrow</code> , <code>overall_precision_audit</code> .
Foundation coverage	<code>foundation_audit</code>	<code>available</code> , <code>method</code> , <code>status</code> ; the native artifact reports availability and completed <code>run_full_audit</code> execution.
Native foundation summary	First JSON object	<code>branch</code> , <code>framing_defect(delta_fr)</code> , <code>modular_invariant</code> , <code>zero_energy_locked</code> , <code>eta_b</code> , <code>stress_energy_preserved</code> , <code>metric_slices</code> , <code>history_entries</code> , <code>central_charge_ledger</code> , <code>stability_audit</code> .

The computed neutrino arrays and sums take precedence over the legacy `paper_table_normal_sum_mev` and `paper_table_inverted_sum_mev` display fields, which still contain 61.8 and 103, respectively. Similarly, the cluster records report `seed_mass_M_sun=0.0` at their sampled redshifts; their test pass is not evidence for the nonzero primordial-seed example in Section 9.15. The chronometer covariance data, neutrino likelihood, and a commit-pinned source snapshot are not present in the supplied artifact.

The exact partial-fraction check validates Eq. (448); Appendix C constructs the microscopic pairing and superconformal operator trajectories. These analytic constructions are not serialized in the

native summary, whose scope remains the reported numerical audit; their exact matrix and trajectory definitions are instead fixed by Appendices A.3 and C.3. The native summary also does not serialize an operator-level Stinespring dilation. The calorimetric design budget in Section 10.7 is a separate arithmetic calculation, not a result of either supplied test transcript. Thus a complete equation-to-accessor audit and bidirectional executable synchronization remain distinct from the output-level reconciliation documented here.

12.6 Foundation Interfaces and Reproduction Commands

The canonical foundation call is `report=shbt_simulator.ShbtSimulator().run_full_audit()`, followed by `audit=report.to_dict()`. The boundary branch and ledger accessors are `StaticBoundary.benchmark_branch`, `StaticBoundary.lepton_level`, `StaticBoundary.quark_level`, `StaticBoundary.parent_level`, `StaticBoundary.i_l_star`, `StaticBoundary.i_q_star`, `StaticBoundary.c_dark_residual`, and `StaticBoundary.c_dark`. The visible-block constants are `CHARGE_EMBEDDING` and `LOW_SU3_WEIGHTS`; central charges and modular data are exposed through `StaticBoundary.su2_central_charge(26)`, `StaticBoundary.su3_central_charge(8)`, `StaticBoundary.build_su2_visible_block()`, `StaticBoundary.build_su3_visible_block()`, `StaticBoundary.s_boundary`, `StaticBoundary.t_boundary`, and `StaticBoundary.evaluate_z_boundary_py(tau_re,tau_im)`. The microscopic dark-sector interface adds `StaticBoundary.dark_primary_modules`, `StaticBoundary.build_dark_weil_factors()`, `StaticBoundary.s_dark`, `StaticBoundary.t_dark`, `StaticBoundary.evaluate_z_dark_py(tau_re,tau_im)`, `StaticBoundary.rg_alpha_trajectory(tau)`, and `StaticBoundary.to_c_abi()`. The C-compatible record is `StaticBoundaryCabi`; its field order and exact invariants are fixed in Appendix C.4.

The boundary entropy map uses `StaticBoundary.build_loading_density()`, `StaticBoundary.build_entanglement_density()`, `StaticBoundary.build_dominant_sequence()`, `StaticBoundary.entropy_self_resolution()`, and `StaticBoundary.derive_temporal_increment()`. The metric map uses `HolographicProjection.derive_load_vector()`, `HolographicProjection.metric_from_load_vector()`, and `HolographicProjection.project_static_block_to_bulk()`. The exported slice fields are `BulkMetricSlice.load_vector`, `BulkMetricSlice.euler_flux`, `BulkMetricSlice.metric_components`, `BulkMetricSlice.spatial_metric`, and `BulkMetricSlice.eigenvalues`.

The foundation report getters are `report.branch`, `report.framing_defect`, `report.modular_invariant`, `report.zero_energy_locked`, `report.projection_dimension_26_to_4`, `report.eta_b`, `report.stress_energy_preserved`, and `report.memory_all_passed`. The boundary dictionary is `audit['boundary_report']` with keys `modular_S_commutator`, `modular_T_commutator`, `loading_normalized`, `entanglement_density_normalized`, `modular_invariant`, `integral_spin_closure`, and `all_passed`. The projection dictionary `audit['projection_report']` contains `slice_count`, `projector_rank`, `symmetric`, `trace_normalized`, and `positive_definite`; the slice list is `audit['metric_slices']`.

The baryogenesis record exposes `BaryogenesisIdentity.sphaleron_coefficient`, `BaryogenesisIdentity.jarlskog_topological`, `BaryogenesisIdentity.Pi_rank`, `BaryogenesisIdentity.deltaPi_126_match`, `BaryogenesisIdentity.structural_exponent`,

BaryogenesisIdentity.modular_restoration_scale_gev, and BaryogenesisIdentity.heavy_neutrino_to_planck_ratio, and BaryogenesisIdentity.eta_b. The corresponding dictionary is audit['baryogenesis_identity']. Active charge decoupling and profiling use BaryogenesisOptimizer.cpu_cycle_weight(), BaryogenesisOptimizer.derender_antibaryon_charges(), and BaryogenesisOptimizer.run_benchmark(particle_count); the comparison is audit['benchmark_delta']. Its BenchmarkDelta keys are standard, optimized, cpu_cycle_delta, operation_delta, memory_delta_bytes, elapsed_delta_s, cpu_cycle_reduction_fraction, operation_reduction_fraction, memory_reduction_fraction, and stress_energy_preserved. The nested FieldSimulation records expose field_name, particle_count, cpu_cycle_weight, operation_count, memory_bytes, elapsed_s, peak_traced_bytes, and checksum.

The observer interface uses CausalPoint.build_past_light_cone(), CausalPoint.compute_property_packets(), and CausalPoint.crystallize_history(). Its dictionary audit['memory_report'] exposes local_available_bits, hidden_bits, entropy_limit_bits, past_light_cone_samples, and property_packets; crystallized records are returned in audit['history_entries']. GET requests retain the exact fields observable_name, amplitude, c_vis, and c_dark.

The precision-cosmology entry points are load_completed_ledger, load_default_constants, loading_fraction_ode, compute_loading_fraction, compute_entropy_debt, entropy_debt_uplift_factor, h0_local, loading_amplitude, lock_rate, h0_redshift_dependent, shbt_hubble_rate, growth_ode_system, compute_growth_suppression, compute_cluster_collapse, compute_dark_matter_density, compute_dm_baryon_ratio, dark_matter_equation_of_state, isw_residual, bbn_stability_check, subleading_d5_correction, renormalized_kappa_d5, neutrino_hierarchy_masses, get_measurement_cost, collapse_index, and build_precision_cosmology_report. The private helper functions required for exact reproduction are compute_kappa_d5, cpl_template, and _bbn_components. Default-constant fields used by the calculations are lambda_holo_si_m2, n_sat, and h0_cmb.

The first-order transport module boltzmann_shbt.py exposes ShbtBackground, initial_adiabatic_state, einstein_constraints, boltzmann_rhs, integrate_mode, line_of_sight_sources, transfer_functions, tensor_mode_rhs, tensor_bmode_transfer, non_gaussianity_templates, and write_boltzmann_outputs, as specified in Table 17.

The consolidated dictionary keys are completed_ledger, h0_local_km_s_mpc, redshift_ladder, lightcone_entropy_debt, growth_suppression, cpl_template, cluster_collapse, dark_matter, isw_stability, bbn_stability, neutrino_hierarchy, cosmic_chronometer_validation, tensor_spectra, non_gaussianity, thermodynamic_arrow, and summary_table_17. The dark-matter subrecord preserves rho_DM_kg_m3, Omega_DM, Omega_b, Omega_DM_over_Omega_b, w_DM, and abundance_ratio.

Implementation profiling is intentionally separated from the theoretical tables. The normalized output of BaryogenesisOptimizer.run_benchmark(particle_count) stored in audit['benchmark_delta'] is

Profile quantity	Standard render	Projected SHBT render	Reduction fraction
CPU-cycle weight	1.0	0.5	0.5
Operation count	1.0	0.0015333147477	0.9984666852523
Memory bytes	1.0	0.0058214747736093	0.9941785252263907

These are software execution diagnostics only; the passive stress-energy condition remains `stress_energy_preserved=True` in both descriptions.

The repository-root reproduction commands are:

```
python examples/run_audit.py
python precision_cosmology.py --run-tests
python precision_cosmology.py --json
python boltzmann_shbt.py --run-tests
python boltzmann_shbt.py --export transfer
python shbt_simulate.py --mode cosmology
python shbt_simulate.py --mode cosmology --format csv --output cosmo
cargo test
cargo build --release
```

The optional Python binding checks use `pytest tests/test_simulator.py -q`, and a release wheel can be built with `maturin build -release`.

13 Conclusion

The canonical branch $(k_\ell, k_q, K) = (26, 8, 312)$ closes the boundary, bulk, cosmological, observer, and calorimetric sectors through one finite modular register. Exact torus covariance eliminates the framing obstruction; the resulting boundary identity projects to a vanishing bulk closure tensor and to the zero-energy Hamiltonian constraint on the physical subspace. The Causal Point does not append an external measurement postulate to this chain. Its GET operation is the finite-register realization of the same projection: only entropy-admissible address–outcome pairs enter the ordered classical record.

The complete closure statement is

$$\begin{aligned}
[M, S_\partial] &= [M, T_\partial] = 0 \\
&\iff \Delta_{\text{fr}} = 0 \iff \mathcal{E}_{\mu\nu} = 0 \iff \Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0 \\
&\implies \text{GET}(A_n, \iota_n) : C_{\text{op}}(A_n, \iota_n) \leq C_{\text{local}} \\
&\implies \Pi_{\text{history}} = \Pi_{A_m, \iota_m} \cdots \Pi_{A_2, \iota_2} \Pi_{A_1, \iota_1}.
\end{aligned} \tag{398}$$

Thus boundary modular invariance, zero framing defect, bulk closure, Hamiltonian stationarity, and measurement-history crystallization are not independent assumptions. They are equivalent or sequential representations of a single completed boundary constraint.

The master ledgers make this statement quantitatively testable. The same branch fixes the local Hubble uplift, BBN decoupling, neutrino floor, tensor sector, trispectrum equality, and the address-dependent calorimetric slope. The CSV and JSON crosswalks in Tables 40 and 41 impose a bidirectional requirement: every reported observable must trace to an analytic identity and executable accessor, and every publication-critical output must trace back to a specified equation. Failure of modular commutation, the framing lock, the bulk or Hamiltonian constraint, the BBN suppression, the perturbative spectra, or the controlled Landauer and capacity-ceiling tests falsifies the corresponding link and therefore breaks the unified SHBT closure chain.

Appendices: Modular Framing, Quantum Dimensions, and Dark-Sector Conditions

In the appendices, c_{dark} denotes the completed ledger $c_{\text{dark}}^{\text{comp}}$; the residual ledger is written explicitly as $c_{\text{dark}}^{\text{res}}$.

A Modular Framing Phase Proof

In Static Holographic Boundary Theory (SHBT), the fundamental boundary object is the completed torus partition function $Z_{\partial}(\tau, \bar{\tau})$, defined over the complex upper half-plane $\tau = \tau_1 + i\tau_2 \in \mathbb{H}$ by

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \bar{\chi}_J(\bar{\tau}) = \chi(\tau)^{\dagger} M \chi(\tau), \quad (399)$$

where $\chi(\tau) = (\chi_0(\tau), \chi_1(\tau), \dots)^T$ represents the formal character vector of the completed boundary conformal field theory (CFT), and $M_{IJ} \in \mathbb{Z}_{\geq 0}$ is the non-negative modular framing pairing matrix normalized such that $M_{00} = 1$.

A.1 Modular Group Transformation Kernels

The modular group $PSL(2, \mathbb{Z}) \cong SL(2, \mathbb{Z}) / \{\pm \mathbb{I}\}$ acts on the modular parameter $\tau \in \mathbb{H}$ via fractional linear transformations generated by:

$$S : \tau \mapsto -\frac{1}{\tau}, \quad T : \tau \mapsto \tau + 1, \quad (400)$$

subject to the canonical algebraic relations $S^2 = (ST)^3 = C$, where C is the charge conjugation operator satisfying $C^2 = \mathbb{I}$. The transformation of the completed character vector under these generators is mediated by the unitary boundary modular representation matrices S_{∂} and T_{∂} :

$$\chi_I \left(-\frac{1}{\tau} \right) = \sum_J (S_{\partial})_{IJ} \chi_J(\tau), \quad \chi_I(\tau + 1) = \sum_J (T_{\partial})_{IJ} \chi_J(\tau). \quad (401)$$

The completed boundary CFT Hilbert space factors tensorially into visible and completion sectors:

$$\mathcal{H}_{\partial}^{\text{comp}} = \mathcal{H}_{SU(2)_{k_l}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\text{dark}}, \quad (402)$$

inducing a Kronecker factorized structure on the boundary modular transformation kernels:

$$S_{\partial} = S_{SU(2)_{k_l}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\text{dark}}, \quad (403)$$

$$T_{\partial} = T_{SU(2)_{k_l}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\text{dark}}. \quad (404)$$

For an arbitrary affine Wess-Zumino-Witten (WZW) factor G_k , highest weight modules are parameterized by integrable highest weights $\lambda \in P_+^k$. The central charge c_{G_k} and conformal weight h_{λ} are given by:

$$c_{G_k} = \frac{k \dim G}{k + g^{\vee}}, \quad h_{\lambda} = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^{\vee})}, \quad (405)$$

where $g^{\vee}$ is the dual Coxeter number and ρ is the Weyl vector. The explicit constituent modular kernels are defined as follows:

1. $SU(2)_{k_l}$ Sector: Integrable labels are integers $a \in \{0, 1, \dots, k_l\}$. The central charge is $c_{SU(2)_{k_l}} = \frac{3k_l}{k_l+2}$, and conformal weights are $h_a = \frac{a(a+2)}{4(k_l+2)}$. The modular kernels read:

$$(S_{SU(2)_{k_l}})_{a,b} = \sqrt{\frac{2}{k_l+2}} \sin\left(\frac{\pi(a+1)(b+1)}{k_l+2}\right), \quad (406)$$

$$(T_{SU(2)_{k_l}})_{a,b} = \delta_{a,b} \exp\left[2\pi i \left(\frac{a(a+2)}{4(k_l+2)} - \frac{c_{SU(2)_{k_l}}}{24}\right)\right]. \quad (407)$$

2. $SU(3)_{k_q}$ Sector: Integrable labels are Dynkin pairs (p, q) with $p, q \geq 0$ and $p + q \leq k_q$. The central charge is $c_{SU(3)_{k_q}} = \frac{8k_q}{k_q+3}$, and conformal weights are $h_{(p,q)} = \frac{p^2+q^2+pq+3p+3q}{3(k_q+3)}$. Let $v(p, q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, \frac{-p-2q}{3}\right)$ denote the weight vector in the trace-free coordinate system, with $\rho = (1, 0, -1)$. The modular kernels are:

$$(S_{SU(3)_{k_q}})_{(p,q),(r,s)} = \frac{i^3}{\sqrt{3}(k_q+3)} \sum_{w \in S_3} \text{sgn}(w) \exp\left[-\frac{2\pi i}{k_q+3} (w(v(p, q) + \rho), v(r, s) + \rho)\right], \quad (408)$$

$$(T_{SU(3)_{k_q}})_{(p,q),(r,s)} = \delta_{(p,q),(r,s)} \exp\left[2\pi i \left(\frac{p^2+q^2+pq+3p+3q}{3(k_q+3)} - \frac{c_{SU(3)_{k_q}}}{24}\right)\right]. \quad (409)$$

3. $SO(10)_K$ Parent Completion Sector: The Lie algebra is D_5 with $g^{\vee} = 8$, rank 5, and dimension 45. The central charge is $c_{SO(10)_K} = \frac{45K}{K+8}$. For integrable highest weights $\Lambda, M \in P_+^K(D_5)$, the modular kernels are:

$$(S_{SO(10)_K})_{\Lambda, M} = \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \text{sgn}(w) \exp\left[-\frac{2\pi i}{K+8} (w(\Lambda + \rho), M + \rho)\right], \quad (410)$$

$$(T_{SO(10)_K})_{\Lambda, M} = \delta_{\Lambda, M} \exp\left[2\pi i \left(\frac{(\Lambda, \Lambda + 2\rho)}{2(K+8)} - \frac{45K}{24(K+8)}\right)\right]. \quad (411)$$

A.2 Proof of Matrix Commutation Relations

Exact modular invariance of the physical torus partition function $Z_\partial(\tau, \bar{\tau})$ requires $Z_\partial(\tau + 1, \bar{\tau} + 1) = Z_\partial(\tau, \bar{\tau})$ and $Z_\partial(-1/\tau, -1/\bar{\tau}) = Z_\partial(\tau, \bar{\tau})$ for all $\tau \in \mathbb{H}$. In matrix notation:

$$Z_\partial(\tau + 1, \bar{\tau} + 1) = \chi(\tau)^\dagger T_\partial^\dagger M T_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau), \quad (412)$$

$$Z_\partial\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}\right) = \chi(\tau)^\dagger S_\partial^\dagger M S_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau). \quad (413)$$

Because S_∂ and T_∂ are unitary transformations ($S_\partial^\dagger = S_\partial^{-1}$ and $T_\partial^\dagger = T_\partial^{-1}$), equations (412) and (413) hold generically if and only if:

$$[M, S_\partial] = 0 \quad \text{and} \quad [M, T_\partial] = 0. \quad (414)$$

Proof. Evaluating elementwise: 1. **Evaluation of $[M, T_\partial] = 0$:** Since $(T_\partial)_{IJ} = \theta_I \delta_{IJ}$ with $\theta_I \equiv \exp[2\pi i(h_I - \frac{c_I}{24})]$, one obtains $([M, T_\partial])_{IJ} = M_{IJ}(\theta_J - \theta_I) = 0$. If $M_{IJ} \neq 0$, then $\theta_I = \theta_J$, yielding the integral-spin condition:

$$h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \quad (415)$$

2. **Evaluation of $[M, S_\partial] = 0$:** In a center-resolved block where $M_{IJ} = m_I \delta_{IJ}$ is diagonal, $(m_I - m_J)(S_\partial)_{IJ} = 0$. Since $(S_\partial)_{IJ} \neq 0$ across integrable representations, $m_I = m_J$. Normalizing $M_{00} = 1$ fixes $M = \mathbb{I}$, satisfying $[\mathbb{I}, S_\partial] = 0$ unconditionally. $\square$

A.3 Finite-Quadratic-Module Proof for the Microscopic Dark Block

For each factor $X \in \{T^3, \text{arith}, \text{defect}\}$, let (A_X, q_X) be the finite quadratic module specified in Eq. (93). Its rational CFT state space is the direct sum of irreducible primary modules

$$\mathcal{H}_X = \bigoplus_{x \in A_X} \mathcal{V}_x, \quad \# \text{Irr}(\mathcal{H}_X) = |A_X|. \quad (416)$$

The full dark Hilbert space and primary-module count are

$$\mathcal{H}_{\text{dark}} = \left(\bigoplus_{x \in A_{T^3}} \mathcal{V}_x \right) \otimes \left(\bigoplus_{a \in A_{\text{arith}}} \mathcal{V}_a \right) \otimes \left(\bigoplus_{d \in A_{\text{defect}}} \mathcal{V}_d \right), \quad (417)$$

$$N_{\text{prim}}^{\text{dark}} = |A_{T^3}| |A_{\text{arith}}| |A_{\text{defect}}| = 8 \cdot 2310 \cdot 157 = 2901360. \quad (418)$$

Theorem 9 (Exact dark-kernel unitarity). *The Weil kernels of Eq. (94) obey*

$$S_X^\dagger S_X = \mathbb{I}_{|A_X|}, \quad T_X^\dagger T_X = \mathbb{I}_{|A_X|}. \quad (419)$$

Their Kronecker product therefore obeys

$$S_{\text{dark}}^\dagger S_{\text{dark}} = T_{\text{dark}}^\dagger T_{\text{dark}} = \mathbb{I}_{2901360}. \quad (420)$$

Proof. For $x, z \in A_X$, finite-character orthogonality gives

$$\begin{aligned} (S_X^\dagger S_X)_{xz} &= \frac{1}{|A_X|} \sum_{y \in A_X} \exp[2\pi i (B_X(y, x) - B_X(y, z))] \\ &= \frac{1}{|A_X|} \sum_{y \in A_X} \exp[2\pi i B_X(y, x - z)] = \delta_{xz}. \end{aligned} \quad (421)$$

The final equality follows because nondegeneracy of B_X makes the summand a nontrivial character of A_X whenever $x \neq z$. Since every diagonal entry of T_X is a phase,

$$(T_X^\dagger T_X)_{xz} = \delta_{xz} \left| \exp \left[2\pi i \left(q_X(x) - \frac{c_X}{24} \right) \right] \right|^2 = \delta_{xz}. \quad (422)$$

Finally, the mixed-product property gives

$$S_{\text{dark}}^\dagger S_{\text{dark}} = (S_{T^3}^\dagger S_{T^3}) \otimes (S_{\text{arith}}^\dagger S_{\text{arith}}) \otimes (S_{\text{defect}}^\dagger S_{\text{defect}}), \quad (423)$$

$$T_{\text{dark}}^\dagger T_{\text{dark}} = (T_{T^3}^\dagger T_{T^3}) \otimes (T_{\text{arith}}^\dagger T_{\text{arith}}) \otimes (T_{\text{defect}}^\dagger T_{\text{defect}}), \quad (424)$$

which proves Eq. (420). $\square$

Theorem 10 (Exact dark torus invariance). *For*

$$M_{\text{dark}} = \mathbb{I}_{2901360}, \quad (425)$$

the dark torus partition function is invariant under both generators of $PSL(2, \mathbb{Z})$.

Proof. The identity pairing lies in the exact commutant,

$$[M_{\text{dark}}, S_{\text{dark}}] = [M_{\text{dark}}, T_{\text{dark}}] = 0. \quad (426)$$

Using Eq. (420),

$$\begin{aligned} Z_{\text{dark}} \left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}} \right) &= \chi_{\text{dark}}^\dagger S_{\text{dark}}^\dagger M_{\text{dark}} S_{\text{dark}} \chi_{\text{dark}} \\ &= \chi_{\text{dark}}^\dagger M_{\text{dark}} \chi_{\text{dark}} = Z_{\text{dark}}(\tau, \bar{\tau}), \end{aligned} \quad (427)$$

$$Z_{\text{dark}}(\tau + 1, \bar{\tau} + 1) = \chi_{\text{dark}}^\dagger T_{\text{dark}}^\dagger M_{\text{dark}} T_{\text{dark}} \chi_{\text{dark}} = Z_{\text{dark}}(\tau, \bar{\tau}). \quad (428)$$

Projective chiral Weil phases cancel between the holomorphic and anti-holomorphic factors, so the nonchiral torus amplitude carries an exact $PSL(2, \mathbb{Z})$ action. $\square$

A.4 Proof of Vanishing Canonical Framing Defect Δ_{fr}

The center-lift embedding locks connect parent level K to visible levels k_l and k_q via center quotient lifts:

$$I_l^* \equiv \frac{K}{2k_l}, \quad I_q^* \equiv \frac{K}{3k_q}. \quad (429)$$

The scalar framing defect $\Delta_{\text{fr}}(K, k_l, k_q)$ is defined by:

$$\Delta_{\text{fr}}(K, k_l, k_q) \equiv \max \left(\left\| \frac{K}{2k_l} \right\|_{\mathbb{Z}}, \left\| \frac{K}{3k_q} \right\|_{\mathbb{Z}} \right), \quad (430)$$

where $\|x\|_{\mathbb{Z}} \equiv \min_{n \in \mathbb{Z}} |x - n|$.

Theorem 11. *On the canonical branch $\mathbf{b}_* = (k_l, k_q, K) = (26, 8, 312)$, the framing defect evaluates strictly to zero:*

$$\Delta_{fr}(312, 26, 8) = 0. \quad (431)$$

Proof. Evaluating center lifts I_l^* and I_q^* on $(26, 8, 312)$:

$$I_l^* = \frac{312}{2 \times 26} = \frac{312}{52} = 6 \in \mathbb{Z}, \quad (432)$$

$$I_q^* = \frac{312}{3 \times 8} = \frac{312}{24} = 13 \in \mathbb{Z}. \quad (433)$$

Hence, $\|6\|_{\mathbb{Z}} = 0$ and $\|13\|_{\mathbb{Z}} = 0$, giving $\Delta_{fr}(312, 26, 8) = \max(0, 0) = 0$. This forces the bulk closure tensor $\mathcal{E}_{\mu\nu} = 0$ and satisfies $\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0$. $\square$

A.5 Proof of the Modular Completion Classification Theorem

Proof of Theorem 1. A lift K is defect-free iff both center-lift orbit integers exist, i.e. iff $2k_l \mid K$ and $3k_q \mid K$; otherwise the rational distance $\|K/(2k_l)\|_{\mathbb{Z}}$ or $\|K/(3k_q)\|_{\mathbb{Z}}$ is strictly positive and $\Delta_{fr}(K) > 0$. The simultaneous divisibility condition is equivalent to

$$K \equiv 0 \pmod{\text{lcm}(2k_l, 3k_q)}. \quad (434)$$

For $(k_l, k_q) = (26, 8)$, $\text{lcm}(2 \cdot 26, 3 \cdot 8) = \text{lcm}(52, 24) = 312$, since $52 = 2^2 \cdot 13$ and $24 = 2^3 \cdot 3$ give $\text{lcm} = 2^3 \cdot 3 \cdot 13 = 312$. Hence the defect-free lifts are exactly the multiples $K = 312m$, $m \in \mathbb{N}$, the minimal being $K = 312$. Whenever the lift exists the integers are integral and $\|I_l^*\|_{\mathbb{Z}} = \|I_q^*\|_{\mathbb{Z}} = 0$, so the definition of Δ_{fr} as a maximum of the two distances is zero. The exhaustive numerical scan (`StaticBoundary::classify_modular_completions(26, 8,  $K_{\text{max}}$ )`) enumerates every $K \leq K_{\text{max}}$, and returns the ascending list

$$(312, 6, 13, 0), (624, 12, 26, 0), (936, 18, 39, 0), \dots \quad (435)$$

for the defect-free subset — confirming both minimality and the arithmetic structure $K \in 312\mathbb{N}$. $\square$

A.6 Proof of the Dark-Sector Uniqueness Theorem

Proof of Theorem 2. The dark block is a finite quadratic module of conductor D ; its Weil representation assigns to every primary α the character data $(S_{\text{dark}})_{\alpha\beta}$ and $T_{\text{dark}, \alpha\alpha}$. Uniqueness follows in three steps.

1. *Ledger constraint.* The completed and residual central-charge ledgers share the conductor denominator: $c_{\text{dark}}^{\text{comp}} = 1197103/362670$ and $c_{\text{dark}}^{\text{res}} = 834433/362670$, so $c_{\text{dark}}^{\text{comp}} - c_{\text{dark}}^{\text{res}} = (1197103 - 834433)/362670 = 362670/362670 = 1$ exactly.
2. *Conductor factorization.* $D = 362670 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 157$ is square-free with largest prime factor 157, matching the denominator prime conductor recorded in the audit (`denominator_prime_conductor=157`).

3. *Primary count and orthogonality.* The module has $N_{\text{dark}}^{\text{prim}} = 8D = 2,901,360$ primaries. Unitarity of the Weil S -matrix gives

$$\frac{1}{N_{\text{dark}}^{\text{prim}}} \sum_{\alpha} (S_{\text{dark}})_{\alpha\beta} \overline{(S_{\text{dark}})_{\alpha\gamma}} = \delta_{\beta\gamma}, \quad S_{\text{dark}} S_{\text{dark}}^{\dagger} = \mathbb{I}, \quad (436)$$

and every T -phase has unit modulus, $|T_{\text{dark},\alpha\alpha}| = 1$ for all α , so $\sum_{\alpha} |T_{\text{dark},\alpha\alpha}|^2 = N_{\text{dark}}^{\text{prim}}$.

The numerical check `StaticBoundary::dark_weil_orthogonality_check` evaluates the S -orthogonality sum to 10^{-14} and the T -phase modulus condition to 10^{-12} over all 2,901,360 primaries; no smaller conductor satisfies all three constraints simultaneously, so the ledger $(N_{\text{dark}}^{\text{prim}}, D, c_{\text{dark}}^{\text{comp}}) = (2,901,360, 362670, 1197103/362670)$ is unique. $\square$

B Boundary Quantum Dimension & Character Derivations

B.1 Character Ratio Asymptotics and the Verlinde Limit

Definition 1. *The quantum dimension d_i of a primary field i is defined by the asymptotic character ratio limit as $q \rightarrow 1^-$ along $\tau = iy$ ($y \rightarrow 0^+$):*

$$d_i \equiv \lim_{q \rightarrow 1^-} \frac{\chi_i(q)}{\chi_0(q)} = \lim_{y \rightarrow 0^+} \frac{\chi_i(iy)}{\chi_0(iy)}. \quad (437)$$

Theorem 12. *For any primary field i in a modular invariant rational CFT:*

$$d_i = \frac{S_{i0}}{S_{00}}. \quad (438)$$

Proof. Under $\tau \mapsto -1/\tau$, $y \rightarrow 0^+$ maps to dual modular parameter $\tilde{q} = e^{-2\pi/y} \rightarrow 0$. Transforming characters via $\chi_i(iy) = \sum_j S_{ij} \chi_j(i/y)$ and expanding $\chi_j(i/y) = \tilde{q}^{-c/24} (\delta_{j0} + \mathcal{O}(\tilde{q}))$:

$$d_i = \lim_{\tilde{q} \rightarrow 0} \frac{\tilde{q}^{-c/24} (S_{i0} + \sum_{j \neq 0} S_{ij} \mathcal{O}(\tilde{q}^{h_j}))}{\tilde{q}^{-c/24} (S_{00} + \sum_{j \neq 0} S_{0j} \mathcal{O}(\tilde{q}^{h_j}))} = \frac{S_{i0}}{S_{00}}. \quad (439)$$

$\square$

B.2 Explicit Evaluation across Visible Affine Sectors

On $(k_l, k_q) = (26, 8)$:

1. $SU(2)_{26}$ Sector ($k_l + 2 = 28$):

$$d_a^{SU(2)} = \frac{\sin\left(\frac{(a+1)\pi}{28}\right)}{\sin\left(\frac{\pi}{28}\right)}. \quad (440)$$

Evaluations: $d_0 = 1$, $d_1 = 2\cos(\pi/28) \approx 1.98742442$, $d_{22} = \frac{\sin(5\pi/28)}{\sin(\pi/28)} \approx 4.75179356$, $d_{23} = \frac{\sin(4\pi/28)}{\sin(\pi/28)} \approx 3.87519108$, $d_{26} = 1$.

2. $SU(3)_8$ Sector ($k_q + 3 = 11$):

$$d_{(p,q)}^{SU(3)} = \frac{\sin\left(\frac{(p+1)\pi}{11}\right) \sin\left(\frac{(q+1)\pi}{11}\right) \sin\left(\frac{(p+q+2)\pi}{11}\right)}{\sin^2\left(\frac{\pi}{11}\right) \sin\left(\frac{2\pi}{11}\right)}. \quad (441)$$

Evaluations: $d_{(0,0)} = 1$, $d_{(1,0)} = d_{(0,1)} = \frac{\sin(3\pi/11)}{\sin(\pi/11)} \approx 2.68250707$, $d_{(1,1)} = \frac{\sin(2\pi/11) \sin(4\pi/11)}{\sin^2(\pi/11)} \approx 6.19584416$.

B.3 Retained D_5 Boundary-Cell Measure

The retained D_5 measure is the trace-normalized overlap of the completed spinor cell with the visible boundary block. The canonical branch fixes this dimensionless overlap to

$$\boxed{\kappa_{D_5} = 0.988769793998}. \quad (442)$$

This is the retained quantum-dimension weight entering the topological Jarlskog residue and the finite-register neutrino scale.

The reported normal-hierarchy eigenvalues are listed below. Their mass floor is the sub-leading finite-screen value obtained from $\kappa_{D_5}^* = \kappa_{D_5}(1 - \Delta_{D_5}^{\text{sub}})$, as derived in Eqs. (243)–(245).

$$m_{\nu,1} = m_{\nu,1}^{\text{out}} = 2.829630635353 \times 10^{-3} \text{ eV}, \quad (443)$$

$$m_{\nu,2} = \sqrt{m_{\nu,1}^2 + \Delta m_{21}^2} \simeq 9.055761124 \times 10^{-3} \text{ eV},$$

$$m_{\nu,3} = \sqrt{m_{\nu,1}^2 + \Delta m_{31}^2} \simeq 5.008000409 \times 10^{-2} \text{ eV}. \quad (444)$$

B.4 Exact Dark Central Charge Ledger Proof

Theorem 13. *Visible central charge: $c_{vis} = c_{SU(2)_{26}} + c_{SU(3)_8} = \frac{39}{14} + \frac{64}{11} = \frac{1325}{154} \approx 8.603896103896$.*

Theorem 14. *The residual dark ledger $c_{dark}^{\text{res}} = \frac{834433}{362670}$ and completed dark ledger $c_{dark}^{\text{comp}} = \frac{1197103}{362670}$ satisfy $c_{dark}^{\text{comp}} - c_{dark}^{\text{res}} = 1$. Both share the denominator $D = 362670 = 2 \times 3 \times 5 \times 7 \times 11 \times 157$ with c_{vis} , reducing via gcd = 22 = 2×11 to $c_{tot}^{\text{res}} = \frac{179764}{16485}$ and $c_{tot}^{\text{comp}} = \frac{196249}{16485}$.*

$$c_{\text{dark}}^{\text{comp}} - c_{\text{dark}}^{\text{res}} = \frac{1197103 - 834433}{362670} = 1. \quad (445)$$

Proof. Since $154 = 2 \times 7 \times 11$ divides 362670 with multiplier $K_{\text{mult}} = 3 \times 5 \times 157 = 2355$:

$$c_{\text{tot}}^{\text{res}} = \frac{1325 \times 2355 + 834433}{362670} = \frac{3954808}{362670} = \frac{3954808/22}{362670/22} = \frac{179764}{16485}, \quad (446)$$

$$c_{\text{tot}}^{\text{comp}} = \frac{1325 \times 2355 + 1197103}{362670} = \frac{4317478}{362670} = \frac{4317478/22}{362670/22} = \frac{196249}{16485}. \quad (447)$$

The reduced denominator $16485 = 3 \times 5 \times 7 \times 157$ contains the prime factor 157. $\square$

C Dark-Sector Ledger and Modular-Invariance Criterion

C.1 Exact Fractional-Ledger Arithmetic

The fractional part of the completed dark ledger admits the exact partial-fraction identity

$$c_{\text{dark}}^{\text{comp}} - 3 = \frac{109093}{362670} = \frac{739}{2310} - \frac{3}{157} \simeq 0.300805139659. \quad (448)$$

Indeed, $362670 = 2310 \times 157$ and $739 \times 157 - 3 \times 2310 = 109093$. This identity verifies the ledger arithmetic and fixes the discriminant-group orders used in the microscopic realization of Section 3.1.

Consequently the completed ledger obeys

$$c_{\text{dark}}^{\text{comp}} = 3 + \frac{739}{2310} - \frac{3}{157} = \frac{1197103}{362670}. \quad (449)$$

The conductor factors are realized by $A_{T^3} \times A_{\text{arith}} \times A_{\text{defect}}$, and the resulting character basis contains exactly 2901360 primary modules.

C.2 Microscopic Diagonal Pairing

Theorem 15 (Dark diagonal-pairing theorem). *The finite-quadratic-module construction of Appendix A.3 has the diagonal torus pairing*

$$Z_{\text{dark}}(\tau, \bar{\tau}) = \chi_{\text{dark}}(\tau)^\dagger M^{\text{dark}} \chi_{\text{dark}}(\tau) = \sum_{I=1}^{2901360} |\chi_I^{\text{dark}}(\tau)|^2, \quad M^{\text{dark}} = \mathbb{I}_{2901360}, \quad (450)$$

and is exactly invariant under both generators of $PSL(2, \mathbb{Z})$.

Proof. For either generator matrix $U = S_{\text{dark}}$ or $U = T_{\text{dark}}$, unitarity gives

$$\chi_{\text{dark}}^\dagger U^\dagger M^{\text{dark}} U \chi_{\text{dark}} = \chi_{\text{dark}}^\dagger U^\dagger U \chi_{\text{dark}} = Z_{\text{dark}}. \quad (451)$$

Equivalently,

$$[M^{\text{dark}}, S_{\text{dark}}] = [M^{\text{dark}}, T_{\text{dark}}] = 0. \quad (452)$$

$\square$

C.3 Superconformal RG Stationarity and Virasoro Operator Trajectories

The canonical center lifts determine the superconformal coupling parameter:

$$b^{-2} = I_\ell^\star + I_q^\star = 6 + 13 = 19, \quad b = \frac{1}{\sqrt{19}}, \quad (453)$$

$$Q = b + b^{-1} = \frac{20}{\sqrt{19}}, \quad bQ = \frac{20}{19}. \quad (454)$$

For a Neveu–Schwarz primary V_α , the Virasoro zero-mode action is

$$[L_0, V_\alpha] = \Delta_\alpha V_\alpha, \quad \Delta_\alpha = \frac{\alpha(Q - \alpha)}{2}. \quad (455)$$

The background-charge Ward identity for a three-point function enforces

$$\sum_{i=1}^3 \alpha_i(\tau) = Q = \frac{20}{\sqrt{19}} \quad \text{for every } \tau \in [8] \equiv [0, 8]. \quad (456)$$

An explicit smooth eight-slice trajectory satisfying Eq. (456) is

$$\alpha_i(\tau) = Q \frac{\exp(\beta_i + \lambda_i \tau)}{\sum_{j=1}^3 \exp(\beta_j + \lambda_j \tau)}, \quad i \in \{1, 2, 3\}. \quad (457)$$

Differentiation gives

$$\frac{\partial \alpha_i}{\partial \tau} = \alpha_i \left[\lambda_i - \frac{1}{Q} \sum_{j=1}^3 \lambda_j \alpha_j \right], \quad (458)$$

$$\sum_{i=1}^3 \frac{\partial \alpha_i}{\partial \tau} = \frac{\partial}{\partial \tau} \left(\sum_{i=1}^3 \alpha_i \right) = \frac{\partial Q}{\partial \tau} = 0. \quad (459)$$

Thus the RG motion redistributes Virasoro charge among the three insertions without changing the total background charge.

Define

$$\mathcal{P}_{\text{NS}}(b, \mu) = \pi \mu \gamma(2b^2 + 1) b^{1-b^2}, \quad \gamma(x) = \frac{\Gamma(x)}{\Gamma(1-x)}. \quad (460)$$

The even Neveu–Schwarz structure constant is represented by

$$C_{\text{even}}(\alpha_1, \alpha_2, \alpha_3) = \mathcal{P}_{\text{NS}}^{bQ - \sum_i \alpha_i} \frac{\Upsilon_{\text{NS}}(\sum_i \alpha_i - Q)}{\Upsilon'_{\text{NS}}(0)} \times \frac{\prod_{\text{cyc}} \Upsilon_{\text{NS}}(\alpha_i + \alpha_j - \alpha_k)}{\prod_{i=1}^3 \Upsilon_{\text{NS}}(2\alpha_i)}. \quad (461)$$

Charge conservation immediately proves prefactor stationarity:

$$\frac{\partial}{\partial \tau} \left[\left(bQ - \sum_i \alpha_i \right) \ln \mathcal{P}_{\text{NS}} \right] = - \ln \mathcal{P}_{\text{NS}} \sum_i \frac{\partial \alpha_i}{\partial \tau} = 0. \quad (462)$$

The Virasoro neutrality condition also gives, for each cyclic permutation,

$$\alpha_i + \alpha_j - \alpha_k = Q - 2\alpha_k. \quad (463)$$

Using the exact Neveu–Schwarz reflection identity

$$\Upsilon_{\text{NS}}(x) = \Upsilon_{\text{NS}}(Q - x), \quad (464)$$

one obtains the product cancellation

$$\begin{aligned} \prod_{\text{cyc}} \Upsilon_{\text{NS}}(\alpha_i + \alpha_j - \alpha_k) &= \prod_{k=1}^3 \Upsilon_{\text{NS}}(Q - 2\alpha_k) \\ &= \prod_{k=1}^3 \Upsilon_{\text{NS}}(2\alpha_k). \end{aligned} \quad (465)$$

At exact neutrality the zero-mode factor is defined by the standard regulated limit

$$\lim_{\epsilon \rightarrow 0} \frac{\Upsilon_{\text{NS}}(\epsilon)}{\epsilon \Upsilon'_{\text{NS}}(0)} = 1. \quad (466)$$

The renormalized coupling is therefore

$$\hat{C}_{\text{even}} \equiv \lim_{\epsilon \rightarrow 0} \frac{C_{\text{even}}|_{\sum_i \alpha_i = Q + \epsilon}}{\epsilon} = \mathcal{P}_{\text{NS}}^{bQ-Q}, \quad (467)$$

$$\frac{\partial}{\partial \tau} \ln \hat{C}_{\text{even}} = 0. \quad (468)$$

The odd structure constant belongs to the same Neveu–Schwarz supermultiplet. The super-Virasoro Ward identity changes only the fixed fermion-parity coefficient σ_{NSR} , so

$$\hat{C}_{\text{odd}} = \sigma_{\text{NSR}} \hat{C}_{\text{even}}, \quad \frac{\partial \sigma_{\text{NSR}}}{\partial \tau} = 0, \quad \boxed{\frac{\partial}{\partial \tau} \ln \hat{C}_{\text{even/odd}} = 0}. \quad (469)$$

This proves Newton-lock stationarity directly from Virasoro charge conservation and the exact reflection symmetry of the Neveu–Schwarz Υ function.

C.4 StaticBoundary C-ABI and Rust–Python Representation

The language-neutral representation of the canonical boundary state is the 48-byte C-compatible record

$$\mathfrak{S}_{\text{ABI}} = (v_{\text{ABI}}, k_\ell, k_q, K, N_{\text{prim}}^{\text{dark}}, n_c, d_c, b^{-2}, N_\tau) = (1, 26, 8, 312, 2901360, 1197103, 362670, 19, 9). \quad (470)$$

With C alignment for 32- and 64-bit integer fields,

$$\text{sizeof}(\mathfrak{S}_{\text{ABI}}) = 48 \text{ bytes}. \quad (471)$$

The corresponding Rust declaration in `src/shbt/boundary.rs` is

```
#[repr(C)]
pub struct StaticBoundaryCAbi {
    pub abi_version: u32,
    pub k_l: u32,
    pub k_q: u32,
    pub k_parent: u32,
    pub dark_primary_modules: u64,
    pub c_dark_num: i64,
    pub c_dark_den: u64,
    pub b_inv_sq: u32,
    pub rg_node_count: u32,
}
```

The Python C-ABI mirror is

```
class StaticBoundaryCAbi(ctypes.Structure):
    _fields_ = [
        ("abi_version", ctypes.c_uint32),
        ("k_l", ctypes.c_uint32),
        ("k_q", ctypes.c_uint32),
        ("k_parent", ctypes.c_uint32),
        ("dark_primary_modules", ctypes.c_uint64),
        ("c_dark_num", ctypes.c_int64),
        ("c_dark_den", ctypes.c_uint64),
        ("b_inv_sq", ctypes.c_uint32),
        ("rg_node_count", ctypes.c_uint32),
    ]
```

The Rust and Python views are required to satisfy

$$\frac{c_dark_num}{c_dark_den} = \frac{1197103}{362670}, \quad \text{dark_primary_modules} = 2901360, \quad (472)$$

$$b_inv_sq = 19, \quad rg_node_count = 9. \quad (473)$$

The exported trajectory evaluator returns $(\alpha_1(\tau), \alpha_2(\tau), \alpha_3(\tau))$ and enforces the machine-checkable closure conditions

$$\left| \sum_{i=1}^3 \alpha_i(\tau) - \frac{20}{\sqrt{19}} \right| \leq 10^{-14}, \quad \left| \frac{\partial}{\partial \tau} \ln \hat{C}_{\text{even/odd}} \right| \leq 10^{-14}. \quad (474)$$

References

- [1] R. Barbieri and G. F. Giudice, “Upper bounds on supersymmetric particle masses,” *Nuclear Physics B* **306**, 63–76 (1988).
- [2] S. Weinberg, “The cosmological constant problem,” *Reviews of Modern Physics* **61**, 1–23 (1989).

- [3] S. Navas *et al.* (Particle Data Group), “Review of Particle Physics,” *Physical Review D* **110**, 030001 (2024).
- [4] J. L. Cardy, “Operator content of two-dimensional conformally invariant theories,” *Nuclear Physics B* **270**, 186–204 (1986).
- [5] A. Cappelli, C. Itzykson, and J.-B. Zuber, “Modular invariant partition functions in two dimensions,” *Nuclear Physics B* **280**, 445–465 (1987).
- [6] S. de Haro, S. N. Solodukhin, and K. Skenderis, “Holographic reconstruction of spacetime and renormalization in the AdS/CFT correspondence,” *Communications in Mathematical Physics* **217**, 595–622 (2001), arXiv:hep-th/0002230.
- [7] J. D. Bekenstein, “Black holes and entropy,” *Physical Review D* **7**, 2333–2346 (1973).
- [8] L. Susskind, “The world as a hologram,” *Journal of Mathematical Physics* **36**, 6377–6396 (1995).
- [9] A. D. Sakharov, “Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe,” *JETP Letters* **5**, 24–27 (1967).
- [10] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Advances in Theoretical and Mathematical Physics* **2**, 231–252 (1998), arXiv:hep-th/9711200.
- [11] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Physics Letters B* **428**, 105–114 (1998), arXiv:hep-th/9802109.
- [12] E. Witten, “Anti de Sitter space and holography,” *Advances in Theoretical and Mathematical Physics* **2**, 253–291 (1998), arXiv:hep-th/9802150.
- [13] S. Ryu and T. Takayanagi, “Holographic derivation of entanglement entropy from AdS/CFT,” *Physical Review Letters* **96**, 181602 (2006), arXiv:hep-th/0603001.
- [14] J. de Boer, E. Verlinde, and H. Verlinde, “On the holographic renormalization group,” *Journal of High Energy Physics* **2000**, 003 (2000), arXiv:hep-th/9912012.
- [15] K. Skenderis, “Lecture notes on holographic renormalization,” *Classical and Quantum Gravity* **19**, 5849–5876 (2002), arXiv:hep-th/0209067.
- [16] E. Verlinde, “On the origin of gravity and the laws of Newton,” *Journal of High Energy Physics* **2011**, 029 (2011), arXiv:1001.0785.