

Static Holographic Boundary Theory: Supplementary Material

September 23, 2026

1 Modular Framing Phase Proof

In Static Holographic Boundary Theory (SHBT), the fundamental boundary object is the completed torus partition function $Z_{\partial}(\tau, \bar{\tau})$, defined over the complex upper half-plane $\tau = \tau_1 + i\tau_2 \in \mathbb{H}$ by

$$Z_{\partial}(\tau, \bar{\tau}) = \sum_{I,J} M_{IJ} \chi_I(\tau) \bar{\chi}_J(\bar{\tau}) = \chi(\tau)^{\dagger} M \chi(\tau), \quad (1)$$

where $\chi(\tau) = (\chi_0(\tau), \chi_1(\tau), \dots)^T$ represents the formal character vector of the completed boundary conformal field theory (CFT), and $M_{IJ} \in \mathbb{Z}_{\geq 0}$ is the non-negative modular framing pairing matrix normalized such that $M_{00} = 1$.

1.1 Modular Group Transformation Kernels

The modular group $PSL(2, \mathbb{Z}) \cong SL(2, \mathbb{Z}) / \{\pm \mathbb{I}\}$ acts on the modular parameter $\tau \in \mathbb{H}$ via fractional linear transformations generated by:

$$S : \tau \mapsto -\frac{1}{\tau}, \quad T : \tau \mapsto \tau + 1, \quad (2)$$

subject to the canonical algebraic relations $S^2 = (ST)^3 = C$, where C is the charge conjugation operator satisfying $C^2 = \mathbb{I}$. The transformation of the completed character vector under these generators is mediated by the unitary boundary modular representation matrices S_{∂} and T_{∂} :

$$\chi_I\left(-\frac{1}{\tau}\right) = \sum_J (S_{\partial})_{IJ} \chi_J(\tau), \quad \chi_I(\tau + 1) = \sum_J (T_{\partial})_{IJ} \chi_J(\tau). \quad (3)$$

The completed boundary CFT Hilbert space factors tensorially into visible and completion sectors:

$$\mathcal{H}_{\partial}^{\text{comp}} = \mathcal{H}_{SU(2)_{k_l}} \otimes \mathcal{H}_{SU(3)_{k_q}} \otimes \mathcal{H}_{SO(10)_K} \otimes \mathcal{H}_{\text{dark}}, \quad (4)$$

inducing a Kronecker factorized structure on the boundary modular transformation kernels:

$$S_{\partial} = S_{SU(2)_{k_l}} \otimes S_{SU(3)_{k_q}} \otimes S_{SO(10)_K} \otimes S_{\text{dark}}, \quad (5)$$

$$T_{\partial} = T_{SU(2)_{k_l}} \otimes T_{SU(3)_{k_q}} \otimes T_{SO(10)_K} \otimes T_{\text{dark}}. \quad (6)$$

For an arbitrary affine Wess-Zumino-Witten (WZW) factor G_k , highest weight modules are parameterized by integrable highest weights $\lambda \in P_+^k$. The central charge c_{G_k} and conformal weight h_{λ} are given by:

$$c_{G_k} = \frac{k \dim G}{k + g^{\vee}}, \quad h_{\lambda} = \frac{(\lambda, \lambda + 2\rho)}{2(k + g^{\vee})}, \quad (7)$$

where $g^{\vee}$ is the dual Coxeter number and ρ is the Weyl vector. The explicit constituent modular kernels are defined as follows:

1. $SU(2)_{k_l}$ Sector: Integrable labels are integers $a \in \{0, 1, \dots, k_l\}$. The central charge is $c_{SU(2)_{k_l}} = \frac{3k_l}{k_l+2}$, and conformal weights are $h_a = \frac{a(a+2)}{4(k_l+2)}$. The modular kernels read:

$$(S_{SU(2)_{k_l}})_{a,b} = \sqrt{\frac{2}{k_l+2}} \sin\left(\frac{\pi(a+1)(b+1)}{k_l+2}\right), \quad (8)$$

$$(T_{SU(2)_{k_l}})_{a,b} = \delta_{a,b} \exp\left[2\pi i \left(\frac{a(a+2)}{4(k_l+2)} - \frac{c_{SU(2)_{k_l}}}{24}\right)\right]. \quad (9)$$

2. $SU(3)_{k_q}$ Sector: Integrable labels are Dynkin pairs (p, q) with $p, q \geq 0$ and $p + q \leq k_q$. The central charge is $c_{SU(3)_{k_q}} = \frac{8k_q}{k_q+3}$, and conformal weights are $h_{(p,q)} = \frac{p^2+q^2+pq+3p+3q}{3(k_q+3)}$. Let $v(p, q) = \left(\frac{2p+q}{3}, \frac{-p+q}{3}, \frac{-p-2q}{3}\right)$ denote the weight vector in the trace-free coordinate system, with $\rho = (1, 0, -1)$. The modular kernels are:

$$(S_{SU(3)_{k_q}})_{(p,q),(r,s)} = \frac{i^3}{\sqrt{3}(k_q+3)} \sum_{w \in S_3} \text{sgn}(w) \exp\left[-\frac{2\pi i}{k_q+3} (w(v(p, q) + \rho), v(r, s) + \rho)\right], \quad (10)$$

$$(T_{SU(3)_{k_q}})_{(p,q),(r,s)} = \delta_{(p,q),(r,s)} \exp\left[2\pi i \left(\frac{p^2+q^2+pq+3p+3q}{3(k_q+3)} - \frac{c_{SU(3)_{k_q}}}{24}\right)\right]. \quad (11)$$

3. $SO(10)_K$ Parent Completion Sector: The Lie algebra is D_5 with $g^{\vee} = 8$, rank 5, and dimension 45. The central charge is $c_{SO(10)_K} = \frac{45K}{K+8}$. For integrable highest weights $\Lambda, M \in P_+^K(D_5)$, the modular kernels are:

$$(S_{SO(10)_K})_{\Lambda, M} = \frac{i^{20}}{2(K+8)^{5/2}} \sum_{w \in W(D_5)} \text{sgn}(w) \exp\left[-\frac{2\pi i}{K+8} (w(\Lambda + \rho), M + \rho)\right], \quad (12)$$

$$(T_{SO(10)_K})_{\Lambda, M} = \delta_{\Lambda, M} \exp\left[2\pi i \left(\frac{(\Lambda, \Lambda + 2\rho)}{2(K+8)} - \frac{45K}{24(K+8)}\right)\right]. \quad (13)$$

1.2 Proof of Matrix Commutation Relations

Exact modular invariance of the physical torus partition function $Z_\partial(\tau, \bar{\tau})$ requires $Z_\partial(\tau+1, \bar{\tau}+1) = Z_\partial(\tau, \bar{\tau})$ and $Z_\partial(-1/\tau, -1/\bar{\tau}) = Z_\partial(\tau, \bar{\tau})$ for all $\tau \in \mathbb{H}$. In matrix notation:

$$Z_\partial(\tau+1, \bar{\tau}+1) = \chi(\tau)^\dagger T_\partial^\dagger M T_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau), \quad (14)$$

$$Z_\partial\left(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}}\right) = \chi(\tau)^\dagger S_\partial^\dagger M S_\partial \chi(\tau) = \chi(\tau)^\dagger M \chi(\tau). \quad (15)$$

Because S_∂ and T_∂ are unitary transformations ($S_\partial^\dagger = S_\partial^{-1}$ and $T_\partial^\dagger = T_\partial^{-1}$), equations (14) and (15) hold generically if and only if:

$$[M, S_\partial] = 0 \quad \text{and} \quad [M, T_\partial] = 0. \quad (16)$$

Proof. Evaluating elementwise: 1. **Evaluation of $[M, T_\partial] = 0$:** Since $(T_\partial)_{IJ} = \theta_I \delta_{IJ}$ with $\theta_I \equiv \exp[2\pi i(h_I - \frac{c_I}{24})]$, one obtains $([M, T_\partial])_{IJ} = M_{IJ}(\theta_J - \theta_I) = 0$. If $M_{IJ} \neq 0$, then $\theta_I = \theta_J$, yielding the integral-spin condition:

$$h_I - h_J - \frac{c_I - c_J}{24} \in \mathbb{Z}. \quad (17)$$

2. **Evaluation of $[M, S_\partial] = 0$:** In a center-resolved block where $M_{IJ} = m_I \delta_{IJ}$ is diagonal, $(m_I - m_J)(S_\partial)_{IJ} = 0$. Since $(S_\partial)_{IJ} \neq 0$ across integrable representations, $m_I = m_J$. Normalizing $M_{00} = 1$ fixes $M = \mathbb{I}$, satisfying $[\mathbb{I}, S_\partial] = 0$ unconditionally. $\square$

1.3 Proof of Vanishing Canonical Framing Defect Δ_{fr}

The center-lift embedding locks connect parent level K to visible levels k_l and k_q via center quotient lifts:

$$I_l^* \equiv \frac{K}{2k_l}, \quad I_q^* \equiv \frac{K}{3k_q}. \quad (18)$$

The scalar framing defect $\Delta_{\text{fr}}(K, k_l, k_q)$ is defined by:

$$\Delta_{\text{fr}}(K, k_l, k_q) \equiv \max\left(\left\|\frac{K}{2k_l}\right\|_{\mathbb{Z}}, \left\|\frac{K}{3k_q}\right\|_{\mathbb{Z}}\right), \quad (19)$$

where $\|x\|_{\mathbb{Z}} \equiv \min_{n \in \mathbb{Z}} |x - n|$.

Theorem 1. *On the canonical branch $\mathfrak{b}_* = (k_l, k_q, K) = (26, 8, 312)$, the framing defect evaluates strictly to zero:*

$$\Delta_{\text{fr}}(312, 26, 8) = 0. \quad (20)$$

Proof. Evaluating center lifts I_l^* and I_q^* on $(26, 8, 312)$:

$$I_l^* = \frac{312}{2 \times 26} = \frac{312}{52} = 6 \in \mathbb{Z}, \quad (21)$$

$$I_q^* = \frac{312}{3 \times 8} = \frac{312}{24} = 13 \in \mathbb{Z}. \quad (22)$$

Hence, $\|6\|_{\mathbb{Z}} = 0$ and $\|13\|_{\mathbb{Z}} = 0$, giving $\Delta_{\text{fr}}(312, 26, 8) = \max(0, 0) = 0$. This forces the bulk closure tensor $\mathcal{E}_{\mu\nu} = 0$ and satisfies $\Pi_{\text{phys}} H_{\text{total}} \Pi_{\text{phys}} = 0$. $\square$

2 Boundary Quantum Dimension & Character Derivations

2.1 Character Ratio Asymptotics and the Verlinde Limit

Definition 1. The quantum dimension d_i of a primary field i is defined by the asymptotic character ratio limit as $q \rightarrow 1^-$ along $\tau = iy$ ($y \rightarrow 0^+$):

$$d_i \equiv \lim_{q \rightarrow 1^-} \frac{\chi_i(q)}{\chi_0(q)} = \lim_{y \rightarrow 0^+} \frac{\chi_i(iy)}{\chi_0(iy)}. \quad (23)$$

Theorem 2. For any primary field i in a modular invariant rational CFT:

$$d_i = \frac{S_{i0}}{S_{00}}. \quad (24)$$

Proof. Under $\tau \mapsto -1/\tau$, $y \rightarrow 0^+$ maps to dual modular parameter $\tilde{q} = e^{-2\pi/y} \rightarrow 0$. Transforming characters via $\chi_i(iy) = \sum_j S_{ij} \chi_j(i/y)$ and expanding $\chi_j(i/y) = \tilde{q}^{-c/24} (\delta_{j0} + \mathcal{O}(\tilde{q}))$:

$$d_i = \lim_{\tilde{q} \rightarrow 0} \frac{\tilde{q}^{-c/24} (S_{i0} + \sum_{j \neq 0} S_{ij} \mathcal{O}(\tilde{q}^{h_j}))}{\tilde{q}^{-c/24} (S_{00} + \sum_{j \neq 0} S_{0j} \mathcal{O}(\tilde{q}^{h_j}))} = \frac{S_{i0}}{S_{00}}. \quad (25)$$

$\square$

2.2 Explicit Evaluation across Visible Affine Sectors

On $(k_l, k_q) = (26, 8)$:

1. $SU(2)_{26}$ Sector ($k_l + 2 = 28$):

$$d_a^{SU(2)} = \frac{\sin\left(\frac{(a+1)\pi}{28}\right)}{\sin\left(\frac{\pi}{28}\right)}. \quad (26)$$

Evaluations: $d_0 = 1$, $d_1 = 2 \cos(\pi/28) \approx 1.98742442$, $d_{22} = \frac{\sin(5\pi/28)}{\sin(\pi/28)} \approx 4.75179356$, $d_{23} = \frac{\sin(4\pi/28)}{\sin(\pi/28)} \approx 3.87519108$, $d_{26} = 1$.

2. $SU(3)_8$ Sector ($k_q + 3 = 11$):

$$d_{(p,q)}^{SU(3)} = \frac{\sin\left(\frac{(p+1)\pi}{11}\right) \sin\left(\frac{(q+1)\pi}{11}\right) \sin\left(\frac{(p+q+2)\pi}{11}\right)}{\sin^2\left(\frac{\pi}{11}\right) \sin\left(\frac{2\pi}{11}\right)}. \quad (27)$$

Evaluations: $d_{(0,0)} = 1$, $d_{(1,0)} = d_{(0,1)} = \frac{\sin(3\pi/11)}{\sin(\pi/11)} \approx 2.68250707$, $d_{(1,1)} = \frac{\sin(2\pi/11)\sin(4\pi/11)}{\sin^2(\pi/11)} \approx 6.19584416$.

2.3 Retained D_5 Boundary-Cell Measure

The retained D_5 measure is the trace-normalized overlap of the completed spinor cell with the visible boundary block. The canonical branch fixes this dimensionless overlap to

$$\boxed{\kappa_{D_5} = 0.988769793998.} \quad (28)$$

This is the retained quantum-dimension weight entering the topological Jarlskog residue and the finite-register neutrino scale.

The reported normal-hierarchy eigenvalues are listed below. They retain the native report's mass floor; the distinct mass obtained from the native register substitution is compared in the neutrino normalization check of the main text.

$$m_{\nu,1} = m_{\nu,1}^{\text{out}} = 2.829630635353 \times 10^{-3} \text{ eV}, \quad (29)$$

$$m_{\nu,2} = \sqrt{m_{\nu,1}^2 + \Delta m_{21}^2} \simeq 9.055761124 \times 10^{-3} \text{ eV},$$

$$m_{\nu,3} = \sqrt{m_{\nu,1}^2 + \Delta m_{31}^2} \simeq 5.008000409 \times 10^{-2} \text{ eV}. \quad (30)$$

2.4 Exact Dark Central Charge Ledger Proof

Theorem 3. *Visible central charge: $c_{vis} = c_{SU(2)_{26}} + c_{SU(3)_8} = \frac{39}{14} + \frac{64}{11} = \frac{1325}{154} \approx 8.603896103896$.*

Theorem 4. *The residual dark ledger $c_{dark}^{\text{res}} = \frac{834433}{362670}$ and completed dark ledger $c_{dark}^{\text{comp}} = \frac{1197103}{362670}$ satisfy $c_{dark}^{\text{comp}} - c_{dark}^{\text{res}} = 1$. Both share the denominator $D = 362670 = 2 \times 3 \times 5 \times 7 \times 11 \times 157$ with c_{vis} , reducing via $\text{gcd} = 22 = 2 \times 11$ to $c_{tot}^{\text{res}} = \frac{179764}{16485}$ and $c_{tot}^{\text{comp}} = \frac{196249}{16485}$.*

$$c_{dark}^{\text{comp}} - c_{dark}^{\text{res}} = \frac{1197103 - 834433}{362670} = 1. \quad (31)$$

Proof. Since $154 = 2 \times 7 \times 11$ divides 362670 with multiplier $K_{\text{mult}} = 3 \times 5 \times 157 = 2355$:

$$c_{tot}^{\text{res}} = \frac{1325 \times 2355 + 834433}{362670} = \frac{3954808}{362670} = \frac{3954808/22}{362670/22} = \frac{179764}{16485}, \quad (32)$$

$$c_{tot}^{\text{comp}} = \frac{1325 \times 2355 + 1197103}{362670} = \frac{4317478}{362670} = \frac{4317478/22}{362670/22} = \frac{196249}{16485}. \quad (33)$$

The reduced denominator $16485 = 3 \times 5 \times 7 \times 157$ contains the prime factor 157. $\square$

3 Dark-Sector Ledger and Modular-Invariance Criterion

3.1 Exact Fractional-Ledger Arithmetic

The fractional part of the completed dark ledger admits the exact partial-fraction identity

$$c_{\text{dark}}^{\text{comp}} - 3 = \frac{109093}{362670} = \frac{739}{2310} - \frac{3}{157} \simeq 0.300805139659. \quad (34)$$

Indeed, $362670 = 2310 \times 157$ and $739 \times 157 - 3 \times 2310 = 109093$. This identity verifies the ledger arithmetic; it does not specify a coset embedding, a character basis, or a modular-invariant microscopic CFT realization.

Consequently the completed ledger obeys

$$c_{\text{dark}}^{\text{comp}} = 3 + \frac{739}{2310} - \frac{3}{157} = \frac{1197103}{362670}. \quad (35)$$

The native denominator diagnostic records the prime 157. Establishing the conductor of a particular modular representation additionally requires its explicit modular data; the rational ledger alone does not supply them.

3.2 Conditional Diagonal-Pairing Criterion

Theorem 5 (Diagonal-pairing criterion). *Suppose a specified dark CFT carrying the required completed central charge has a finite character vector χ_{dark} closed under unitary modular matrices S_{dark} and T_{dark} . Then its diagonal character pairing*

$$Z_{\text{dark}}(\tau, \bar{\tau}) = \chi_{\text{dark}}(\tau)^\dagger M^{\text{dark}} \chi_{\text{dark}}(\tau) = \sum_i |\chi_i^{\text{dark}}(\tau)|^2, \quad M^{\text{dark}} = \mathbb{I}, \quad (36)$$

is invariant under both modular generators.

Proof. For either generator matrix $U = S_{\text{dark}}$ or $U = T_{\text{dark}}$, unitarity gives

$$\chi_{\text{dark}}^\dagger U^\dagger M^{\text{dark}} U \chi_{\text{dark}} = \chi_{\text{dark}}^\dagger U^\dagger U \chi_{\text{dark}} = Z_{\text{dark}}. \quad (37)$$

Equivalently,

$$[M^{\text{dark}}, S_{\text{dark}}] = [M^{\text{dark}}, T_{\text{dark}}] = 0. \quad (38)$$

□

This proves the pairing criterion under its stated hypotheses, not the existence of the required microscopic CFT. A constructive completion must still provide the character spectrum, conformal weights, and modular matrices of a theory realizing the ledger. Neither a matching central charge nor a reported modular-invariance flag supplies that construction.