

FROM THE AUTHOR

Physics at a Glance contains all the physics material you require for any of the major GCSE examination boards. It begins with the theory of two major ideas in physics, force and energy. We discover that for anything useful to happen there must be a transfer of energy; and then describe that transfer, by waves, electrically, thermally and by nuclear processes, in more depth. To conclude many applications of physics are explored. Not all the material covered may be relevant to your course and you should ask your teacher or use your examination specification to find out which parts you can leave out.

Many examinations only test a small range of topics encouraging you just to learn the bits you need for your examination and then move on. To be successful at physics it is important to try to make connections between important ideas and, therefore, you will find the same ideas appearing a number of times. This is to help you learn physics by reviewing earlier ideas as you examine a wide range of applications.

The book's visual presentation encourages you to use the mind mapping type approach in your revision, which many learners find helpful as this is often how the brain organizes information. It is intended that the book gives you the 'big picture' while a companion traditional textbook can fill in the detail.

Physics is a mathematical science so some of the questions require you to carry out a calculation. Many of these are of the 'show that' type where an approximate answer is given, so that you can check that you are able to reach the correct solution for yourself. It is vital to show how you got to the solution by showing all your calculations. There are always marks for this and is a good habit to develop. Many questions are quite straightforward, but there a couple designed to make you think, sometimes quite hard about the physics. Tackling these, and persisting until you are successful, will develop real understanding of physics.

The GCSE specifications also require you to understand 'How Science Works'. There is a page midway through the book devoted to these ideas together with examples and questions throughout designed to develop your ability to address these issues in context. I hope you enjoy using *Physics at a Glance* and your GCSE Physics course.

T. Mills

Copyright © 2009 Manson Publishing Ltd

ISBN: 978-1-84076-106-1

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the written permission of the copyright holder or in accordance with the provisions of the Copyright Act 1956 (as amended), or under the terms of any licence permitting limited copying issued by the Copyright Licensing Agency, 33–34 Alfred Place, London WC1E 7DP, UK.

Any person who does any unauthorized act in relation to this publication may be liable to criminal prosecution and civil claims for damages.

A CIP catalogue record for this book is available from the British Library.

For full details of all Manson Publishing Ltd titles please write to:
Manson Publishing Ltd, 73 Corringham Road, London NW11 7DL, UK.
Tel: +44(0)20 8905 5150
Fax: +44(0)20 8201 9233
Website: www.mansonpublishing.com

Project manager: Ruth Maxwell, Clair Chaventré
Design, illustration, and layout: Cathy Martin, Presspack Computing Ltd

Printed by Replika Press Pvt Ltd, Haryana, India

PHYSICS

at a Glance

Tim Mills, BSc
Head of Physics
Brampton College
London

Illustrations by
Cathy Martin










 MANSON
PUBLISHING

CONTENTS

Electrical circuit symbols	4	Electrical cells, alternating and direct current	53
FUNDAMENTAL CONCEPTS			
Forces and Motion	5	Diodes, rectification and capacitors	54
Measuring and describing motion	5	Mains electricity and wiring	55
Motion graphs	6	Electrical safety	56
Equations of motion	8	Electron beams	57
Describing forces	9	Magnetic Fields	58
Balanced forces – Newton's first law	10	Magnetic fields and the Earth's magnetic field	58
Unbalanced forces – Newton's second law	11	Electromagnetism and the motor effect	59
Gravitational forces	12	Thermal Energy	60
Terminal velocity	13	Heat and temperature – what is the difference?	60
Projectiles	14	Temperature scales	60
Newton's third law	15	Specific and latent heat	61
Momentum and force (Newton's laws revisited)	16	Heat transfer 1 – conduction	62
Momentum conservation and collisions	17	Heat transfer 2 – convection	62
Motion in circles and centripetal forces	19	Heat transfer 3 – radiation	63
Moments and stability	20	Reducing energy wastage in our homes	64
Energy	21	Kinetic model of gases	65
Types of energy and energy transfers	21	Gas laws	66
Energy conservation	22	Radioactivity	67
Work done and energy transfer	23	Atomic structure	67
Power	24	What is radioactivity?	67
Gravitational potential energy and kinetic energy	25	A history of our understanding of the atom	68
Energy calculations	26	Background radiation	69
Efficiency and the dissipation of energy	27	Three types of nuclear radiation	70
TRANSFER OF ENERGY			
Waves	28	Radioactive decay and equations	71
Describing waves	28	N/Z curve	72
Wave speed	29	Fundamental particles	73
Electromagnetic waves	30	Half-life	74
How electromagnetic waves travel	31	Is radiation dangerous?	76
Absorption, reflection and transmission of electromagnetic waves	32	Nuclear fission	77
The Earth's atmosphere and electromagnetic radiation	33	Nuclear fusion	78
Uses of electromagnetic waves, including laser light	34	APPLICATIONS OF PHYSICS	
Dangers of electromagnetic waves	35	How science works	79
Reflection, refraction and total internal reflection	36	The Supply and Use of Electrical Energy	80
Refractive index and dispersion	38	Examples of energy transformations involving electrical devices and the impact of electricity on society	80
Diffraction and interference	39	What influences the energy resources we use?	81
Polarization and photon model of light	40	Electricity generation (electromagnetic induction)	82
Seismic waves and the structure of the Earth	41	How power stations work	84
Sound waves	42	The transformer	85
Electrical Energy	43	The national grid	86
Static electricity	43	The environmental impact of electricity generation	87
Electric currents	44	Renewable energy resources	88
Potential difference and electrical energy	45	Calculating the cost of the electrical energy we use	90
Energy transfers in series and parallel circuits	46	The motor and dynamo	91
Resistance	47	Logic gates	92
Electrical measurements and Ohm's Law	48	Electricity and the human body	94
Power in (Ohmic) electrical circuits	49	Transport	95
Properties of some electrical components	50	Stopping distances	95
Potential dividers	52	Road safety	96
		Waves and Communications	97
		Using waves to communicate	97
		Analogue and digital signals	98
		AM/FM radio transmission	99

Satellite orbits and their uses	100	Telescopes and types of radiation used to learn about the Universe	115
Images and ray diagrams	101	The motion of objects in the sky	116
Mirrors and lenses, images	102	Exploring space	117
Optical fibres	104	Forces in the Solar System	118
Ultrasound and its applications	105	The structure of the Universe	119
Uses of electron beams	106	The Sun	120
Beams of light – CDs and relativity	107	Stars and their spectra	121
Radioactivity	108	The life story of a star	122
How is nuclear radiation used in hospitals?	108	How did the Solar System form?	124
Other uses of radioactivity	109	The expanding Universe	125
Radioactive dating	110		
Nuclear power and weapons	111		
Radioactive waste	112		
Our Place in the Universe	113		
Geological processes	113		
The Solar System	114		
		APPENDICES	
		Formulae	127
		INDEX	128

ELECTRICAL CIRCUIT SYMBOLS

Conductors crossing (no connection)		Cell	
Conductors joined		Battery	
Switches			
Open		Closed	
Ammeter			
Voltmeter		Power supply (a.c.)	
		Power supply (d.c.)	
		Transformer	
Fixed resistor		Light emitting diode	
Variable resistor		Lamp	 
Potential divider		Loudspeaker	
Thermistor		Microphone	
Light dependent resistor (LDR)		Motor	
Diode		Generator	
Logic gates			
NOT		Fuse	
AND		Earth connection	
NAND			

FUNDAMENTAL CONCEPTS


FORCES AND MOTION Measuring and Describing Motion

$$\text{Speed (m/s)} = \frac{\text{distance (m)}}{\text{time (s)}}$$

Velocity is speed in a given direction (an example of a **vector** – it has size and direction)


$$\text{Average speed (m/s)} = \frac{\text{total distance (m)}}{\text{time taken (s)}}$$

Instantaneous speed is speed at a given time



$$\text{Acceleration (m/s}^2) = \frac{\text{change in velocity (m/s)}}{\text{time taken (s)}}$$

Ticker tape – 1 dot every 1/50th second
1/10 second



Constant speed – equally spaced dots.

Measure distance for 5 dots, time taken was 1/10th second.

Acceleration – dots get further apart

Length of a 5-tick is proportional to the speed.

SCALAR – size only

SPEED – rate of change of position.


VELOCITY – speed in a given direction.

ACCELERATION – rate of change of velocity (usually taken as increasing, but can be either).

DECELERATION – rate of decrease of velocity.

VECTOR – size and direction

Measuring speed



$$\text{Speed in gate} = \frac{\text{length of interrupt card}}{\text{time beam blocked}}$$

$$\text{Acceleration} = \frac{\text{change in speed between gates}}{\text{time between gates}}$$


$$\text{Average speed} = \frac{\text{distance between gates}}{\text{time between gates}}$$

Questions


1. A toy train runs round a circular track of circumference 3 m. After 30 s, it has completed one lap.
 - a. What was the train's average speed?
 - b. Why is the train's average velocity zero?
 - c. The train is placed on a straight track. The train accelerated uniformly from rest to a speed of 0.12 m/s after 10 s. What was its acceleration?
 - d. Describe three different ways of measuring the train's average speed and two different ways of measuring the train's instantaneous speed.
 - e. How could light gates be used to measure the train's acceleration along a 1 m length of track?
2. Explain the difference between a scalar and vector. Give an example of each.
3. A car leaks oil. One drip hits the road every second. Draw what you would see on the road as the car accelerates.

FORCES AND MOTION Motion Graphs

$$\text{Gradient} = \frac{\text{change in distance (m)}}{\text{change in time (s)}} = \frac{\Delta d}{\Delta t} = \text{speed (m/s)}$$




$$\text{Gradient} = \frac{\text{change in velocity (m/s)}}{\text{change in time (s)}} = \frac{\Delta v}{\Delta t} = \text{acceleration (m/s}^2)$$





Questions

- Copy and complete the following sentences:
 - The slope of a distance – time graph represents _____
 - The slope of a velocity – time graph represents _____
 - The area under a velocity – time graph represents _____
- Redraw the last four graphs from p7 for an object that is decelerating (slowing down).
- Sketch a distance–time graph for the motion of a tennis ball dropped from a second floor window.
- Sketch a velocity-time graph for the motion of a tennis ball dropped from a second floor window. Take falling to be a negative velocity and bouncing up to be a positive velocity.



FORCES AND MOTION Equations of Motion




Questions

Show ALL your working.

- What quantities do the variables x , u , v , a , and t each represent?
- Write a list of three equations which connect the variables x , u , v , a , and t .
- A car accelerates from 10 m/s to 22 m/s in 5 s. Show that the acceleration is about 2.5 m/s².
- Now show the car in (3) travelled 80 m during this acceleration:
 - Using the formula $v^2 = u^2 + 2ax$.
 - Using the formula $x = ut + \frac{1}{2}at^2$.
- A ball falls from rest. After 4 s, it has fallen 78.4 m. Show that the acceleration due to gravity is 9.8 m/s².
- Show that $x = \frac{1}{2}(u+v)(v-u)/a$ rearranges to $v^2 = u^2 + 2ax$.
- A ball thrown straight up at 15 m/s, feels a downward acceleration of 9.8 m/s² due to the pull of the Earth on it. How high does the ball go before it starts to fall back?


FORCES AND MOTION

Describing Forces



Free body diagrams

E.g. Sliding box



Simple diagrams to show all the forces acting *on* an object.


properly called

Normal contact force

Push of one surface on another at right angles to the surface.
Due to atoms in each surface being slightly squashed together and pushing back

Weight – shorthand for ‘the gravitational pull of a planet or moon (usually the Earth) on an object’

E.g. pendulum




Resultant force – a single force that can replace all the forces acting on a body and have the same overall effect as all the individual forces acting together.


It is the sum of all the individual forces taking their directions into account.

E.g. rocket

Acceleration (double-headed arrow not attached to object)




Parallel forces – add



Resultant = push of wind on boat – drag of water on boat

Perpendicular forces – Pythagoras Theorem



Questions

- List three effects forces can have.
- Explain what the term ‘resultant force’ means.
- To describe a force fully, what three pieces of information should be recorded?
- Copy and add arrows to these diagrams to show all the forces (and their directions) acting on:
 - A netball flying through the air.
 - A jet ski.



- A cyclist freewheeling down a hill.
- A child on a swing pushed by an adult.


- Calculate the resultant force in the following cases:

- 3 N ← → 5 N
- 7 N ← → 4 N ↑ 10 N
- 3 N ↑ 4 N
- 4 N ↑ 8 N ← → 5 N
- 1 N ↑ 3 N ← → 7 N ↓ 4 N



FORCES AND MOTION


Balanced Forces – Newton's First Law




Why do moving objects seem to slow down?

On Earth objects move:


Over solid surfaces



In or over water



In the atmosphere



In all cases, resistive forces act to oppose motion. Therefore, unless a force is applied to balance the resistive force the object will slow down. In space, there are no resistive forces and objects will move at constant speed in a straight line unless another force acts.

Newton's First Law of Motion:


- If the resultant force acting on a body is zero, it will remain at rest or continue to move at the same speed in the same direction.
- If the resultant force acting on a body is not zero, it will accelerate in the direction of the resultant force.

Questions

1. In which of the following situations is the resultant force zero? Explain how you decided.
 - A snooker ball resting on a snooker table.
 - A car accelerating away from traffic lights.
 - A ball rolling along level ground and slowing down.
 - A skier travelling down a piste at constant speed.
 - A toy train travelling round a circular track at constant speed.
2. A lift and its passengers have a weight of 5000 N. Is the tension in the cable supporting the lift:

- Greater than 5000 N, ii. Less than 5000 N,
- iii. Exactly 5000 N when:
 - The lift is stationary?
 - Accelerating upwards?
 - Travelling upwards at a constant speed?
 - Decelerating whilst still travelling upwards?
 - Accelerating downwards?
 - Travelling downwards at constant velocity?
 - Decelerating while still travelling downward?
- Explain why all objects moving on Earth will eventually come to rest unless another force is applied?

FORCES AND MOTION Unbalanced Forces – Newton's Second Law



We define the Newton as the force needed to accelerate a 1 kg mass at 1 m/s². Therefore, we can write:

$$\text{Force (N)} = \text{mass (kg)} \times \text{acceleration (m/s}^2\text{)}.$$

Newton's Second Law



$$\text{Resultant force} = 2800 \text{ N}$$

$$\text{Acceleration} = \text{force/mass} = 2800 \text{ N}/1000 \text{ kg} = 2.8 \text{ m/s}^2.$$




$$\text{Resultant force} = \text{mass} \times \text{acceleration}$$

$$= 100 \text{ kg} \times 0.3 \text{ m/s}^2$$

$$= 30 \text{ N}$$

$$\text{Resultant force} = \text{driving force} - 20 \text{ N}$$

$$= 50 \text{ N}$$



Questions

- Calculate:
 - The force needed to accelerate a 70 kg sprinter at 6 m/s².
 - The acceleration of a 10 g bullet with 2060 N explosive force in a gun barrel.
 - The mass of a ship accelerating at 0.09 m/s² with a resultant thrust of 6 400 000 N from the propellers.
- An underground tube train has mass of 160 000 kg and can produce a maximum driving force of 912 000 N.
 - When accelerating in the tunnel using the maximum driving force show the acceleration should be 5.7 m/s².
 - In reality, the acceleration is only 4.2 m/s². Hence show the resistive forces on the train are 240 000 N.
- Explain why towing a caravan reduces the maximum acceleration of a car (two reasons).
- A football made of concrete would be weightless in deep space. However, it would not be a good idea for an astronaut to head it. Why not?

FORCES AND MOTION Gravitational Forces

A gravitational field is a region of space where objects with mass feel forces. Since we live in the Earth's gravitational field, gravitational forces are very common to us.

We give the gravitational force between a mass and the Earth a special name, weight.

Weight always acts towards the centre of the Earth.

The force per kilogram of mass is a good way of measuring the strength of the Earth's gravitational field.

If an object is in free fall, the only force on it is weight (difficult in practice because of air resistance).


$$W = mg$$

10 m/s^2 Hence, acceleration due to free fall is equivalent to gravitational field strength.

$$g = 10 \text{ N/kg}$$

Earth

The mass of an object is the same everywhere. The weight depends on the gravitational field strength.



$$\text{Weight (N)} = \text{mass (kg)} \times \text{gravitational field strength (N/kg)}$$

Applying Newton's second law



$$\text{Resultant force} = \text{mass} \times \text{acceleration}$$

$$\Rightarrow \text{Then acceleration } (\text{m/s}^2) = \text{weight (N)} / \text{mass (kg)}.$$

However, this is also the definition of gravitational field strength.

You can think of this as the weight increasing to compensate for the increased mass so all objects fall at the same rate, independent of their mass.

All masses fall with an acceleration of about 10 m/s^2 (providing there are no resistive forces).




Questions

1. Near the surface of the Earth, what are the values of:
 - The acceleration due to free fall?
 - The gravitational field strength?
2. What are the weights on the Earth of:
 - A book of mass 2 kg?
 - An apple of mass 100 g?
 - A girl of mass 60 kg?
 - A blade of grass of 0.1 g?
3. What would the masses and weights of the above objects be on the moon? (Gravitational field strength on the moon = 1.6 N/kg).
4. 6400 km above the surface of the Earth a 1 kg mass has a weight of 2.5 N. What is the gravitational field strength here? If the mass was dropped, and started falling towards the centre of the Earth, what would its initial acceleration be?
5. Write a few sentences to explain the difference between mass and weight.

FORCES AND MOTION

Terminal Velocity




Questions

- What happens to the size of the drag force experienced by an object moving through a fluid (e.g. air or water) as it speeds up?
 - What force attracts all objects towards the centre of the Earth?
 - Why does a car need to keep its engine running to travel at constant velocity?
 - A hot air balloon of weight 6000 N is released from its mooring ropes.
 - The upward force from the hot air rising is 6330 N. Show the initial acceleration is about 0.5 m/s^2 .
- This acceleration gradually decreases as the balloon rises until it is travelling at a constant velocity. Explain why.
 - A mass of 100 kg is thrown overboard. What will happen to the balloon now?
 - Sketch a velocity-time graph for the whole journey of the balloon as described in parts a-c.
 - Explain why the following are likely to increase the petrol consumption of a car:
 - Towing a caravan.
 - Adding a roof rack
 - Driving very fast.


FORCES AND MOTION

Projectiles



The secret is to consider the velocity of the projectile to be made up of horizontal and vertical velocities, which can be considered separately.



A true projectile only has one force, weight, acting on it when it is in flight.



Examples:




Questions

1. In an ideal world how many forces act on a projectile, and what are they?
2. State the value of the vertical acceleration of a projectile.
3. Explain why the horizontal acceleration of a projectile is zero. What assumption has to be made?
4. Explain why a firework rocket cannot be analysed as a projectile with the methods shown here.
5. A ball is kicked so it has a velocity of 15.59 m/s horizontally and 9.0 m/s vertically.

- Show that the resultant velocity of the ball has a magnitude of 18.0 m/s.
- Show that the ball takes 0.92 s to reach its maximum height above the ground.
- For how long in total is the ball in the air and how far along the ground will it travel?
- Show the maximum height the ball reaches is 4.1 m.
- What will the magnitude of its resultant velocity be when it hits the ground? Hint: no calculation needed.

FORCES AND MOTION Newton's Third Law

Whenever an object experiences a force it always exerts an equal and opposite force on the object causing the force.



Questions

1. Explain what is meant by the term 'normal contact force'.
2. A jet engine in an aircraft exerts 200 000 N on the exhaust gases. What force do the gases exert on the aircraft?
3. Describe the force that forms a Third Law pair with the following. In each case, draw a diagram to illustrate the two forces:
 - a. The push east of the wind on a sail.
 - b. The push left of a bowstring on an arrow.
 - c. The frictional push south of a train wheel on a rail.
 - d. The normal contact force downwards of a plate on a table.


- e. The attraction right of the north magnetic pole of a bar magnet on a south magnetic pole of a different magnet.
4. Why are the following not Third Law pairs? (There may be more than one reason for each.)
 - a. The weight of a mug sitting on a table; the normal contact force of the tabletop on the mug.
 - b. The weight of the passengers in a lift car; the upward tension in the lift cable.
 - c. The weight of a pool ball on a table; the horizontal push of the cue on the ball.
 - d. The attraction between the north and south magnetic poles of the same bar magnet.
5. Explain why it is very difficult (and dangerous) to ride a bicycle across a sheet of ice.

FORCES AND MOTION Momentum and Force (Newton's Laws revisited)

Momentum helps to describe how moving objects will behave.


$$\text{Momentum (kgm/s)} = \text{mass (kg)} \times \text{velocity (m/s)}$$

Momentum is a vector. It has size and direction (the direction of the velocity).




Hence, an alternative version of Newton's Second Law

If a resultant force acts on a body free to move a change in momentum occurs equal to the product of the force and the time for which it acts.



Also consider



Questions


1. What units do we use to measure momentum and impulse? (2 answers)
2. Calculate the momentum of:
 - A 55 kg girl running at 7 m/s north.
 - A 20 000 kg aircraft flying at 150 m/s south.
 - A 20 g snail moving at 0.01 m/s east.
3. What is the connection between force and change in momentum?
4. What is the change in momentum in the following cases:
 - A 5 N force acting for 10 s?
 - A 500 N force acting for 0.01 s?
5. What force is required to:
 - Accelerate a 70 kg athlete from 0 to 9 m/s in 2 s?
 - Accelerate a 1000 kg car from rest to 26.7 m/s in 5 s?
 - Stop a 10 g bullet travelling at 400 m/s in 0.001 s?
6. What would be the effect on the force needed to change momentum if the time the force acts for is increased?
7. A 2564 kg space probe is to be accelerated from 7.7 km/s to 11.0 km/s. If it has a rocket motor that can produce 400 N of thrust, for how long would it need to burn assuming that no resistive forces act? Why might this not be practical? How else might the space probe gain sufficient momentum (see p113 for ideas)?

FORCES AND MOTION Momentum Conservation and Collisions

Law of Conservation of Momentum:

Momentum cannot be created or destroyed but can be transferred from one object to another when they interact.

There are no exceptions to this. It is applied to analyse interactions between objects, which can be classified as:



FORCES AND MOTION Momentum Conservation and Collisions (continued)

The calculation of the force exerted on the bullet and the ball would work equally well if the force on the bat or the rifle were calculated. The size of the force would be the same, but in the opposite direction according to Newton's Third Law. Again using $F\Delta t = mv - mu$.

Force of ball on bat

$$F \times 0.02 \text{ s} = 2 \text{ kg} \times (-13.76 \text{ m/s}) - 2 \text{ kg} (-20 \text{ m/s})$$


$$F = 624 \text{ N} \text{ (positive, to the right).}$$

Force of bullet on gun

$$F \times 0.002 \text{ s} = (3 \text{ kg} \times 1.33 \text{ m/s}) - (3 \text{ kg} \times 0 \text{ m/s})$$

$$F = 2000 \text{ N} \text{ (positive, to the right).}$$

These calculations show that the force involved depends on.




Questions

- When a raindrop hits the ground where does its momentum go?
- Why do boxers wear padded gloves?
- A squash ball is hit against a wall and bounces off. An equal mass of plasticine is thrown at the same wall with the same speed as the ball, but it sticks on impact. Which exerts the larger force on the wall and why?
- A golfer swings a 0.2 kg club at 45 m/s. It hits a stationary golf ball of mass 45 g, which leaves the tee at 65 m/s.
 - What was the momentum of the club before the collision?
 - What was the momentum of the ball after the collision?
 - Hence, show that the club's velocity is about 30 m/s after the collision.
 - If the club is in contact with the ball for 0.001 s, what is the average force the club exerts on the ball?
- A 1.5 kg air rifle fires a 1 g pellet at 150 m/s. What is the recoil velocity of the rifle? Show that the force exerted by the rifle on the pellet is about 70 N if the time for the pellet to be fired is 0.0021 s.
- Assume that the average mass of a human being is 50 kg. If all 5.5×10^9 humans on Earth stood shoulder to shoulder in one place, and jumped upward at 1 m/s with what velocity would the Earth, mass 6×10^{24} kg recoil?
- Two friends are ice-skating. One friend with mass 70 kg is travelling at 4 m/s. The other of mass 60 kg travelling at 6 m/s skates up behind the first and grabs hold of them. With what speed will the two friends continue to move while holding onto each other?

FORCES AND MOTION Motion in Circles and Centripetal Forces


Object moving in circular path.




$$\text{Centripetal acceleration (m/s}^2) = \frac{\text{[velocity (m/s)]}^2}{\text{radius (m)}} \\ a = v^2/r$$

$$\text{Centripetal force} = \frac{\text{mass (kg)} \times \text{acceleration (m/s}^2)}{\text{radius (m)}} \\ F = mv^2/r = \frac{\text{mass (kg)} \times \text{[velocity (m/s)]}^2}{\text{radius (m)}}$$


Centripetal force is not a force in its own right – it must be *provided* by another type of force.



Frictional push sideways of road on tyres.



Normal contact force on clothes in washing machine drum provides centripetal force.




Questions

- What force provides the centripetal force in each of these cases?
 - The Earth moving in orbit around the Sun.
 - Running around a sharp bend.
 - A child on a swing.
- Explain how a passenger on a roundabout at a funfair can be moving at constant speed around the circle and yet accelerating. In what direction is the acceleration?
- What is the centripetal acceleration of, and force on, the following?
 - A wet sweater of mass 1 kg, spinning in a washing machine drum of radius 35 cm, moving at 30 m/s.
 - A snowboarder of mass 70 kg travelling round a half pipe of radius 6 m at 5 m/s.
- The Earth has a mass of 6×10^{24} kg. Its orbit radius is 1.5×10^{11} m and the gravitational attraction to the Sun is 3.6×10^{22} N.
 - Show that the circumference of the Earth's orbit is about 9.5×10^{11} m.
 - Show that the Earth's speed around the Sun is about 30 000 m/s.
 - Therefore, show that the time to orbit the Sun is about 3×10^7 s.
 - Show that this is about 365 days.
- On a very fast rotating ride at a funfair, your friend says that they feel a force trying to throw them sideways out of the ride. How would you convince your friend that actually they are experiencing a force pushing *inwards*? You should refer to Newton's First and Third Laws in your explanation.

FORCES AND MOTION


Moments and Stability

A moment (or torque) is the turning effect of a force.



A body will not rotate if there is no resultant moment.

Moment (Nm) = Force (N) \times perpendicular distance from line of action of the force to the axis of rotation (m).



Rotation neither raises, nor lowers, the centre of mass.


Questions

1. What is the moment in each of the diagrams below?



2. If the forces in question 1 acted at 60° to the spanner rather than 90° would the moment be greater, the same as, or less than that calculated in question 1? Explain.


3. What are the missing forces or distances in the diagrams below?




4. A letter P is cut from thin cardboard. Explain how to locate its centre of mass.

5. The following letters are cut from a thick plank of wood. W, P, O, I, H, L, U. If stood upright in their normal positions, which are in stable equilibrium, which unstable, and which neutral? Which letter would you expect to be easiest to topple and why?

ENERGY Types of Energy and Energy Transfers



Energy transfers can be shown on simple diagrams.



Many transducers have a number of output energies. Sometimes we ignore some of these if they are insignificant.

E.g. Candle
Chemical energy (in wax) Heat Light } (In flame)

E.g. Electric motor
Electrical energy (in wires) Kinetic energy
Heat energy
Sound energy (very small so probably ignored)

Some transfers are cyclical
E.g. pendulum

Gravitational potential Kinetic Gravitational potential

Questions

1. Nuclear energy is stored in the nucleus of atoms. Make a list of the other types of energy that can be stored giving an example of each.
2. What is a transducer? Make a list of five transducers that might be found in a home and the main energy change in each case.
3. Draw an energy transfer diagrams for the following showing the main energy transfers in each case:

- a. Electric filament light bulb.
 - b. Solar cell.
 - c. Electric kettle.
 - d. Loudspeaker.
 - e. Mobile phone 'charger'.
 - f. Clockwork alarm clock.
 - g. Playground swing.
 - h. Bungee jumper.
 - i. Petrol engine.
 - j. Microphone.
4. What provides the energy input for the human body? List all types of energy that the body can transfer the energy input into.


ENERGY Energy Conservation

Probably the most important idea in Physics is the Principle of Conservation of Energy, which states:

Energy cannot be created or destroyed. It can only be transformed from one form to another form.

This means that the total energy input into a process is the same as the total energy output.

We can use a more sophisticated energy transfer diagram, called a *Sankey diagram*, to show this.




The majority of the rest of this book explores this in more detail.

Questions

- State the Principle of Conservation of Energy.
- What units is energy measured in?
- Explain the difference between energy transformations and energy transfers. Suggest four ways energy can be transferred.
- A TV set uses 25 J of energy each second. If 15 J of energy is converted to light and 2 J is converted to sound, how much energy is converted to heat, assuming this is the only other form of energy produced?
- The motor in a toy train produces 1 J of heat energy and 2 J of kinetic energy every second. What must have been the minimum electrical energy input per second? If the train runs uphill and the electrical energy input stays the same, what would happen to its speed?
- Use the following data to draw a Sankey diagram for each device:
 - Candle (chemical energy in wax becomes heat energy 80% and light 20%).
 - Food mixer (electrical energy supplied becomes 50% heat energy in the motor, 40% kinetic energy of the blades, and 10% sound energy).
 - Jet aircraft (chemical energy in fuel becomes 10% kinetic energy, 20% gravitational potential energy, and 70% heat).

ENERGY Work Done and Energy Transfer



Whenever something useful happens, energy must be transferred but how can we measure energy? The only way to measure energy directly is by considering the idea of *work done*.



Questions

1. Copy and complete:
‘Work is done when a ? moves an object. It depends on the size of the ? measured in ? and the ? the object moves measured in ?. Whenever work is done, an equal amount of ? is transferred. The unit of energy is the ?. Work is calculated by the formula: work = ? × distance moved in the ? of the ?.’
2. I push a heavy box 2 m along a rough floor against a frictional force of 20 N. How much work do I do? Where has the energy come from for me to do this work?
3. A parachute exerts a resistive force of 700 N. If I fall 500 m, how much work does the parachute do?
4. A firework rocket produces a constant thrust of 10 N.
 - a. The rocket climbs to 150 m high before the fuel is used up. How much work did the chemical energy in the fuel do?
 - b. Explain why the chemical energy stored in the fuel would need to be much greater than the work calculated in (a).
 - c. The weight of the empty rocket and stick is 2.5 N. How much work has been done against gravity to reach this height?
 - d. The answers to parts (a) and (c) are not the same, explain why.

ENERGY Power




Power is the number of Joules transferred each second.

The unit of power is the Joule per second, called the Watt, W.

$$\text{Power (W)} = \frac{\text{energy transferred (J)}}{\text{time taken (s)}}$$

Calculating power. Non-mechanical:

- Find out total (heat, light, electrical) energy transferred
- Find out how long the energy transfer took
- Use the formula above





Bulb is switched on for 20 s.

Compare these: imagine how tired you would get if you personally had to do all the work necessary to generate all the electrical power your house uses.

$$\begin{aligned} \text{Power} &= \frac{\text{energy transferred}}{\text{time taken}} \\ &= \frac{1200 \text{ J}}{20 \text{ s}} \\ &= 60 \text{ W} \end{aligned}$$

Slow conversion of electrical to heat and light energy.



Power is the rate of energy conversion between forms.


'Rate' means how quickly something happens.

Energy transferred = work done, so


$$\text{Power (W)} = \frac{\text{work done (J)}}{\text{time taken (s)}}$$

Mechanical:

(i.e. where a force moves through a distance)



- Calculate the work done = force (N) × distance (m)
- Find out how long the work took to be done
- Use the formula above



$$\begin{aligned} \text{Work done} &= 300 \text{ N} \times 0.5 \text{ m} \\ &= 150 \text{ J per lift} \end{aligned}$$


$$\text{Total work done} = 20 \times 150 \text{ J} = 3000 \text{ J}$$

$$\begin{aligned} \text{Power} &= \frac{\text{work done}}{\text{time taken}} = \frac{3000 \text{ J}}{60 \text{ s}} \\ &= 50 \text{ W} \end{aligned}$$

Questions

- A kettle converts 62,000 J of electrical energy into heat energy in 50 s. Show its power output is about 1,200 W.
- A car travels at constant velocity by exerting a force of 1,025 N on the road. It travels 500 m in 17 s. Show that its power output is about 30 kW.
- The power to three electrical devices is as follows: energy efficient light bulb, 16 W; the equivalent filament bulb, 60 W; a TV on standby, 1.5 W.
 - How many more Joules of electrical energy does the filament bulb use in one hour compared to the energy efficient bulb?
- When I bring my shopping home, I carry two bags, each weighing 50 N up two flights of stairs, each of total vertical height 3.2 m. I have a weight of 700 N.
 - How much work do I do on the shopping?
 - How much work do I do to raise my body up the two flights of stairs?
 - If it takes me 30 s to climb all the stairs, show that my power output is about 170 W.

ENERGY Gravitational Potential Energy and Kinetic Energy



Questions


1. Make a list of five objects that change their gravitational potential energy.
2. Using the diagram above calculate the kinetic energy of the car and the lorry.
3. How fast would the car have to go to have the same kinetic energy as the lorry?
4. The mass of the lift and the passengers in the diagram is 200 kg. Each floor of the building is 5 m high.
 - a. Show that the gravitational potential energy of the lift when on the eighth floor is about 80 000 J.
 - b. How much gravitational potential energy would the lift have when on the third floor? If one passenger of mass 70 kg got out on the third floor, how much work would the motor have to do on the lift to raise it to the sixth floor?
 - c. What is the gravitational potential energy of a 0.5 kg ball 3 m above the surface of the Moon where the gravitational field strength is about 1.6 N/kg?
5. A coin of mass 10 g is dropped from 276 m up the Eiffel tower.
 - a. How much gravitational potential energy would it have to lose before it hits the ground?
 - b. Assuming all the lost gravitational potential energy becomes kinetic energy, how fast would it be moving when it hit the ground?
 - c. In reality, it would be moving a lot slower, why?

ENERGY Energy Calculations

GPE = gravitational potential energy KE = kinetic energy

All energy calculations use the *Principle of Conservation of Energy*.

E.g. Bouncing ball



Air resistance is ignored

$$\text{GPE at } h_1 = \text{KE at bottom} \quad \text{Elastic potential at bottom} = \text{KE leaving floor} = \text{GPE at } h_2$$

Thermal energy in deformed ball

Conservation of energy

GPE at top of bounce = KE at bottom of bounce

$$mg\Delta h = \frac{1}{2}mv_1^2$$


$$v_1 = \sqrt{2gh}$$

GPE at top of skydive

Work against friction

As small as possible to prevent injury

Roller Coaster



Time to reach top of track

$$= \text{GPE gain / power of motor} = mg h_1 / \text{power}$$

The time will be greater than this as some electrical energy is converted to KE and does work against friction.


$$\text{GPE on leaving plane} = mg\Delta h$$

GPE at top is not equal to KE at bottom as some GPE was transferred to work against friction (air resistance).

GPE = KE + work against friction

$$mg\Delta h = \frac{1}{2}mv^2 + F \times \Delta h$$

$$F = \frac{mg\Delta h - \frac{1}{2}mv^2}{\Delta h}$$



$$\text{KE at bottom} = \frac{1}{2}mv^2$$

Questions Take $g = 9.8 \text{ m/s}^2$.

1. At the start of a squash game, a 44 g ball is struck by a racquet and hits the wall at 10 m/s.
 - a. Show its KE is about 2 J.
 - b. The ball rebounds at 8 m/s. Calculate the loss in KE.
 - c. Where, and into what form, has this energy been transferred?
2. An aerobatics aircraft of mass 1000 kg is stationary on a runway. Its take off speed is 150 m/s.
 - a. Show that the KE of the aircraft at take off is about $11 \times 10^6 \text{ J}$

b. The maximum thrust of the engines is 20 000 N. Show the aircraft travels over 500 m along the runway before it lifts off.

- c. Give two reasons why the runway will actually need to be considerably longer.
- d. The aircraft climbs to a height of 1000 m. Show it gains about $10 \times 10^6 \text{ J}$.


- e. If the aircraft takes 5 minutes to reach this height, show the minimum power of the engine must be about 33 kW.
- f. Why must this be the minimum power?
- g. The aircraft then flies level at 200 m/s. What is its KE now?
- h. The pilot cuts the engine and goes into a vertical dive as part of the display. When the plane has dived 500 m what is the maximum KE the plane could have gained?
- i. Hence, what is the maximum speed the plane could now be travelling at?
- j. In reality, it will be travelling slower, why?

ENERGY Efficiency and the Dissipation of Energy

If energy is conserved, why do we talk about 'wasting energy'?

Usually when energy is transferred only a proportion of the energy is converted to a useful form, the remainder is converted to other less useful forms of energy, often heat.

E.g. Light bulb



Useful – the type of energy we want from a light bulb.


This energy is not useful – light bulbs are not designed to be heaters. We say this energy has been 'wasted'.

We say that this energy, that is not useful, is 'wasted'.


The proportion of the total energy transferred that is useful is called the *efficiency* of the system.

$$\text{Efficiency (\%)} = \frac{\text{useful energy output (J)}}{\text{total energy input (J)}} \times 100\%$$

The greater the percentage of energy transformed to a useful form, the more efficient the device.




E.g. Power generation



Notice that most of the wasted energy ends up as heat.

Heat energy naturally spreads out into the environment.





Questions

- An electric motor on a crane raises 50 kg of bricks 10 m. If the energy supplied to the motor was 16 000 J show that the motor is about 30% efficient.
- A rollercoaster has 250 000 J of GPE at the top of the first hill. At the bottom of the first hill, the coaster has 220 000 J of KE. Where did the rest of the energy go, and what is the overall efficiency of the GPE to KE conversion?
- A ball of mass 30 g falls from 1.5 m and rebounds to 0.8 m. Show that the efficiency of the energy transformation is about 50%. Why do you not need to know the mass of the ball?
- A car engine is about 20% efficient at converting chemical energy in petrol. If a car of mass 1000 kg has to climb a hill 50 m high, how much chemical energy will be required? Why in reality would substantially more chemical energy be needed than the value you calculated?
- A filament light bulb produces a lot of waste heat. Explain why this waste heat energy cannot be put to other uses very easily.
- What are the main sources of energy wastage in:
 - A vacuum cleaner?
 - A motor car?




TRANSFER OF ENERGY

WAVES Describing Waves



Questions

1. Identify the measurements a, b and c in the following diagrams:
 - a. Amplitude
 - b. Wavelength
 - c. Frequency
2. Write a sentence to define each of the following terms:
 - a. Wavelength.
 - b. Frequency.
 - c. Amplitude.
3. Give one similarity and one difference between a longitudinal and transverse wave and give an example of each.
4. For each of particles a, b, and c in the diagram decide if the particle is moving up, moving down, or is momentarily stationary.




WAVES Wave Speed

The speed of a wave is given by the equation


$$\text{Wave speed (m/s)} = \text{frequency (Hz)} \times \text{wavelength (m)}$$

Here is how to see why



Examples

Water Wave:



$$\begin{aligned} \text{frequency} &= \frac{\text{speed}}{\text{wavelength}} \\ &= \frac{5 \text{ m/s}}{2 \text{ m}} \\ &= 2.5 \text{ Hz} \end{aligned}$$

Light Wave:

$$\begin{aligned} \text{Speed of light} &= 3 \times 10^8 \text{ m/s} \\ \text{frequency} &= 5 \times 10^{14} \text{ Hz}, \\ \text{wavelength} &= \frac{\text{speed}}{\text{frequency}} = \frac{3 \times 10^8 \text{ m/s}}{5 \times 10^{14} \text{ Hz}} \\ &= 6 \times 10^{-7} \text{ m} \end{aligned}$$

Common speeds:


Speed of light = 3×10^8 m/s (300 000 000 m/s)

Speed of sound \approx 340 m/s (in air at room temperature)

Wave speeds can also be calculated by

$$\text{Wave speed (m/s)} = \frac{\text{distance travelled (m)}}{\text{time taken (s)}}$$

E.g. Sonar




N.B. Remember time is the out and back time.

Questions

1. Calculate the speed of the following waves:
 - A water wave of wavelength 1 m and frequency 2 Hz.
 - A water wave of wavelength 3 m and frequency 0.4 Hz.
2. Rearrange the formula $\text{wave speed} = \text{frequency} \times \text{wavelength}$ to read:
 - wavelength = _____. b. frequency = _____.
3. Calculate the frequency of a sound wave of speed 340 m/s and wavelength:
 - 2 m.
 - 0.4 m.
4. Calculate the wavelength of a light wave of speed 300 000 000 m/s and frequency:
 - 4.62×10^{14} Hz.
 - 8.10×10^{14} Hz.
5. Calculate the speed of the following waves. Why might we say that all of these waves belong to the same family?
 - Wavelength 10 m, frequency = 3×10^7 Hz.
 - Wavelength 4×10^{-3} m, frequency 7.5×10^{10} Hz.
 - Wavelength 6×10^{-10} m, frequency 5×10^{17} Hz.
6. In the sonar example above, the echo takes 0.3 s to return from the sea floor. If the sea is 225 m deep, show that the speed of sound in seawater is about 1500 m/s.
7. A radar station sends out radiowaves of wavelength 50 cm and frequency 6×10^8 Hz. They reflect off an aircraft and return in 4.7×10^{-5} s. Show that the aircraft is about 7 km from the radar transmitter.

WAVES Electromagnetic Waves

Electromagnetic waves, like all waves transfer energy. They also have the following properties in common.




Questions

- State three properties all electromagnetic waves have in common.
- Calculate the wavelength of electromagnetic waves of the following frequencies:
 - 5×10^9 Hz.
 - 5×10^{14} Hz.
 - 5×10^{15} Hz.
- What part of the electromagnetic spectrum does each of these waves come from?
- Calculate the frequencies of electromagnetic waves of the following wavelengths:
 - 1 m.
 - 1×10^{-5} m.
 - 5×10^{-8} m.
- What part of the electromagnetic spectrum does each of these waves come from?
- List the electromagnetic spectrum in order of increasing energy.
- Which has the longest wavelength, red or blue light? List the colours of the visible spectrum in order of increasing frequency.


WAVES How Electromagnetic Waves Travel

What is 'waving' in an electromagnetic wave?

It is formed from linked oscillating electric and magnetic fields, hence the name.



Electromagnetic radiation from a point source (e.g. a star, lamp filament) obeys the inverse square law.



$$\text{Intensity (W/m}^2\text{)} = \frac{\text{power (W)}}{\text{area (m}^2\text{)}}$$

After travelling a distance $2r$, it will be spread over a sphere of surface area $4A$. (Intensity, $\frac{P}{4A}$)

Laser light Stands for
'Light Amplification by Stimulated Emission of Radiation'

Only one wavelength from a given source. Waves form a narrow beam with very little spreading so the light is very intense.

Laser light

All waves travel together in step (in phase). { Peaks and troughs always coincide.

vs.

Ordinary light


Waves travel in all directions. Waves do not travel in step.

Generally a mixture of wavelengths from a source.

Questions

- What is waving in an electromagnetic wave?
- A 60 W light bulb can be considered a point source of light. What is the intensity of the light?
 - 1 m from the bulb when it has spread through a sphere of area 12.6 m^2 ?
 - 2 m from the bulb when it has spread through a sphere of area 50.3 m^2 ?
 - Suggest what the intensity would be 3 m from the bulb.
- The intensity of the Sun's radiation at the Earth is about 1400 W/m^2 . Jupiter is about five times further from the Sun. Show that the intensity of the Sun's radiation here is about 56 W/m^2 .
- Suggest three differences between laser light and ordinary light from a lamp.

WAVES Absorption, Reflection, and Transmission of Electromagnetic Waves



Whether a wave is absorbed, reflected, or transmitted depends on the type of radiation and the material. Here are some examples.

Radiation	Metals	Glass	Living Tissue	Water
Radiowaves	Absorbed by aerials, but otherwise reflected	Transmitted	Transmitted	Reflected
Microwaves	Reflected, e.g. satellite dishes and inside of microwave ovens	Transmitted	Transmitted except 12 cm wavelength which is absorbed by water in the tissues	12 cm wavelength absorbed, otherwise transmitted
Infrared	Absorbed by dull/black surfaces, reflected by shiny ones	Transmitted/reflected depending on wavelength	Absorbed	Absorbed
Visible light	Absorbed by dull/black surfaces, reflected by shiny ones	Transmitted	Some wavelengths absorbed, some reflected – giving the tissue a distinctive colour	Transmitted
Ultraviolet	Absorbed	Absorbed	Absorbed and causes ionization	Absorbed
X-rays	Partially absorbed and partially transmitted. The denser the material the more is absorbed		Partially absorbed and partially transmitted. The denser the tissue the more is absorbed	Transmitted
Gamma rays			Transmitted	Transmitted

Questions

- Define the following and give an example of a type of radiation and material that illustrates each:
 - Transmission.
 - Reflection.
 - Absorption.
- Suggest three possible results of the absorption of electromagnetic radiation by a material.
- Copy and complete the table below (look ahead to p33 and 34 if you need help).

Sending signals to mobile phones. Cooking. Aerials. Broadcasting. Suntans. Sterilization. Medical X-rays. Mirrors. Walls of a microwave oven.

	Transmission	Absorption	Reflection
Radiowave			Round the globe broadcasting by bouncing off the ionosphere
Microwave			
Infrared		Cooking	
Visible light	Lenses		
Ultraviolet			
X-ray			
Gamma ray			

WAVES The Earth's Atmosphere and Electromagnetic Radiation

Electromagnetic waves either




pass straight through
the atmosphere

or



are absorbed by
molecules in the atmosphere

or



are scattered by
molecules in the atmosphere





Type of radiation	Effect of the atmosphere	Potential uses	Potential problems
Radiowaves	Generally pass straight through, except some long wavelengths will be reflected by a layer called the ionosphere, high in the atmosphere	Carrying messages over long distances. Bouncing radiowaves off the ionosphere allows them to reach receivers out of the line of sight	
Microwaves	Pass through all parts of the atmosphere	Send information to and from satellites in orbit; send information to and from mobile phones; radar	
Infrared	Absorbed by water vapour and other gases such as carbon dioxide (present in small amounts) and methane (present in minute amounts)	<p>Humans are increasing the amount of greenhouse gases in the atmosphere. Some scientists think this is causing the Earth to warm up. Possible consequences are ...</p> <ul style="list-style-type: none"> • Rising sea levels due to melting of the polar ice caps • Extreme weather conditions occurring more often • Loss of farmland (too wet, dry) 	Infrared is emitted by all warm surfaces including the Earth's surface. Some is lost into space but some is absorbed by gases (water, carbon dioxide) in the atmosphere warming it. This is called the <i>Greenhouse effect</i> and those gases that absorb infrared, greenhouse gases. Too high a concentration of greenhouse gases leads to <i>global warming</i>
Visible light	 Passes through clear skies. Blue light is scattered more than red light giving blue skies during the day and red skies at dawn and dusk Randomly scattered from water vapour in clouds	Provides plants with energy for photosynthesis and hence all living things with food. Warms the Earth's surface	 Molecules in atmosphere Red Evening (sun low in the sky) Sunlight Blue Earth Midday (sun overhead)
Ultraviolet	Absorbed by ozone gas high in the atmosphere (the ozone layer)	Ozone layer protects plants and animals from exposure to too much ionizing ultraviolet radiation from the Sun which would harm them	Ozone layer is being destroyed by chemical reactions with man-made gases
X-rays and gamma rays	Absorbed by the atmosphere		

Questions

1. Which types of electromagnetic radiation pass straight through the atmosphere, which are scattered, and which are absorbed?
2. What is the Greenhouse effect? Suggest why the concentration of carbon dioxide in the atmosphere has been rising for the last 200 years. Suggest three consequences of global warming.
3. Why are cloudy nights generally warmer than when there are clear skies?
4. If the polar ice caps melt, will the Earth's surface absorb more or less radiation from the Sun? Hence will this increase or decrease the rate of global warming?
5. How is the ozone layer helpful to humans and why should we be concerned about a hole in it?

WAVES Uses of Electromagnetic Waves, Including Laser Light

There is an almost limitless range of uses for electromagnetic waves. The selection below gives a flavour of some of the more common.

Type of radiation	
Radiowaves	Broadcasting (long, medium, and shortwave radio, TV [UHF]) (see pages 97, 99). Emergency services communications
Microwaves	<p>Microwaves are strongly absorbed by water molecules making them vibrate violently. This can be used to heat materials (e.g. food) containing water.</p> <p>Microwave energy penetrates more deeply than infrared so food cooks more quickly</p>  <p>Microwaves bounce off the metal walls until absorbed by the food</p> <p>Food must be rotated to ensure all parts are cooked evenly</p>  <p>Sending signals to and from mobile phones or orbiting satellites (see p97)</p>
Infrared	<p>Fibre-optic cables (see p104)</p> <p>Remote controls</p> <p>Toasters and ovens</p> <p>Infrared cameras for looking at heat loss from buildings, night vision, and searching for trapped people under collapsed buildings</p>
Visible light	<p>Seeing and lighting</p> <p>Laser light</p> <p>Surveying, as laser beams are perfectly straight</p>  <p>Fibre-optic cables (see p104)</p> <p>To read CDs, DVDs, and barcodes in shops (see p107)</p> <p>Eye surgery (can be used to 'weld' a detached retina back into place on the back of the eyeball)</p> <p>Retina</p>
Ultraviolet	<p>Can be produced by passing electrical current through mercury vapour</p> <p>If the tube is coated with a fluorescent chemical this absorbs the ultraviolet radiation and emits visible light</p> <p>Fluorescent strip lights</p>  <p>Ultraviolet radiation produced</p> <p>Mercury vapour</p> <p>Electric current</p> <p>Security markers use fluorescent chemicals, which glow in ultraviolet radiation but are invisible in visible light</p> <p>Used for tanning lamps in sun beds</p> <p>Washing powder contains fluorescent chemicals to make clothes look 'whiter than white'</p>
X-rays	Absorption depends on density of the material so can be used to take shadow picture of bones in bodies or objects in luggage (see p108)
Gamma rays	Used to kill cancerous cells Sterilize hospital equipment and food

Questions

- Write a list of all the things you use electromagnetic radiation for during a typical day.
- Food becomes hot when the molecules in it vibrate violently. Suggest one similarity and one difference between how this is achieved in a microwave oven and in a conventional thermal oven.
- Group the uses listed in (1) under the headings:
 - 'Electromagnetic waves used to communicate'.
 - 'Electromagnetic waves used to cause a change in a material'.
 - 'Electromagnetic waves used to gather information'.

WAVES Dangers of Electromagnetic Waves

When electromagnetic radiation is absorbed by the body, it deposits its energy. The more energy deposited, the greater the potential for damage. This depends on the type of radiation, its intensity, and time for which the body is exposed to it.

To reduce the hazard from electromagnetic waves you can reduce the time of exposure, reduce the intensity (for example by moving away from the source or using a lower power source), or by the use of a physical barrier to absorb the radiation.


Type of radiation	Hazard	How to reduce hazard
Non-ionizing. These are a lower hazard		
Radiowaves	Minimal. These generally pass straight through the body and carry little energy	
Microwaves	<p>Low intensity radiation from mobile phones and their transmitter masts may be a health risk, but the evidence is inconclusive</p> <p>Microwaves used in ovens causes a heating effect in water, which would therefore damage water-containing cells</p>	<i>Reduce time of exposure:</i> reduce phone usage <i>Reduce intensity:</i> use a hands free kit to reduce exposure <i>Physical barrier:</i> microwave ovens have metal case and grille over the door to prevent microwaves escaping
Infrared	Absorbed infrared can lead to cell damage, which we call a burn	<i>Reduce time of exposure and intensity:</i> the body has a natural defence mechanism of instinctively moving away from sources of infrared that are uncomfortably hot
Visible light	Only laser light presents a significant hazard	<i>Reduce exposure:</i> never look into the beam <i>Physical barrier:</i> most laser products, especially if high intensity, have the beam shielded
Ionizing – able to break molecules into smaller parts (ions) which may go on to be involved in further (possibly harmful) chemical reactions. If these molecules are in the cells of the body the ions can cause changes to the DNA of the cell causing it to divide and grow incorrectly. This is called <i>cancer</i>		
Ultraviolet	<p>Absorption may cause cell mutations (particularly in skin) which can lead to cancer</p> <p>Sunburn</p>	<i>Physical barrier:</i> sun cream and sun block contain chemicals that strongly absorb ultraviolet providing a barrier between the radiation and the skin Wear clothing <i>Reduce time of exposure:</i> avoid excessive sunbathing or tanning treatment
X-rays	Some absorbed and some transmitted. Absorbed radiation may cause cell mutations leading to cancer	<i>Reduce time of exposure:</i> limit number of X-rays you are exposed to (but sometimes the medical benefits outweigh the potential risks) <i>Physical barrier:</i> health workers use lead shielding to reduce their exposure
Gamma rays	High enough energy to directly kill cells (radiation burns), or to cause cancerous cell mutation	<i>Physical barrier:</i> gamma rays from nuclear power plants are shielded from the outside by thick layers of lead, steel, and concrete <i>Reduce time of exposure:</i> nuclear industry workers have their exposure times carefully monitored and controlled

Questions


1. Suggest three ways that exposure to harmful electromagnetic waves can be reduced.
2. What is the difference between ionizing and non-ionizing radiation?
3. A parent is worried about the possible health risks of a child using a mobile phone while sunbathing in swimwear on a very sunny day. What advice would you give them?

WAVES Reflection, Refraction and Total Internal Reflection


Waves reflect off a plane surface.




Waves travel at different speeds depending on the media they are travelling in.



If the waves meet the boundary at an angle . . .




Think about cars on a road, if they slow down they get closer together but the number of cars passing each second stays the same.



The material light passes through is called the *medium*.


If the speed of light is different in two different media, it also refracts.




As light slows down it changes direction towards the normal (angle of incidence, i , > angle of refraction, r).

As light speeds up it changes direction away from the normal (angle of incidence, i , < angle of refraction, r).



Real and apparent depth





Questions

1. Explain how to decide whether a wave changes direction towards or away from the normal. In what special situation is there no change in direction?
2. Copy and complete the diagram right to show the paths of the two rays through and out the bottom of the glass block.
3. Copy and complete the diagrams right to show what happens to light incident on a glass/air surface when the angle of incidence is less than, equal to, or above the critical angle.
4. Explain, with the aid of a diagram, why waves meeting a boundary where they slow down at an angle change direction.
5. Complete the path of the light ray through this prism and suggest a use for such a prism.




WAVES Refractive Index and Dispersion


When light travels from a vacuum (or air since it makes very little difference to the speed) into another medium, it is slowed down. The amount of slowing is expressed by the ratio:

$$\frac{\text{Speed of light in vacuum (m/s)}}{\text{Speed of light in medium (m/s)}} = \text{refractive index, } n$$

Small refractive index





Large refractive index



Therefore, the bigger the refractive index the greater the change in direction of the light wave as it passes into the medium.

Hence Snell's Law



$$\text{Refractive index } n, = \frac{\sin i}{\sin r}$$



The bigger the refractive index the more the light is slowed as it passes into the medium.

Increasing
refractive
index

Decreasing
refractive
index



Dispersion

Red light has the longest wavelength and is slowed the least. It changes direction the least.


Light slows as it enters the prism and is refracted towards the normal.

White light contains a mixture of wavelengths (colours).

Blue light has the shortest wavelength and is slowed the most. It changes direction the most.

Light speeds up as it leaves the prism and is refracted away from the normal. Again, red light changes direction the least and blue the most.


Spectrum (rainbow) produced on a screen.




Total internal reflection

1. Light must change direction away from the normal so must be going from high to low refractive index.
2. Angle of incidence must be greater than the critical angle.

Small refractive index



Large refractive index



The higher the refractive index of the material, the greater the change of direction away from the normal and therefore, the lower its critical angle.


The critical angle, c , can be calculated from the ratio of the refractive indices either side of the boundary.

$$\text{Sin (critical angle)} = \frac{\text{refractive index of second material}}{\text{refractive index of first material}}$$

$$\text{Sin } c = \frac{n_r}{n_i}$$

Questions


1. Which colour, blue or red, is slowed most as it enters a glass prism?
2. Copy the water droplet and complete the diagram to show how the drop splits the white light into colours. Show the order of these colours on your diagram.
3. The speed of light in a vacuum is 3×10^8 m/s. Show that:
 - a. The refractive index of water is about 1.3 given the speed of light in water is 2.256×10^8 m/s.
 - b. The speed of light in diamond is about 1.2×10^8 m/s given its refractive index is 2.42.
4. The refractive index of glass is about 1.52. A ray of light enters a glass block at 25° to the normal. Show that it continues through the block at about 16° .
5. What is the critical angle for light travelling from water, refractive index 1.33, to air, refractive index 1.00? Why is it not possible to calculate a critical angle for light travelling from air into water?



WAVES Diffraction and Interference

Both diffraction and interference are properties of waves. The fact that all electromagnetic waves display both effects is strong evidence for them having a wave nature.

Diffraction – the spreading out of wave energy as it passes through a gap or past an obstacle.





Light has a very short wavelength (about 5×10^{-7} m), so needs very small gap sizes for diffraction to be noticeable.

Interference – when two waves meet, their effects add.


When two waves arrive in step, they reinforce each other and this is called *constructive interference*. For light the result would be bright and for sound, loud.

When two waves arrive out of step they cancel out and this is called *destructive interference*. For light this would be dark and for sound, quiet.



Questions

1. The speed of sound in air is about 340 m/s. Calculate wavelength of the note 'middle C', frequency = 256 Hz. Hence, explain why a piano can be heard through an open doorway, even if the piano itself cannot be seen.
2. A satellite dish behaves like a gap with electromagnetic waves passing through. Explain why the dish sending the signal to a satellite should have a much wider diameter than the wavelength of the waves, whereas a dish broadcasting a signal from a satellite over a wide area should have the same diameter as the wavelength of the waves.
3. The diagram shows plan view of a harbour. The wavelength of the waves arriving from the sea is 10 m.
 - a. How long is length x ?
 - b. How many waves fit in the length E_1 to B?
 - c. How many waves fit in the length E_2 to B?
 - d. Therefore, will the waves arrive in or out of step at the buoy, B? Hence, describe the motion of a boat tied to it.
 - e. If the wavelength increased to 20 m how would your answers to b-d change?



WAVES Polarization and the Photon Model of Light

Electric field

Magnetic field

Direction of wave motion

Electromagnetic waves are made from an electric and magnetic field oscillating at right angles to each other and to the direction of wave motion.

Symbols

Unpolarized

Electric field oscillates in all possible planes.

Polarized

Electric field oscillates in one plane only.

The arrow shows the plane of oscillation of the electric field.

A beam of electromagnetic radiation is made up of many waves. If the electric field in all the waves only oscillates in one plane (so the magnetic field only oscillates in one plane too), the waves are said to be *polarized*.

A mechanical wave on a string can be polarized by passing it through a frictionless slot; only the oscillation in one plane passes through.

All produce unpolarized light. It contains waves with all possible polarizations.

Polaroid is a material that contains molecules that behave like the slot for visible light.

SUN

Polaroid

Overall, light intensity is reduced.

Only one polarization can pass through.

Polaroid sunglasses help to block this and reduce glare.

Light is polarized by reflection, e.g. from water.

① Electromagnetic radiation always delivers energy in packets (or quanta). These have been called photons.

② The energy carried by each photon is directly proportional to the frequency of the electromagnetic wave.

③ Hence, the electromagnetic spectrum is also a scale of increasing photon energy.

④ The total energy delivered by a beam of electromagnetic radiation (its intensity) depends on ...

Gamma

X-ray

Ultraviolet

Visible

Infrared

Microwave

Radiowave

Increasing Photon Energy

Continuous wave

Absorbing surface

Photons

Energy delivered

Time

Energy delivered

Time

Number of photons arriving per second

Energy each photon carries

Area over which the photons are spread

Diagram illustrating the Photon Model of Light. It shows a continuous wave hitting an absorbing surface, which emits photons. A graph of energy delivered vs. time shows a continuous curve for a continuous wave and a step-like curve for photons. A graph of energy delivered vs. time for photons shows a constant energy per photon multiplied by the number of photons. The total energy is the product of energy per photon, number of photons, and area. The electromagnetic spectrum is shown on the right, with increasing photon energy from radio waves to gamma rays.


Questions

- What do we mean by a polarized wave? Draw a diagram to illustrate your answer.
- Reflected light from a lake in summer is horizontally polarized. Which orientation of light should the Polaroid material in sunglasses allow to pass if the glasses are to cut down glare from the lake?
- What is a photon?
- What type of radiation delivers more energy per photon, X-rays or radiowaves?
- Suggest why X-rays and gamma rays can knock electrons out of atoms (ionize them) but visible light and infrared cannot. What effect might this have on the human body?
- The photons in a beam of electromagnetic radiation carry 4×10^{-17} J each. If 1×10^{18} photons arrive each second over a 2 m^2 area what is the total energy arriving per m^2 ?

WAVES Seismic Waves and the Structure of the Earth

Earthquakes occur when stresses build up at fault lines where the Earth's tectonic plates are moving past each other. The energy stored can be suddenly released as the plates shift, sending out a shock or seismic wave.

Epicentre – the point directly above the focus on the Earth's surface.



Focus – point at which stress between plates is released.

Crust (solid) – maximum 70 km thick.

Mantle (very viscous – behaves almost as a solid).

S wave shadow zone. S waves cannot travel through the molten outer core so this area forms a 'shadow' of the Earth's core. Seismometers in this region do not detect S waves.

Diameter 12 735 km
P waves only

Not to scale

P wave shadow zone – caused by refraction as the P waves enter and leave the core. Seismometers here pick up no waves.



The distance to the epicentre can be estimated by the time lag between the P and S waves arriving.

These seismic waves come in two types
P (primary) waves **S (secondary) waves**

Faster (about 10 km/s)
Longitudinal
Travel through solids and liquids

Cause most damage as they make buildings move side to side

Slower (about 6 km/s)
Transverse
Travel through solids only
Cause less damage as they make buildings move up and down



Outer core (molten, iron and nickel).

Inner core (solid, due to great pressure. Iron and nickel).

Waves follow a curved path as the density of the Earth steadily increases as you go into the Earth. Therefore, their speed changes and they are steadily refracted.


Sudden change in direction at join between layers, because the density of the material suddenly changes so there is a sudden change in wave speed and hence direction by refraction (see p36).

Questions

1. What is the difference between an Earthquake's epicentre and its focus?
2. Draw a labelled diagram of the layers in the Earth. If the crust is a maximum of 70 km thick, what percentage of the total radius of the Earth is made up of crust?
3. Write down two similarities and three differences between P and S waves.
4. Explain how scientists know that the outer core of the Earth is molten.
5. Here is a seismometer trace for an earthquake:
 - a. Which trace, X or Y, shows the arrival of the S waves and which the P waves?
 - b. If the speed of the P waves is 10 km/s and they took 150 s to arrive, how far away was the earthquake?
 - c. If the speed of the S waves is 6 km/s, how long should they take to arrive?
 - d. Hence, what is the time interval t marked on the graph?




WAVES Sound Waves



Questions

- What causes sound? Explain how the sound from a loudspeaker reaches your ear.
- Explain why sound cannot travel in a vacuum.
- Use the formula speed = frequency × wavelength to calculate the range of wavelengths of sound the human ear can hear in air where the speed of sound is about 340 m/s.
- Why does sound travel faster in solids than in gases?
- What does the pitch of a sound wave depend on?
- What does the loudness of a sound wave depend on?
- What is a harmonic?
- Copy this waveform and add:
 - A waveform of twice the frequency but the same amplitude.
 - A waveform of half the amplitude but the same frequency.
 - A waveform of the same amplitude and frequency but of a higher quality.




ELECTRICAL ENERGY Static Electricity

The positively charged nucleus is orbited by negatively charged electrons. These do not escape because opposite charges attract.

All materials are made of atoms

Normally the number of positive and negative charges is equal in each atom




Measured in Coulombs, C.

Static electricity is formed when electrical charges are trapped on an insulating material that does not allow them to move. You can charge up a material by . . .

Like charges repel
Unlike charges attract

1) Friction

Wool duster Electrons (negative charge) rubbed off the wool onto the polythene rod.




Clothing and aircraft can be charged by friction.




Both plane and fuel tanker are connected to Earth before fuelling to prevent any sparks.

2) Induction



Antistatic sprays contain a conducting chemical to avoid the build up of charge.




E.g.

Danger of explosions – flammable vapours or dust ignited by sparks – earthing needed in these environments

Uses and dangers of static electricity

Spraying paint and pesticides




Negatively charged metal object.
Attraction
As drops all have the same charge, they repel giving a fine spray.

Negative charge induced on leaf by positive drops.
Positively charged droplets are attracted to leaf.

When large particles are collected on the plates, they fall down the chimney due to their weight.

Prevents dirty smoke entering the atmosphere.

Electrostatic precipitator




Questions

1. Complete the following sentences. Like charges _____, unlike (opposite) charges _____.
2. What is the unit of electrical charge?
3. What is the difference between an insulator and a conductor?
4. Why would it not be possible to charge a copper rod by rubbing it, no matter how furiously you rubbed?
5. Explain in as much detail as possible how a balloon rubbed on a woolly jumper sticks to a wall.
6. Make a list of all the examples of static electricity in action mentioned in this page. Divide your list into cases where static electricity is useful, where it is a nuisance, and where it is dangerous. Try to add your own examples to the list.


ELECTRICAL ENERGY Electric Currents

An electric current is a flow of charged particles.


Solids




Graphite



Liquids



Gases



What do all these have in common?

- Three conditions for an electric current to flow:
1. There must be charge carriers (electrons or ions).
 2. They must be free to move.
 3. There must be a potential difference to repel them from one side and attract them to the other.

Current flows between plates.


Current (in Amps) is the rate of flow of charge; the number of Coulombs of charge flowing past a point per second.

$$\text{Current (A)} = \frac{\text{Charge (C)}}{\text{time (seconds, s)}}$$

1 Amp = 1 coulomb per second

In equations we usually use I for current and Q for charge. Hence $I = Q/t$.


Electric current is always measured with an ammeter, always placed in series.



Electron current $- \rightarrow +$.


Current rules

1. The current is the same all the way round a series circuit.
Current is **not** used up.




These rules mean that charge is conserved. It does not 'pile up' anywhere in the circuit.

2. The current flowing into a junction = current flowing out.



Questions


1. Why must ionic solids be molten or dissolved to conduct an electric current?
2. In a circuit 4 C of charge passes through a bulb in 2.5 s. Show that the current is 1.6 A.
3. An ammeter in a circuit shows a current of 1.2 A.
 - a. The current flows for 2 minutes. Show the total charge passing through the ammeter is 144 C.
 - b. How long would it take 96 C to pass through the ammeter?
4. In the following circuit, how many Amps flow through the battery?
5. The laws of circuit theory were all worked out in the 1800s. The electron was discovered in 1897. Discuss why we have conventional direct current flowing from positive to negative, when we know that the electrons actually flow from negative to positive.



ELECTRICAL ENERGY Potential Difference and Electrical Energy

What actually happens in an electric circuit?

We can use a model to help us understand what is happening.



We can measure the energy difference between the loaded lorries going into the bulb and the empty ones leaving it using a voltmeter. The voltmeter is connected *across* the bulb to measure how much energy has been transferred to the bulb by comparing the energy (Joules) carried by the lorries (Coulombs) before and after the bulb. *Each Volt represents one Joule transferred by one Coulomb*. The proper name of this is potential difference (because the current has more potential to do work before the bulb than after it) but is often called the voltage.

Questions


1. Why the ammeter readings are the same all the way round a series circuit.
2. Why the total current flowing into a junction is the same as the total current flowing out.
3. Why all the bulbs in a parallel circuit light at full brightness.
4. Why the bulbs get dimmer as you add more in a series circuit.
5. Why the cell goes 'flat' more quickly if you add more bulbs in parallel.

6. Should a 'flat' battery be described as discharged or de-energized? Discuss.
7. This model cannot explain all the features of a circuit. Try to explain:
 - a. How the lorries know to save some energy for the next bulb in a series circuit.
 - b. Whether it takes time for the first full lorries to reach the bulb and make it light up.
 - c. Whether there are full lorries left in the wires when you take the circuit apart.

ELECTRICAL ENERGY Energy Transfers in Series and Parallel Circuits

1 Volt = 1 Joule of energy per Coulomb of charge

The voltage of a cell is a measure of how many Joules of chemical energy are converted to electrical energy per Coulomb of charge passing through it.



Potential difference is the energy given to a device *by* each Coulomb of charge passing through it.

Voltage (sometimes called electromotive force or emf for short) is the energy transferred *to* each Coulomb of charge passing through a source of electrical energy.

The potential difference across a component is a measure of how many Joules are converted from electrical energy to other forms of energy per Coulomb of charge passing through the component.


A bulb converts electrical energy to thermal and light energy.
A motor converts electrical energy to kinetic energy.
A resistor converts electrical energy to thermal energy.
A loudspeaker converts electrical energy to sound energy.

As energy cannot be created or destroyed all the electrical energy supplied by the cell must be converted into other forms of energy by the other components in the circuit.

N.B. Voltmeters connected in *parallel*.

This means that in a *series* circuit the sum of the voltages across the components must equal the voltage across the cell.

The current is the *same* through all components, the potential difference is *shared* between components.



In a *parallel* circuit, each Coulomb of charge only passes through one component before returning to the cell. Therefore, it has to give all the energy it carries to that component. Therefore, the potential difference across each component is the same as the potential difference of the cell.


Potential difference is the *same* across all components, current is *shared* between components.

Questions


- What is a Joule per Coulomb more commonly called?
- A cell is labelled 9 V, explain what this means.
- Explain whether or not voltage splits at a junction in a circuit.
- A 1.5 V cell is connected in series with a torch bulb. The bulb glows dimly. Explain why adding another cell, in series, will increase the brightness of the bulb.
- Considering the same bulb as in question 2, adding a second cell in parallel with the first will make no difference to the brightness. Why not?
- When making electrical measurements we talk about the *current through* a component, but the *voltage across* a component, explain why.
- Try to write down a formula relating voltage, energy, and charge.

ELECTRICAL ENERGY Resistance



Resistance is a measure of how much energy is needed to make something move or flow.



High resistance



Low resistance



We *define* resistance as the ratio of the potential difference across a component to the current flowing through it.

$$\text{Resistance } (\Omega) = \frac{\text{potential difference (V)}}{\text{current (A)}}$$


Unit = Ohms, Ω

i.e. if we have a high resistance then a bigger push is needed to push the current round the circuit.
Note that this is not Ohm's Law, just the definition of resistance.

What causes resistance in wires?

In the lorry analogy on p45, the lorry had to use some energy (fuel) to move along the roads (wires). This represents the resistance of the wires.

Wires have resistance because the electrons moving through the wire bump into the positive metal ions that make up the wire.



The electrons give some of their kinetic energy to the metal ions, which makes them vibrate so electrical energy is converted to thermal energy and the wire gets warm.

The same process happens in a resistor, but the materials are chosen to increase the number of collisions making it more resistant to charge flow.

Factors affecting resistance

Length of wire – the longer the wire the more collisions each electron will make.


Width of wire – the wider the wire the more electrons can flow at any one time, hence you get a bigger current for the same potential difference, so a lower resistance.

Temperature – the hotter the wire the more the metal ions vibrate and so the more likely electrons are to collide.

Material of wire – this affects size and layout of the metal ions and so the number of collisions the electrons make.


Questions

- Show that a resistor with 5 V across it and 2 A flowing through it has a resistance of 2.5Ω .
- A 12Ω resistor has 2.4 V across it. Show that the current flowing is 0.2 A.
- A lamp has a resistance of 2.4Ω and 5 A flows through it. Show the potential difference is 12 V.
- The potential difference across the lamp in (3) is doubled. What would you expect to happen to a. the filament temperature, b. the resistance, c. the current?
- In the following circuit, which resistor has the largest current flowing through it?
- Why do many electronic devices, e.g. computers, need cooling fans?



ELECTRICAL ENERGY Electrical Measurements and Ohm's Law

Experimental technique for measuring resistance



Finding the resistance of a component for a given current

Use meter readings and the formula Resistance (Ω) = potential difference (V) / current (A) to find the resistance.


Datalogging could be used to do this

Use voltmeter and ammeter sensors. Computer software automatically plots voltage vs. current as the variable resistor is altered.

$$\begin{aligned} \textcircled{3} \text{ Gradient of graph} &= \frac{\text{change in } y}{\text{change in } x} \\ &= \frac{\text{change in potential difference, V}}{\text{change in current, A}} \\ &= \text{Resistance, } \Omega \end{aligned}$$

The steeper the line the greater the resistance.

N.B. Beware, sometimes these graphs are plotted with the axes reversed. Then the resistance is not the gradient, it is $1/\text{gradient}$.




Finding the resistance of a component for a range of currents

- ① Record voltmeter and ammeter readings for many different settings of the variable resistor in a table.

Voltage across component (Volts)	Current through component (Amps)

- ② Plot a graph like this



Ohm's Law

A component where the current is directly *proportional* to the voltage is said to obey Ohm's Law and is called *ohmic*.


This means:

1. A graph of V vs. I is a straight line through the origin.
2. $V = I \times R$ where R is constant whatever the value of the current or voltage.

Note that the definition of resistance applies to all components; they are only ohmic if their resistance does not change as the current changes.

Questions

1. Calculate the gradients of the three lines in the graph above and confirm they have the resistances shown.
2. 1.5 A flows in a 1 m length of insulated wire when there is a potential difference of 0.3 V across it.
 - a. Show its resistance is 0.2Ω .
 - b. If 0.15 A flows in a reel of this wire when a potential difference of 3 V is placed across it, show that the length of the wire on the reel is 100 m.
3. Current and voltage data is collected from a mystery component using the method above. When plotted the graph looks like this:
Is the resistance of the component increasing, decreasing, or staying the same as the current increases?



ELECTRICAL ENERGY Power in (Ohmic) Electrical Circuits

The general definition of power is : Power (W) = $\frac{\text{energy transferred (J)}}{\text{time taken (s)}}$

We also know that 1 Volt = 1 Joule per Coulomb.

As a formula we represent this as potential difference (Volts) = $\frac{\text{energy transferred (Joules)}}{\text{charge passing (Coulombs)}}$

We also know that current (Amps) = $\frac{\text{charge passing (Coulombs)}}{\text{time taken (seconds)}}$


Then current \times voltage = $\frac{\text{charge passing}}{\text{time taken}} \times \frac{\text{energy transferred}}{\text{time taken}} = \frac{\text{energy transferred}}{\text{time taken}} = \text{Power}$

Also as voltage = current \times resistance then

$$\text{Power} = \text{current}^2 \times \text{resistance}$$

$$P = I^2 R$$

Mains appliances use 230 V and always have a power rating.




We can use this information to calculate the current that flows through them when working normally.

Device	Voltage (V)	Power (W)
Filament bulb	230	60
Energy efficient lamp	230	9
Kettle	230	1500
Microwave oven	230	1600
Electric cooker	230	1000–11 000
TV set	230	30


Questions

- Redraw the circuit using standard circuit symbols adding voltmeter to measure the potential difference across the lamp and an ammeter to measure the current through it.
 - The voltmeter reads 6 V. How many Joules of energy are transferred per Coulomb?
 - The ammeter reads 2 A. How many Coulombs pass through the lamp each second?
 - Hence, how many Joules per second are transferred to the lamp?
 - If the voltmeter now reads 12 V and the ammeter still reads 2 A then how many Joules are transferred to the lamp each second?
- A 1.5 V cell is used to light a lamp.
 - How many Joules does the cell supply to each Coulomb of electric charge?
 - If the current in the lamp is 0.2 A, how many Coulombs pass through it in 5 s?
 - What is the total energy transferred in this time?
 - Hence, show the power of the lamp is 0.3 W.
- A 6 V battery has to light two 6 V lamps fully. Draw a circuit diagram to show how the lamps should be connected across the battery. If each draws a current of 0.4 A when fully lit, explain why the power generated by the battery is 4.8 W.
- Use the data in the table above to show that the current drawn by an 'energy efficient' lamp is over 6 times less than the current drawn by a normal filament bulb.
- Show that a 60 W lamp with a potential difference of 240 V across it has a resistance of 960 Ω .




ELECTRICAL ENERGY Properties of Some Electrical Components

A graph of voltage against current (or vice versa) for a component is called its characteristic. This circuit can be used to measure the characteristic of component X.



The cell can be reversed to measure negative values of current and voltage.


N.B. Check carefully whether current or voltage is plotted on the x axis.




All the following components are *non-ohmic* as their resistance is not independent of the current flowing through them.

1. Filament lamp

Symbol




As the current increases, the filament wire in the bulb becomes hotter.



Eventually it glows red, and then white, hot.

Increasing current in the wire means the electrons make more collisions with fixed ions in the wire.




Electrons find it harder to move; you need a greater potential difference to drive the same current so the resistance increases.

2. Diode

Symbol

A diode only allows current to flow in one direction.

Forward bias – very low resistance, high current flow



Reverse bias – extremely high resistance, negligible current flow


In the reverse direction, the current is almost negligible until very large voltages are reached when the diode may fail.

Diodes are made from two types of semiconductor

n-type – excess of negative charges



Small gradient




The forward current ‘turns on’ at about 0.5 V and very large currents can be achieved by 0.7 V when there is almost no resistance in the diode.


p-type – shortage of negative charge. The spaces left are called holes and behave as positive charges.

Electrons will flow $n \rightarrow p$ and holes $p \rightarrow n$. Therefore, the diode will conduct when the n-type end is negative and the p-type end is positive.

3. Light dependent resistor (LDR)



4. Thermistor




Notice that this is the opposite behaviour to a wire, whose resistance increases with increasing temperature.

Thermistors are used to control circuits that need to respond to temperature changes, e.g. to switch off a kettle.


Questions

1. Draw the circuit symbols for: a. A filament lamp. b. An LDR. c. A thermistor. d. A diode.
2. Sketch a graph of current against voltage for a filament lamp. Explain in terms of the motion of electrons through the filament the shape of the graph.
3. Show that a thermistor with a potential difference of 3 V across it and a current of 0.2 A flowing through it has a resistance of 15Ω . If the temperature of the thermistor was raised, what would you expect to happen to its resistance?
4. Show that an LDR with a potential difference of 1.5 V across it and a current of $7.5 \times 10^{-3} \text{ A}$ (7.5 mA) flowing through it has a resistance of 200Ω . If the LDR is illuminated with a brighter light, with the same potential difference across it what would you expect to happen to the current flowing in it and why?
5. Sketch a graph of current against voltage (both positive and negative values) for a diode. Use it to explain why a diode only passes current in one direction.
6. Consider the following circuits. In which circuit will the ammeter show the greatest current?
7. A student plans to use a thermistor to investigate how the temperature of the water in a kettle varies with time after it is switched on.
 - a. Draw a circuit involving an ammeter and voltmeter the student could use.
 - b. Explain how they would use the ammeter and voltmeter readings together with a graph like the one printed above on this page, to find the temperature of the water at any given time.




ELECTRICAL ENERGY Potential Dividers

Two resistors in series form a potential divider.




The current in both resistors is the same as they are in series. The resistor with the greater value will take more voltage to drive the current through it so has the greater potential difference across it.



The potential difference of the cell, V_{in} , is divided between the two resistances in the ratio of their resistances.

The output voltage can be calculated using the formula:

$$V_{out} = V_{in} \times \left(\frac{R_2}{R_1 + R_2} \right)$$





Slider divides the resistance into two separate resistors, R_1 and R_2 . The position of the slider determines the relative size, hence ratio, of the two resistances.

If one of the resistors is a thermistor, the output voltage depends on temperature.

Thermistors have a high resistance in the cold.


Questions

- Use the formula above to calculate V_{out} if R_1 and R_2 in the circuit provided have the following values:
 - $R_1 = 10 \Omega$, $R_2 = 20 \Omega$.
 - $R_1 = 20 \Omega$, $R_2 = 10 \Omega$.
 - $R_1 = 1 \text{ k}\Omega$, $R_2 = 5 \text{ k}\Omega$.
 - $R_1 = 1.2 \text{ k}\Omega$, $R_2 = 300 \Omega$.
- For each of the pairs of resistors in question 1, decide whether R_1 or R_2 has the greater potential difference across it.
- An LDR has a resistance of 1000Ω in the light and $100\,000 \Omega$ in the dark. In the circuit, the variable resistor is set to 5000Ω . Calculate V_{out} in the light and in the dark. If the resistance of the variable resistor is reduced, will the values of V_{out} increase or decrease?
- Draw a potential divider circuit where the output rises as the temperature rises. Suggest a practical application of this circuit.



ELECTRICAL ENERGY Electric Cells, Alternating and Direct Current

Electron flow allows chemicals inside the cell to react.



Electric cells convert stored chemical energy into electrical energy.


Chemicals only react while current is flowing.

Eventually the reaction is complete and no further stored chemical energy can be released from the cell.

Circuit symbols




Battery (many cells)



A number of cells joined are a battery.

Current produced by cells only ever flows one-way.


Conventional current – positive to negative



Electron current – negative to positive

Both these currents are *direct currents* (or d.c.). They flow consistently in one direction.

Compare this with mains electricity.



In the UK, the frequency is 50 Hz (cycles per second).

Alternating currents are widely used because:

- They are easier to generate (p83–85).
- They are easier to distribute (p86–7) than direct currents.

Many devices need direct current to work so alternating current often has to be converted to direct current (p54).


Peak value of supply potential difference = 325 V. But, because it only has the peak value for a short time the supply transfers an equivalent power to a 230 V direct current supply.

Questions


1. Sketch a labelled graph of the variation of supply potential difference with time (for 10 seconds) for alternating current of frequency 2 Hz, and peak value 3 V. Add to the graph a line showing the output from a battery of terminal potential difference 2 V.
2. The capacities of two cells are AA = 1.2 Amp-hours and D = 1.4 Amp-hours. How long will each cell last when supplying:
 - a. A current of 0.5 A to a torch bulb?
 - b. 50 mA to a light emitting diode?
3. Some people claim that battery powered cars do not cause any pollution. A battery is just a store of electrical energy so where do battery-powered cars really get their energy from? Hence, are they really non-polluting, or is the pollution just moved elsewhere?
4. Draw up a table of advantages and disadvantages of batteries compared to mains electricity. Consider relative cost, how they are used, potential power output, and impact on the environment.

ELECTRICAL ENERGY Diodes, Rectification and Capacitors



Although alternating current is easier to generate and distribute, many appliances, especially those with microchips, need direct current. The process of converting alternating to direct current is called *rectification*.



A better result is obtained with a bridge circuit.



A capacitor is a component that can store charge.



As the potential difference starts to fall it releases the charge. This helps to keep the current and potential difference up and we say the direct current is smoothed.

Questions

1. What is a diode?
 - a. Complete the graphs in the circuit below to show the effect of the diode.
 - b. Why is the output an example of direct current? Why do we say it is 'unsmoothed'?
 - c. If the diode were reversed what would be the effect, if any, on the direct current output?
2. What name do we give a device that stores charge?
3. Explain the difference between full wave rectification and half-wave rectification. Illustrate your answer with voltage-time graphs.
4. Draw a circuit that produces full wave rectification. Show how the current flows through the circuit.



ELECTRICAL ENERGY Mains Electricity and Wiring




N.B. Never inspect any part of mains wiring without first switching off at the main switch next to the electricity meter.

The UK mains electricity supply is alternating current varying between +325 V and -325 V with a frequency of 50 Hz. It behaves as the equivalent of 230 V direct current.

Earth connection is connected to the Earth, usually by a large metal sheet buried in the ground.

Live connection varies between +325 V and -325 V with respect to neutral.



Neutral connection stays at 0 V with respect to the Earth.

The Earth acts as a vast reservoir of charge; electrons can flow into it or out of it easily. Therefore, its potential is 0 V.

Touching the live wire is dangerous because if you are also connected to Earth, electrons can flow across the potential difference between Earth and live, through you. This will give you a shock.

When the live is at +325 V electrons flow from neutral to live.

When the live is at -325 V electrons flow from live to neutral.

Ensure wires are connected firmly, with no stray metal conductors.

Three pin plug

Ensure cable grip is tightened, so if the cable is pulled, the conductors are not pulled out of their sockets.

Most appliances use three core cables.

Therefore, switches and fuses are always placed in the live wire.

Yellow/green wire – Earth wire. This can never become live, as it will always conduct electrons into or out of the Earth, so its potential will always be 0 V. All metal cased appliances must have an Earth connection.

Brown wire, live.
Blue wire, neutral.

Needed to complete the circuit.

Inner insulation to separate the conductors.

Outer insulation to protect the conductors from damage.

The current drawn by an appliance can be calculated using the equation:

$$\text{Current} = \frac{\text{Power}}{\text{voltage}} = \frac{\text{Power}}{230 \text{ V}}$$

Power ratings can be found on the information label on the appliance.


Questions

- What colours are the following electrical wires: live, neutral, Earth?
- My kettle has a power output of 1 kW and my electric cooker 10 kW. What current will each draw? Why does the cooker need especially thick connecting cables?
- Some countries use 110 V rather than 230 V for their mains supply. Suggest how the thickness of the conductors in their wiring would compare to the conductors used in the UK. How will this affect the cost of wiring a building? What advantages does using a lower voltage have?
- Study this picture of a three-pin plug – how many faults can you find?
- Placing a light switch in the neutral wire will not affect the operation of the light but could make changing a bulb hazardous. Why?



ELECTRICAL ENERGY Electrical Safety

N.B. Never inspect any part of mains wiring without first switching off at the main switch next to the electricity meter.




Questions

- Choose (from 3 A, 5 A, and 13 A) the most appropriate fuse for the following:
 - An electric iron of power output 800 W.
 - A table lamp of power output 40 W.
 - A washing machine of total power 2500 W.
- Explain why a fuse must always be placed in the live wire.
- Explain why a double insulated appliance does not need an Earth wire, but does need a fuse.
- The maximum current that can be safely drawn from a normal domestic socket is 13 A. At my friend's house, I notice a 2.5 kW electric fire, an 800 W iron, and three 100 W spot lamps all connected to a single socket. What advice should I give my friend? Use a calculation to support your answer.
- While using my electric lawnmower I cut the flex, and the live wire comes into contact with the damp grass. An r.c.d will make this wire safe very quickly. What is an r.c.d and how does it work in this case?

ELECTRICAL ENERGY Electron Beams

Not all electric currents flow in wires. It is possible to produce a beam of electrons travelling through a vacuum.




Questions


- Describe the process of thermionic emission. Why is it important that the electron beam be produced in a vacuum?
- What would happen to the kinetic energy of the electrons produced by an electron gun if the potential difference between the heated filament and the accelerating anode was increased?
- What would happen to the charge transferred per second (the current) in the electron beam if the heater temperature was increased but the accelerating potential was not changed? Would the kinetic energy of the electrons change?
- Given that the charge of one electron is 1.6×10^{-19} C, show that the kinetic energy of an electron in the beam is 3.2×10^{-17} J when the accelerating potential is 200 V.
- If the current in the electron beam is 2 mA, show that the number of electrons boiled off the filament each second is about 1.3×10^{16} [charge on the electron = 1.6×10^{-19} C].
- Use the answers to questions 4 and 5 to show that the total energy delivered by the beam per second (i.e. its power) is 0.4 W.
- An electron beam passes through two charged plates as shown in the diagram. What would be the effect on the deflection of:
 - Increasing the potential difference across the deflecting plates?
 - Decreasing the accelerating voltage across the electron gun?

MAGNETIC FIELDS Magnetism and the Earth's Magnetic Field

A magnetic field is a region of space in which magnets and magnetic materials feel forces. The only magnetic materials are iron, steel, nickel, and cobalt. We represent magnetic fields by drawing magnetic field lines.



The Earth's magnetic field




Questions

1. What is a magnetic field? Make a list of three properties of magnetic field lines.
2. Make a list of the four magnetic materials. How could you test an unknown material to discover whether it is one of the four in the list?
3. Using a magnet how would you tell if a piece of steel was magnetized or un-magnetized?
4. If the Earth's magnetic field were to disappear, it would be very bad news for our health. Explain why. (You might need to look at p69.)
5. Why might a magnetic compass not work very well close to the North or South Pole?

MAGNETIC FIELDS Electromagnetism and The Motor Effect

A current carrying wire produces a magnetic field around it.


Plotting compasses line up with the field.




The magnetic field of a wire can be made to interact with another magnetic field to produce a *catapult field*, which exerts a force on the wire.

Circular field lines (getting further apart as the field gets weaker further from the wire).


A coil (or solenoid) produces a magnetic field through and around it.




Right Hand Grip Rule



The more turns on the coil, a bigger current and adding a soft iron core all strengthen the magnetic field.




Fields cancel out as they are in opposite directions.



Magnetic field lines try to get as short as possible by pushing the wire out of the field and removing the distortion.

The directions of the external field, current, and the resulting force (and hence motion of the wire) are remembered by Fleming's Left Hand Rule.



Reversing either current or field will reverse direction of the force.


If the current is parallel to the external magnetic field the two magnetic fields are at right angles to each other and cannot interact so no force is produced.

Size of the force can be increased by:

- Using a larger current
- Using a stronger external field

Questions

1. In what ways are the fields around a bar magnet and around a long coil (solenoid) similar and in what ways are they different?
2. What would happen to the direction of the magnetic field lines around a wire, or through a coil, if the current direction reverses?
3. Make a list of five uses for an electromagnet and suggest why electromagnets are often more useful than permanent magnets.
4. What happens to the direction of the force on a current carrying wire if both the field and current directions are reversed?
5. Copy the diagrams (right) and add an arrow to show the direction of the force on the wire.




Thermal Energy Heat and Temperature – What is the Difference?

All energy ultimately ends up as heat. In most energy transfers, a proportion ends up as heat energy, and often this is not useful. Sometimes we want to encourage heat transfers, in cooking for example, and sometimes discourage them, in preventing heat losses from your home for example. Therefore, understanding heat energy and how it is transferred is important.


Are heat and temperature the same thing?

We define heat energy as the total kinetic energy of the particles in a substance (in Joules).

Identical kettles, both switched on for the same time.




Same electrical energy supplied to both. Therefore, the water gains the same amount of heat energy. Therefore, the total kinetic energy of the particles in both kettles will be the same.



We have a special name for this average; we call it *temperature*.

Experience tells us that kettle B is hotter than A. This means that the particles in B have a higher average kinetic energy than those in A. This is reasonable because the same amount of energy is spread over fewer particles in B than in A.

Temperature differences tell us how easily heat is transferred. The bigger the temperature difference between an object and its surroundings the more easily heat will be transferred.




Questions

- Explain how a bath of water at 37°C can have more heat energy than an electric iron at 150°C.
- A red-hot poker placed in a small beaker of water will make the water boil, but placed in a large bucket of water the temperature of the water only rises a few degrees, why?
- Which should lose heat faster, a mug of tea at 80°C in a fridge at 5°C, or the same mug of tea at 40°C, placed in a freezer at -10°C?

Temperature (in Kelvin) is proportional to the average kinetic energy of the particles.

Temperature Scales




Questions

- Convert the following into Kelvin: 42°C, 101°C, -78°C, -259°C.
- Convert the following into $^{\circ}\text{C}$: 373K, 670K, 54K, 4K.

Thermal Energy Specific and Latent Heat

When an object cools, it transfers heat to its surroundings.

Consider



This tells us that 1 kg of aluminium has less heat energy stored in it than 1 kg of water, so the average kinetic energy (temperature) of the particles when mixed is less. We say aluminium has a lower specific heat capacity than water.

Since temperature is proportional to the average kinetic energy of the particles we are actually measuring the energy needed to increase the average kinetic energy of the particles by a set amount. This will depend on the structure of the material, i.e. what it is made of and whether it is a solid, liquid, or gas. Therefore, *all materials have their own specific heat capacities*.

Latent heat is a measure of the energy needed to completely melt or boil 1 kg of a material.


Energy (J) = mass (kg) × specific latent heat (J/kg)

$$\Delta E = m \times slh$$

Units J/kg

Specific latent heat depends on the strength and number of intermolecular bonds between molecules, so depends on the material and its state.

Heat energy used to raise average kinetic energy of molecules, therefore temperature rises



Temperature of water / aluminium when they come to equilibrium <50°C (and no heat has been lost from the container).

Specific heat capacity is a measure of how much heat energy 1 kg of a material can hold, defined as:


The energy needed to be supplied to raise the temperature of 1 kg of a material by 1K.

$$\text{Units J/kgK}$$

Energy supplied (J) = mass (kg) × specific heat capacity (J/kgK) × temperature change (K).

$$\Delta E = m \times shc \times \Delta T$$

Energy transferred is used to break bonds between molecules, not to increase their kinetic energy (temperature).




Questions

- What happens to the average kinetic energy of the particles in material when the temperature rises?
- A pan of boiling water stays at 100°C until all the water has evaporated. Why?
- Explain why adding ice to a drink cools it down.
- Given that specific heat capacity of water = 4200 J/kgK and of steam = 1400 J/kgK and that the specific latent heat for melting ice is 334 000 J/kg and for boiling water = 2 260 000 J/kg show that if the graph in the text above represents 2.5 kg of water:
 - The energy supplied between a and b is 835 000 J.
 - The energy supplied between b and c is 1 050 000 J.
 - The energy supplied between c and d is 5 650 000 J.
- A student finds that it takes 31 500 J to heat a 1.5 kg block of aluminium from 21°C to 44°C. Show that the specific heat capacity of aluminium is about 900 J/kgK.


Thermal Energy Heat Transfer 1 – Conduction

Conduction is the transfer of thermal energy from a high temperature region to low temperature region by the transfer of kinetic energy between particles in a material.



Heat Transfer 2 – Convection

Convection happens in *fluids* because the particles can move and carry heat energy away from the source.
Hot fluids have a lower density than cool fluids.




Questions – conduction

- Using the idea of particles explain why metals are such good conductors of heat and why air is a bad conductor.
- Air is often trapped, for example between layers of clothing, to reduce conduction. Make a list of five places where air is trapped to prevent conduction.
- Stuntmen can walk (quickly) across a bed of burning coals without injury, yet briefly touching a hot iron causes a painful burn, why?


Questions – convection

- Explain the result opposite using the ideas of conduction and convection:
- Why is the heating element at the bottom of a kettle?
- Suggest what causes currents in the oceans (in detail).




Thermal Energy Heat Transfer 3 – Radiation

Thermal radiation is the transfer of heat energy by (infrared) electromagnetic waves (see p30 and 32).




The greenhouse effect



Temperature rises due to the reabsorption of the trapped heat radiation. This leads to global warming.


Questions

1. By which method of heat transfer does the heat from Sun reach Earth? How can you tell?
2. Why are solar panels fixed to roofs and designed to heat water painted black?
3. Why are many teapots made of shiny steel?
4. Explain why it is important to reduce the amount of carbon dioxide we pump into the atmosphere.
5. Look at the following diagram of a thermos flask and explain why:
 - a. There is a vacuum between the walls of the flask.
 - b. The walls of the flask are shiny.
 - c. The drink stays hotter longer if the stopper is put in.
 - d. Liquid nitrogen (boiling point 77K, -196°C) stays as a liquid in the flask for a long time, but rapidly boils and evaporates if poured out.



Thermal Energy Reducing Energy Wastage in Our Homes

Reducing our demand for energy is as important in reducing greenhouse gas emissions as finding renewable energy resources. Although energy is conserved, we often convert it to forms that are not useful. For example:



To reduce our demand for energy there are many practical steps we can take, many involving the reduction of heat transfer. However, householders must also consider payback time.

$$\frac{\text{Cost of installing}}{\text{annual saving}} = \text{payback time in years.}$$



Many of these methods rely on trapping pockets of air. Air, because it is a bad conductor, and trapped, to prevent it convection.

Loft insulation made of fibreglass or mineral wool. Annual saving £180–220 and 1500 kg CO₂.


Hot water tank jackets – air trapped in fibres. Annual saving £10–20, 50 kg CO₂.

Energy saving light bulb. Annual saving £9, 40 kg CO₂. Produces much less waste heat.

A condensing boiler condenses hot waste gas from the burnt fuel making it give up its latent heat rather than pumping it straight into the atmosphere.

Not leaving appliances on standby. Annual saving £20, 40 kg CO₂.

Turn down thermostats – the smaller the temperature difference between inside and outside the slower the rate of loss of thermal energy. 1°C reduction has an annual saving of £50 and 300 kg CO₂.



Saving energy also reduces carbon dioxide emissions because carbon dioxide is a waste product of burning any fossil fuel, either directly such as gas in a boiler, or indirectly to generate electricity in a power station.



(Data correct (2006) Energy Saving Trust www.est.org.uk.)

Questions

1. A householder could spend £230 on loft insulation that would save £180 in fuel bills each year, or they could spend £75 on draughtproofing and save £20 each year. Which would you recommend they do and why?
2. Why do we talk about wasting energy when physics tells us energy is conserved?
3. Which of the energy saving measures above are free?
4. The annual savings quoted above are both in terms of money and CO₂ saved. Which do you consider to be more important and why?

Thermal Energy Kinetic Model of Gases

The kinetic model of gases is the name we give to the idea that a gas is made up of microscopic particles moving randomly, colliding with each other and the walls of the container.




Questions


- Explain carefully why heating a gas in a sealed container raises the pressure. Why might this not be very safe?
- A motorist checks his tyre pressures before a long journey. At the end of the journey, he notices his tyres are warm and that their pressure has risen, why?
- A toy balloon containing helium is released accidentally by a child. The balloon rises high into the atmosphere where the air pressure is a lot lower. Eventually it bursts. Why?
- When you breathe in your lung volume increases. What happens to the air pressure in your lungs, and why does air rush in through your nose?
- When petrol burns in a car engine it gets very hot very quickly. Why does this force the cylinder out?
- When carbon dioxide is released rapidly from a fire extinguisher, it makes the nozzle get very cold, why?

THEMAL ENERGY Gas Laws


Measuring the variation of gas pressure with temperature:



It would make sense to start a temperature scale here – we call this the Kelvin scale.



Measuring the variation of gas pressure with volume:



Questions

- The pressure of air in a sealed container at 22°C is 105 000 Pa. The temperature is raised to 85°C. Show that the new pressure is about 130 000 Pa assuming that the volume of the container remains constant.
- A bubble of air of volume 2 cm³ is released by a deep-sea diver at a depth where the pressure is 420 000 Pa. Assuming the temperature remains constant show that its volume is 8 cm³ just before it reaches the surface where the pressure is 105 000 Pa.
- A sealed syringe contains 60 cm³ of air at a pressure of 105 000 Pa and at 22°C. The piston is pushed in rapidly until the volume is 25 cm³ and the pressure is 315 000 Pa. Show that the temperature of the gas rises to about 95°C.
- When a star forms a gas cloud in space is attracted together by gravity compressing it. As the volume of the gas reduces what happens to its pressure and hence temperature?

RADIOACTIVITY Atomic Structure

All atoms have the same basic structure:


In all atoms the number of protons = number of electrons. This makes atoms uncharged, or *neutral*.

Naming atoms:

Atomic (proton) number Z = number of protons in the nucleus

Mass (nucleon) number A = total number of protons plus neutrons in the nucleus


Symbol for the element A_Z^X



What is Radioactivity?

Some elements give out random bursts of radiation. Each individual nucleus can only do this once, and when it has happened, it is said to have decayed. As even a tiny sample of material contains billions of atoms, many bursts of radiation can be emitted before all the nuclei have decayed.

Ionizing – it can knock electrons out of other atoms.




Elements that behave like this are called *radioactive*.

We can measure the radioactivity as the number of decays (and, therefore, bursts of radiation emitted) per second.

1 decay per second = 1 Becquerel, Bq

Each element has a unique number of protons. Therefore, the atomic number uniquely identifies the element.

Some atoms of the same element have different numbers of neutrons.



E.g. all these atoms are carbon as they all have 6 protons, but they have different numbers of neutrons. They are called isotopes of carbon.

Isotopes are always the same element, i.e. same atomic number but have different numbers of neutrons (and so mass number).

The relative masses of protons, neutrons, and electrons and their relative electric charges are:

	Mass	Charge
Proton	1	+1
Neutron	1	0
Electron	$\frac{1}{1870}$	-1

Questions

1. Copy and complete the table.

	No. of protons	No. of electrons	No. of neutrons
Carbon $^{12}_6\text{C}$			
Barium $^{137}_{56}\text{Ba}$			
Lead $=^{208}\text{Pb}$		82	125
Iron ^{56}Fe	26		
Hydrogen $\frac{1}{1}\text{H}$			
Helium $\frac{4}{2}\text{He}$			
Helium $\frac{3}{2}\text{He}$			
Element X A_Z^X			


2. a. Draw a diagram to show all the protons and neutrons in the nuclei of $^{35}_{17}\text{Cl}$ and $^{37}_{17}\text{Cl}$.
b. What word do we use to describe these two nuclei?
c. Why is there no difference in the way the two types of chlorine atoms behave in chemical reactions?
d. If naturally occurring chlorine is 75% $^{35}_{17}\text{Cl}$ and 25% $^{37}_{17}\text{Cl}$ explain why on a periodic table it is recorded as $^{35.5}_{17}\text{Cl}$?
3. What is a Becquerel?
4. If ionizing radiation knocks electrons out of atoms, will the ions left behind be positively or negatively charged? Why?
5. Explain what you understand by the term 'radioactive element'.

RADIOACTIVITY A History of Our Understanding of the Atom

In 1803, John Dalton noted that chemical compounds always formed from the same ratio of elements, suggesting particles were involved. He called these atoms from the Greek, meaning indivisible.

J.J. Thomson (1897) discovered the electron, a particle that could be knocked out of an atom. He suggested a 'plum pudding' model of the atom.


Rutherford, Geiger, and Marsden investigated this in 1910. They decided to probe the nucleus further with alpha particles. These are particles with two positive charges, which they considered to be like little bullets.




1. Detector detects the alpha particles that have travelled through the foil. It can be moved to any angle round the foil so that the number of alpha particles in any direction can be recorded.

3. A very small number were turned through angles greater than 90°.

5. Rutherford proposed the nuclear model.



2. The majority of alpha particles travelled through the foil with very little change in direction.



4. Plum pudding model cannot explain this since as the positive and negative charges were reasonably evenly distributed no alpha particles should get scattered through large angles.


Summary

The alpha scattering experiment proves that:

1. Atoms have massive, positively charged nuclei.
 2. The majority of the mass of the atom is the nucleus.
 3. Electrons orbit outside of the nucleus.
- Most of the atom is empty space.

Bohr further developed the atomic model by suggesting that the electrons were arranged in energy levels around the nucleus.

To move up a level it has to absorb precisely the right amount of energy from an electromagnetic wave.



Kinetic energy is transferred to potential energy in the electric field round the nucleus as the alpha particle does work against the repulsive force. This is returned to kinetic energy on leaving the region near the nucleus.

- The larger the charge on the nucleus the greater was the angle of scatter.
- The thicker the foil the greater the probability that an alpha particle passes close to a nucleus.
- Slower alpha particles remain in the field around the nucleus for longer – increases the angle of scattering.

If an electron moved down a level, it has to get rid of some energy in the form of an electromagnetic wave.


Questions

1. List the main conclusions of the alpha scattering experiment.
2. What evidence did Thomson have for the plum pudding model?
3. Suggest why the alpha scattering apparatus has to be evacuated (have all the air taken out of it).
4. Suggest why the gold foil used in the alpha scattering experiment needs to be very thin.
5. The diameter of an atom is about 10^{-10} m and of a gold nucleus 10^{-14} m. Show that the probability of directly hitting a nucleus with an alpha particle is about 1 in 108. What assumptions have you made?

RADIOACTIVITY Background Radiation

Radioactive elements are naturally found in the environment and are continually emitting radiation. This naturally occurring radiation is called *background radiation*, which we are all exposed to throughout our lives.


Background radiation comes from a number of sources. (Note that these are averaged across the population and may differ for different groups, for example depending on any medical treatment you may have, or whether you make many aeroplane flights.)



(Source: Health Protection Agency, 2006.)


One of the major sources of background radiation is radon gas. This is produced by minute amounts of uranium, which occurs naturally in rocks, and is present in all parts of the country. It disperses outdoors so is only a problem if trapped inside a building. Exposure to high levels of radon can lead to an increased risk of lung cancer.

Since we all inhale radon throughout our lives it accounts for about half our annual radiation dose in the UK.



Maps courtesy of the Health Protection Agency and the British Geological Survey
For more information go to:
www.ukradon.co.uk

Geological conditions in some areas produce higher than average radon concentrations as shown in the map.




Questions

1. Make a list of sources of background radiation.
2. Give at least two reasons why the percentages shown above in the sources of background radiation are only averages and will differ for different people.
3. On average what percentage of the total background radiation is man-made?
4. Should we worry about background radiation?

RADIOACTIVITY Three Types of Nuclear Radiation

There are three types of radiation emitted by radioactive materials. They are all emitted from unstable *nuclei*:



Name	Identity		Mass	Charge	
Alpha (α)	Helium nucleus	 ${}^4_2 \text{He}$	4	+2	Massive and highly charged. Therefore, interacts strongly with other matter causing ionization, and loses energy rapidly. Easily stopped and short range
Beta (β)	Fast moving electron ejected from the <i>nucleus</i> . Note that it is not an atomic orbital electron		$\frac{1}{1870}$	-1	Nearly $8000 \times$ less massive than alpha and only half the charge. Therefore, does not interact as strongly with other matter causing less ionization, and loses energy more gradually. Harder to stop and has a longer range
Gamma (γ)	Electromagnetic wave		0	0	No mass or charge so only weakly interacts with matter. Therefore, very difficult to stop


Questions

- Describe the differences between alpha, beta, and gamma radiation. What materials will stop each one?
- Alpha and beta particles are deflected in both electric and magnetic fields but gamma is not. Explain why. Why are alpha and beta deflected in opposite directions?
- A student has a radioactive source. When the source is placed 1 cm in front of a GM tube connected to a ratemeter it counts 600 counts per minute.
 - Moving the source back to 10 cm the count drops to 300 counts per minute.
 - Replacing the source at 1 cm and inserting 2 mm thickness of aluminium foil gives 300 counts per minute.
 - Moving the source back to 5 cm and inserting 2 cm of lead gives 150 counts per minute.
 Explain how you know what type(s) of radiation the source emits.
- Many smoke alarms contain a small radioactive source emitting alpha particles. This is inside an aluminium box, and placed high on a ceiling. Use the properties of alpha particles to explain why smoke alarms do not pose any health risk.

RADIOACTIVITY Radioactive Decay and Equations


Most nuclei never change; they are stable. Radioactive materials contain unstable nuclei. These can break up and emit radiation. When this happens, we say the nucleus has *decayed*. The result for alpha and beta decay is the nucleus of a different element. For gamma decay, it is the same element but it has less energy.

Alpha decay




Mass number decreases by 4 (2 protons + 2 neutrons lost). Atomic number decreases by 2 (2 protons lost).

Atomic number



Or




Beta decay





Neutron becomes a proton and electron.

Daughter nucleus has one more proton than the parent so the atomic number increases by one.

Overall number of protons plus neutrons is unchanged so the mass number does not change.




Or




Proton becomes a neutron and a positron (an anti-electron with all the same properties as an electron but the opposite charge).

Daughter nucleus has one less proton than the parent so the atomic number decreases by one.

Overall number of protons plus neutrons is unchanged so the mass number does not change.



Or



Gamma decay

Often after either alpha or beta decay the nucleons have an excess of energy. By rearranging the layout of their protons and neutrons, they reach a lower energy state and the excess energy is emitted in the form of a gamma ray.



Rules for nuclear equations

The total mass number must be the same on both sides of the equation.

The total atomic number on both sides of the equation must be the same.

The total charge must be the same on both sides of the equation.

Questions

Copy and complete the following nuclear equations:

1. ${}_{84}^{215}\text{Po} \rightarrow {}_{82}^{211}\text{Pb} + \underline{\quad}$.
2. ${}_{90}^{228}\text{Th} \rightarrow \underline{\quad}\text{Ra} + \frac{4}{2}\alpha$.
3. ${}_{82}^{214}\text{Pb} \rightarrow {}_{83}^{214}\text{Bi} + \underline{\quad}$.
4. ${}_{8}^{15}\text{O} \rightarrow {}_{7}^{15}\text{N} + \underline{\quad}$.

$$5. \underline{\quad}\text{Si} \rightarrow {}_{13}^{27}\text{Al} + {}_{+1}^0e^-$$

$$6. \underline{\quad}\text{U} \rightarrow {}_{90}^{238}\text{Th} + \frac{4}{2}\alpha$$


$$7. \underline{\quad}\text{As} \rightarrow \underline{\quad}\text{Se} + {}_{-1}^0e^-$$

$$8. \underline{\quad}\text{Ac} \rightarrow {}_{87}^{227}\text{Fr} + \underline{\quad}$$

RADIOACTIVITY N/Z Curve

Nuclei have positive charge due to the protons in them. All the protons repel, so why does the nucleus not explode?


There is another force acting called the *strong nuclear force*. This acts between all nucleons, both protons and neutrons.



Plotting the number of protons vs. number of neutrons in stable nuclei gives this graph.

For elements where $Z > 80$ these decay by α decay.




Alpha particles consist of two protons and two neutrons. Therefore, the atomic number falls by two and the mass number by four.



N.B. Remember alpha particle is ${}_{2}^{4}\text{He}$.



These isotopes need to gain protons and lose neutrons to move towards the line of stability. They have too much strong nuclear force and not enough electrostatic force. β^- decay allows this to happen.

A neutron turns into a proton and an electron. The equations for this process are:





N.B. remember the beta particle is an electron.

E.g.



E.g.




Questions

- Explain why proportionately more neutrons are needed in larger nuclei?
- Using the graph above, calculate the ratio $Z:N$ when $Z = 6$ and when $Z = 80$. Comment on your answer. Why does the line on the graph curve away from the line $Z = N$?
- What type of decay occurs in isotopes with too much strong nuclear force? How do these changes help the nucleus to become more stable?
- Repeat question 3 for isotopes with too much electrostatic force.
- Nuclei do not contain electrons, so where does the electron emitted from a nucleus in beta-minus decay come from?
- Balance the equation ${}_{6}^{11}\text{C} \rightarrow {}_{7}^{11}\text{B} + \underline{\quad}$. (Hint: are there too many protons or too many neutrons in the carbon nucleus?), hence will β^+ or β^- decay occur?

RADIOACTIVITY Fundamental Particles

A fundamental particle is one that cannot be split into anything simpler.

The word atom means 'indivisible' because scientists once thought atoms were fundamental particles.



Similar experiments to Rutherford's alpha scattering using electrons fired at protons and neutrons reveals that they are made up of smaller particles - **quarks**.

Scientists now think that quarks, together with electrons and **positrons** are examples of fundamental particles.



There are actually six types of quark given odd names. They also have fractional charges as shown below.

Up	Charge	Charm	Charge	Top	Charge
u	$+2/3$	c	$+2/3$	t	$+2/3$
Down	Charge	Strange	Charge	Bottom	Charge
d	$-1/3$	s	$-1/3$	b	$-1/3$

An example of antimatter.
All particles have antiparticles; they are identical in mass but opposite in charge. Our Universe is made of matter. Antimatter is made in particle accelerators or as the result of some nuclear processes such as beta-plus decay.


Normally we are not allowed fractional charges, but quarks never occur on their own, only in combinations that add up to a whole charge.

Protons and neutrons are made of just two types of quark, the up and the down. Other particles have to be created in special machines called particle accelerators.



Beta decay

In beta decay, one of the up quarks changes to a down quark or *vice versa*.




Questions

- What is meant by the statement 'an electron is a fundamental particle'?
- How many different types of quark make up protons and neutrons?
- What quarks are found in a neutron?
- Describe the changes in quarks when a proton decays to a neutron by beta-plus decay.
- What is antimatter?

RADIOACTIVITY Half-Life

Most types of nuclei never change; they are stable. However, radioactive materials contain unstable nuclei. The nucleus of an unstable atom can break up (decay) and when this happens, it emits radiation.

A nucleus of a different element is left behind.




As time goes by radioactive materials contain fewer and fewer unstable atoms and so become less and less radioactive and emit less and less radiation.

There is no way of predicting when an individual nucleus will decay; it is a completely random process. A nucleus may decay in the next second or not for a million years. This means it is impossible to tell how long it will take for all the nuclei to decay.

Like throwing a die, you cannot predict when a six will be thrown. However, given a very large number of dice you can estimate that a certain proportion, $\frac{1}{6}$ th, will land as a six.

We define *activity* as the number of nuclei that decay per second (N.B. 1 decay per second = 1 Bq). The time it takes for the activity of a radioactive material to halve (because half of the unstable nuclei that were originally there have decayed) is called the **half-life**.



We see the activity falling as there are fewer nuclei available to decay. However, note that the time taken to halve is independent of the number of nuclei, in this case 2 seconds. Half-lives are unique to each individual isotope and range from billions of years to fractions of a second.

The half-life of a radioactive isotope is formally defined as:

'The time it takes for half the nuclei of the isotope in a sample to decay, or the time it takes for the count rate from a sample containing the isotope to fall to half its initial level.'

Calculations

1. Numerically e.g. a radioisotope has an activity of 6400 Bq and a half-life of 15 mins.

$$\text{After 15 mins the activity will be } \frac{6400 \text{ Bq}}{2} = 3200 \text{ Bq.}$$

$$\text{After 30 mins the activity will be } \frac{3200 \text{ Bq}}{2} = 1600 \text{ Bq.}$$

$$\text{After 45 mins the activity will be } \frac{1600 \text{ Bq}}{2} = 800 \text{ Bq.}$$

$$\text{After 1 hour the activity will be } \frac{800 \text{ Bq}}{2} = 400 \text{ Bq.}$$

Alternatively, consider the number of half-lives, e.g. $1\frac{1}{2}$ hrs = 6×15 mins = 6 half-lives.

$$\text{Therefore activity} = \frac{\text{original activity}}{(2 \times 2 \times 2 \times 2 \times 2 \times 2)}$$


$$(\text{i.e. divide by 2, six times}) = \frac{\text{original activity}}{2^6}$$

$$\text{In general, activity} = \frac{\text{original activity}}{2^{\text{no. of half-lives}}}$$

Therefore after 6 half-lives, in this case,


$$\text{activity} = \frac{6400 \text{ Bq}}{2^6} = 100 \text{ Bq.}$$

2. Graphically A graph of activity vs. time can be plotted from experimental measurements. We must remember to subtract the background count from the actual count to find the count due to the source alone. We call this the *corrected count rate*.



Results:

As long as the sample is large enough, this curve is smooth, because although the process is random, probability tells us half the atoms will decay in a certain time, but not which.



Nuclear radiation never completely dies away, but eventually drops to a negligible level, close to the background. At this point, a source is considered safe. Consideration of half-life therefore, has importance when considering which isotopes to use for various applications and the disposal of radioactive waste – see section on applications of radioactivity.

Questions

1. What is the activity of a radioactive source?
2. Write down a definition of half-life. Suggest why we can measure the half-life of a substance, but not its 'full life' (i.e. the time for all the atoms to decay).
3. ^{99}Tc (Technetium) has a half-life of 6 hrs. A sample of technetium has an initial count rate of 128 000 Bq
 - i. What will the count rate be after: a. 6 hrs? b. 18 hrs?
 - ii. How many hours will it take the count rate to fall to: a. 32 000 Bq? b. 8000 Bq? c. 1000 Bq?
4. A student has a sample of ^{137}Ba (Barium). They record the count rate every 60 s and record the following results:



Time in seconds	0	60	120	180	240	300	360	420	480	540	600	660	720
Count rate (decays/s)	30.8	23.8	18.4	14.2	11.1	8.7	6.9	5.4	4.4	3.5	2.9	2.4	2.0

The background count rate, with no source present, was 0.8 counts per second.

- a. Copy the table and include a row for the corrected count rate.
- b. Draw a graph of count rate vs. time and use it to show that the half-life is approximately 156 s.
- c. Do you think this isotope would present significant disposal problems, why or why not?
5. A student has a sample of radioactive material. In one lesson the activity recorded was 2000 Bq. The next day, at the same time, the count rate was just over 500 Bq. Which of the following isotopes is the sample most likely to be?
 - a. ^{135}I (iodine) half-life = 6.7 hrs.
 - b. ^{87}Sr (strontium) half-life = 2.9 hrs.
 - c. ^{42}K (potassium) half-life = 12.5 hrs.
 - d. ^{187}W (tungsten) half-life = 24 hrs.

RADIOACTIVITY Is Radiation Dangerous?

All nuclear radiation is ionizing. It can knock electrons out of atoms, or break molecules into bits. If these molecules are part of a living cell, this may kill the cell.




Questions

1. Explain which type of radiation is most harmful:
 - a. Outside the body.
 - b. Inside the body.
2. Explain the difference between contamination and irradiation. Which would you consider a more serious problem?
3. How does nuclear radiation cause damage to living tissues?
4. What is a Sievert?
5. Explain three precautions you should take if you had to handle a low activity radioactive source.

RADIOACTIVITY Nuclear Fission

Nuclear fission is the splitting of an atomic nucleus.


A large parent nucleus, such as ^{235}U , splits into two smaller daughter nuclei, of approximately equal size. This process also releases energy (heat) which can be used to generate electricity (see p111). Normally, this will happen spontaneously but can be speeded up by inducing fission.




If uranium were burned chemically to uranium oxide, it would release about 4500 J/g. The equivalent energy release from nuclear fission is 8.2×10^{10} J/g.

The daughter products themselves are radioactive because they still tend to be neutron rich (i.e. lying above the N/Z curve), and decay, releasing more thermal energy and nuclear radiation. They have a wide range of half-lives. These factors need to be taken into account when considering their disposal, (see p112).

Fuel rods of uranium or plutonium.



Control rods absorb neutrons before they can cause further fissions.



Lowering the control rods absorbs more neutrons and slows the reaction, raising the control rods speeds it up.

Questions


- Balance this equation, a fission reaction of uranium producing the daughter nuclei barium and krypton.

$$^{92}_{\Lambda}\text{U} + {}_0^1\text{n} \rightarrow {}_{56}^{136}\text{Ba} + {}_{36}^{90}\text{Kr} + 2 {}_0^1\text{n}$$
- In what form is the majority of the energy released by a nuclear reaction?
- Why do the products of fission reactions need careful handling?
- How do the control rods in a reactor control the rate of the nuclear reaction?
- For a stable chain reaction, neither speeding up nor slowing down, suggest how many neutrons from each fission should go on to cause a further fission.
- Use the data above to show that the energy released from the fission of 1 g of ^{235}U is about 20 million times as much as when the same gram is burnt in oxygen to form uranium oxide.

RADIOACTIVITY Nuclear Fusion


Nuclear fusion is the joining of two light nuclei to form a heavier nucleus. It is the process by which energy is released in stars.

In the nucleus, the **STRONG NUCLEAR FORCE** attracts protons and neutrons together; it is stronger than the **ELECTROSTATIC REPULSION** between the protons but it is a very short-range force.



This means that the gas containing the nuclei has to be very hot, dense, and under high pressure.


The gas is so hot that none of the electrons now orbits the nuclei. This is called plasma.




This is very difficult to do on Earth as this plasma would melt any container. Confining plasma is a major area of research because for the same mass of fuel, fusion of hydrogen to helium releases much more energy than fission and is the reaction occurring in the core of stars. We have a plentiful supply of hydrogen in water on Earth and the products are not polluting.

To fuse two nuclei they must be brought very close together so the strong nuclear force can bind their protons and neutrons together.

To do this you have to overcome the electrostatic repulsion between the nuclei.



Therefore, the nuclei have to travel very fast so they have a lot of kinetic energy to do work against the repulsive force.



When the nuclei join, energy is released as the kinetic energy of the product nucleus.


The nucleus formed has less mass than the total mass of the nuclei that fused to create it. The missing mass (or mass defect) has been converted to energy by Einstein's famous relationship

$$\Delta E = \Delta m c^2$$

ΔE = energy released in J

Δm = mass loss in kg

c = speed of light = 3×10^8 m/s



Scientists still have not achieved the process under control. They can do it where the reaction is explosive, in a hydrogen bomb. Some scientists once claimed they could do fusion at room temperature, but no one has been able to repeat this.

Key

- H^+ proton
- Deuterium nucleus ($1n + 1p$)
- $^3_2 He$ nucleus
- ⊕ Positron (β^+ particle)


Questions

1. Explain the differences between nuclear fission and fusion.
2. What are the two forces that must be kept in balance in a stable nucleus?
3. What is plasma?
4. Why does fusion require such high temperatures and what problems may occur as a result?
5. Explain why scientists are working hard to achieve controlled fusion on Earth.
6. A helium-4 nucleus is only 99.3% of the mass of the 4 hydrogen nuclei from which it was formed. The other 0.7% of its mass is converted into energy. Use Einstein's equation $\Delta E = \Delta m c^2$ to show that the energy released from the fusion of 1 kg of hydrogen nuclei, is about 6.3×10^{14} J (c = speed of light = 3×10^8 m/s).

APPLICATIONS OF PHYSICS

How Science Works


The remainder of this book outlines some of the ways that these ideas have been put to use. The link between these two aspects is how science works.




THE SUPPLY AND USE OF ELECTRICAL ENERGY

Examples of Energy Transformations Involving Electrical Devices and the Impact of Electricity on Society

Electricity supplies the majority of the energy we use in our daily lives. It is clean and very easy to control. Most houses contain many appliances that work by transforming electricity into other forms.



The majority of people in the developed world have access to a cheap, reliable electricity supply. The majority of people in the developing world do not. The fact that we have electricity so easily available has had a huge impact on all aspects of our lives and society. Here is a flavour.




Questions

- One hundred years ago open coal fires heated many homes. Now electric heaters heat many houses. Suggest some reasons why.
- Draw an energy flow diagram for three more electrical devices found in your home not shown above.
- Make a list, with justification of each, of one positive and one negative impact of electricity socially, politically, environmentally and economically not shown above.


THE SUPPLY AND USE OF ELECTRICAL ENERGY

What Influences the Energy Resources We Use?

Electricity provides the majority of the energy needs of the UK. The demand for electricity is predicted to continue to rise. Electricity is a secondary energy source; another (primary) energy source is needed to generate it.




The following pages outline how electricity is generated from these resources, but how do we decide what resources to use? There are a huge number of questions to be answered, and many of the answers may be contradictory.




Questions

1. Make a list of energy resources we use to generate electricity and divide your list into the renewable and non-renewable resources.
2. The type of energy resources that the UK should use to generate electricity in the future is very controversial. Why do you think this is? Do you think that there are any 'right' answers to the question, 'What energy resources should the UK use in the future?'



Questions

1. A wire is moved at right angles to a magnetic field. What would happen to the size of the potential difference across the wire if:
 - a. The wire was moved faster?
 - b. The magnet was moved instead of the wire, but it was moved at the same speed as the wire?
 - c. A weaker magnetic field was used?
 - d. The wire stopped moving?
 - e. Two magnets were used end to end so more wire was in the field?
 - f. The wire moved from a north pole to a south pole along the magnetic field lines?
2. When pushing a magnet into a coil how could you make the size of the induced potential difference bigger (3 ways)? How could you reverse the direction of the potential difference (2 ways)?
3. When generating electricity by induction where does the energy that is converted into electrical energy come from?




Questions (continued)

4. List five ways the output of an alternating current generator can be increased.
5. The mains electricity in the UK alternates through 50 complete cycles per second. What does this tell us about the rate of rotation of the generators in power stations in the UK?
6. Suggest two differences between the simple generator shown above and the generators used to generate mains electricity.
7. Why is the potential difference produced by a generator zero twice every revolution?
8. Draw a labelled diagram of an alternating current generator and use it to explain why the current it produces is alternating.

THE SUPPLY AND USE OF ELECTRICAL ENERGY How Power Stations Work

Electricity is very useful energy source because it is easy to distribute and control. However, it is a *secondary energy source* because another primary energy source has to be used to generate it. In conventional power stations, that energy source is either fossil fuels (coal, oil, or natural gas) or nuclear energy stored in uranium or plutonium (see p77 and p111). Increasingly renewable energy resources (see p88 and p89) are also being used.

Here we focus on conventional power stations.




Power stations are situated in the country to reduce pollution in towns. They are often close to rivers or the sea for their cooling water supply.

Questions

1. Name energy sources used to generate electricity in thermal power stations.
2. Draw an energy flow diagram for a coal-fired power station. Start with chemical energy in the coal and end with electrical energy produced.
3. What is combined heat and power?
4. Why are thermal power stations built near rivers or the sea?
5. What is the typical efficiency of conversion of chemical energy to electricity in a thermal power station? To what form of energy is most of the chemical energy converted?

THE SUPPLY AND USE OF ELECTRICAL ENERGY The Transformer



Questions


1. Copy and complete the following table, giving answers to the nearest whole number:

Transformer	Primary turns	Secondary turns	Primary voltage	Secondary voltage
A		120	240	12
B	625	10 000		400 000
C	20 000		11 000	240
D	2180	1000	240	

- Which transformers are step-up and which are step-down?
- Explain why a transformer needs AC not DC current to work.
 - Remember that electrical power = current × voltage.
 - If a transformer is supplied with 0.2 A at 240 V, what is the input power?
 - Assuming the transformer is 100% efficient, what is the output power?
 - If the ratio $N_p:N_s = 2400:60$ what is the output voltage?
 - Hence, what current can be drawn from it?
 - Many people leave mobile phone chargers, containing transformers, plugged in when not in use. The primary coil is connected to the mains, but no current is drawn from the secondary coil by the phone since it is not connected.
 - How, and from where, does the charger still waste energy?
 - Even though the energy wasted is small, why should people be encouraged to unplug chargers when not in use?
 - DC electricity is more useful for many applications, but the mains electricity is supplied as AC. Suggest why.

THE SUPPLY AND USE OF ELECTRICAL ENERGY The National Grid

Electricity is supplied from power stations to consumers by a 'national grid' of interconnected cables and transformers. They allow energy to be sent where it is needed anywhere in the country, and diverted around any faults that develop.




Questions

- Suggest two reasons for a 'national grid' to supply electricity, rather than each town having its own power station.
- Assuming the super-grid power lines operate at 400 000 V, calculate the ratio $N_p:N_s$ for each of the transformers in the diagram above.
- Why do we use very high voltages to distribute electricity when a lower voltage would be a lot safer?
- Step-down transformer B (above) has an output of 300 A at 132 000 V. what is the current flowing into it assuming the input voltage is 400 000 V and it is 100% efficient?
- Explain (using a formula) the statement, 'Doubling the current in a wire, quadruples the energy loss from it as heat'.
- Draw up a table of advantages and disadvantages of underground vs. overground cables.

THE SUPPLY AND USE OF ELECTRICAL ENERGY

All electricity generation has some impact on the environment.




Questions


1. Make a list of 5 ways you could reduce electricity consumption in your house.
2. In the UK in 2007 there are 1200 wind turbines producing a total of 772 megawatts.
 - a. On average how many megawatts does one turbine produce?
 - b. All the fossil fuel power stations in the UK combined produce about 60 000 megawatts. How many turbines would be needed to replace all the fossil fuel power stations?
 - c. Solar cells produce free electricity without any pollution. Suggest some reasons why they are not very widely used in Britain.
3. The environmental impact of electricity generation is an international problem. Give three examples from above where the impact on the environment could affect more than just the country generating the electricity.
4. Some people say the destruction of a wildlife habitat to build a new dam is not justified. If the dam replaced a coal-fired power station do you agree or not? Justify your argument.

THE SUPPLY AND USE OF ELECTRICAL ENERGY Renewable Energy Resources

Renewable energy resources are those that are *not used up* like fossil fuels. They can be used on a large scale, mainly to generate electricity, or for individual buildings either to provide heating or to generate electricity. All of these resources have advantages and disadvantages. To use renewable resources effectively a combination of different resources must be used, both on a national and local scale.




The carbon dioxide produced is absorbed by new plants grown to replace those harvested. Overall, it is a carbon neutral process.




Heating and alternative fuels

Large south facing windows provide passive solar heating. Warmed contents of the building radiate long wavelength infrared that cannot pass through the glass.



Some systems used curved mirrors to focus the Sun's heat energy onto a pipe containing water, whatever its position in the sky. Solar collectors must face the Sun; south in the northern hemisphere. Ideally, they should move to track the sun.

Solar heating supplements rather than replaces existing water and space heating. It depends on how sunny it is but can reduce domestic carbon dioxide emissions by 400 kg per year.



Questions

- What is our most significant source of energy on Earth?
- Look at the map of wind farms in the UK. The most common wind direction in the UK is from the southwest. Scotland, Wales and northern England are hilly. Hence, explain why there are very few wind farms in southeast England.
- Summarize all the information in this chapter in a table as shown.

Source of renewable energy	How it works (you might include a diagram here)	Advantages	Disadvantages or problems


You should be able to include at least eight separate rows.

- For each of the following, why might people object to having them built near their local area? How might you persuade them to accept the proposal?
 - A wind farm of twenty large windmills.
 - A hydroelectric power scheme involving flooding a valley by building a dam across it.
 - Building a barrage across a river estuary to generate tidal power.
- UK receives 40% of Europe's total available wind energy but only generates 0.5% of its power from it. Discuss some of the possible reasons why.

THE SUPPLY AND USE OF ELECTRICAL ENERGY

Energy is the ability to do *work*. Electrical energy is a convenient way to distribute it to houses, shops, schools, offices, and factories. Like any product, electricity has a *cost* and the more you use the more you pay. Therefore, the amount used has to be measured.

Whenever electric charge flows through an appliance, the energy it carries is given up to the appliance.



The number of Joules used by an appliance every second = power.

1 Joule every second = 1 Watt.

1000 Joules per second = 1 kilowatt, kW.

$$\text{Electrical power (W)} = \text{current (A)} \times \text{voltage (V)}$$

Consider a typical device such as a kettle. 4.35 A flows when it is connected to the 230 V mains supply.

Switched on for 2 mins to boil some water.


If a 1 kW appliance is switched on for 1 hour it uses $1000 \text{ W} \times 3600 \text{ s} = 3.6 \text{ million Joules}$
We call this a kilowatt-hour or kWh.

$$\text{Electrical power} = 4.35 \text{ A} \times 230 \text{ V} = 1000 \text{ W} = 1 \text{ kW}$$

$$\text{Electrical energy} = 1000 \text{ W} \times 2 \text{ mins} \times 60 \text{ s} = 120\,000 \text{ J}$$

Electricity companies call this a unit.

Calculating the Cost of the Electrical Energy We Use



$$\text{Electrical energy (kWh)} = \text{power (kW)} \times \text{time (h)}$$


N.B. Be careful with the units: power must be in kilowatts not watts.

Energy efficiency

Remember Efficiency = $\frac{\text{useful energy output}}{\text{total energy input}}$

Higher efficiency means:

- Cheaper bills.
- Less pollution.



The cost of the electricity will be
Number of kilowatt-hours × cost per kilowatt-hour

Some people can buy cheaper electricity during the night (midnight–8am).

Advantages

Can be used to heat water or 'storage' heaters overnight, which give out their heat during the day.

Disadvantages

You need to use over 20% of your electricity at night, to be cost effective as the daytime rate is usually more expensive than normal. The heat in the water or heaters may have dissipated by the evening – just when you need it most.

Questions


1. Calculate how many Joules each of the following uses:
 - a. A 100 W light bulb on for 10 s.
 - b. A 500 W TV on for 30 s.
 - c. A 1 kW kettle on for 60 s.
 - d. A 2 kW fire on for 5 mins.
 - e. A 1 kW iron on for 1 hour.
2. Is a kWh a unit of power or energy? How many Joules are there in a kWh? How many kWh is 14.4 million Joules?
3. Copy and complete the following table:

Appliance	Current (A)	Voltage (V)	Power (kW)	Time (hours)	Energy (Joules)	Units used (kWh)	Cost of electricity at 10 p per unit
Storage heater		230	2	2		$2 \times 4 = 8$	$8 \times 10 = 80$
Cooker	26	230		2			
CD player	0.048	230		2.0			
Kettle		230	2			0.2	
Iron	2	230		1.2			
Fridge		230	0.12			2.88	
Lamp		230	0.06		432 000		

4. Give two advantages of buying more energy efficient devices. Where can you look to find energy efficiency information when shopping for new household appliances?
5. Why do you think electricity companies offer cheap electricity overnight?

THE SUPPLY AND USE OF ELECTRICAL ENERGY The Motor and Dynamo

The motor effect from p59 can be used to make a practical electric motor.



Questions


1. How can an electric motor be made more powerful?
2. What would happen to a motor if there was no way of reversing the current direction every half turn? How does a split ring commutator avoid this situation?
3. What are the energy changes in an electric motor? Therefore, why are electric motors not 100% efficient?
4. A motor can lift a weight of 20 N through 3 m in 10 s. If the current flowing is 1.79 A when the voltage of the electricity supply is 12 V, show that the motor is about 30% efficient.
5. What is a dynamo? Explain how it works in as much detail as possible using the ideas from this page and p82–83.

THE SUPPLY AND USE OF ELECTRICAL ENERGY Logic Gates

A logic gate is a circuit that can make decisions depending on the signals it receives.

The input signal for a logic gate can either be high (about 5 V) or low (about 0 V). The high input is always denoted by 1, and low input by 0. Signals between these values are not counted. The gate's output is either high or low depending on whether the input signals are high or low.

Potential divider circuits are used to provide the input voltage for a logic gate that can be either high or low depending on the conditions.



$$V_{\text{out}} = R_1 / (R_1 + R_2) \times \text{supply p.d.}$$


If R_2 is a variable resistor the temperature or light level at which V_{out} becomes large or small enough to trigger a high or low input at the logic gate can be set. This allows the light level or temperature that the sensor will respond to, to be set.

Name	Symbol	Truth table Input	Truth table Output
OR Output high (1) when either input (A or B) is high (1)		0 0 1 0 0 1 1 1	0 1 1 1
AND Output high (1) when both inputs (A and B) are high (1)		0 0 1 0 0 1 1 1	0 0 0 1
NOR Output high (1) when neither input (A nor B) is high (1)		0 0 1 0 0 1 1 1	1 0 0 0
NAND Output high (1) unless both inputs (A and B) are high (1)		0 0 1 0 0 1 1 1	1 1 1 0
NOT Reverse the input		0 1	1 0

Reversing the position of the LDR or thermistor with the fixed resistor, reverses the output.


Logic gates can be used to activate a certain output when required input conditions are met. This can be shown in a *block diagram*.

E.g. A courtesy light switches on in a car when either driver or passenger door, or both, are opened.



Designing a logic system:

1. Draw up a block diagram with logic gates




E.g. A washing machine door must only unlock if the drum is not turning and there is no water in the drum. It should stay locked until the programme has finished, although this should have a manual override.

2. Draw a truth table and check it gives the desired output.


W	D	OR output	OR output	AND output	NOR output
0	0	0	0	1	1
1	0	1	1	0	0
0	1	1	1	0	0
1	1	1	1	0	0

W	D	AND output	OR output
0	0	1	0
1	0	0	1
1	1	0	1

Latching circuits Sometimes an output is needed which does not change even if the input is removed, e.g. an alarm that continues to ring even if the input that triggered it is removed.




A low signal at both inputs does not change the output. This circuit is called a *bistable* – it has two stable states.




Questions

- What do the numbers '1' and '0' represent in a truth table?
- A microwave oven must not start unless the door is closed and the timer is set. Draw a block diagram with a suitable logic gate for this, and include the truth table.
- In the following circuit, the resistance of the thermistor (R_1) at 100°C is $1.2 \text{ k}\Omega$. What resistance should the variable resistor R_2 be set to so $V_{\text{out}} = 5 \text{ V}$ when the temperature reaches 100°C ?
- What is a relay and where are they used? Draw a labelled diagram.
- In a greenhouse, automatic shades should be drawn if the soil around the plants becomes too dry and if the light level or the temperature rises too much. Draw a suitable block diagram using logic gates and give its truth table.
- Draw a truth table for the circuit shown:
- A water pipe may burst if the temperature drops below freezing. Draw a suitable block diagram using two logic gates for a system that will shut off the water to a house if the temperature falls below freezing and not switch it back on until it is reset by a plumber who has inspected the pipes for damage. Draw a truth table for your system.



THE SUPPLY AND USE OF ELECTRICAL ENERGY Electricity and the Human Body

The body sends electrical signals, via nerves from the brain to stimulate muscles.




The heart is a muscle and the signals that control it come, not from the brain via nerves, but from some special cells in the heart, called the sinoatrial node. This means that the heart works automatically without us having to think about it.

Potentials detected on the surface of the body as they are transmitted from the heart by conducting body tissues.

Defibrillation

Fibrillation occurs when a patient's heart does not beat rhythmically but quivers. Blood is not pumped and the patient will soon die.



All the heart muscles contract strongly.
Usually when they relax, they settle back into a regular rhythm of beating.

Care must be taken not to shock the operator.


Pacemakers

- Cells producing stimulus to the heart stop functioning.
- Electrical devices are fitted producing tiny, but regular, shocks to the heart.
- It is placed under the skin and a wire fed through a vein to the heart.
- Fitted with long-life batteries.

① Blood arrives from the lungs and the rest of the body and collects in the atria.



Potential difference at electrodes / mV




Questions

- Using the figures above show:
 - That the energy delivered to the heart during defibrillation is about 300 J.
 - That the resistance of the body is about 150Ω .
- When electrodes are attached to a patient for an ECG or defibrillation, a conducting paste or pad is applied between the skin and the electrode. Why?
- Often when a person receives an electric shock muscles contract violently, sometimes even breaking bones. Why do electric shocks often have this effect?
- A normal resting heartbeat is about 70 beats per minute. Draw an ECG trace to scale showing 5 beats. A patient who has tachycardia has a fast heart rate. Add this ECG trace to your original and label it. A patient who has arrhythmia has an irregular heartbeat. Add a third trace showing this.

TRANSPORT Stopping Distances


Reaction time is the time it takes a driver to see a hazard, recognize they need to take action, decide on the action, and use the vehicle's controls.

E.g. a child runs into the road, the driver sees the child and considers braking or swerving. They decide to brake and press the brake pedal. Normally all these stages happen in about 1 second.



When braking, the brakes do work against friction.

The total stopping distance of the vehicle = thinking distance + braking distance.



This converts the kinetic energy of the vehicle into heat energy.


Hence kinetic energy = work done against friction

$$\frac{1}{2} (\text{mass of vehicle}) \times (\text{velocity of vehicle})^2 = \text{brake force} \times \text{distance travelled while braking.}$$

$$\frac{1}{2} mv^2 = F \times d.$$

$$\text{Therefore minimum braking distance} = \frac{1}{2} mv^2/F$$

Braking distance is proportional to the mass of the vehicle assuming the same brake force. Therefore, large vehicles need brakes that can exert a larger force.



Questions


- Write a list of factors that affect braking distance.
- As the speed of a vehicle increases what happens to the size of the brake force needed to stop it in a certain distance?
- Explain why each of the following is a driving offence that the police might stop you for in terms of their effect on the thinking distance or braking distance of a vehicle. The first one has been done as an example:
 - Driving faster than the speed limit. Answer – increases both thinking distance and braking distance so a vehicle is less likely to be able to stop in the distance the driver can see to be clear.
 - Having bald tyres.
 - Driving under the influence of alcohol.
 - Using a mobile phone while driving.
- Copy and complete the following table:

Mass of vehicle (kg)	Initial velocity (m/s)	Maximum braking force (N)	Minimum braking distance (m)
1000	13.3 (= 30 mph)	6500	
1000	31.1 (= 70 mph)	6500	
5000	13.3 (= 30 mph)	6500	

Explain why the braking distances you calculated are minimum distances.

TRANSPORT Road Safety

Most car safety features are designed to reduce the force of any collision on the passengers, which reduce the injuries they may suffer.




Questions

- Using the words 'force', 'deceleration', 'momentum', 'energy' in your answers explain:
 - How motorcycle and cycle helmets made of a material that deforms on impact with a hard surface, reduce the severity of head injuries to the rider in a collision.
 - Why crash barriers alongside roads are often made of deformable material like steel tube rather than a rigid material like concrete.
 - An escape lane (a pit full of deep sand) is provided at the bottom of steep hills for drivers to steer into if their brakes fail.
- A car passenger of mass 70 kg is travelling at 13.3 m/s (30 mph). Show that their momentum is 931 kgm/s. In a collision, they hit the dashboard and stop in 0.01 s. Show that the force exerted is about 93 kN. In another collision, the passenger is cushioned by an air bag and stops in 0.1 s. Show the force is now only 9.3 kN.
- A car of mass 1000 kg travelling at 13.3 m/s hits a brick wall and stops in 0.1 s. Calculate the deceleration. What force is exerted on the car by the wall? The same car is now fitted with a crumple zone and stops in 0.5 s. What force is exerted by the wall now?
- Explain how antilock brakes (ABS) can help to reduce stopping distances when a driver brakes hard.

WAVES AND COMMUNICATIONS Using Waves to Communicate

The long distance transmission of information, other than by a message on paper or via an electrical signal in a wire, relies on using waves.




Questions


1. Name four types of electromagnetic waves used to send messages. Suggest why the other types of electromagnetic radiation are unsuitable.
2. Explain three ways radiowaves can be used to send messages over long distances. Use diagrams to help your explanation.
3. i. Use the formula $wave\ speed = frequency \times wavelength$ to calculate the wavelength of radiowaves of frequency:
a. 3 MHz (3×10^6 Hz). b. 1800 MHz (1.8×10^9 Hz). c. 30 GHz (3×10^{10} Hz).
ii. Explain which of the above frequencies would be most useful for:
a. Diffracting around large obstacles like hills. b. Sending to a satellite using a dish.
c. Mobile telephone communication.
4. A signal is to be sent from the UK to America across the Atlantic. Explain:
a. Why a signal sent by a ground wave would be very weak by the time it reached America.
b. Why the ionosphere is needed if the signal is to be sent by a sky wave.
c. Why a satellite is needed if the signal is to be sent by a space wave.

WAVES AND COMMUNICATIONS Analogue and Digital Signals

Communication signals are either




Or



Computers work with binary code, a series of high or low voltages representing 1 and 0. They are particularly good at processing digital signals.

Conversion:


Sample too often and the process is too slow.
Sample too few times and not enough information is available to reconstruct the original signal.



Advantages of digital signals:


1) Less noise

Noise is any unwanted interference picked up by the signal.



With analogue signals it is hard to remove noise.


Digital signals are still clearly 1s and 0s even with noise.



When decoded the noise is removed.

Therefore, digital radio and TV have better sound and picture quality.

2) More information can be transmitted at once. Signals can be interleaved, (called multiplexing).




Therefore, lots of information can be sent on one signal. Hence, this is why you can have so many digital TV and radio stations but relatively few analogue ones.

Questions

1. Use diagrams to illustrate the difference between a digital and an analogue signal.
2. When listening to a radio station a hissing sound is heard. What is likely to have caused this and is the signal most likely to have been analogue or digital?
3. Morse code is transmitted as a series of pulses of electricity in a wire or flashes of light representing dots and dashes. Explain whether it is an analogue or digital signal.
4. How are analogue signals converted to digital?
5. What is multiplexing?
6. Explain two advantages of digital signals compared to analogue.
7. When signals are amplified, noise is also amplified. Why is this less of a problem for digital signals?

WAVES AND COMMUNICATIONS AM/FM Radio Transmission

When you tune to a given radio or TV station, you select a particular frequency of radiowave to be received. This wave is called a *carrier wave*, but how is the message added to the carrier wave? There are two methods by which the carrier wave is *modulated* (or varied) by the message signal.




Questions

1. What is a carrier wave?
2. What do you understand by the term 'modulation' in the context of radiowaves?
3. What do the abbreviations AM and FM stand for?
4. Use diagrams to explain the difference between AM and FM radio transmissions.
5. Which type of transmission, AM or FM suffers less from noise?
6. Can two different national radio stations covering the whole of the UK use the same carrier wave frequency? What about two local stations?

WAVES AND COMMUNICATIONS Satellite Orbits and Their Uses

Satellites are objects that orbit larger objects in space. They can be natural, like the moon orbiting the Earth or artificial (man-made).




The gravitational attraction between the two objects provides the necessary centripetal force to keep the satellite moving in a circular path (see p19). Without this force, satellites would move off in a straight line into space. The gravitational attraction provides the centripetal acceleration towards the centre of the Earth that causes the satellite's direction of motion to change continuously.

$$\text{Since centripetal force} = \frac{\text{mass} \times \text{velocity}^2}{\text{radius}}$$

launching a satellite involves getting it to the right height, at the right speed.

Orbital period increases with height



Newton used a thought experiment to try to illustrate this.

1. Little gunpowder in cannon. Low initial speed and shot falls to Earth quickly.
2. More gunpowder. Higher initial speed and shot falls further before hitting ground.
3. Given enough gunpowder, the shot will orbit because it is travelling fast enough that the ground is 'falling' away under it at the same rate the shot is accelerating towards it.

Satellites in a low orbit make many orbits of the Earth per day. If they orbit over the poles and the Earth spins beneath them, they can observe many different parts of the Earth's surface each day.

Useful for:

- Weather forecasting.
- Imaging the Earth's surface (e.g. for spying, to monitor crops or pollution).

Communications – as they stay in the same place in the sky, satellite dishes only need to point in one fixed position.

Used for
Satellite stays over the same point, as the Earth and satellite are turning at the same rate.


Questions

1. State and explain two reasons why satellite orbit period increases with height above the Earth.
2. Using diagrams state and explain as many differences as possible between geostationary and polar orbits.
3. For each type of orbit, geostationary and polar:
 - a. State a use for a satellite in that orbit.
 - b. Explain why that orbit is used.
4. A geostationary satellite orbits 36 000 km above the surface of the Earth. The radius of the Earth is 6400 km.
 - a. How many hours does it take a geostationary satellite to orbit the Earth? What is this in seconds?
 - b. Show that the circumference of the satellite's orbit is about 270×10^6 m.
 - c. Hence show that its orbital speed is about 3080 m/s.
 - d. Use the formula centripetal force = mass × velocity² / radius to find the resultant force on a 10 kg satellite.
 - e. What provides this resultant force?

WAVES AND COMMUNICATIONS Images and Ray Diagrams



Light follows straight lines, or rays, from a source of light to an observer unless it is reflected, by a mirror, or refracted, by a lens, on route.

Mirrors and lenses come in a variety of shapes to manipulate the light rays in various useful ways. Ray diagrams help us to understand their effects.



Rays show the direction the light waves are travelling in. Light rays always travel in straight lines (as light waves travel in straight lines) except when reflected or refracted when they change direction.


An image is formed at a point where the light rays from an object appear to come from, had their direction not been changed by a mirror or lens.



Greater than 1 – *magnified* (image larger than object).

$$\text{Magnification} = \frac{\text{image height}}{\text{object height}}$$

Smaller than 1 – *diminished* (image smaller than the object).



Questions


1. Make a list of three properties of an image that describe the 'nature of an image'.
2. State three differences between a real and virtual image.
3. Is the image in a plane (flat) mirror real or virtual?
4. What is a light ray?
5. What is the formula for magnification? If the magnification of a lens is less than 1, would the image be larger or smaller than the object?
6. A tree has a height of 20 m. In a photograph, it has a height of 20 cm. What is the magnification?
7. A letter 'T' in a book has a height of 5 mm. When viewed through a magnifying glass with a magnification of 1.9, how high will it appear?

WAVES AND COMMUNICATIONS Mirrors and Lenses, Images

Mirrors

(1) Plane (flat)

Nature of image
Virtual
Upright
Same size as object




Law of reflection (applies to all mirrors):

Angle of incidence, i = angle of reflection, r

(2) Concave – curving in (like a cave)

Brings light to a focus so is a *converging* mirror.

Equal angles of incidence and reflection




Diffuse reflection from a rough surface – no image formed.

Nature of images

Object	Image
Beyond C	Between C and F Real Inverted Diminished
At C	At C Real Inverted Same size
Between C and F	Beyond C Real Inverted Magnified
Closer than F	Virtual Upright Magnified

(3) Convex – bulges out

Spreads light rays out so is a *diverging* mirror.




Rules for drawing ray diagrams for concave mirrors

Ray from the object

1. Parallel to optic axis – reflects through F .
2. To centre of mirror is reflected, forming equal angles with optic axis.
3. Through F is reflected parallel to the optic axis.

Lenses




The more powerful a lens, the greater the change in direction of the light rays, and therefore the closer the focus is to the centre line of the lens.

$$\text{Power of lens (dioptrē)} = 1/\text{focal length (m)}$$


The more curved the surface the greater the refraction of the light. Therefore, fat lenses have short focal lengths and are more powerful.

1. Convex (converging) lens



Object	Image	Uses
	Further than 2F Between F and 2F Real Inverted Diminished	Camera: convex lens focuses light from a distant object to form a diminished image on the film close to the lens
	Between F and 2F Further than 2F Real Inverted Magnified	Projector: convex lens focuses light from a nearby object to form an enlarged image on a distant screen
	Closer than F Upright Virtual Magnified	Magnifying glass

2. Concave (diverging) lens





Questions

- Describe what we mean by the term 'focal point'.
- Draw the shapes of convex and concave mirrors and lenses. Show with ray diagrams which will bring parallel light waves to a focus, and which will diverge them.
- What three rays are drawn in a ray diagram for:
 - A convex lens?
 - A concave mirror?
- Does a powerful lens have a short or long focal length? What unit is the power of a lens measured in?
- A lens has a focal length of 0.1 m. What is its power?
- Draw a ray diagram for an object placed at 2F from a convex lens and at F from a convex lens.
- Draw a ray diagram for a camera and a projector; include the object, image, and lens.

WAVES AND COMMUNICATIONS Optical Fibres

An optical fibre is a thin strand of very clear glass through which visible light or infrared radiation can be guided.



Questions


- Copy and complete the following diagram as accurately as possible showing the path of the light along the fibre-optic cable. What can you say about the size of the pairs of angles a and b , and x and y ?
- What types of electromagnetic radiation are commonly used with fibre optics?
- Outline some benefits of using fibre optics rather than copper wires for sending messages.
- The light in a fibre optic gradually gets less intense as it travels along the fibre due to impurities in the glass absorbing some of the light energy. What is the electrical equivalent of this?
- What is an endoscope? Suggest two possible uses for one.
- Suggest why doctors often prefer to see inside people using an endoscope rather than carrying out an operation to open up the patient.



WAVES AND COMMUNICATIONS Ultrasound and its Applications


Ultrasound is a sound wave with a frequency of greater than 20 000 Hz. This is above the upper limit of hearing for humans, so we cannot hear it, although in all other respects it behaves in exactly the same manner as normal sound.

Ultrasound can be used to detect the distance between the boundaries of two objects.



Questions

- Is ultrasound a longitudinal or transverse wave? How is ultrasound different to normal sound?
- The speed of ultrasound in soft tissue is 1540 m/s. The oscilloscope trace shows the returning pulses. How far below the surface of the body was pulse A and pulse B reflected?
- Suggest two reasons why ultrasound may be preferable to X-rays for medical examinations.
- Explain how ultrasound could be used to locate the depth below the skin of a cyst (fluid filled pocket) in an organ.
- Suggest one use of ultrasound in medicine and one in industry other than for making images of hidden objects.




WAVES AND COMMUNICATIONS Uses of Electron Beams


Review p57. Particularly note . . .

1. Electron beams are produced by 'boiling' electrons off a heated filament (thermionic emission). The hotter the filament the more electrons are produced.
2. The electrons are accelerated across a potential difference to increase their kinetic energy.
Kinetic energy = electronic charge (1.6×10^{-19} C) \times accelerating voltage

Cathode ray tubes – used in computer monitors, TVs, and oscilloscopes.





X-ray tubes:




Electrons gain kinetic energy as they accelerate towards the anode. Increasing the potential difference between filament and anode gives electrons more kinetic energy on impact, so higher energy (or 'harder') more penetrating X-rays are produced.

Kinetic energy of electrons on impact
1% X-ray energy
99% thermal energy



Questions


1. The diagram shows the X and Y plates in an oscilloscope viewed end on. In each case which of the dots shown (a, b, or c) correctly shows the position of the beam falling on the screen?
2. How many lines are there on a TV screen? Explain how the electron beam is made to move across the screen.
3. Describe three ways that the tungsten anode in an X-ray tube is kept cool.
4. What adjustment to an X-ray tube produces X-rays that are more penetrating?
5. An X-ray tube accelerates an electron through a potential difference of 40 000 000 V. (Charge on the electron = 1.6×10^{-19} C)
 - Show that its kinetic energy when it hits the anode is about 6.4×10^{-12} J.
 - If 1.6×10^{15} electrons hit the anode, show the total energy they deliver is about 10 kJ.
 - If this energy is delivered in about 0.2 s what is the power of the tube?
 - What percentage of the energy above is converted to X-ray energy and hence explain why the tungsten anode needs to be cooled?
 - Explain what effect increasing the filament temperature would have on the number of X-rays produced in an X-ray tube.



WAVES AND COMMUNICATIONS Beams of Light – CDs and Relativity


Einstein's theory of relativity is one of the most creative and challenging ideas in physics, while reading the information from a CD is a very straightforward application of physics. Yet they both involve ideas about beams of light.

A beam of laser light reads the information stored on a CD (or DVD).



Relativity

This theory makes some weird predictions about how we measure length and time when moving very fast relative to another object.




Questions


1. Laser beams can be made very narrow and do not spread out much. Why is this necessary for reading a CD as described above?
2. If you shake a CD player while playing a disc the music can be interrupted or skip a section. Using the above description try to explain why.
3. What is a thought experiment?
4. What predictions did Einstein make from his thought experiments?
5. Suggest three ways Einstein's predictions have been tested.

RADIOACTIVITY How is Nuclear Radiation Used in Hospitals?

Stable nuclei are bombarded with protons. These unstable proton-rich nuclei decay by beta-plus emission with short half-lives. They emit positrons.




This can be used to find out how well the drug moves round the body and how well organs of interest are working, or if they contain a tumour.



X-rays are high frequency, short wavelength, electromagnetic waves. They are ionizing so can damage cells. Exposure to them needs to be limited.

The benefits of the use of X-rays to diagnose medical problems often outweigh any cell damage caused.

X-rays expose photographic film and bones show up as a shadow.



Progress of the drug can be tracked by a radiation detector outside the body.

Monitoring the flow of the radioisotope over time can tell doctors about how organs are working.

Gamma emitter made into a drug and injected.

Alpha emitters cannot be used as they would not pass through the body and are highly ionizing so would cause a lot of cell damage.

Radioisotope attached to a drug that is absorbed by an organ of interest, e.g. a kidney.

Blockage – radioisotope will not pass and no radiation detected in this area outside body.

Short half-life used so radiation does not stay in the body too long.


Cancerous cells are those where the DNA has been damaged and grow and divide uncontrollably.

Ionizing radiation can damage the DNA in cells.

If the radiation dose is small, the cells may be able to repair themselves.

Large doses of radiation can kill cells. This can be used to kill cancerous cells.

Source of gamma rays




Radiotherapy may not be successful if the tumour is very large. In this case radiotherapy may be used to reduce suffering. This is called *palliative care*.

Questions

- Which types of radiation, alpha, beta, or gamma can pass through flesh?
- Why do the radioisotopes injected into patients always have short half-lives?
- What absorbs X-rays better, flesh or bone?
- What does PET stand for? Describe how it works, for example to identify the location of a cancerous tumour.
- The thyroid gland stores iodine. How could injecting a patient with radioactive iodine-131 allow a doctor to find out how well the thyroid gland is working?
- Ionizing radiation can cause the DNA in cells to mutate and cause cancer. Therefore, why can we also use ionizing radiation as a treatment for cancer?
- Why is the source of gamma rays in radiotherapy rotated around the patient?
- All ionizing radiation causes damage to the body. How do doctors justify exposing patients to it?

RADIOACTIVITY Other Uses of Radioactivity




Questions

- Explain whether an alpha, beta, or gamma source is most useful for the following and why:
 - Smoke alarms.
 - Detecting aluminium foil thickness in a factory.
 - Following the flow of oil along a pipe.
- Should a radioactive material with a long or short half-life be chosen for the following and why?
 - Smoke alarm.
 - Tracer in an oil pipe.
 - Thickness detection in a factory.
- Many people are concerned about the effect on their health of radioactive sources. How would you address the following concerns?
 - 'I don't have a smoke alarm as I do not want a radioactive source in my house.'
 - 'I am concerned that irradiated food might be radioactive.'

RADIOACTIVITY Radioactive Dating

Carbon dating




Measuring the activity of a sample of ancient materials that were once living and comparing the activity to a living sample can give a fairly accurate indication of when the ancient material was alive.


(It works for plant or animal material because animals eat plants and absorb carbon-14 from them.)

Dating of rocks

Many rocks contain traces of radioactive uranium. This decays to stable lead with a half-life of 4.5 billion years.



Assuming that there was no lead in the rock when it was formed the ratio of uranium to lead gives an approximate age for the rock.



Questions

- The graph shows the radioactive decay of carbon-14.
 - Use the graph to calculate the half-life of carbon-14. What does carbon-14 decay into?
 - A wooden post from an archaeological dig produces 150 counts/min. Wood from an identical species of tree currently alive gives 600 counts/min. How long ago did the wood from the archaeological dig die?
 - What assumption have you made in the above calculation?
- Two samples of rock are analysed. The ratio of 238-uranium to 206-lead are as follows:
Sample A: uranium to lead 5:1 Sample B: uranium to lead 7:1.
Which rock is older and how do you know? What assumption have you made?
- The age of the Earth is thought to be about 4.5 billion years. Why can there be no rock in which the number of lead nuclei formed from the decay of uranium outweighs the number of uranium nuclei remaining?


RADIOACTIVITY Nuclear Power and Weapons

See p77 for a description of the nuclear fission process and the nuclear reactor.

Large amounts of radiation is produced by a radioactive core (gamma rays, alpha and beta particles, high-energy neutrons).

Extra neutrons absorbed by reactor casing – this makes them neutron rich and radioactive (by beta decay, see p72).


Thick lead and concrete surrounds the reactor to prevent this radiation escaping.



Uranium is the fuel for most commercial nuclear reactors. It is obtained by mining uranium ore.

The nuclear debate

(fallout) – spreads contamination over a wide area. The bomb works like a nuclear reactor, but there are no control rods to keep the reaction in check so the chain reaction proceeds explosively.




Questions

- Write out a list of energy changes in a nuclear power station starting from nuclear energy stored in uranium fuel and ending with electrical energy in the wires leading from the generator.
- Where does the fuel for a nuclear power station come from and what has to happen to it before it can be used?
- The energy released by 1 kg of ^{235}U is about 8×10^{13} J. Show that this could light a 60 W light bulb for about 42 thousand years.
- Using the diagram of a nuclear power plant above explain:
 - Why is the reactor surrounded by a thick layer of concrete and lead?
 - Why is the pressure vessel made of steel?
 - Why are the pipes in the heat exchangers coiled up?
- Nuclear weapons cause damage to living things in three ways – what are they?
- 'Nuclear power damages the environment and should be banned.' Give arguments in favour and against this statement.

RADIOACTIVITY Radioactive Waste

- Sources:
- Nuclear fission power stations (p111).
 - Industrial users of radioactivity (p109).
 - Hospitals and other medical establishments (p108).
 - Laboratories.
 - Decommissioned nuclear weapons.
- These wastes should be disposed of in a way that does not significantly increase the naturally occurring background level of radiation around the disposal site.




Questions

- What are the three classifications of nuclear waste?
- What types of materials make up low-level waste?
- What is the main constituent of intermediate level waste?
- What constitutes high-level waste and why is this generally hot?
- What happens to low-level waste?
- What happens to intermediate level waste?
- What happens to high-level waste?
- Why are spent fuel rods left in cooling ponds for 3 months after use?
- You are responsible for finding a site for a new managed underground radioactive waste store.
 - What features would you look for in identifying a suitable site?
 - What concerns might local residents have?
 - How might you go about addressing these concerns?

OUR PLACE IN THE UNIVERSE Geological Processes

Theory of plate tectonics




Questions

- What is the difference between a constructive and destructive plate boundary?
- Explain why the majority of earthquakes and volcanoes occur near plate boundaries.
- Give three pieces of evidence mentioned above in support of the idea of plate tectonics.
- Why did people find it difficult to accept Wegener's ideas?
- What is the name of the process that causes the material in the mantle to circulate and drag the plates along?
- Describe how the magnetization of the rocks of the oceanic crust could be used to show that the ocean is growing wider over millions of years.
- Describe and explain the differences between the collision of two continental plates compared to a continental and an oceanic plate.

OUR PLACE IN THE UNIVERSE The Solar System

Oort cloud – objects made of ice and dust orbiting the Sun far beyond Pluto. Sometimes these fall in towards the Sun and become a comet due to minor gravitational disturbances.



The planets orbit the Sun in elliptical (slightly squashed circle) orbits. The Sun is at one focus of the ellipse.

Quantity	Mercury	Venus	Earth	Mars	Jupiter	Saturn	Uranus	Neptune	Pluto
Mean distance from sun (orbit radius), million km	57.9	108	150	228	778	1430	2870	4500	5900
Time to orbit the Sun, years	0.241	0.615	1.00	1.88	11.9	29.5	84.0	165	248
Orbital speed, km/s	47.9	35.0	29.8	24.1	13.1	9.64	6.81	5.43	4.74
Equatorial diameter, km	4880	12 100	12 800	6790	143 000	120 000	51 800	49 500	?3000
Mass (Earth = 1)	0.0558	0.815	1.000	0.107	318	95.1	14.5	17.2	?0.010
Density g/cm ³	5.600	5.200	5.520	3.950	1.310	0.704	1.210	1.670	?
Moons	0	0	1	2	16	17	15	2	1
Typical surface temperature, °C	167	457	14	-55	-153	-185	-214	-225	-236
Atmosphere	None	Carbon dioxide	Nitrogen, oxygen	Carbon dioxide	Hydrogen, helium	Hydrogen, helium	Hydrogen, helium	Hydrogen, helium	None

Questions


- Which of the following orbit the Sun directly and which orbit planets?
Comets, moons, asteroids, artificial satellites, planets.
- Explain why the density of Jupiter, Saturn, Uranus, and Neptune is a lot less than that of Mercury, Venus, Earth, and Mars.
- Using the data in the table show that:
 - The circumference of the Earth's orbit is 942 million km.
 - The time the Earth takes to orbit the Sun is 31.6×10^6 s. c. That 31.6×10^6 s = 1 year.
- Plot a graph of surface temperature vs. distance from the Sun. State and explain any trend you see. One planet is anomalous, which is it and give a scientific explanation for why it does not fit the trend?
- Explain why the speed of a comet decreases as it moves away from the Sun.

OUR PLACE IN THE UNIVERSE

Telescopes and Types of Radiation Used to Learn About the Universe

Everything we know about space outside the solar system comes from analyzing the electromagnetic radiation collected from space by telescopes.

Different objects in space emit different wavelengths.




Questions

1. Make a list of the advantages and disadvantages of space telescopes compared to ground based telescopes.
2. Why do you think optical telescopes that collect visible light are often placed on mountains whilst radio telescopes can be at sea level?
3. Will the image in a refracting telescope be upright or inverted? Use a ray diagram to illustrate your answer. Suggest two advantages of having a very large objective lens and explain why there is a limit on how big the objective lens can be.
4. The aperture of a reflecting telescope is 0.7 m in diameter and it collects light of wavelength of about 0.00000055 m. Its objective mirror has a focal length of 0.4 m and its eyepiece a focal length of 1.50 cm. The diameter of the Jodrell Bank radio telescope dish is 76.2 m and the wavelengths it collects are around 1 m.
 - a. What is the angular magnification of the reflecting telescope?
 - b. Which telescope would you expect to suffer the most from diffraction?
5. Suggest at least three reasons why astronomers need to work together in international groups.

OUR PLACE IN THE UNIVERSE The Motion of Objects in the Sky


The Moon orbits the Earth once every 27.3 days. It also takes 27.3 days to rotate once so it always presents the same side to the Earth.

The phases of the Moon take about 29 days to complete a cycle because the Earth has changed its position during the cycle as it orbits the Sun.





During one sidereal day, the Moon completes $\frac{1}{27}$ th of an orbit. For the Moon to appear in the same place in the sky the Earth will have to turn $\frac{1}{27}$ th of a revolution further. This takes about an extra 49 minutes. Therefore, moonrise occurs once every 24 hrs 49 minutes.

Different stars are seen in different seasons.



Lunar eclipse
Moon in Earth's shadow
Eclipses do not happen once each month because the Moon's orbit is tilted so is usually above or below the direct line between the Earth and Sun.



The sun appears to take $(23 \text{ hrs } 56 \text{ minutes} \times 0.986^\circ)/360^\circ = 4 \text{ minutes}$ longer to move from east to west across the sky and back to its starting point. This makes a solar day 4 minutes longer (i.e. 24 hrs exactly) than a sidereal day.


Some planets are visible with the naked eye (Mercury, Venus, Mars, Jupiter, and Saturn). They follow complicated paths in the sky because they too are moving around the Sun, at different speeds relative to the Earth. The term planet means 'wandering star'.

Questions

- Do most objects in the sky appear to move east to west or west to east? Which objects do not always follow this pattern?
- What is the difference between a solar day and a sidereal day?
- What is the difference between a lunar eclipse and a solar eclipse?
- Why do we not have an eclipse once a month?
- Why does the Moon appear at slightly different places in the sky each night at the same time?

OUR PLACE IN THE UNIVERSE Exploring Space

The Universe is vast. The light from the nearest star takes 4.2 years to reach Earth. If we could build a spaceship to travel at the speed of light the round trip journey time would be as follows.



Questions


- What are the three main ways scientists find out about the Universe?
- Copy and complete the following table to summarize the advantages and disadvantages of two ways of exploring the solar system.

	Manned spaceflight	Unmanned robotic probes
Advantages		
Disadvantages		

- It is proposed to send astronauts to Mars. Apart from the journey time of a couple of years, what other considerations are necessary when designing a spacecraft to make the journey?
- 'Exploring Space is a waste of money that would be better spent on giving aid to people who live in poverty.' Do you agree or disagree with this statement? Give some explanation to try to convince somebody to support your view.
- Explain why manned missions outside the solar system are very unlikely.

OUR PLACE IN THE UNIVERSE Forces in the Solar System

All masses exert gravitational attractions on all other masses.




This centripetal force is provided by the gravitational attraction between the planet and the Sun.

Satellites and moons orbiting a planet also need centripetal force acting towards the planet at the centre of their orbit. It is provided by the gravitational attraction between the satellite or moon and the planet.


From p19 Centripetal force = mass × velocity²/radius of orbit.

Therefore, to stay in orbit at a particular distance from a larger body, a smaller body must travel at a particular speed in order that the centripetal force required is exactly provided by the available gravitational attraction between the masses.



This gravitational attraction is:

- Proportional to the product of the two masses ($F \propto M \times m$)
- Inversely proportional to the square of the distance between their centres of mass ($F \propto 1/r^2$).




Mathematics


Orbital circumference = $2\pi r$
Orbital period = T
Therefore orbital speed,
 $v = 2\pi r/T$

Orbiting body

Centripetal force provided by gravitational attraction is the net resultant force.



As force and distance travelled are always perpendicular, no work is done (N.B. remember work = force × distance in the direction of the force). Therefore, the body does not need any energy to be transferred to stay in orbit.




Questions

1. A planet orbits the Sun. What would happen to the size of its gravitational attraction to the Sun if:
 - a. It doubled in mass but stayed in the same orbit?
 - b. It stayed the same mass but moved to an orbit twice the distance from the Sun?
2. What happens to the orbital period of a planet as you move away from the Sun? Does the table on p114 confirm this? Give two reasons why the orbital period varies in this way.
3. A geostationary satellite has a mass of 5 kg and an orbit radius of 42×10^6 m.
 - a. Show that its orbit circumference is about 260×10^6 m.
 - b. Given that its orbital period is 86 400 s, show that its orbital speed is about 3000 m/s.
 - c. Therefore, show that the centripetal acceleration is about 0.2 m/s^2 .
 - d. Explain why the satellite's weight in this orbit is about 1 N.

OUR PLACE IN THE UNIVERSE

The Structure of the Universe

The Universe consists of:
Planets, moons, asteroids, and comets.



Galaxies can have many different shapes.



Our Sun is in an arm of a spiral galaxy we call the Milky Way. We see this as a band of stars across the night sky as we look through the plane of our galaxy.



The nearest star to Earth is about 4.2 light years away. Since the light has taken 4.2 years to reach Earth we see the star as it was 4.2 years ago.

Each galaxy is separated from its neighbours by many times its own diameter so, as the Universe contains billions of galaxies, it is unimaginably vast.

Astronomers have detected planets around some nearby stars. If even a small proportion of the stars in the rest of the Universe have planets, some astronomers think it is very likely that there is life elsewhere in the Universe.

Like the solar system, galaxies are held together by gravity.

Some astronomers are looking for signals sent by intelligent life from elsewhere in the Universe. This is called the 'search for extraterrestrial intelligence' or SETI.

Questions


- List the following objects in order of size: galaxy, planet, star, and comet.
- What force is responsible for holding galaxies together?
- A galaxy is 100 000 light years from Earth. When we look at the galaxy through a telescope, we are seeing it as it was 100 000 years ago. Explain why.
- If the nearest star is 4 light years away, show it would take a rocket travelling at 11 km/s (the speed needed to just escape the Earth) about 109 000 years to get there. (Speed of light = 3×10^8 m/s.)
- Suggest why astronomers find it so difficult to detect planets around stars other than the Sun.

OUR PLACE IN THE UNIVERSE The Sun

For many years, scientists could not work out the source of energy for the Sun. Some thought the energy was released as the Sun shrank in size releasing gravitational potential energy. Others thought it was a chemical reaction like coal burning in a fire. However, geologists knew that the age of the Earth was about 5000 million years old and none of these ideas would provide enough energy to keep the Sun's energy output at the observed rate for anything like that long.

We now know that the Sun is about 4600 million years old and its energy comes from nuclear fusion (see p78 for more details). In the core, under extreme pressure and temperature, hydrogen nuclei are forced together to form helium nuclei releasing vast amounts of energy. There is enough fuel for another 5000 million years.

Einstein's famous relation $\Delta E = \Delta m c^2$, shows this enormous energy release, ΔE , comes at the expense of a small overall loss in the mass of the particles, Δm , linked by the speed of light $c = 3 \times 10^8$ m/s. Inside the Sun, 600 million tonnes of hydrogen are converted in nuclear fusion reactions every second, and 4 million tonnes of this is converted into energy.



Light can exert a force on particles of matter. We do not usually notice this because the force is extremely small. However, with the amount of light in the Sun the pressure is significant and stops the outer layers collapsing under gravity.

Questions

- What provides the energy for the Sun?
- How can you guess the temperature of a star simply by looking at it? (N.B. Never look directly at the Sun.)
- What problems can solar flares cause on Earth?
- If the Sun loses 4 million tonnes (4×10^9 kg) every second, use $\Delta E = \Delta m \times (3 \times 10^8 \text{ m/s})^2$ to calculate the energy output of the Sun per second, i.e. its power.
- Show that if the Sun has a diameter 109× that of the Earth, its volume is over 1 million times greater.

OUR PLACE IN THE UNIVERSE Stars and Their Spectra


Analyzing the light from stars can tell us . . .

- How bright they are (and therefore how far away they are).
- What elements they contain.
- How fast they are moving.


1. More advanced physics tells us the electrons in Rutherford's model of the atom should lose energy and spiral into the nucleus.



2. Clearly, this does not happen, atoms are stable. Neils Bohr suggested that the electrons could only have certain energy levels, like the rungs on a ladder.



All hot bodies, like stars, produce a continuous spectrum of wavelengths.



Brightness can be used to find the distance to the star (by the inverse square law [see p31]).

This is an absorption line spectrum as certain wavelengths have been absorbed by the outer layers of the star.

Absorption spectra can be thought of as like a fingerprint, or barcode, uniquely identifying the element responsible.

The missing colours are unique to the atoms that absorbed them. Each element produces its own unique *line spectrum*.

The line spectrum identifies what elements are present in the outer layers of the star.

Questions

1. What piece of apparatus could be used to split starlight into its spectrum?
2. Explain how scientists can estimate the temperature of a star by analyzing the light received on Earth from it.
3. Why are there black lines present in the spectra of stars? What information does the position of the lines give us?
4. Here are the line spectra of two stars together with the line spectra of some common elements measured in a laboratory. Which elements are present in each star?

a.



b.



Calcium

Iron

Hydrogen


Magnesium

OUR PLACE IN THE UNIVERSE The Life Story of a Star

A cloud of gas in space is called a *nebula*.

It is made mainly of hydrogen gas, but there may be small amounts of other elements and dust.

Black holes sometimes attract matter from a neighbouring star. This spirals into the black hole, like water down a plughole. This releases X-rays that astronomers can use to detect their presence.



1. The gas tends to be quite cool, preventing it dispersing into space.

2. Sometimes the density in parts of the nebula increases.


Billions of years

3. Other molecules are gravitationally attracted to this denser region.

4. The molecules moving towards this region gain kinetic energy (like a ball falling to Earth). Therefore, as the nebula contracts the gas molecules gain kinetic energy. Kinetic energy of a collection of gas molecules is usually referred to as heat energy.


Another way of thinking about this is to consider that the gravitational attraction is compressing the gas, and from the gas laws (p66) the temperature rises proportionally to the pressure.

Fusion
Nuclei are moving so fast that when they collide their electrostatic repulsion cannot keep them apart and they join.



Vast amounts of energy released.

5. The nebula becomes a protostar.



Temperature rise to 10⁶

6. The nebula begins to heat up.

This massive explosion releases so much energy that the outer layers are blown off and their nuclei are ripped apart and reformed in every possible way, making the whole range of elements in the periodic table.

Eventually can be imploded

Supernova

Briefly as bright as a whole galaxy.


This is a region of space where, because a very large mass has been compressed into a tiny (possibly zero) volume, gravity is so strong not even light can travel fast enough to escape from it.

If the mass of the neutron star is large enough even it will collapse under gravity to form a black hole.

Neutron star

The remaining core is formed entirely of neutrons. It is a neutron star.

The matter here is incredibly dense; 1 cm³ has a mass of about 10 000 000 tonnes.




Questions

- What is an interstellar gas cloud called?
- Outline the history of the Sun from its formation to its current state.
- What process provides the energy to make stars shine and stop them collapsing under gravity?
- Outline what will happen to the Sun when it runs out of hydrogen fuel in its core.
- What type of star will end in supernova and what might happen to the debris from this explosion?
- What is a black hole? Can we see them?
- The early universe only contained hydrogen. Where did all the other elements we see around us come from?


OUR PLACE IN THE UNIVERSE How Did the Solar System Form?

We saw on p122 that the Sun began to form when a nebula (of gas and dust) collapsed under gravity. The centre of the nebula began to heat up until about 4500 million years ago, when the temperature was high enough, fusion started, and the Sun became a star.

- Orbiting around the newly formed Sun was the remains of the gas and dust from the nebula.




- The debris' gravitational attraction to the Sun kept it in orbit, but the pressure from the radiation escaping from the Sun pushed the lighter gases into a larger orbit, leaving more massive dust particles orbiting closer to the Sun.



There is evidence for radiation pressure because comets tails, formed from gas like the gas in the early solar system, always point away from the Sun as the gas is blown away by the radiation from the Sun.

- The dust particles collided with each other and began to collect into larger clumps. These grew as they collected more dust into rocks, which eventually joined to form planets.




Some astronomers think the asteroid belt, between Mars and Jupiter, might be the remains of planets that collided, perhaps due to the influence of Jupiter's very strong gravitational field.



The heat generated by these collisions melted the rocks allowing the young planets to form into spheres before they cooled down.

Craters on other planets are evidence for these collisions, which still continue. We call the small rocks asteroids. Plate tectonics covers up craters on Earth, but there are still some impact craters to be seen. Scientists think these may explain some extinction processes such as that of the dinosaurs.

The gases further out in the solar system also collected together to form the gas planets.



Pluto does not fit this pattern. It is suggested that it has been captured by the Sun's gravity and did not form in the solar system.

Questions



- What force is responsible for keeping the planets in orbit around the Sun?
- Explain why the rocky planets are found close to the Sun, whilst the gas planets are found further away.
- What evidence is there that the planets formed by collisions between lumps of dust and rock?
- The explanation of how the solar system formed is just a theory. Suggest why scientists have found it difficult to get evidence to support the theory.

OUR PLACE IN THE UNIVERSE The Expanding Universe


Consider a moving source of waves.

This is the *Doppler effect*.

The change in pitch as an ambulance siren passes is an example of this.




The light from hydrogen in the galaxy looks redder than the same light from stationary hydrogen on Earth. This *red shift* tells us the galaxy is moving away. The *bigger* the red shift the *faster* the galaxy is *moving away*.



This implies the Universe is *expanding*. Consider the galaxies like dots on an inflating balloon.


All the galaxies move away from all the other galaxies. The Earth is *not* at the centre of the Universe!



Be careful – the Universe is not expanding into empty space like this balloon. It is the space *between* the galaxies that is growing, like the rubber of the balloon stretching *between* the dots.

The Universe must have started from an incredibly hot and dense point and exploded outwards. This expansion continues to this day and is called the 'Big Bang Theory of the Universe.'

OUR PLACE IN THE UNIVERSE The Expanding Universe – Further Evidence



The age of the Universe can be calculated by measuring how far galaxies have moved and their speed.

The calculated age assumes that the Universe has expanded at a steady rate.

If the expansion is slowing due to gravity, the actual age is less than this.


Our best estimate of the age of the Universe is

14 thousand million years

Some scientists disagree because they think some galaxies are older than this.

As gravity acts between all masses, it slows down the expansion of the Universe. Just like a ball thrown upwards slows down.


Size of Universe



Other theories

Oscillating Universe

Steady State



Which of fates 1, 2, or 3 will occur depends on the total mass of the Universe and the rate of expansion.

We need to calculate the mass of the Universe.

But we think the majority of the mass of the Universe is 'dark matter', which we cannot see.

Measuring this and the exact rate of expansion is very hard so the age and fate of the Universe remain very controversial.

Questions

1. What is the Doppler effect? Suggest where you might be able to observe the Doppler effect in everyday life.
2. If a galaxy was moving towards the Earth, how would the light received from it be affected? What if it was moving away?
3. Explain what led Hubble to propose that the Universe is expanding.
4. What was the Big Bang? Suggest two pieces of evidence for this theory of the Universe.
5. Outline three possible fates for the Universe. What two factors will dictate which outcome actually occurs?
6. Suggest some reasons why scientists are uncertain about the age and the fate of the Universe.
7. Make a list of three controversial facts in this topic. Explain why they are controversial. If possible, suggest some data scientists could collect to try to settle the dispute.

FORMULAE

The lists below bring together all the formulae in the book.

FORCES AND MOTION

Speed (m/s) = distance (m) / time (s)	$s = d/t$
Average speed (m/s) = total distance travelled (m) / total time taken (s)	$s = d/t$
Acceleration (m/s^2) = change in velocity (m/s) / time taken (s)	$a = \Delta v/\Delta t$
Equations of motion for uniformly accelerated motion	
$v = u + at$	
$x = ut + \frac{1}{2}at^2$	
$v^2 = u^2 + 2ax$	
v = final velocity (m/s)	
u = initial velocity (m/s)	
a = acceleration (m/s^2)	
t = time taken (s)	
x = distance travelled (m)	
Force (N) = mass (kg) \times acceleration (m/s^2)	$F = ma$
Weight (N) = mass (kg) \times acceleration due to gravity (m/s^2)	$W = mg$
Weight (N) = mass (kg) \times gravitational field strength (N/kg)	$W = mg$
Density (kg/m^3) = mass (kg) / volume (m^3)	$D = m/V$
Pressure (N/m^2 or Pa) = force (N) / area (m^2)	$P = F/A$
Momentum (kgm/s) = mass (kg) \times velocity (m/s)	
Impulse (Ns) = Force (N) \times time force acts for (s) = change in momentum (kgm/s)	$F\Delta t = mv - mu$
Centripetal acceleration (m/s^2) = [velocity (m/s)] 2 / radius (m).	$a = v^2/r$
Centripetal force = mass (kg) \times acceleration (m/s^2) = mass (kg) \times [velocity (m/s)] 2 / radius (m)	$F = mv^2/r$
Orbital speed (m/s) = orbit circumference (m) / orbital period (s)	$v = 2\pi r/T$
Moment (Nm) = Force (N) \times perpendicular distance from line of action of the force to the axis of rotation (m).	

Principle of moments

Sum of anticlockwise moments = sum of clockwise moments when in equilibrium.

Energy

Work done = force (N) \times distance moved in the direction of the force (m).	$w.d. = F \times d$
Power (W) = energy transferred (J) / time taken (s).	$P = E/t$
Energy transferred = work done	
Gravitational potential energy transferred (J) = mass (kg) \times gravitational field strength (N/kg) \times change in height (m)	$GPE = mg\Delta h$
Kinetic energy (J) = $\frac{1}{2}$ mass of object (kg) \times [speed (m/s) 2]	$KE = \frac{1}{2}mv^2$
Efficiency (%) = useful energy output (J) / total energy input (J) \times 100%.	
Nuclear energy	
Energy released (J) = change in mass (kg) \times [speed of light (m/s) 2]	$\Delta E = \Delta mc^2$

Waves

Wave speed (m/s) = frequency (Hz) \times wavelength (m).	$v = f\lambda$
Intensity (W/m^2) = power (W) / area (m^2).	$I = P/A$
Refractive index, n = speed of light in vacuum (m/s) / speed of light in medium (m/s)	$n = c/v$
Snell's Law	
Refractive index n_r = sin (angle of incidence) / sin (angle of refraction)	$n_r = \sin i / \sin r$
sin (critical angle) = refractive index of second material / refractive index of first material	
	$\sin c = n_r / n_i$

Magnification = image height / object height	
Power of lens (dioptrre) = 1/focal length (metres)	
Angular magnification = focal length of objective lens / focal length of eyepiece lens	

Electricity

Current (A) = charge passing (C) / time taken (s).	$I = Q/t$
Potential difference (V) = energy transferred (J) / charge passing (C).	$V = E/Q$
Resistance (Ω) = potential difference (V) / current (A)	$R = V/I$
Power (W) = [current (A)] 2 \times resistance (Ω)	$P = I^2R$
Power (W) = current (A) \times voltage (V)	$P = IV$
Power (W) = [voltage (V)] 2 / resistance (Ω)	$P = V^2/R$
Electrical energy (kWh) = power (kW) \times time (h)	
Kinetic energy of an electron (J) = charge on the electron (C) \times potential difference (V)	$KE = e \times V$
Transformer formula	
Primary voltage (V) / secondary voltage (V) = No. of turns on primary / No. of turns on secondary.	$V_p/V_s = N_p/N_s$

Thermal physics

Kelvin \rightarrow $^{\circ}\text{C}$ = (temperature / K) - 273	
$^{\circ}\text{C} \rightarrow$ Kelvin = (temperature / $^{\circ}\text{C}$) + 273	
Energy supplied (J) = mass (kg) \times specific heat capacity (J/kg K) \times temperature change (K)	$\Delta E = m \times s.h.c. \times \Delta T$
Energy (J) = mass (kg) \times specific latent heat (J/kg)	$E = mL$
Pressure (Pa) / temperature (Kelvin) = constant.	$P/T = \text{constant.}$
Pressure (Pa) \times volume (m^3) = constant.	$PV = \text{constant}$

Units

Length - metres, m	
Time - seconds, s	
Mass - kilogram, kg	
Speed or velocity - metres per second, m/s	
Acceleration - metres per second 2 , m/s 2	
Force - Newton, N	
Momentum - kilogram metre per second, kgm/s	
Impulse - Newton second, Ns	
Moment - Newton metre, Nm	
Density - kilograms per metre 3 , kg/m 3	
Pressure - Newton per metre 2 , N/m 2 (equivalent to 1 Pascal, Pa)	
Work done - Newton metre, Nm	
Power - Watt, W	
Energy - Joule, J (equivalent to one Newton metre, Nm)	
Frequency - Hertz, Hz	
Wavelength - metre, m	
Intensity - Watts per metre 2 , W/m 2	
Power of lens - dioptrre	
Current - Amps, A	
Charge - Coulombs, C	
Potential difference - Volts, V	
Resistance - Ohms, Ω	
Electrical energy - Joules, J (or kiloWatt-hours, kWh. 1 kWh = 3.6×10^6 J)	
Temperature - Kelvin, K or Celsius, $^{\circ}\text{C}$.	
Specific heat capacity - Joules per kilogram per Kelvin, J/kg K	
Specific latent heat - Joules per kilogram, J/kg	

INDEX

- absolute zero 60, 65
absorption spectra 121
 hydrogen gas 125
acceleration 5, 8, 10, 11
action potential 94
air resistance 13
airbags 96
alpha decay 72
alpha particles 68, 70, 71, 72, 76, 109
alpha scattering
 experiment 68
ammeter 44, 45
amplitude 28
Amps 49
analogue signals 98
antimatter 73
asteroids 114, 124
atomic structure 67, 73
 history of understanding 68
Becquerel 67
beta decay 73
beta particles (electrons) 72, 76, 109, 121
Big Bang 126
Big Crunch 126
biomass 89
black dwarf 123
black holes 122
brakes, antilock 96
braking distance 95, 96
brushes 91
camera 103
capacitor 54
carbon dating 110
carbon dioxide 33, 63, 89
cathode ray tubes 106
CDs, reading 107
Celsius scale 60
centripetal force 19, 100, 118
chemical energy 21
coil (solenoid) 59
collisions 17–18
comets 114
communication signals, digital 98
communication signals, analogue 98
computer 98
computer monitors 106
conduction (heat) 62
conductor, electrical 82
Conservation of energy 22, 26
Conservation of
 momentum 18
convection 62
cosmic rays 69
Coulomb 43, 45
crash barriers 96
craters, impact 124
critical angle 37
crumple zone 96
deceleration 10, 96
defibrillation 94
digital signals 98
diode 50, 54
distance-time graphs 6–7
DNA 76, 108
Doppler effect 105, 125
drag 13
dynamo 91
Earth
 geological processes 113
 gravitational field 12
 magnetic field 58, 113
 place in solar system 114
 structure 41, 113
Earth wire 55, 56
earthquakes 41
efficiency 27
elastic potential 21
electric current
 conditions for 44
 definition 44
electric motor 59, 91
electric shock 55, 56
electrical appliances 49
 electricity use 90
 hazards 56
 safety 55, 56
electrical circuits 45
 parallel 46
 series 46
electrical components 50–1
 non-ohmic 50–1
 ohmic 48
 resistance 47–8
electrical energy
 resistance 47–8
 transformation 80
electrical wires 55
electricity
 alternating current 53
 calculating costs 90
 direct current 53
 efficiency 90
 human body 94
 impact of life/society 80
 mains and wiring 55
 power 49, 90
 ‘rechargeable’ cells 53
 rectification 54
 safety 55, 56
 static 43
 units 90
electricity generation 82–3
 environmental impacts 87
 power stations 84
 renewable energy
 sources 88–9
electricity supply, national grid 86
electrocardiogram (ECG) 94
electromagnetic induction 82
electromagnetic spectrum 30
electromagnetic waves 30–1
 absorption 32
 dangers of 35
 and Earth’s atmosphere 33
 frequency 28, 36
 ionizing 35, 67
 polarization 40
 properties 30
 reflection 32, 102
 transmission 32
travel 31
uses of 34
electromagnetism 59
electron beams 57, 106
electrons 67, 72, 76, 109
 discovery 68
 energy levels 68, 121
electrostatic attraction 19
endoscope 104
energy
 calculations 26
 conservation 22, 26
 dissipation 27
 Sun 120
 transfer of 21, 23
 transformation 22
 types 21
energy resources 81
 influences on use 81
non-renewable 81
power stations 84
renewable 81, 88–9
energy transfer
 efficiency 27
 waves 28–9
energy wastage, reducing 64
equilibrium 20
evidence, scientific 79
explosion 17
filament lamp 50
fire risk, electricity 56
focal length 102
forces
 balanced (Newton’s First Law) 10
 effects of 9
 types of 9
 unbalanced (Newton’s Second Law) 11
fossil fuels 81, 87
free body diagrams 9

magnifying glass 103
 mantle 41, 113
 Mars 114, 116, 124
 mass 11, 12
 medicine
 nuclear radiation 34, 108
 optical fibres 104
 ultrasound 105
 medium 36
 Mercury 114, 116
 microwaves 30, 32, 33, 34
 dangers of 35
 uses of 34, 97
 Milky Way 119
 mirrors 102
 moment 20
 momentum 16, 96
 momentum conservation 17–18
 Moon 116
 moons 114
 morse code 97
 motion
 in circles 19
 equations of 8
 graphs 6–7
 motor effect 59, 91
 ‘multiplexing’ 98, 104
 muscle cells 94
 mutations 76
 national grid 86
 nebula 122
 Neptune 114
 neutrons 67, 73
 Newton's First Law 10, 13, 16
 Newton's Second Law 11, 12, 16, 25
 Newton's Third Law 15, 16, 18
 normal contact forces 15
 nuclear debate 111
 nuclear decay 71, 74–5
 nuclear energy 21
 nuclear equations 71
 nuclear fission 77
 nuclear fusion 78, 120, 122
 nuclear power 111
 nuclear power station 87
 nuclear reactor 77
 nuclear weapons 111
 Ohm 47
 Ohm's Law 48
 Oort cloud 114
 optical fibres 104
 orbits 118
 ozone layer 33
 pacemakers 94
 parallel circuit 46
 Pascal 66
 peer review 79
 photons 40
 physics, aims of 79
 planets 114, 116, 119
 characteristics 114
 formation 124
 plasma 78, 122
 plate tectonics 113
 plug, three-pin electric 55
 Pluto 114, 124
 plutonium 111
 polaroid 40
 pollution 81, 87
 positron emission tomography (PET) 108
 positrons 73
 potential difference 45, 46, 82
 potential divider 52
 power 24
 calculating 24
 definition 24
 electrical 49
 power stations
 environmental impacts 87
 working of 84
 pressure (gases) 66
 prisms 37, 38
 probes, space 117
 projectiles 14
 projector 103
 protons 67, 73
 Pythagoras' Theorem 14
 quarks 73
 radiation, ionizing 35, 67, 76, 108
 radiation (heat transfer) 63
 radio telescope 115
 radio transmission 99
 radioactive dating 110
 radioactive decay 71, 74–5
 radioactive materials, half-life 74, 109
 radioactivity 67
 alpha 70
 background 69
 beta 70
 contamination 76
 dangers and safety
 precautions 76
 gamma rays 30, 32, 33, 70, 76, 108, 109
 N/Z curve 72
 nuclear debate 111
 uses of 34, 108–12
 radioisotope tracers 108, 109
 radiotherapy 108
 radiowaves 30, 32, 33, 34, 35, 97
 carrier 99
 radon gas 69
 ray diagrams 101, 102
 reaction of a surface 15
 rectification 54
 red giant 123
 red shift 125
 reflection 36, 102
 refraction 36, 103
 refractive index 38
 relativity theory 107
 renewable energy 81, 88–9
 resistance, electrical 47–8, 86
 resistive forces 10, 13
 resistors 51, 52
 resultant force 9, 10, 16
 Right Hand Grip Rule 59
 Right Hand Rule 82
 road safety 96
 Sankey diagram 22
 satellites
 communication 97
 orbits 100, 118
 Saturn 114, 116
 science, methods 79
 seatbelts 96
 seismic waves 41
 series circuit 46
 sidereal day 116
 Sieverts 76
 signals, ‘multiplexing’ 98, 104
 smoke alarm 109
 Snell's Law 38
 solar collectors 89
 solar day 116
 solar eclipse 116
 solar energy 88, 89
 solar flares 120
 solar system 114
 forces 118
 formation 124
 solenoid (coil) 59
 sound, speed of 29
 sound waves 42
 space exploration 117
 space telescopes 115
 specific heat capacity 61
 spectrum
 electromagnetic 30
 stars 121
 speed 5
 stability 20
 stars 119
 life story 122–3
 spectra 121
 structure 120
 static electricity 43
 sterilization 109
 stopping distances 95
 strong nuclear forces 72, 77
 Sun
 age of 120, 123
 structure 120
 surface, reaction of 15
 tectonic plates 41, 113
 telescopes 115
 temperature 60, 65
 and pressure 66
 scales 60
 thermal energy 21, 60–6, 62
 reducing wastage 64
 transfer 62–3
 thermionic emission 57
 thermistor 51, 52
 thickness measurement 109
 thought experiments 100, 107
 tidal energy 87, 89
 total internal reflection 37, 38, 104
 trajectory 14
 transducer 21
 transformers 85, 86
 truth table 92, 93
 TVs 106
 ultrasound 105
 ultraviolet (UV) light 30, 31
 dangers of 35
 uses of 34
 Universe
 age of 126
 collecting data 115, 117
 expansion 125–6
 fates of 126
 structure 119
 uranium 77, 111
 Uranus 114
 vector 5
 vehicles
 safety features 96
 stopping distances 95
 velocity 5, 16, 17
 terminal 13
 Venus 114, 116
 voltage (electromotive force/emf) 45, 46, 49
 Watts 24, 49
 wavelength 28, 29
 waves
 description 28
 diffraction 39
 dispersion 38
 frequency 28, 36
 interference 39
 longitudinal 28, 42
 reflection 36, 102
 refraction 36, 103
 seismic 41
 sound 42
 speed 29, 36
 total internal reflection 37
 transverse 28
 see also electromagnetic waves
 Wegener's theory 113
 weight 9, 12
 white dwarf star 123
 wind power 87, 88
 work done 23
 X-ray tubes 106
 X-rays 30, 32, 33
 dangers of 35
 uses of 34, 108