

# PLAcER: Plastic Accumulation Estimation using R

*Davi Castro-Tavares, Esteban Acevedo-Trejos & Jailson F. Moura*

*2019-08-27*

## Contents

Overview . . . . .	1
Installation . . . . .	1
Usage . . . . .	1
References . . . . .	2

## Overview

### Installation

To install simply type in the R console

```
install.packages('placer')
```

### Usage


Load data containing observations of presence and absence of plastic debris.

species	location	country	latitude	longitude	year	debris_presence
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes
hydroprogne_caspia	ile_aux_ouiseaux	Senegal	13.62795	-16.63338	2018	yes

Calculate the bootstrap prevalence probability of plastic based on a specific number of samples. Then from the resampled data the 95% confidence intervals (CI) are computed. To quantify the prevalence probability and the CI simply provide the presence/absence data, the maximum number of sample sizes to test (default 300), and the number of bootstrap simulations (default 1000) as:

```
ternsci<-placer::plastic.ci(cterns$debris_presence, 300, 1000)
```

Plot the plastic preevalence probability and the confidence interval as function of sample size:



## References