

1. We consider the training examples shown in the following table for a binary classification problem.

Instance	$a_1$	$a_2$	$a_3$	Target Class
1	T	T	1	+
2	T	T	6	+
3	T	F	5	-
4	F	F	4	+
5	F	T	7	-
6	F	T	3	-
7	F	F	8	-
8	T	F	7	+
9	F	T	5	-

- a) Calculate the respective changes in the Gini index value when  $a_1$  and  $a_2$  are used for partitioning the training set.

The original Gini index is  $1 - (\frac{4}{9})^2 - (\frac{5}{9})^2 = 0.494$

After splitting on  $a_1$ , the Gini index becomes

$$\frac{4}{9}[1 - (\frac{3}{4})^2 - (\frac{1}{4})^2] + \frac{5}{9}[1 - (\frac{1}{5})^2 - (\frac{4}{5})^2] = 0.344$$

As a result, the change in Gini index is

$$\Delta G(a_1) = 0.494 - 0.344 = 0.15.$$

After splitting on  $a_2$ , the Gini index becomes

$$\frac{5}{9}[1 - (\frac{2}{5})^2 - (\frac{3}{5})^2] + \frac{4}{9}[1 - (\frac{2}{4})^2 - (\frac{2}{4})^2] = 0.489$$

As a result,

$$\triangle G(a_2) = 0.494 - 0.489 = 0.005.$$

b) Calculate the respective changes in the classification error when

$a_1$  and  $a_2$  are used for partitioning the training set.

The original classification error is  $1 - \max(\frac{4}{9}, \frac{5}{9}) = \frac{4}{9}$

After splitting on  $a_1$ , the classification error becomes

$$\frac{4}{9}[1 - \max(\frac{3}{4}, \frac{1}{4})] + \frac{5}{9}[1 - \max(\frac{1}{5}, \frac{4}{5})] = \frac{2}{9}$$

As a result, the change in classification error is

$$\triangle E(a_1) = 4/9 - 2/9 = 2/9.$$

After splitting on  $a_2$ , the classification error becomes

$$\frac{5}{9}[1 - \max(\frac{2}{5}, \frac{3}{5})] + \frac{4}{9}[1 - \max(\frac{2}{4}, \frac{2}{4})] = \frac{4}{9}$$

As a result,

$$\triangle E(a_2) = 4/9 - 4/9 = 0.$$

c) For  $a_3$ , which is a continuous attribute, compute the information gain for every possible split. What is the best threshold for splitting the set of attribute values?

We consider the different possible split points for  $a_3$  as follows:

$a_3$	Class label	Split point	Entropy	Info gain
1	+	2.0	0.848	0.143
3	-	3.5	0.989	0.002
4	+	4.5	0.918	0.073
5	-	5.5	0.984	0.007
5	-			
6	+	6.5	0.973	0.018
7	+	7.5	0.889	0.102
7	-			
8	-			

The best split for  $a_3$  occurs when the split point is equal to 2.