mun. trenepalers.com

## CAMBRIDGE INTERNATIONAL EXAMINATIONS GCE Advanced Level

## MARK SCHEME for the May/June 2013 series

## 9701 CHEMISTRY

9701/53

Paper 5 (Planning, Analysis and Evaluation), maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9701	53

Question		ion	Expected Answer	
1	(a)	(i)	(Solubility will) decrease	1
			Dissolving/reaction is exothermic so reaction shifts left (owtte).	1
			Increase negates both marks.	
			<b>Allow:</b> Variations in the wording but the word exothermic or heat evolved or the reverse process must be included.	
		(ii)	Axes are correctly labelled <b>AND</b> graph is a curve/straight line showing a decrease in solubility with temperature. (ignore units)	1
			Graph goes through the point 25 on temperature scale and 5 on solubility scale <b>AND</b> goes from 0 to 100 °C	1
			Allow ecf from (i) prediction.	
	(b)		(i) temperature (increase)	
			(ii) solubility (of chlorine)	1
	(c)	1	Pipette (5,10, 20, 25, 50 cm <sup>3</sup> ), burette (25, 50 or 100 cm <sup>3</sup> ) both required for mark.	1
		2	Starch indicator <b>AND</b> blue/blue-black <b>AND</b> colourless/opaque.	1
		3	Concentration of $Cl_2 = 0.0704 \text{ mol dm}^{-3}$ .	1
		4	Calculates $M_r$ of Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> .5H <sub>2</sub> O as 248.2 <b>AND</b> calculates mass with unit required for a solution of stated concentration and volume. (Allow any concentration)	1
		5	Mass and volume used must produce a solution twice as concentrated as the chlorine solution (ecf from $Cl_2$ ).	1
		6	Describes making of solution in volumetric flask which must include: dissolving, making up to mark.	1
		7	Titration is repeated to achieve concordant titration results/average titre, 'concordant' not required if meaning clear.	1
		8	Calculates moles $Cl_2$ in titration from $0.5 \times$ moles thiosulfate in titre and therefore concentration <b>AND</b> concentration of $Cl_2$ in mol dm <sup>-3</sup> in aqueous chlorine.	1
			Allow any explanation which covers these points, calculations involving concentrations or moles to mass and concentration in $gdm^{-3}$ , or any formula that would produce a correct answer e.g. mv / n = mv / n	

	GCE AS/A LEVEL – May/June 2013	9701	53
(d)	Chlorine <b>OR</b> iodine are harmful	to akin/ayaa aiya	1
Wear a mask/use a fume cupboard/for iodine if harmful to skin/eyes given, allow resistant gloves/goggles		n, <b>1</b>	
			[Total: 1

Syllabus

Paper

Mark Scheme

Page 3

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2013	9701	53

2 (a)	151.9 <b>AND</b> 18.0	151.9 <b>AND</b> 18.0			1	
(b)	Columns are headed with a label, an expression and units as below.				1	
	Mol of FeSO <sub>4</sub> AN	Mol of FeSO <sub>4</sub> <b>AND</b> mol of H <sub>2</sub> O are correct to 3 sig. figs.				1
	ECF incorrect M	r. ECF the use	of incorrect expr	ressions into data	а.	
	D	E	F	G		
	FeSO <sub>4</sub> (C – A) / g	H₂0 B – C / g	FeSO <sub>4</sub> (C – A) / 151.9 OR D / 151.9 mol OR mole	H <sub>2</sub> O (B – C) / 18 OR E / 18 mol OR mole		
	1.00	0.83	0.00658	0.0461		
	1.31	1.00	0.00862	0.0556		
	1.30	1.08	0.00856	0.0600		
	1.39	1.16	0.00915	0.0644		
	1.50	1.24	0.00987	0.0689		
	1.63	1.35	0.0107	0.0750		
	1.78	1.48	0.0117	0.0822		
	1.84	1.53	0.0121	0.0850		
	1.95	1.62	0.0128	0.0900		
	2.03	1.76	0.0134	0.0978		
(c)		<i>x</i> -axis labelled 'mol of FeSO <sub>4</sub> ' and <i>y</i> -axis 'mol H <sub>2</sub> O' <b>AND</b> plotted points cover at least half the grid in both directions.			1	
	Allow a correct le	Allow a correct letter from the table as a label.				
	All 10 points plotted correctly.				1	
	Best fit straight	Best fit <b>straight</b> line drawn.				1

Page 5	ge 5 Mark Scheme		Paper
	GCE A LEVEL – May/June 2013	9701	53

	1		Г	
(d)	Points 2 and 10 circled. (The circled points in the reasons.)	nts must be unambiguously referred	1	
Point 2 (mass of crucible 15.10) Not all the water had be sulfate crystals <b>OR</b> anhydrous FeSO <sub>4</sub> absorbed some wimpurity that does not decompose.			1	
	Allow water loss is low(er) (than expecte	ed).		
	Point 10 (mass of crucible = 15.01) The anhydrous FeSO <sub>4</sub> had decomposed <b>OR</b> prior to heating the crucible/original sample was wet and water removed on heating <b>OR</b> contained an impurity which decomposed/was removed on heating.			
	Allow some mass lost (spits out) on heat	ting.		
(e)	Appropriately drawn lines on the graph.		1	
	Correctly read values from the graph.		1	
	(Figures from the table allowed if no construction lines drawn providing graph drawn does actually go through the points used.)			
	Correctly calculated value of the slope given to 2 or more sig. figs up to calculator value and using the <b>candidate's</b> figures <b>AND</b> no units given.		1	
(f)	Most of the points are on the line <b>OR</b> only a few points are not on the line <b>OR</b> there are only a few anomalies.		1	
(g) (i)	FeSO <sub>4</sub> .7H <sub>2</sub> O (ecf on slope in <b>(e)</b> )		1	
(ii)	The gradient/slope is the ratio of (moles) of H <sub>2</sub> O:FeSO <sub>4</sub> (is 7 or 7:1).		1	
		[Tota	l: 15]	
	1	1		