



CANDIDATE NAME

CENTRE NUMBER

Cambridge International Examinations

Cambridge International Examinations Cambridge International Advanced Subsidiary and Advanced Level			
	CANDIDATE		

CHEMISTRY

9701/23

Paper 2 Structured Questions AS Core

May/June 2014

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials:

Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 9 printed pages and 3 blank pages.



Answer **all** the questions in the spaces provided.

1	(a)	Define the term <i>mole</i> .			
			[1		
	(b)	100	cm ³ of a gaseous hydrocarbon, C_xH_y , was reacted with 100 cm ³ of oxygen gas, an excess.		
		The	e final volume of the gaseous mixture was 95 cm ³ .		
		This gaseous mixture was treated with concentrated, aqueous sodium hydroxide to absorb the carbon dioxide present. This reduced the gas volume to 75 cm ³ .			
		All gas volumes were measured at 298 K and 100 kPa.			
		(i)	Write an equation for the reaction between sodium hydroxide and carbon dioxide.		
			[1		
		(ii)	Calculate the volume of carbon dioxide produced by the combustion of the hydrocarbon.		
			volume of CO ₂ produced = cm ³ [1		
		(iii)	Calculate the volume of oxygen used up in the reaction with the hydrocarbon.		
			volume of O_2 used = cm ³ [1		
		<i>(</i> ,)	-		
		(iv)	Use your answers to (b)(ii) and (b)(iii) , together with the initial volume of hydrocarbon, to balance the equation below.		
			$C_xH_y +O_2 \rightarrowCO_2 + zH_2O$ [2]		
		(v)	Deduce the values of x , y and z in the equation in (iv).		
			<i>X</i> =		
			<i>y</i> =		
			z =[3		
			[o		

(c)	Another hydrocarbon, \mathbf{W} , with the formula C_4H_8 , reacts with hydrogen bromide, HBr, to give two products \mathbf{X} and \mathbf{Y} . \mathbf{X} and \mathbf{Y} are structural isomers of molecular formula C_4H_9Br .			
	Reaction of \boldsymbol{X} with aqueous alkali production of \boldsymbol{X} with aqueous alkali production dichromate (VI).	uces an alcohol, Z , that has no reaction with acidified		
	(i) Give the structures and names of the	ne compounds W , X , Y , and Z		
	W	X		
	Y	Z		
		[4]		
	(ii) When W reacts with hydrogen brom	nide, more X than Y is produced. Explain why.		

[Total: 15]

.....[2]

	ble of a hydrated double salt, $Cu(NH_4)_x(SO_4)_2.6H_2O$, was boiled with an excess of sodiide. Ammonia was given off.	um
	nmonia produced was absorbed in $40.0\mathrm{cm^3}$ of $0.400\mathrm{moldm^{-3}}$ hydrochloric acid. The result n required $25\mathrm{cm^3}$ of $0.12\mathrm{moldm^{-3}}$ sodium hydroxide to neutralise the excess acid.	ing
(a) W	rite the ionic equation for the reaction between ammonium ions and hydroxide ions.	
(b) (i)	Calculate the amount, in moles, of hydrochloric acid in 40.0 cm ³ of 0.400 mol dm ⁻³ solution	
(ii)	Calculate the amount, in moles, of sodium hydroxide needed to neutralise the exceading acid. This will be equal to the amount of hydrochloric acid left in excess.	[1] ess
(iii)	Calculate the amount, in moles, of hydrochloric acid that reacted with ammonia.	[1]
(iv)	Calculate the amount, in moles, of ammonium ions in the sample of the double salt.	[1]
(v)	The sample contained 0.413 g of copper. Use this information and your answer to (iv) calculate the value of x in $Cu(NH_4)_x(SO_4)_2.6H_2O$.	[1]) to
(vi)	Calculate the $M_{\rm r}$ of Cu(NH ₄) _x (SO ₄) ₂ .6H ₂ O.	[2]
		[1]

[Total: 8]

BLANK PAGE

3	Nitı	roger	n dioxide, NO ₂ , can enter the atmosphere in a variety of ways.
	(a)	(i)	State one natural and one man-made source of atmospheric NO ₂ .
			natural
			man-made
			[1]
		(ii)	Write an equation to show how NO ₂ leads to the formation of nitric acid in acid rain.
			[1]
		(iii)	Use equations to illustrate the catalytic role of NO_2 in the formation of sulfuric acid in acid rain.
			[3]
	(b)	Nitr	ogen dioxide exists in equilibrium with dinitrogen tetroxide, N ₂ O ₄ .
			$2NO_2(g) \rightleftharpoons N_2O_4(g)$
		esta	Omol of dinitrogen tetroxide was sealed in a container at 350 K. After equilibrium had been ablished the total pressure was 140 kPa and the mixture of gases contained 1.84 mol of trogen tetroxide.
		(i)	Give the expression for the equilibrium constant, K_p , for this equilibrium.
			K_{p} =
			[1]
		(ii)	Calculate the number of moles of NO ₂ present at equilibrium.
			[1]
		(iii)	Calculate the total number of moles of gas present at equilibrium and hence the mole fraction of each gas present at equilibrium.

(iv)	Calculate the partial pressure of each gas present at equilibrium.
	[2]
(v)	Calculate the value of the equilibrium constant, K, at 350 K.
(-)	Calculate the value of the equilibrium constant, K_p , at 350 K. Give your answer to three significant figures and include the units.
	$K_p = \dots$
	units =
	[2]
	[Total: 13]

4

The halogens and their compounds have a wide variety of uses and the chemical and physical properties of the elements show regular patterns related to their positions in Group VII.

(a)	Chlorine, bromine and iodine all react with hydrogen.			
	(i)	State the trend in the reactivities of the halogens with hydrogen.		
		[1]		
	(ii)	Explain this trend in terms of bond energies.		
		[2]		
(b)	In th	ne laboratory it is not very convenient to prepare hydrogen halides from their elements.		
	Нус	drogen halides can be prepared from their salts.		
	(i)	Write an equation for the reaction of calcium chloride, ${\rm CaC}l_2$, with concentrated sulfuric acid.		
		[1]		
	(ii)	Explain why hydrogen iodide is not prepared in this way.		
		[1]		
	(iii)	When potassium bromide, KBr, reacts with concentrated sulfuric acid, sulfur dioxide, SO_2 , is produced. State what you would see and write an equation for this reaction.		
		[3]		

(c)	(1)	secondary or tertiary.	isomers of C₄H ₉ Br and identify each as primary		
			[4]		
	(ii)	Name the isomer of C ₄ H ₉ Br that contains structures of the two optical isomers.	a chiral centre and draw the three-dimensiona		
		name			
		structures			
			[3]		
(d)	Aqueous silver nitrate solution was added to separate tubes containing chloroethane, bromoethane and iodoethane. The tubes were heated in a water bath.				
	A yellow precipitate appeared first in the tube containing iodoethane, followed by a cream precipitate in the tube containing bromoethane and finally a white precipitate appeared in the tube containing chloroethane.				
	Exp	plain these observations.			
			[2]		

(e)	(i)	Give the full name of the mechanism for the reaction between aqueous sodium hydroxic and bromoethane.	de		
		[2]		
	(ii)	Complete the diagram below to represent this mechanism. Include all necessary curly arrows, partial charges and lone pairs.			
		H H H			
		HO ⁻			
		[.	2]		
(f)	In th	the past, CFCs such as CF_3Cl were widely used as refrigerants.			
	(i)	State a property of CFCs which makes them suitable for use as refrigerants.			
		[1]		
	(ii)				
		Explain your answer.			

[Total: 24]

[2]

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.