

www.xiremenabers.com



| CANDIDATE<br>NAME |  |                     |  |  |
|-------------------|--|---------------------|--|--|
| CENTRE<br>NUMBER  |  | CANDIDATE<br>NUMBER |  |  |

CHEMISTRY 9701/43

Paper 4 Structured Questions

May/June 2011

2 hours

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

### **Section A**

Answer all questions.

### Section B

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

| For Examiner's Use |  |  |
|--------------------|--|--|
| . O. =Xuiii        |  |  |
| 1                  |  |  |
| 2                  |  |  |
| 3                  |  |  |
| 4                  |  |  |
| 5                  |  |  |
| 6                  |  |  |
| 7                  |  |  |
| 8                  |  |  |
| Total              |  |  |

This document consists of 16 printed pages and 4 blank pages.



## Section A

For Examiner's Use


Answer all questions in the spaces provided.


| 1   | (a) | Hyd   | rogen fluoride, HF, b  | ehaves as a weak acid i              | in water, with $K_a = 5.6 \times 10^{-4} \text{mol dr}$ | n <sup>–3</sup> . |
|-----|-----|-------|--|--------------------------------------|---|-------------------|
|     |     | Cald  | culate the pH of a 0.0   | $050\mathrm{moldm^{-3}}$ solution of | HF.   |                   |
|     |     |       |  |                                      |   |                   |
|     |     |       |  |                                      |   |                   |
|     |     |       |  |                                      | pH =  | [2]               |
|     | (b) |       | eous ammonia and ride.   | hydrogen fluoride react              | together to give solid ionic ammor                      | nium              |
|     |     |       | NH <sub>3</sub> (g) + HF   | $f(g) \rightleftharpoons NH_4F(s)$   | $\Delta H = -147 \text{kJ} \text{mol}^{-1}$             |                   |
|     |     | (i)   | What type of reaction  |                                      |   |                   |
|     |     |       |  |                                      |   |                   |
|     |     | (ii)  | Draw dot-and-cross compounds involved  |                                      | only) describing the bonding in the t                   | hree              |
|     |     |       |  |                                      |   |                   |
|     |     |       |  |                                      |   |                   |
|     |     |       | NH <sub>3</sub>  | HF                                   | NH <sub>4</sub> F                                       |                   |
| (ii |     | (iii) |  |                                      | and state where in the compound o                       | each              |
|     |     |       | There are <b>three</b> types of bonding in $NH_4F$ . Give the names of each of the three types, and state where in the compound each |                                      |   |                   |
|     |     |       |  |                                      |   |                   |
|     |     |       |  |                                      |   |                   |
|     |     |       |  |                                      |   |                   |

|     | (iv)     | The reaction between $\mathrm{NH_3}$ and HF is reversible. What conditions of temperature and pressure would favour the <b>reverse</b> reaction, i.e. the dissociation of $\mathrm{NH_4F?}$ Explain your answer.   | For<br>Examiner's<br>Use |
|-----|----------|--|--------------------------|
|     |          |  |                          |
|     |          | [9]  |                          |
| (c) | on tair, | by commercial copper and brass polishes contain ammonia. The tarnish that forms the surface of copper is often copper sulfide, CuS. In the presence of $\rm O_2$ from the NH $_3$ can combine with this copper sulfide to produce the soluble cuprammonium ate, $\rm [Cu(NH_3)_4]SO_4$ . |                          |
|     | (i)      | Construct an equation for this reaction.   |                          |
|     | (ii)     | State the colour of cuprammonium sulfate solution.   |                          |
|     | (iii)    | Describe what you would see if a solution of cuprammonium sulfate was diluted with water. Explain your answer.   |                          |
|     |          | [3]  |                          |
| (d) | hyd      | en sulfuric acid is added to $Cu^{2+}(aq)$ , no colour change occurs, but when concentrated rochloric acid is added to $Cu^{2+}(aq)$ , the solution turns yellow-green. The solution erts to its original colour when it is diluted with water.  |                          |
|     |          | gest the type of reaction occurring with $HCl(aq)$ , suggest what is formed during the ction, and write an equation for the change.  |                          |
|     |          |  |                          |
|     |          | [3]  |                          |
|     |          | [Total: 17]  |                          |

2 (a) (i) On the following grids, plot points showing the variation in the named property of the Group IV elements. Your points should show for each element, whether the melting point/electrical conductivity is 'high', 'medium' or 'low'. The point for silicon has already been plotted in each case.

For Examiner's Use





Suggest explanations of these trends in terms of the structure and bonding of the Group IV elements.

| melting point           |      |     |
|-------------------------|------|-----|
|                         | <br> |     |
| electrical conductivity | <br> |     |
|                         | <br> |     |
|                         | <br> | [6] |

(b) Choose one reaction to illustrate each of the following statements. Write an equation for each of your chosen reactions, and describe what you would see as the reaction is carried out.

| (i)   | PbO is more stable than PbO <sub>2</sub> .            |
|-------|---|
|       |   |
|       |   |
| (ii)  | CO is easily oxidised to CO <sub>2</sub> .            |
|       |   |
|       |   |
| (iii) | Aqueous SnCl <sub>2</sub> is a useful reducing agent. |
|       |   |
|       |   |

[Total: 10]

[4]

| <ul> <li>(b) If the charge on the electron, the A<sub>r</sub> and the valency of copper are known, the value of the Avogadro number can be determined experimentally. This is done by passing a known current for a known time through a copper electrolysis cell, and weighing the mass of copper deposited onto the cathode.</li> <li>(i) Draw a diagram of suitable apparatus for carrying out this experiment.  Label the following: power supply (with + and - terminals); anode; cathode; and ammeter.  State the composition of the electrolyte.</li> <li>The following are the results obtained from one such experiment.  current passed through the cell = 0.500 A  time current was passed through cell = 30.0 min  initial mass of copper cathode = 52.243 g  final mass of copper cathode = 52.542 g</li> <li>(ii) Use these data and relevant information from the Data Booklet to calculate a value of L to 3 significant figures.</li> </ul> |     |         | te the relationship between the Faraday constant, $F$ , the charge on the electron, $e$ , the Avogadro number, $L$ .  |
|--|-----|---------|---|
| Label the following: power supply (with + and – terminals); anode; cathode; and ammeter.  State the composition of the electrolyte.  The following are the results obtained from one such experiment.  current passed through the cell = 0.500 A  time current was passed through cell = 30.0 min  initial mass of copper cathode = 52.243 g  final mass of copper cathode = 52.542 g  (ii) Use these data and relevant information from the Data Booklet to calculate a value   | (b) | of that | he charge on the electron, the $A_r$ and the valency of copper are known, the value he Avogadro number can be determined experimentally. This is done by passing nown current for a known time through a copper electrolysis cell, and weighing the |
| current passed through the cell = 0.500 A time current was passed through cell = 30.0 min initial mass of copper cathode = 52.243 g final mass of copper cathode = 52.542 g  (ii) Use these data and relevant information from the Data Booklet to calculate a value   |     | (i)     | Label the following: power supply (with + and - terminals); anode; cathode; and ammeter.  |
| current passed through the cell = 0.500 A time current was passed through cell = 30.0 min initial mass of copper cathode = 52.243 g final mass of copper cathode = 52.542 g  (ii) Use these data and relevant information from the Data Booklet to calculate a value   |     |         |   |
|  |     |         | current passed through the cell = 0.500 A<br>time current was passed through cell = 30.0 min<br>initial mass of copper cathode = 52.243 g   |
|  |     | (ii)    |   |
|  |     |         |   |
| <i>L</i> =[9]  |     |         |   |

**(c)** Use relevant information from the *Data Booklet* to identify the substances formed at the anode and at the cathode when aqueous solutions of the following compounds are electrolysed.

For Examiner's Use

| compound          | product at anode | product at cathode |
|-------------------|------------------|--------------------|
| AgF               |                  |                    |
| FeSO <sub>4</sub> |                  |                    |
| MgBr <sub>2</sub> |                  |                    |

[5]

[Total: 15]

**4 (a)** Polyvinyl acetate, PVA, is a useful adhesive for gluing together articles made from wood, paper or cardboard. The monomer of PVA is ethenyl ethanoate, **B**.

For Examiner's Use

PVA is formed from **B** by the process of addition polymerisation.

(i) Draw a section of the PVA molecule containing at least 2 monomer molecules, and identify clearly the repeat unit.

The ester **B** can be hydrolysed in the usual way, according to the following equation.

$$CH_3$$
  $O$   $+ H_2O$   $CH_3$   $O$   $+$   $CH_3$   $O$   $+$   $C(C_2H_4O)$ 

(ii) Use this information to suggest a possible structure for **C** and draw it in the box above.


When substance  $\bf C$  is extracted from the product mixture, it is found that it does **not** decolourise  $Br_2(aq)$ , but it **does** form a pale yellow precipitate with alkaline aqueous iodine.

(iii) Suggest a structure for C that fits this new information.

| (iv) | Suggest a confirmatory test for the functional group in the structure you have drawn in (iii). Your answer should include the reagent you would use and the observation you would make. |
|------|---|
|      |   |
|      |   |

[6]

(b) The following diagram represents a section of another polymer.





- (i) On the above formula draw brackets, [ ], around the atoms that make up the repeat unit of this polymer.
- (ii) Name the functional group in polymer **D**.

- (iii) Suggest and draw the structure of the monomer, **E**, that could form this polymer.
- (iv) What *type of polymerisation* is involved in making polymer **D** from its monomer?

What is the relationship between the repeat unit of polymer D and the repeat unit of PVA?

- (c) Monomer **E** exists as two stereoisomers. Heating either isomer with  $Al_2O_3$  gives a mixture of two unsaturated carboxylic acids **F** and **G**, which are stereoisomers of each other.
  - (i) Name the *type of stereoisomerism* shown by compound **E**.

.....

(ii) Suggest structures for **F** and **G**, and name the type of stereoisomerism they show.


type of isomerism ......[4]

- -

| 5 (a)  | Describe a           | nd explain how the acidi  | ties of ethanol and phenol compare to that of water.   |
|--------|----------------------|---|--|
|        |                      |   |  |
|        |                      |   |  |
|        |                      |   |  |
|        |                      |   |  |
|        |                      |   |  |
|        |                      |   |  |
|        |                      |   | [4]  |
| (b)    | phenol. Inc          | he following equations s<br>lude reaction conditions<br>curs write <i>no reaction</i> i | howing <b>all</b> the products of each of these reactions of where appropriate in the boxes over the arrows. If no n the products box. |
|        |                      |   |  |
| OH<br> |                      |   |  |
|        |                      |   |  |
|        | + Na                 | -   |  |
|        |                      |   |  |
|        |                      |   |  |
| OH<br> |                      |   |  |
|        | NI OII               |   |  |
|        | + NaOH               |   |  |
|        |                      |   |  |
|        |                      |   |  |
| OH<br> |                      |   |  |
|        |                      |   |  |
|        | + CH <sub>3</sub> CC | D <sub>2</sub> (1)  |  |
|        |                      |   |  |
|        |                      |   |  |
| OH     |                      |   |  |
|        | + Br <sub>2</sub>    |   |  |
|        | . 5.2                | -   |  |
|        |                      |   |  |
|        |                      |   | [5]  |

**(c)** The analgesic drug paracetamol can be synthesised from phenol by the following route. Suggest reagents and conditions for the each of three steps, and suggest the structure of the intermediate **H**. Write your answers in the boxes provided.

For Examiner's Use



[Total: 13]

[4]

## **Section B**

For Examiner's Use

Answer all questions in the spaces provided.

6


|     |       | s are protein molecules that are highly efficient in catalysing specific chemical s in living organisms.  |
|-----|-------|---|
| (a) |       | work in tissues, enzyme molecules generally need to be water-soluble. What does tell you about the nature of the side-chains on the exterior of the molecules?  |
|     | ••••• | [4]   |
|     | ••••• | [1]   |
| (b) | enz   | symes function by a substrate molecule interacting with a particular part of the yme known as the 'active site'. The substrate is converted into products that are then ased, to be replaced by another substrate molecule. |
|     | (i)   | Describe briefly the primary, secondary and tertiary structures of an enzyme.   |
|     |       |   |
|     |       |   |
|     |       |   |
|     |       |   |
|     |       |   |
|     |       |   |
|     | (ii)  | The activity of an enzyme depends upon the tertiary structure of the protein molecule. Explain how the tertiary structure produces an effective active site.  |
|     |       |   |
|     | (iii) | Give <b>two</b> conditions that can <b>reduce</b> the activity of an enzyme, explaining the reason in each case.  |
|     |       | I   |
|     |       |   |
|     |       |   |
|     |       | II  |
|     |       |   |
|     |       | [6]   |

(c) An individual enzyme operates best at a specific pH. Different enzymes operate best under conditions of different pH. Three enzymes involved in the digestion of food are amylase, pepsin and trypsin.

For Examiner's Use

- Amylase, found in saliva, hydrolyses starch to a mixture of glucose and maltose under approximately neutral conditions.
- Pepsin hydrolyses proteins to peptides in the acid conditions of the stomach.
- Trypsin continues the hydrolysis of peptides to amino acids in the mildly alkaline conditions of the small intestine.

The graph below shows the activity of two of the three enzymes mentioned above.



- (i) Label each peak shown with the name of the enzyme responsible, either amylase, pepsin or trypsin.
- (ii) On the axes above, sketch the graph that the third enzyme would produce, and label it with the name of that enzyme.

[3]

[Total: 10]

7 The technique of DNA fingerprinting has been one of the most important developments in biochemical analysis in recent times. It has enabled enormous advances to be made in forensic science, medicine and archaeology.

For Examiner's Use

(a) The table shows different stages in the production of a genetic fingerprint. Use the numbers 1 to 6 to put the stages in the correct sequence in the blank column.

| stages | process                        | correct sequence<br>(numbers) |
|--------|--------------------------------|-------------------------------|
| Α      | place samples on agarose gel   |                               |
| В      | use polymerase chain reaction  |                               |
| С      | label with radioactive isotope |                               |
| D      | extract DNA                    |                               |
| Е      | use restriction enzyme         |                               |
| F      | carry out electrophoresis      |                               |

[3]


| /L\ | \ \O_= = = f | the stages | 0 0 0 10 110 |            |         |        |
|-----|--------------|------------|--------------|------------|---------|--------|
| m   | i Une oi     | The stades | anove us     | ses a radi | oacnvei | SOIODE |
|     |              |            |              |            |         |        |

| (i) | What icotopo is used?  |  |
|-----|------------------------|--|
| (I) | vynal isolobe is used? |  |

| 1 | (ii) | Why   | is | this | isotone | chosen? |
|---|------|-------|----|------|---------|---------|
| ٨ | ,    | VVIIV | 13 | นแง  | 1301000 | CHOSCHE |

| , 10 11110 10010   10 011000111 |     |
|---------------------------------|-----|
|                                 |     |
|                                 |     |
|                                 |     |
|                                 | [2] |

(c) The following DNA fingerprints were taken from a family of mother, father and four children.



|     | (i)  | Are all of the children related to the mother? State the evidence for your answer.  |
|-----|------|---|
|     | (ii) | Which child is unlikely to be related to the father? State the evidence for your answer.  |
|     |      |   |
| (d) | DN   | [2] A fingerprinting has been successfully used in archaeological investigations.   |
|     | (i)  | Ancient writings were often made on goatskins. Over the centuries these have often become broken into fragments, making reconstruction of the writings almost impossible. |
|     |      | Suggest how the use of DNA fingerprinting might be able to identify which fragments came from a particular skin.  |
|     |      |   |
|     |      |   |
|     |      |   |
|     |      |   |
|     | (ii) | Apart from the examples of human remains and goatskins, state one other material that could be investigated using this technique.   |
|     |      |   |
|     |      | [3]   |
|     |      | [Total: 10]   |

|   |     | 16   |
|---|-----|--|
| 8 Nanotechnology is a fast-developing area of science based on the ability to m materials of very small dimensions. |     |  |
|   | (a) | On the scale shown in metres, mark the upper and lower limits of the range of sizes for nanoparticles.   |
|   |     | $ \begin{array}{cccccccccccccccccccccccccccccccccccc$  |
|   | (b) | One of the most commonly recognised nanoparticles is the 'buckyball', a spherical form of carbon containing 60 carbon atoms. It has been referred to as the third allotrope of carbon. |
|   |     |  |
|   |     | Diamond and graphite are two other allotropes of carbon. Suggest what is meant by the term <i>allotrope</i> .  |
|   |     |  |
|   |     | [2]  |
|   | (c) | Nanoparticles are used to deliver drugs within cells. Suggest what property of nanoparticles enables them to be used in this way. Explain your answer.                                 |

| (d) | d) Copper is an important metal that has been used for thousands of years. The proble today is that most of the ores rich in copper compounds have been used up. A centuago ores containing >2% of copper by mass would have been worked; today's minhave to operate at much lower percentages, down to 0.5% of copper by mass. |   |  |
|-----|---|---|--|
|     | (i)   | By what <i>type of reaction</i> is the copper present in the ore converted to copper metal?   |  |
|     | One   | of the main ores of copper contains the mineral <i>chalcopyrite</i> , CuFeS <sub>2</sub> .  |  |
|     | (ii)  | Calculate the percentage of copper by mass in <i>chalcopyrite</i> .   |  |
|     | (iii)   | If the ore contains 2% of <i>chalcopyrite</i> by mass, calculate the mass of copper which can be produced from each tonne of ore.   |  |
|     | (iv)  | Certain bacteria are able to extract copper from the 'spoil' heaps of previously mined copper ore. These bacteria are sprayed onto the spoil heaps in an aqueous solution and the resulting solution containing iron(II) sulfate and copper(II) sulfate is collected in tanks.  Suggest how the copper could be recovered as metal. |  |
|     |   |   |  |
|     |   | [4]<br>[Total: 10]  |  |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.