

6.S091 Problem Set 2

Suyeol Yun
6.S091: Causality

January 24, 2023

Problem 1


Plotting Regression Residuals [1 point]

(a)

$$\hat{\beta}_{12} = 2.0032613606359835$$

(b)

$$\hat{\beta}_{21} = 0.4853818665988653$$



Causal Direction Inference [1 point]

(c)

The plot explains the relationship between ε_1 and ε_2 . Since $\varepsilon_1 \perp\!\!\!\perp \varepsilon_2$, we have to choose the SCM which shows such independence. Therefore, the data is more likely to be generated from SCM M^a .

Problem 2

Partial correlation [2 points]

(a)

$$\hat{\rho}(X_1, X_4, \emptyset) = 0.18515108160562416$$

(b)

$$\hat{\rho}(X_1, X_4, \{X_2, X_3\}) = 0.0093340865561515$$

Fisher's z-transformation [1 point]

(c)

$$\hat{z}(X_1, X_4, \{X_2, X_3\}) = 0.9332023767104407$$

p-values [1 point]

(d)

$$\text{compute_pvalue}(\text{pcalg_samples}, 1, 4, [2, 3]) = 0.35071548783635986$$

Skeleton phase [2 points]

(e)

$$\text{pcalg_skeleton}(\text{samples}[: 500], 0.2) = 10$$

(f)

$$\text{pcalg_skeleton}(\text{samples}[: 500], 0.001) = 10$$

Orientation phase [2 points]

(g)

$X_1 \rightarrow X_3 \leftarrow X_2$ is the only unshielded collider in the output of $\text{pcalg_orient}(\text{estimated_skeleton}, \text{estimated_separator_function})$