Input data

- nurses $n \in \{1, \dots, N\}$, days $d \in \{1, \dots, D\}$, shifts $s \in \{1, \dots, S\}$
- week numbers $w \in \{0, \dots, W\}, W = \lfloor D/7 \rfloor$
- nurses demand per day and shift $demand_{d,s} \in \mathbb{N}$
- workhours lower and upper limit per nurse $minhours_n \in \mathbb{N}$, $maxhours_n \in \mathbb{N}$
- vacation requests $VR \subset N \times D$
- preferred companions $PC \subset N \times N$
- unpreferred companions $UC \subset N \times N$
- preferred slots $PS \subset N \times D \times S$
- unpreferred slots $US \subset N \times D \times S$

Optimization variables: schedule for each nurse, day and shift

$$schedule_{n,d,s} \in \{0,1\}$$

and interactions of nurses (weekends 2)

$$interaction_{n,n',d,s} \in \{0,1\}$$

and "worked in given weekend" indicators

$$weekend_{n,w} \in \{0,1\}$$

and "fraction of contract fulfilled" proportions, as well as lower and upper bound for those

$$\alpha_n = \frac{\frac{24}{S} \sum_{d,s} schedule_{n,d,s}}{workhours_n}, \ \alpha_{min}, \ \alpha_{max}$$

Reward function

$$\begin{split} & \lambda_{PC} \cdot \sum_{(n,n') \in PC} \sum_{d,s} interaction_{n,n',d,s} \\ & - \lambda_{UC} \cdot \sum_{(n,n') \in UC} \sum_{d,s} interaction_{n,n',d,s} \\ & + \lambda_{PS} \cdot \sum_{(n,d,s) \in PS} schedule_{n,d,s} \\ & - \lambda_{US} \cdot \sum_{(n,d,s) \in US} schedule_{n,d,s} \end{split}$$

 $-(\alpha_{max} - \alpha_{min})$ (tight alpha gap iff work distributed proportionately to wokrhours specified)

Constraints

$$\forall_{d,s} \sum_{n} schedule_{n,d,s} = demand_{d,s} \quad (\text{demand is met})$$

$$\forall_{n} \ minhours_{n} \leq \frac{24}{S} \sum_{d,s} schedule_{n,d,s} \leq maxhours_{n} \quad (\text{workhours limits are not exceeded})$$

$$\forall_{n,d} \sum_{s} schedule_{n,d,s} \leq 1 \quad (\text{max 1 shift per day})$$

$$\forall_{n,w} \sum_{d=7w+1}^{\min(7(w+1),D)} schedule_{n,d,S} \leq 6 \quad (\text{max 6 night shifts per week})$$

$$\forall_{n,d} \ schedule_{n,d,s} + schedule_{n,d+1,1} \leq 1 \quad (\text{can't continue past midnight})$$

$$\forall_{(n,d)\in VR} \ \forall_{s} \ schedule_{n,d,s} = 0 \quad (\text{vacations are respected})$$

$$\forall_{n,n',d,s} \ interaction_{n,n',d,s} \leq schedule_{n,d,s} \quad (\text{interactions 1})$$

$$\forall_{n,n',d,s} \ interaction_{n,n',d,s} \leq schedule_{n',d,s} \quad (\text{interactions 2})$$

$$\forall_{n,n',d,s} \ interaction_{n,n',d,s} \geq schedule_{n,d,s} + schedule_{n',d,s} - 1 \quad (\text{interactions 3})$$

$$\forall_{n,w} \ weekend_{n,w} \geq \frac{1}{2S} \sum_{s} schedule_{n,w+6,s} + schedule_{n,w+7,s} \quad (\text{weekends 1})$$

$$\forall_{n,w} \ weekend_{n,w} \leq \sum_{s} schedule_{n,w+6,s} + schedule_{n,w+7,s} \quad (\text{weekends 2})$$

$$\forall_{n,w} \ weekend_{n,w} \leq \alpha_{n} \leq \alpha_{max} \quad (\text{alphas with bounds})$$