

Lagrange Interpolation & The Chinese Remainder Theorem

Tsz-Wo Sze

2025 · 8 · 10

Abstract

In this article, we present an application of Lagrange interpolation to solve the puzzle below. Then, we discuss the similarity between Lagrange interpolation and the Chinese remainder theorem.

$$\begin{aligned} 37\#21 &= 928, \\ 77\#44 &= 3993, \\ 123\#17 &= 14840, \\ 71\#6 &= ? \end{aligned}$$

1 Lagrange Interpolation

Theorem 1.1 (Lagrange Interpolation). *For $1 \leq i \leq n$, let (a_i, b_i) be n distinct pairs of numbers and c_i be n not necessarily distinct numbers. By Lemma 1.2 below, we may translate (a_i, b_i) 's and assume*

$$a_i b_j - a_j b_i \neq 0 \quad \text{for all } j \neq i. \quad (1.1)$$

Define the Lagrange polynomial

$$f(x, y) := \sum_{1 \leq i \leq n} c_i \prod_{1 \leq j \leq n, j \neq i} \frac{xb_j - a_j y}{a_i b_j - a_j b_i}. \quad (1.2)$$

Then, f is a degree d homogeneous polynomial with $d < n$ and $f(a_i, b_i) = c_i$ for all i .

Lemma 1.2. *For $1 \leq i \leq n$, let (a_i, b_i) be n distinct pairs of numbers. Suppose $a_{i_0} b_{j_0} - a_{j_0} b_{i_0} = 0$ for some $j_0 \neq i_0$. By translating a_i 's and b_i 's, let*

$$A_{i,s} := a_i + s \quad \text{and} \quad B_{i,t} := b_i + t.$$

There exists integers $1 \leq s \leq n$ and $1 \leq t \leq n(n-1)/2$ such that $A_{i,s} B_{j,t} - A_{j,s} B_{i,t} \neq 0$ for all $j \neq i$.

Proof. If $a_i \neq 0$ for all i , set $s = 0$. Otherwise, there are at most $n-1$ non-zero a_i 's. Choose $1 \leq s \leq n$ such that $s \neq -a_i$ for $1 \leq i \leq n$. Then, $A_i \neq 0$ for all i . We may assume $a_i \neq 0$ for all i .

If $a_{i_0} = a_{j_0} \neq 0$, then $b_{i_0} = b_{j_0}$ and these two pairs are not distinct. We may assume $a_{i_0} \neq a_{j_0}$. Then,

$$a_{i_0} B_{j_0,t} - a_{j_0} B_{i_0,t} = (a_{i_0} - a_{j_0})t \neq 0 \quad \text{for } 1 \leq t \leq n(n-1)/2.$$

If $a_i B_{j,t_0} - a_j B_{i,t_0} = 0$ for some $(i, j) \neq (i_0, j_0)$, then $a_i B_{j,t_1} - a_j B_{i,t_1} = (a_i - a_j)(t_1 - t_0) \neq 0$ for any $t_1 \neq t_0$. Since there are exactly $n(n-1)/2 - 1$ pairs of i, j other than i_0, j_0 , the lemma follows. $\square$

1.1 Solving The Puzzle

The puzzle mentioned in the abstract has $n = 3$. Using the notation $a_i \# b_i = c_i$, let $a_1 = 37, b_1 = 21, c_1 = 928, a_2 = 77, b_2 = 44, c_2 = 3993, a_3 = 123, b_3 = 17, c_3 = 14840$. Then,

$$\begin{aligned} a_1 b_2 - a_2 b_1 &= 37 \cdot 44 - 77 \cdot 21 &= 11, \\ a_1 b_3 - a_3 b_1 &= 37 \cdot 17 - 123 \cdot 21 &= -1954, \\ a_2 b_3 - a_3 b_2 &= 77 \cdot 17 - 123 \cdot 44 &= -4103; \end{aligned}$$

$$\begin{aligned} x \# y &= \frac{c_1(xb_2 - a_2y)(xb_3 - a_3y)}{(a_1b_2 - a_2b_1)(a_1b_3 - a_3b_1)} + \frac{c_2(xb_1 - a_1y)(xb_3 - a_3y)}{(a_2b_1 - a_1b_2)(a_2b_3 - a_3b_2)} + \frac{c_3(xb_1 - a_1y)(xb_2 - a_2y)}{(a_3b_1 - a_1b_3)(a_3b_2 - a_2b_3)} \\ &= 928 \frac{(44x - 77y)(17x - 123y)}{(11)(-1954)} + 3993 \frac{(21x - 37y)(17x - 123y)}{(-11)(-4103)} + 14840 \frac{(21x - 37y)(44x - 77y)}{(1954)(4103)} \\ &= x^2 - y^2. \end{aligned}$$

Finally, $71\#6 = 71^2 - 6^2 = 5005$.

1.2 The Idea Behind Lagrange Interpolation

The Lagrange polynomial (1.2) looks complicated but the idea behind it actually is simple. Given n distinct pairs (a_i, b_i) and n numbers c_i , construct a homogeneous polynomial f satisfying $f(a_i, b_i) = c_i$. For discussion purpose, set $n = 3$. Let

$$f(x, y) := c_1\delta_1(x, y) + c_2\delta_2(x, y) + c_3\delta_3(x, y),$$

where δ_i 's are homogeneous polynomials with the following property:

$$\delta_i(x, y) := \begin{cases} 1 & , \text{ if } x = a_i \text{ and } y = b_i, \\ 0 & , \text{ if } x = a_j \text{ and } y = b_j \text{ for } j \neq i. \end{cases} \quad (1.3)$$

Then, $f(a_i, b_i) = c_i$ for $i = 1, 2, 3$ as desired. The remaining task is to construct δ_i 's.

Take δ_1 as an example. Since we want to have $\delta_1(a_2, b_2) = 0$ and $\delta_1(a_3, b_3) = 0$, define it as

$$\delta_1(x, y) := g_2(x, y)g_3(x, y)k_1,$$

where $g_i(x, y) := xb_i - a_iy$ and k_1 is a constant. Note that $g_i(a_i, b_i) = 0$ for all i . Then,

$$\delta_1(x, y) = \begin{cases} g_2(a_1, b_1)g_3(a_1, b_1)k_1 & , \text{ if } x = a_1 \text{ and } y = b_1, \\ 0 & , \text{ if } x = a_j \text{ and } y = b_j \text{ for } j \neq 1. \end{cases}$$

By assumption (1.1), we have $g_j(a_i, b_i) \neq 0$ for all $j \neq i$. In order to have $\delta_1(a_1, b_1) = 1$, set $k_1 := k_{1,2}k_{1,3}$, where $k_{i,j} := 1/g_j(a_i, b_i)$ for all $j \neq i$. Therefore,

$$\delta_1(x, y) = g_2(x, y)g_3(x, y)k_{1,2}k_{1,3} = \frac{(xb_2 - a_2y)(xb_3 - a_3y)}{(a_1b_2 - a_2b_1)(a_1b_3 - a_3b_1)}$$

satisfies property (1.3). Of course, δ_2 and δ_3 can be constructed similarly.

2 Chinese Remainder Theorem

Theorem 2.1 (Chinese Remainder Theorem). *Let $a_1, \dots, a_n > 1$ be n pairwise coprime integers. Let $c_1, \dots, c_n$ and m be integers such that*

$$m \equiv c_1 \pmod{a_1}, \quad \dots, \quad m \equiv c_n \pmod{a_n}. \quad (2.1)$$

Then,

$$m \equiv \sum_{1 \leq i \leq n} c_i \prod_{1 \leq j \leq n, j \neq i} a_j k_{i,j} \pmod{A}, \quad (2.2)$$

where $A := \prod_{1 \leq i \leq n} a_i$ and $k_{i,j}$'s are integers such that $k_{i,j} \equiv a_j^{-1} \pmod{a_i}$ for $j \neq i$.

2.1 The Idea Behind The Chinese Remainder Theorem

Interestingly, the Chinese remainder theorem equation (2.2) looks similar to the Lagrange polynomial (1.2). Indeed, the ideas behind them are essentially the same. As before, we set $n = 3$ in the following discussion. Let

$$m := c_1\delta_1 + c_2\delta_2 + c_3\delta_3,$$

where δ_i 's are integers such that

$$\delta_i \bmod a_j \equiv \begin{cases} 1 & , \text{ if } j = i, \\ 0 & , \text{ if } j \neq i. \end{cases} \quad (2.3)$$

Then, m satisfies conditions (2.1) as desired. The remaining task is to construct δ_i 's.

Take δ_1 as an example. Since we want to have $\delta_1 \equiv 0 \pmod{a_2}$ and $\delta_1 \equiv 0 \pmod{a_3}$, define it as

$$\delta_1 := a_2a_3k_1,$$

for some integer k_1 . Then,

$$\delta_1 \bmod a_j \equiv \begin{cases} a_2a_3k_1 & , \text{ if } j = 1, \\ 0 & , \text{ if } j \neq 1. \end{cases}$$

By the assumption of a_i 's being pairwise coprime, the multiplicative inverses of $a_2, a_3 \pmod{a_1}$ exist and they are respectively $k_{1,2}, k_{1,3}$. In order to have $\delta_1 \equiv 1 \pmod{a_1}$, set $k_1 := k_{1,2}k_{1,3}$. Therefore,

$$\delta_1 = a_2a_3k_{1,2}k_{1,3}$$

satisfies property (2.3). Of course, δ_2 and δ_3 can be constructed similarly.