

Partial Differential Equations

Problem Set 1: First order equations and shock waves

- (Conservation) Let $u = u(x, t)$ denote the density per unit length of some substance, which fulfils the conservation law without any sources. Show that, provided the flux q vanishes at infinity and u is x -integrable, the total amount of u in space is constant in time.
- Solve the following problems. Draw their characteristics on the $x - t$ plane and sketch the solution $u(x, t)$ for several times.

$$a) \begin{cases} u_t + 10u_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \frac{1}{1+x^2} & x \in \mathbb{R}, \end{cases}$$

$$b) \begin{cases} u_t + 2tu_x = -u, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = e^{-x^2}, & x \in \mathbb{R}. \end{cases}$$

$$c) \begin{cases} u_t - x^2u_x = \sin u, & t > 0, \quad x > 0; \\ u(x, 0) = 2 \arctan x, & x \geq 0, \end{cases}$$

$$d) \begin{cases} u_t + 2tu_x = xtu, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = x, & x \in \mathbb{R}. \end{cases}$$

- (Signaling problem) Suppose that a lighthouse is located at $x = 0$ and constantly sends its signals with the intensity $\psi(t)$.

(a) As you know, light travels through any medium with a constant speed c . Let $u = u(x, t)$ denote the light intensity at (x, t) . Find $u(x, t)$ for $x > 0$ and $t \in \mathbb{R}$ (lighthouse has been sending its signals for ever).

(b) Solve the above problem but now assume that the light is absorbed by clouds and other aerosols with a rate proportional to its intensity.

- (Initial-boundary value problems) Sometimes we have to solve a problem where both the initial and boundary data is given. Consider the following equations.

$$a) \begin{cases} u_t + c u_x = 0, & t > 0, \quad x > 0, \quad c > 0; \\ u(x, 0) = \phi(x), & x > 0; \\ u(0, t) = \psi(t), & t > 0, \quad \phi(0) = \psi(0) = 0, \end{cases}$$

$$b) \begin{cases} u_t + tu_x = -u^2, & t > 0, \quad x > 0; \\ u(x, 0) = x, & x > 0; \\ u(0, t) = \sin t, & t > 0, \end{cases}$$

What conditions have to be fulfilled for the a) case to have a solution with $c < 0$?

- (Quasi-linear equations) Solve the following problems and sketch the characteristics.

$$a) \begin{cases} u_t + \ln u u_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = e^x, & x \in \mathbb{R}, \end{cases}$$

$$b) \begin{cases} u_t + u_x = u^2, & t \in \mathbb{R}, \quad x > 0; \\ u(0, t) = \psi(t), & t \in \mathbb{R}. \end{cases}$$

$$c) \begin{cases} u_t + u^2 u_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = x, & x \geq 0, \end{cases}$$

$$d) \begin{cases} u_t + u^{-1} u_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \frac{1}{1+x}, & x \in \mathbb{R}. \end{cases}$$

- Solve the following full quasi-nonlinear problem.

$$a) \begin{cases} u_t + uu_x = -u, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = -x, & x \in \mathbb{R}. \end{cases}$$

$$b) \begin{cases} u_t + uu_x = 2t, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = x, & x \in \mathbb{R}. \end{cases}$$

- (Shock waves) Find the solutions of given problems. Draw the characteristics and a shock wave trajectory.

$$a) \begin{cases} u_t + u^2 u_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 2, & x < 0; \\ 1, & 0 > 0, \end{cases} \end{cases}$$

$$b) \begin{cases} u_t + 2uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 3, & x < 0; \\ 2, & x > 0, \end{cases} \end{cases}$$

8. (*Shock fitting*) Analyse the following shock-fitting problems. Draw the solution $u(x, t)$ for several times.

$$\text{a) } \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 1, & x < 0; \\ -1, & 0 < x < 1; \\ 0, & x > 1, \end{cases} \end{cases} \quad \text{b) } \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 1, & x < 0; \\ 1 - x, & 0 < x < 1; \\ 0, & x > 1. \end{cases} \end{cases}$$

9. (*Paint*) The paint flows down the wall and has a thickness $u(x, t)$. As we derived, the governing equation has the form

$$u_t + u^2 u_x = 0, \quad t > 0, \quad x \in \mathbb{R}.$$

Solve the paint flow problem with the given initial profile

$$u(x, 0) = \begin{cases} 0, & x < 0 \text{ or } x > 1; \\ 1, & 0 < x < 1. \end{cases}$$

10. (*Car traffic*) Let $u = u(x, t)$ be a density of cars per unit length on a road at time t . Define the traffic flux $q(x, t)$ as the number of cars per unit time passing through a fixed x at time t .

- (a) Argue that the conservation of cars leads to the equation

$$u_t + q_x = 0.$$

- (b) Define the car velocity as $v = q/u$ (as in the example with convection). A simple model of velocity distribution assumes that there exists a critical density at which cars stop (traffic jam)

$$v(u) = v_m \left(1 - \frac{u}{u_c} \right).$$

From this, find q and plug it in the conservation of cars to obtain the governing equation.

- (c) (*Red light turns green*) Solve the car flow problem for a case where the red light at $x = 0$ turns green at $t = 0$, i.e.

$$u(x, 0) = \begin{cases} u_c, & x < 0; \\ 0, & x > 0. \end{cases}$$

- (d) (*Green light turns red*) Now, a red light suddenly turns on and the traffic with a density u_0 suddenly have to stop at $x = 0$.

$$u(x, 0) = u_0 < u_c, \quad x \in \mathbb{R}, \quad u(0, t) = u_c, \quad t > 0.$$

11. (*Continuation of the discontinuity*) Solve the following equation

$$u_t + uu_x = 0, \quad t > 0, \quad x \in \mathbb{R},$$

for each of the two initial conditions

$$\text{a) } u(x, 0) = \Phi_1(x) := \begin{cases} 1, & x \leq 0; \\ 1 - \frac{x}{\epsilon}, & 0 < x < \epsilon; \\ 0, & x \geq \epsilon, \end{cases} \quad \text{b) } u(x, 0) = \Phi_2(x) := \begin{cases} 0, & x \leq 0; \\ \frac{x}{\epsilon}, & 0 < x < \epsilon; \\ 1, & x \geq \epsilon. \end{cases}$$

In both cases examine the limit $\epsilon \rightarrow 0$.

12. (*Flood hydrograph*) Imagine a long and narrow river. We can introduce a curvilinear variable x which denotes the position measured along its bed. Moreover, let $A = A(x)$ be the wetted cross-section at x . Conservation of mass along with Manning's Law states that

$$A_t + A^m A_x = 0, \quad m > 0.$$

Assume also that $A(x, 0) = A_0(x)$.

- Find a solution of the above equation given in an implicit form.
 - A flood can arise as a result of a sudden rainfall at some point x . To model this situation assume that $A_0(x) = \delta(x)$, where δ is the Dirac delta¹. Draw the characteristics.
 - Find the shock wave. You do not have to solve the equation in order to do that.
 - A flood hydrograph is a graph of the flux at some fixed point x_0 . Hydrologists use this tool a lot. Draw the hydrograph by yourself.
13. (*Blow-up time*) For a general initial condition it is not usually straightforward to determine the shock wave development. It is, however, possible to find the first time of a blow-up. Consider

$$\begin{cases} u_t + c(u)u_x = 0, & t > 0, \quad x \in \mathbb{R}, \\ u(x, 0) = \phi(x) \end{cases},$$

where $c > 0$ and $\phi > 0$ have an opposite monotonicity (say, $c' > 0$ and $\phi' < 0$).

- Pick any characteristic $X = X(t)$ and define $P(t) := u_x(X(t), t)$. Compute P' in terms of u and its derivatives.
- Use the differential equation for u in order to get rid of the second derivatives in the expression for P' .
- Solve the obtained equation for P and whence, for u_x .
- Define $F(\xi) := c(\phi(\xi))$ and show that the first time t_b for which u_x becomes infinite is

$$t_b = \frac{1}{\max_{\xi} |F'(\xi)|}.$$

14. Find the first time of a blow-up for the following problems.

$$\begin{aligned} \text{a)} \quad & \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = e^{-x^2} \end{cases} & \text{b)} \quad & \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \frac{1}{\cosh x^2}. \end{cases} \\ \text{c)} \quad & \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 2 - x^2, & x < 1; \\ 1, & x > 1. \end{cases} \end{cases} & \text{d)} \quad & \begin{cases} u_t + uu_x = 0, & t > 0, \quad x \in \mathbb{R}; \\ u(x, 0) = \begin{cases} 1, & x \leq 0; \\ \cos x, & 0 < x \leq \frac{\pi}{2}; \\ 0, & x > \frac{\pi}{2}. \end{cases} \end{cases} \end{aligned}$$

Łukasz Płociniczak

¹If you do not feel comfortable with distributions, think about some narrow and high spike of unit mass such as $\frac{1}{\epsilon} \chi_{(-\epsilon/2, \epsilon/2)}(x)$ or $\frac{1}{\epsilon} \varphi(x/\epsilon)$, where φ is any integrable function with $\int |\varphi| dx = 1$.