Respiratory Rate Determination by Non-Invasive Means

Stephen Su, 844503, COMP90072

1 Introduction

1.1 Background

(Penzel et al., 2000).


2 The data

2.1 Data source


(Goldberger et al., 2000).

2.2 Sample plot

10-second sample of subject a01



8-hour R-R intervals of subject a01



```
interval <- rr_interval(data$voltage)
range(interval) ## Range of R-R intervals

#> [1] 0.13 1.89
60 / range(interval) ## Range of instantaneous heart rate per minute


#> [1] 461.53846 31.74603
```


abnormal_interval <- abnormal_rr(data\$voltage, max_bpm = 130, min_bpm = 40)
length(abnormal_interval) ## Number of abnormal R-R intervals

#> [1] 58
length(interval) ## Total number of heart beats


#> [1] 29941
invisible(map(seq_len(58), function(i) print(plot_abnormal_rr(data, i))))


Recorded abnormal R-R interval of 0.22s







Recorded abnormal R-R interval of 0.31s







Recorded abnormal R-R interval of 0.31s







Recorded abnormal R-R interval of 1.89s







Recorded abnormal R-R interval of 1.51s







Recorded abnormal R-R interval of 1.55s







Recorded abnormal R-R interval of 1.56s







Recorded abnormal R-R interval of 1.52s







Recorded abnormal R-R interval of 1.54s







Recorded abnormal R-R interval of 1.6s







Recorded abnormal R-R interval of 1.51s







Recorded abnormal R-R interval of 1.51s







Recorded abnormal R-R interval of 1.6s







Recorded abnormal R-R interval of 1.53s







Recorded abnormal R-R interval of 1.64s







Recorded abnormal R-R interval of 1.55s







Recorded abnormal R-R interval of 1.51s







Recorded abnormal R-R interval of 1.53s







Recorded abnormal R-R interval of 1.57s







Recorded abnormal R-R interval of 1.54s







Recorded abnormal R-R interval of 1.52s







Recorded abnormal R-R interval of 1.65s







Recorded abnormal R-R interval of 1.58s







Recorded abnormal R-R interval of 1.63s







Recorded abnormal R-R interval of 1.51s







Recorded abnormal R-R interval of 1.65s







Recorded abnormal R-R interval of 1.71s







Recorded abnormal R-R interval of 1.51s





Recorded abnormal R-R interval of 1.52s





References

- Goldberger, A. L., Amaral, L. A., Glass, L., Hausdorff, J. M., Ivanov, P. C., Mark, R. G., Mietus, J. E., Moody, G. B., Peng, C.-K., & Stanley, H. E. (2000). Physiobank, physiotoolkit, and physionet. *Circulation*, 101(23), e215–e220.
- Penzel, T., Moody, G. B., Mark, R. G., Goldberger, A. L., & Peter, J. H. (2000). The apnea-ecg database. Computers in Cardiology 2000, 255–258.