

Some recent work in proof mining

Thomas Powell
University of Bath

(jww Franziskus Wiesnet)

MINISYMPOSIUM ON PROOF AND COMPUTATION IN MATHEMATICS

DMV Annual Meeting (held online)
14 September 2020

These slides are available at
<https://t-powell.github.io/talks>

Disclaimer...

This is a presentation of work in progress.

Not all definitions, theorems etc. are in their final form.

Contractive mappings

In this talk, X is a Banach space.

Let $E \subseteq X$ and consider a mapping $T : E \rightarrow X$. Then T is contractive if $\forall x, y \in E$:

$$x \neq y \Rightarrow \|Tx - Ty\| < \|x - y\|$$

A *modulus* of contractivity for T (cf. [Kohlenbach and Oliva, 2003]) is a function $\tau : (0, \infty) \rightarrow (0, \infty)$ such that $\forall x, y \in E$:

$$\|x - y\| \geq \varepsilon \Rightarrow \|Tx - Ty\| + \tau(\varepsilon) \leq \|x - y\|$$

Contractive mappings with a modulus generalise certain notions from the literature e.g. *weakly contractive* mappings:

A mapping T is weakly contractive if there exists a continuous and strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that $\forall x, y \in E$:

$$\|Tx - Ty\| \leq \|x - y\| - \psi(\|x - y\|)$$

Contractive mappings

In this talk, X is a Banach space.

Let $E \subseteq X$ and consider a mapping $T : E \rightarrow X$. Then T is contractive if $\forall x, y \in E$:

$$x \neq y \Rightarrow \|Tx - Ty\| < \|x - y\|$$

A *modulus* of contractivity for T (cf. [Kohlenbach and Oliva, 2003]) is a function $\tau : (0, \infty) \rightarrow (0, \infty)$ such that $\forall x, y \in E$:

$$\|x - y\| \geq \varepsilon \Rightarrow \|Tx - Ty\| + \tau(\varepsilon) \leq \|x - y\|$$

Contractive mappings with a modulus generalise certain notions from the literature e.g. *weakly contractive* mappings:

A mapping T is weakly contractive if there exists a continuous and strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that $\forall x, y \in E$:

$$\|Tx - Ty\| \leq \|x - y\| - \psi(\|x - y\|)$$

Contractive mappings

In this talk, X is a Banach space.

Let $E \subseteq X$ and consider a mapping $T : E \rightarrow X$. Then T is contractive if $\forall x, y \in E$:

$$x \neq y \Rightarrow \|Tx - Ty\| < \|x - y\|$$

A *modulus* of contractivity for T (cf. [Kohlenbach and Oliva, 2003]) is a function $\tau : (0, \infty) \rightarrow (0, \infty)$ such that $\forall x, y \in E$:

$$\|x - y\| \geq \varepsilon \Rightarrow \|Tx - Ty\| + \tau(\varepsilon) \leq \|x - y\|$$

Contractive mappings with a modulus generalise certain notions from the literature e.g. *weakly contractive* mappings:

A mapping T is weakly contractive if there exists a continuous and strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that $\forall x, y \in E$:

$$\|Tx - Ty\| \leq \|x - y\| - \psi(\|x - y\|)$$

Contractive mappings

In this talk, X is a Banach space.

Let $E \subseteq X$ and consider a mapping $T : E \rightarrow X$. Then T is contractive if $\forall x, y \in E$:

$$x \neq y \Rightarrow \|Tx - Ty\| < \|x - y\|$$

A *modulus of contractivity* for T (cf. [Kohlenbach and Oliva, 2003]) is a function $\tau : (0, \infty) \rightarrow (0, \infty)$ such that $\forall x, y \in E$:

$$\|x - y\| \geq \varepsilon \Rightarrow \|Tx - Ty\| + \tau(\varepsilon) \leq \|x - y\|$$

Contractive mappings with a modulus generalise certain notions from the literature e.g. *weakly contractive mappings*:

A mapping T is weakly contractive if there exists a continuous and strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that $\forall x, y \in E$:

$$\|Tx - Ty\| \leq \|x - y\| - \psi(\|x - y\|)$$

Contractive mappings

In this talk, X is a Banach space.

Let $E \subseteq X$ and consider a mapping $T : E \rightarrow X$. Then T is contractive if $\forall x, y \in E$:

$$x \neq y \Rightarrow \|Tx - Ty\| < \|x - y\|$$

A *modulus of contractivity* for T (cf. [Kohlenbach and Oliva, 2003]) is a function $\tau : (0, \infty) \rightarrow (0, \infty)$ such that $\forall x, y \in E$:

$$\|x - y\| \geq \varepsilon \Rightarrow \|Tx - Ty\| + \tau(\varepsilon) \leq \|x - y\|$$

Contractive mappings with a modulus generalise certain notions from the literature e.g. *weakly contractive mappings*:

A mapping T is weakly contractive if there exists a continuous and strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that $\forall x, y \in E$:

$$\|Tx - Ty\| \leq \|x - y\| - \psi(\|x - y\|)$$

Convergence theorems for weakly contractive mappings

The convergence properties of weakly contractive mappings are widely studied, and explicit rates of convergence are typically provided. For example

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n - 1))$$

where Ψ denotes the function

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

In this project, we are interested in generalisations of weakly contractive mappings for which

- convergence to a fixpoint is harder to prove
- rates of convergence are not given/known.

Convergence theorems for weakly contractive mappings

The convergence properties of weakly contractive mappings are widely studied, and explicit rates of convergence are typically provided. For example

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n - 1))$$

where Ψ denotes the function

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

In this project, we are interested in generalisations of weakly contractive mappings for which

- convergence to a fixpoint is harder to prove
- rates of convergence are not given/known.

Convergence theorems for weakly contractive mappings

The convergence properties of weakly contractive mappings are widely studied, and explicit rates of convergence are typically provided. For example

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n - 1))$$

where Ψ denotes the function

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

In this project, we are interested in generalisations of weakly contractive mappings for which

- convergence to a fixpoint is harder to prove
- rates of convergence are not given/known.

Example: Total asymptotically weakly contractive mappings

The following definition is due to [Alber et al., 2006].

A mapping $T : E \rightarrow X$ is *total asymptotically weakly contractive* if there exist nonnegative sequences of reals $\{k_n^{(1)}\}$ and $\{k_n^{(2)}\}$ with $k_n^{(1)}, k_n^{(2)} \rightarrow 0$ together with a pair of continuous and strictly increasing functions $\psi, \phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = \psi(0) = 0$ such that $\forall x, y \in E$:

$$\|T^n x - T^n y\| \leq \|x - y\| + k_n^{(1)} \phi(\|x - y\|) - \psi(\|x - y\|) + k_n^{(2)}$$

NOTE. In the case that $k_n^{(1)} = k_n^{(2)} = 0$ we reobtain the class of ordinary weakly contractive mappings.

Cf. Theorem 4.1 of [Alber et al., 2006]

Let $T : E \rightarrow X$ be a total asymptotically weakly contractive mapping with some fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha_n\}$ is some sequence of reals in $(0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Then $\|x_n - q\| \rightarrow 0$.

Example: Total asymptotically weakly contractive mappings

The following definition is due to [Alber et al., 2006].

A mapping $T : E \rightarrow X$ is *total asymptotically weakly contractive* if there exist nonnegative sequences of reals $\{k_n^{(1)}\}$ and $\{k_n^{(2)}\}$ with $k_n^{(1)}, k_n^{(2)} \rightarrow 0$ together with a pair of continuous and strictly increasing functions $\psi, \phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = \psi(0) = 0$ such that $\forall x, y \in E$:

$$\|T^n x - T^n y\| \leq \|x - y\| + k_n^{(1)} \phi(\|x - y\|) - \psi(\|x - y\|) + k_n^{(2)}$$

NOTE. In the case that $k_n^{(1)} = k_n^{(2)} = 0$ we reobtain the class of ordinary weakly contractive mappings.

Cf. Theorem 4.1 of [Alber et al., 2006]

Let $T : E \rightarrow X$ be a total asymptotically weakly contractive mapping with some fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha_n\}$ is some sequence of reals in $(0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Then $\|x_n - q\| \rightarrow 0$.

Example: Total asymptotically weakly contractive mappings

The following definition is due to [Alber et al., 2006].

A mapping $T : E \rightarrow X$ is *total asymptotically weakly contractive* if there exist nonnegative sequences of reals $\{k_n^{(1)}\}$ and $\{k_n^{(2)}\}$ with $k_n^{(1)}, k_n^{(2)} \rightarrow 0$ together with a pair of continuous and strictly increasing functions $\psi, \phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = \psi(0) = 0$ such that $\forall x, y \in E$:

$$\|T^n x - T^n y\| \leq \|x - y\| + k_n^{(1)} \phi(\|x - y\|) - \psi(\|x - y\|) + k_n^{(2)}$$

NOTE. In the case that $k_n^{(1)} = k_n^{(2)} = 0$ we reobtain the class of ordinary weakly contractive mappings.

Cf. Theorem 4.1 of [Alber et al., 2006]

Let $T : E \rightarrow X$ be a total asymptotically weakly contractive mapping with some fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha_n\}$ is some sequence of reals in $(0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Then $\|x_n - q\| \rightarrow 0$.

Example: Total asymptotically weakly contractive mappings

The following definition is due to [Alber et al., 2006].

A mapping $T : E \rightarrow X$ is *total asymptotically weakly contractive* if there exist nonnegative sequences of reals $\{k_n^{(1)}\}$ and $\{k_n^{(2)}\}$ with $k_n^{(1)}, k_n^{(2)} \rightarrow 0$ together with a pair of continuous and strictly increasing functions $\psi, \phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = \psi(0) = 0$ such that $\forall x, y \in E$:

$$\|T^n x - T^n y\| \leq \|x - y\| + k_n^{(1)} \phi(\|x - y\|) - \psi(\|x - y\|) + k_n^{(2)}$$

NOTE. In the case that $k_n^{(1)} = k_n^{(2)} = 0$ we reobtain the class of ordinary weakly contractive mappings.

Cf. Theorem 4.1 of [Alber et al., 2006]

Let $T : E \rightarrow X$ be a total asymptotically weakly contractive mapping with some fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha_n\}$ is some sequence of reals in $(0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Then $\|x_n - q\| \rightarrow 0$.

An abstract formulation of asymptotic contractivity

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \|A_n x - A_n y\| + \tau(\varepsilon) \leq \|x - y\| + \delta$$

EXAMPLE. If $T : E \rightarrow X$ is total asymptotically weakly contractive w.r.t. $\{k_n^{(1)}\}$, $\{k_n^{(2)}\}$, ϕ and ψ and f, g are rates of convergence for $k_n^{(1)} \rightarrow 0$, $k_n^{(2)} \rightarrow 0$ respectively, then $\{T^n\}$ is asymptotically contractive in the above sense with

- $\tau(\varepsilon) := \psi(\varepsilon)$
- $\sigma(\delta, K) := \max \left\{ f \left(\frac{\delta}{2\phi(K)} \right), g \left(\frac{\delta}{2} \right) \right\}$

However, our abstract formulation covers a number of other notions in the literature.

An abstract formulation of asymptotic contractivity

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \|A_n x - A_n y\| + \tau(\varepsilon) \leq \|x - y\| + \delta$$

EXAMPLE. If $T : E \rightarrow X$ is total asymptotically weakly contractive w.r.t. $\{k_n^{(1)}\}$, $\{k_n^{(2)}\}$, ϕ and ψ and f, g are rates of convergence for $k_n^{(1)} \rightarrow 0$, $k_n^{(2)} \rightarrow 0$ respectively, then $\{T^n\}$ is asymptotically contractive in the above sense with

- $\tau(\varepsilon) := \psi(\varepsilon)$
- $\sigma(\delta, K) := \max \left\{ f \left(\frac{\delta}{2\phi(K)} \right), g \left(\frac{\delta}{2} \right) \right\}$

However, our abstract formulation covers a number of other notions in the literature.

An abstract formulation of asymptotic contractivity

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \|A_n x - A_n y\| + \tau(\varepsilon) \leq \|x - y\| + \delta$$

EXAMPLE. If $T : E \rightarrow X$ is total asymptotically weakly contractive w.r.t. $\{k_n^{(1)}\}$, $\{k_n^{(2)}\}$, ϕ and ψ and f, g are rates of convergence for $k_n^{(1)} \rightarrow 0$, $k_n^{(2)} \rightarrow 0$ respectively, then $\{T^n\}$ is asymptotically contractive in the above sense with

- $\tau(\varepsilon) := \psi(\varepsilon)$
- $\sigma(\delta, K) := \max \left\{ f \left(\frac{\delta}{2\phi(K)} \right), g \left(\frac{\delta}{2} \right) \right\}$

However, our abstract formulation covers a number of other notions in the literature.

An abstract formulation of asymptotic contractivity

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \|A_n x - A_n y\| + \tau(\varepsilon) \leq \|x - y\| + \delta$$

EXAMPLE. If $T : E \rightarrow X$ is total asymptotically weakly contractive w.r.t. $\{k_n^{(1)}\}$, $\{k_n^{(2)}\}$, ϕ and ψ and f, g are rates of convergence for $k_n^{(1)} \rightarrow 0$, $k_n^{(2)} \rightarrow 0$ respectively, then $\{T^n\}$ is asymptotically contractive in the above sense with

- $\tau(\varepsilon) := \psi(\varepsilon)$
- $\sigma(\delta, K) := \max \left\{ f \left(\frac{\delta}{2\phi(K)} \right), g \left(\frac{\delta}{2} \right) \right\}$

However, our abstract formulation covers a number of other notions in the literature.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

How to establish convergence of asymptotically contractive mappings

Take the following simple version of asymptotic contractivity:

$$\|T^n x - T^n y\| \leq \|x - y\| - \psi(\|x - y\|) + k_n$$

where ψ satisfies the usual properties and $k_n \rightarrow 0$. Suppose that $Tq = q$ and that

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

for $\sum \alpha_n = \infty$. We now prove that $\|x_n - q\| \rightarrow 0$.

STEP 1. We observe that

$$\begin{aligned}\|x_{n+1} - q\| &= \|(1 - \alpha_n)x_n + \alpha_n T^n x_n - q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n \|T^n x_n - T^n q\| \\ &\leq (1 - \alpha_n) \|x_n - q\| + \alpha_n (\|x_n - q\| - \psi(\|x_n - q\|) + k_n) \\ &= \|x_n - q\| - \alpha_n \psi(\|x_n - q\|) + \alpha_n k_n\end{aligned}$$

STEP 2. Prove more generally that whenever some sequence $\{\mu_n\}$ of nonnegative reals satisfies the recurrence

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n k_n$$

where $k_n \rightarrow 0$, then also $\mu_n \rightarrow 0$.

A key combinatorial lemma

The following is similar to Lemma 3.4 of [Kohlenbach and Powell, 2020], but there are nevertheless a number of important differences.

Theorem

Let $\{\mu_n\}$ be a sequence of nonnegative reals bounded above by L . Suppose that $\{\alpha_n\}$ is a sequence in $(0, \alpha]$ and $r : \mathbb{N} \times (0, \infty) \rightarrow \mathbb{N}$ a function satisfying

$$\sum_{n=k}^{r(k,x)} \alpha_n > x \quad \text{for all } k \in \mathbb{N}, \quad x \in (0, \infty).$$

Finally, let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a monotone function with $\psi(0) = 0$ and $\psi(t) > 0$ for $t > 0$ and suppose there exists some $N : (0, \infty) \rightarrow \mathbb{N}$ such that for any $\delta > 0$ we have

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n \delta$$

for all $n \geq N(\delta)$.

Then $\mu_n \rightarrow 0$ with rate of convergence

$$\Phi_{L,\alpha,r,\psi,N}(\varepsilon) := r \left(N \left(\min \left\{ \psi \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\} \right), 2 \int_{\varepsilon/2}^L \frac{dt}{\psi(t)} \right)$$

A key combinatorial lemma

The following is similar to Lemma 3.4 of [Kohlenbach and Powell, 2020], but there are nevertheless a number of important differences.

Theorem

Let $\{\mu_n\}$ be a sequence of nonnegative reals bounded above by L . Suppose that $\{\alpha_n\}$ is a sequence in $(0, \alpha]$ and $r : \mathbb{N} \times (0, \infty) \rightarrow \mathbb{N}$ a function satisfying

$$\sum_{n=k}^{r(k,x)} \alpha_n > x \quad \text{for all } k \in \mathbb{N}, \quad x \in (0, \infty).$$

Finally, let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a monotone function with $\psi(0) = 0$ and $\psi(t) > 0$ for $t > 0$ and suppose there exists some $N : (0, \infty) \rightarrow \mathbb{N}$ such that for any $\delta > 0$ we have

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n \delta$$

for all $n \geq N(\delta)$.

Then $\mu_n \rightarrow 0$ with rate of convergence

$$\Phi_{L,\alpha,r,\psi,N}(\varepsilon) := r \left(N \left(\min \left\{ \psi \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\} \right), 2 \int_{\varepsilon/2}^L \frac{dt}{\psi(t)} \right)$$

A key combinatorial lemma

The following is similar to Lemma 3.4 of [Kohlenbach and Powell, 2020], but there are nevertheless a number of important differences.

Theorem

Let $\{\mu_n\}$ be a sequence of nonnegative reals bounded above by L . Suppose that $\{\alpha_n\}$ is a sequence in $(0, \alpha]$ and $r : \mathbb{N} \times (0, \infty) \rightarrow \mathbb{N}$ a function satisfying

$$\sum_{n=k}^{r(k,x)} \alpha_n > x \quad \text{for all } k \in \mathbb{N}, \quad x \in (0, \infty).$$

Finally, let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a monotone function with $\psi(0) = 0$ and $\psi(t) > 0$ for $t > 0$ and suppose there exists some $N : (0, \infty) \rightarrow \mathbb{N}$ such that for any $\delta > 0$ we have

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n \delta$$

for all $n \geq N(\delta)$.

Then $\mu_n \rightarrow 0$ with rate of convergence

$$\Phi_{L,\alpha,r,\psi,N}(\varepsilon) := r \left(N \left(\min \left\{ \psi \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\} \right), 2 \int_{\varepsilon/2}^L \frac{dt}{\psi(t)} \right)$$

A key combinatorial lemma

The following is similar to Lemma 3.4 of [Kohlenbach and Powell, 2020], but there are nevertheless a number of important differences.

Theorem

Let $\{\mu_n\}$ be a sequence of nonnegative reals bounded above by L . Suppose that $\{\alpha_n\}$ is a sequence in $(0, \alpha]$ and $r : \mathbb{N} \times (0, \infty) \rightarrow \mathbb{N}$ a function satisfying

$$\sum_{n=k}^{r(k,x)} \alpha_n > x \quad \text{for all } k \in \mathbb{N}, \quad x \in (0, \infty).$$

Finally, let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a monotone function with $\psi(0) = 0$ and $\psi(t) > 0$ for $t > 0$ and suppose there exists some $N : (0, \infty) \rightarrow \mathbb{N}$ such that for any $\delta > 0$ we have

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n) + \alpha_n \delta$$

for all $n \geq N(\delta)$.

Then $\mu_n \rightarrow 0$ with rate of convergence

$$\Phi_{L,\alpha,r,\psi,N}(\varepsilon) := r \left(N \left(\min \left\{ \psi \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\} \right), 2 \int_{\varepsilon/2}^L \frac{dt}{\psi(t)} \right)$$

A special case

Consider the simple case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

for all $n \in \mathbb{N}$, which is all that is usually required for ordinary weakly contractive mappings.

A rate of convergence in this case is just

$$\Phi_{L,r,\psi}(\varepsilon) := r \left(0, 2 \int_{\varepsilon/2}^{\mu_0} \frac{dt}{\psi(t)} \right)$$

Let us now try to find some function $f : \mathbb{N} \rightarrow (0, \infty)$ such that $n \in \mathbb{N}$ we have

$$\Phi(f(n)) = n \quad \text{and thus} \quad \mu_n \leq f(n).$$

This f would form a rate of convergence from the analyst's - as opposed to proof theorists' - point of view.

A special case

Consider the simple case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

for all $n \in \mathbb{N}$, which is all that is usually required for ordinary weakly contractive mappings.

A rate of convergence in this case is just

$$\Phi_{L,r,\psi}(\varepsilon) := r \left(0, 2 \int_{\varepsilon/2}^{\mu_0} \frac{dt}{\psi(t)} \right)$$

Let us now try to find some function $f : \mathbb{N} \rightarrow (0, \infty)$ such that $n \in \mathbb{N}$ we have

$$\Phi(f(n)) = n \quad \text{and thus} \quad \mu_n \leq f(n).$$

This f would form a rate of convergence from the analyst's - as opposed to proof theorists' - point of view.

A special case

Consider the simple case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

for all $n \in \mathbb{N}$, which is all that is usually required for ordinary weakly contractive mappings.

A rate of convergence in this case is just

$$\Phi_{L,r,\psi}(\varepsilon) := r \left(0, 2 \int_{\varepsilon/2}^{\mu_0} \frac{dt}{\psi(t)} \right)$$

Let us now try to find some function $f : \mathbb{N} \rightarrow (0, \infty)$ such that $n \in \mathbb{N}$ we have

$$\Phi(f(n)) = n \quad \text{and thus} \quad \mu_n \leq f(n).$$

This f would form a rate of convergence from the analyst's - as opposed to proof theorists' - point of view.

A special case

Consider the simple case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

for all $n \in \mathbb{N}$, which is all that is usually required for ordinary weakly contractive mappings.

A rate of convergence in this case is just

$$\Phi_{L,r,\psi}(\varepsilon) := r \left(0, 2 \int_{\varepsilon/2}^{\mu_0} \frac{dt}{\psi(t)} \right)$$

Let us now try to find some function $f : \mathbb{N} \rightarrow (0, \infty)$ such that $n \in \mathbb{N}$ we have

$$\Phi(f(n)) = n \quad \text{and thus} \quad \mu_n \leq f(n).$$

This f would form a rate of convergence from the analyst's - as opposed to proof theorists' - point of view.

Reformulating the rate of convergence

Assuming that $r(k, x)$ returns the least n with $\sum_{i=k}^n \alpha_i > x$, then $\Phi(f(n)) = n$ implies that

$$\sum_{i=0}^{n-1} \alpha_i \leq 2 \int_{f(n)/2}^{\mu_0} \frac{dt}{\psi(t)} < \sum_{i=0}^n \alpha_i$$

and so in particular

$$\frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \leq \Psi(\mu_0) - \Psi(f(n)/2)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Therefore

$$f(n) \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

Reformulating the rate of convergence

Assuming that $r(k, x)$ returns the least n with $\sum_{i=k}^n \alpha_i > x$, then $\Phi(f(n)) = n$ implies that

$$\sum_{i=0}^{n-1} \alpha_i \leq 2 \int_{f(n)/2}^{\mu_0} \frac{dt}{\psi(t)} < \sum_{i=0}^n \alpha_i$$

and so in particular

$$\frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \leq \Psi(\mu_0) - \Psi(f(n)/2)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Therefore

$$f(n) \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

Reformulating the rate of convergence

Assuming that $r(k, x)$ returns the least n with $\sum_{i=k}^n \alpha_i > x$, then $\Phi(f(n)) = n$ implies that

$$\sum_{i=0}^{n-1} \alpha_i \leq 2 \int_{f(n)/2}^{\mu_0} \frac{dt}{\psi(t)} < \sum_{i=0}^n \alpha_i$$

and so in particular

$$\frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \leq \Psi(\mu_0) - \Psi(f(n)/2)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Therefore

$$f(n) \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

Reformulating the rate of convergence

Assuming that $r(k, x)$ returns the least n with $\sum_{i=k}^n \alpha_i > x$, then $\Phi(f(n)) = n$ implies that

$$\sum_{i=0}^{n-1} \alpha_i \leq 2 \int_{f(n)/2}^{\mu_0} \frac{dt}{\psi(t)} < \sum_{i=0}^n \alpha_i$$

and so in particular

$$\frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \leq \Psi(\mu_0) - \Psi(f(n)/2)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Therefore

$$f(n) \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

A direct comparison with known bounds

In other words, in the case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

we have $\mu_n \rightarrow 0$ where

$$\mu_n \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Compare with:

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n-1))$$

where we define

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

A direct comparison with known bounds

In other words, in the case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

we have $\mu_n \rightarrow 0$ where

$$\mu_n \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Compare with:

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n-1))$$

where we define

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

A direct comparison with known bounds

In other words, in the case where

$$\mu_{n+1} \leq \mu_n - \alpha_n \psi(\mu_n)$$

we have $\mu_n \rightarrow 0$ where

$$\mu_n \leq 2\Psi^{-1} \left(\Psi(\mu_0) - \frac{1}{2} \sum_{i=0}^{n-1} \alpha_i \right)$$

for $\Psi(s) := \int^s \frac{dt}{\psi(t)}$.

Compare with:

cf. Theorem 3.1 of [Alber and Guerre-Delabriere, 1997]

Let $T : E \rightarrow X$ be a weakly contractive mapping w.r.t ψ , with a fixed point $q \in E$. Let the sequence $\{x_n\}$ satisfy $x_{n+1} = Tx_n$. Then $\|x_n - q\| \rightarrow 0$, and moreover we have the estimate

$$\|x_n - q\| \leq \Psi^{-1}(\Psi(\|x_0 - q\|) - (n - 1))$$

where we define

$$\Psi(s) = \int^s \frac{dt}{\psi(t)}$$

A new quantitative result: Example 1 (apologies in advance...)

Theorem

Suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$ is bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi_{\tau, \sigma, \alpha, r, g, L}(\varepsilon) := r \left(\max \{ \sigma(\delta, K), g(\delta) \}, \frac{2L}{\tau(\varepsilon/2)} \right)$$

where δ is given by

$$\delta := \frac{1}{4} \min \left\{ \tau \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\}$$

A new quantitative result: Example 1 (apologies in advance...)

Theorem

Suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$ is bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi_{\tau, \sigma, \alpha, r, g, L}(\varepsilon) := r \left(\max \{ \sigma(\delta, K), g(\delta) \}, \frac{2L}{\tau(\varepsilon/2)} \right)$$

where δ is given by

$$\delta := \frac{1}{4} \min \left\{ \tau \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\}$$

A new quantitative result: Example 1 (apologies in advance...)

Theorem

Suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$ is bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi_{\tau, \sigma, \alpha, r, g, L}(\varepsilon) := r \left(\max \{ \sigma(\delta, K), g(\delta) \}, \frac{2L}{\tau(\varepsilon/2)} \right)$$

where δ is given by

$$\delta := \frac{1}{4} \min \left\{ \tau \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\}$$

A new quantitative result: Example 1 (apologies in advance...)

Theorem

Suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$ is bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi_{\tau, \sigma, \alpha, r, g, L}(\varepsilon) := r \left(\max \{ \sigma(\delta, K), g(\delta) \}, \frac{2L}{\tau(\varepsilon/2)} \right)$$

where δ is given by

$$\delta := \frac{1}{4} \min \left\{ \tau \left(\frac{\varepsilon}{2} \right), \frac{\varepsilon}{\alpha} \right\}$$

Further results

There are many related notions of asymptotic contractivity which can be approached in a similar way.

Let X be a uniformly smooth space, and $J : X \rightarrow X^*$ the single valued normalized duality mapping satisfying

$$\langle x, Jx \rangle = \|x\|^2 = \|Jx\|^2$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing operation.

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \langle A_n x - A_n y, J(x - y) \rangle + \tau(\varepsilon) \leq \|x - y\| + \delta$$

This generalises the notion of a d -weakly contractive mapping, studied in e.g. [Chidume et al., 2002].

Further results

There are many related notions of asymptotic contractivity which can be approached in a similar way.

Let X be a uniformly smooth space, and $J : X \rightarrow X^*$ the single valued normalized duality mapping satisfying

$$\langle x, Jx \rangle = \|x\|^2 = \|Jx\|^2$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing operation.

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \langle A_n x - A_n y, J(x - y) \rangle + \tau(\varepsilon) \leq \|x - y\| + \delta$$

This generalises the notion of a d -weakly contractive mapping, studied in e.g. [Chidume et al., 2002].

Further results

There are many related notions of asymptotic contractivity which can be approached in a similar way.

Let X be a uniformly smooth space, and $J : X \rightarrow X^*$ the single valued normalized duality mapping satisfying

$$\langle x, Jx \rangle = \|x\|^2 = \|Jx\|^2$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing operation.

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \langle A_n x - A_n y, J(x - y) \rangle + \tau(\varepsilon) \leq \|x - y\| + \delta$$

This generalises the notion of a d -weakly contractive mapping, studied in e.g. [Chidume et al., 2002].

Further results

There are many related notions of asymptotic contractivity which can be approached in a similar way.

Let X be a uniformly smooth space, and $J : X \rightarrow X^*$ the single valued normalized duality mapping satisfying

$$\langle x, Jx \rangle = \|x\|^2 = \|Jx\|^2$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing operation.

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \langle A_n x - A_n y, J(x - y) \rangle + \tau(\varepsilon) \leq \|x - y\| + \delta$$

This generalises the notion of a d -weakly contractive mapping, studied in e.g. [Chidume et al., 2002].

Further results

There are many related notions of asymptotic contractivity which can be approached in a similar way.

Let X be a uniformly smooth space, and $J : X \rightarrow X^*$ the single valued normalized duality mapping satisfying

$$\langle x, Jx \rangle = \|x\|^2 = \|Jx\|^2$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing operation.

We say that a sequence $\{A_n\}$ of mappings with $A_n : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$ if for all $x, y \in E$ and any $\varepsilon, \delta, K > 0$ we have

$$\|x - y\| \in [\varepsilon, K] \wedge n \geq \sigma(\delta, K) \Rightarrow \langle A_n x - A_n y, J(x - y) \rangle + \tau(\varepsilon) \leq \|x - y\| + \delta$$

This generalises the notion of a d -weakly contractive mapping, studied in e.g. [Chidume et al., 2002].

A new quantitative result: Example 2 (even more apologies in advance...)

Theorem

Let X be uniformly smooth, and suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r and $\alpha_n \rightarrow 0$ with rate of convergence f . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$, $\|q\|$ and $\{\|x_n - A_n x_n\|\}$ are all bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi(\varepsilon) := r \left(\max \left\{ \sigma \left(\frac{\delta}{6}, K \right), f \left(\frac{\delta}{6} \right), g \left(\frac{\omega_X((2 + \alpha)L, \delta/6L)}{L} \right) \right\}, \frac{2L}{\tau(\varepsilon/\sqrt{2})} \right)$$

where δ is given by

$$\delta := \frac{1}{2} \min \left\{ \tau \left(\frac{\varepsilon}{\sqrt{2}} \right), \frac{\varepsilon^2}{\alpha} \right\}$$

and ω_X is a modulus of uniform continuity on bounded subsets for J .

A new quantitative result: Example 2 (even more apologies in advance...)

Theorem

Let X be uniformly smooth, and suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r and $\alpha_n \rightarrow 0$ with rate of convergence f . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$, $\|q\|$ and $\{\|x_n - A_n x_n\|\}$ are all bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi(\varepsilon) := r \left(\max \left\{ \sigma \left(\frac{\delta}{6}, K \right), f \left(\frac{\delta}{6} \right), g \left(\frac{\omega_X((2 + \alpha)L, \delta/6L)}{L} \right) \right\}, \frac{2L}{\tau(\varepsilon/\sqrt{2})} \right)$$

where δ is given by

$$\delta := \frac{1}{2} \min \left\{ \tau \left(\frac{\varepsilon}{\sqrt{2}} \right), \frac{\varepsilon^2}{\alpha} \right\}$$

and ω_X is a modulus of uniform continuity on bounded subsets for J .

A new quantitative result: Example 2 (even more apologies in advance...)

Theorem

Let X be uniformly smooth, and suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r and $\alpha_n \rightarrow 0$ with rate of convergence f . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$, $\|q\|$ and $\{\|x_n - A_n x_n\|\}$ are all bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi(\varepsilon) := r \left(\max \left\{ \sigma \left(\frac{\delta}{6}, K \right), f \left(\frac{\delta}{6} \right), g \left(\frac{\omega_X((2 + \alpha)L, \delta/6L)}{L} \right) \right\}, \frac{2L}{\tau(\varepsilon/\sqrt{2})} \right)$$

where δ is given by

$$\delta := \frac{1}{2} \min \left\{ \tau \left(\frac{\varepsilon}{\sqrt{2}} \right), \frac{\varepsilon^2}{\alpha} \right\}$$

and ω_X is a modulus of uniform continuity on bounded subsets for J .

A new quantitative result: Example 2 (even more apologies in advance...)

Theorem

Let X be uniformly smooth, and suppose that the sequence of mappings $\{A_n\} : E \rightarrow X$ is asymptotically d -contractive with moduli $\tau : (0, \infty) \rightarrow (0, \infty)$ and $\sigma : (0, \infty) \rightarrow \mathbb{N}$. Let $\{x_n\}$ be some sequence satisfying

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n$$

where $\{\alpha\}$ is a sequence of reals in $(0, \alpha]$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$ with rate of divergence r and $\alpha_n \rightarrow 0$ with rate of convergence f . Finally, suppose that $q \in E$ satisfies

$$\|A_n q - q\| \rightarrow 0$$

with rate of convergence g , and that $\{\|x_n - q\|\}$, $\|q\|$ and $\{\|x_n - A_n x_n\|\}$ are all bounded above by L .

Then $\|x_n - q\| \rightarrow 0$ with rate of convergence

$$\Phi(\varepsilon) := r \left(\max \left\{ \sigma \left(\frac{\delta}{6}, K \right), f \left(\frac{\delta}{6} \right), g \left(\frac{\omega_X((2 + \alpha)L, \delta/6L)}{L} \right) \right\}, \frac{2L}{\tau(\varepsilon/\sqrt{2})} \right)$$

where δ is given by

$$\delta := \frac{1}{2} \min \left\{ \tau \left(\frac{\varepsilon}{\sqrt{2}} \right), \frac{\varepsilon^2}{\alpha} \right\}$$

and ω_X is a modulus of uniform continuity on bounded subsets for J .

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Conclusion

- There are many interesting classes of weakly contractive mappings, with corresponding convergence results.
- In simple cases, explicit rates of convergence are given.
- We consider abstract formulations of mappings of weakly contractive type, and provide general rates of convergence (which in simple cases match up with those in the literature).
- This is work in progress - by the end we will hopefully have several metatheorems which in each case generalise and unify a class of existing convergence theorems.

THANK YOU!

Alber, Y., Chidume, C., and Zegeye, H. (2006).

Approximating fixed points of total asymptotically nonexpansive mappings.

Fixed Point Theory and Applications, 2006:1687–1812.

Alber, Y. and Guerre-Delabriere, S. (1997).

Principle of weakly contractive maps in Hilbert spaces.

In Gohberg, I. and Lyubich, Y., editors, *New Results in Operator Theory and its Applications*, volume 98, pages 7–22.

Chidume, C., Zegeye, H., and Aneke, S. J. (2002).

Approximation of fixed points of weakly contractive nonself maps in Banach spaces.

Journal of Mathematical Analysis and Applications, 270:189–199.

Kohlenbach, U. and Oliva, P. (2003).

Proof mining: a systematic way of analyzing proofs in mathematics.

Proceedings of the Steklov Institute of Mathematics, 242:136–164.

Kohlenbach, U. and Powell, T. (2020).

Rates of convergence for iterative solutions of equations involving set-valued accretive operators.

Computers and Mathematics with Applications, 80:490–503.