GROUP 1 | Project Cycle 2: One-Page Summary

YRBS_2007.csv • landscape PDF

Research Questions

1) Is the smoking proportion different from 0.20?
2) Is the mean BMIPCT different from 65.0?

Data and Variables + Method

Behavior: CurrentCigaretteUse
Continuous: BMIPCT
One-sample z/t tests with 95% CI

EDA + Extension

Four figures included: smoking bar chart, BMIPCT histogram, age vs smoking, and BMIPCT by sleep code

Proportion Analysis: CurrentCigaretteUse



Key Results

n = 13,323
$\hat{p}$ = 0.1943
95% CI = [0.1876, 0.2010]
p = 0.1015

Mean Analysis: BMIPCT



Key Results

n = 13,062
$\bar{x}$ = 64.8207
95% CI = [64.3487, 65.2926]
p = 0.4564

Extension: smoking rate by age code



EDA Highlight

From age code 3 to 7,
smoking rate increases
from 0.10 to 0.27.

Extension: BMIPCT by sleep code



EDA Highlight

Group means vary, but boxplots overlap heavily across sleep codes.

Decision: Fail to reject H_0 for both analyses. The smoking proportion is not significantly different from 0.20, and the mean BMIPCT is not significantly different from 65.0.