Causal dimension reduction for environmental health



Wake Forest University BDS Faculty Candidate Talk

Thomas Hsiao, Ph.D.
Postdoctoral Fellow
November 4, 2026

Emory University
Rollins School of Public Health
Department of Biostatistics and Bioinformatics

Environmental mixtures analysis

How can we quantify the causal effect of complex environmental exposures?

- Air pollution mixtures ($PM_{2.5}$, NO_2, O_3, ...)
- PFAS compounds (PFOA, PFOS, PFNA, ...)
- Metals (Pb, Cd, As, Hg, ...)

All involve multiple, highly correlated, continuous exposures



EDITORIAL • Environ Health Perspect. 2013 Jan 1;121(1):a6–a8. doi: 10.1289/ehp.1206182

Unraveling the Health Effects of Environmental Mixtures: An NIEHS Priority

Danielle J Carlin [1], Cynthia V Rider [1], Rick Woychik [1], Linda S Birnbaum [1]

• Author information • Article notes • Copyright and License information

PMCID: PMC3553446 PMID: 23409283

Traditionally, toxicological studies and human health risk assessments have focused primarily on single chemicals. However, people are exposed to a myriad of chemical and nonchemical stressors every day and throughout their lifetime. Some recent events that highlight the need to understand these complex exposures and their role in the etiology of disease include the *Deepwater Horizon* oil spill, the earthquake in Japan and subsequent Fukushima nuclear disaster, and the unknown environmental and health effects of hydraulic

Standard setup of the observed data

Observed data: n-vectors $O_1, ..., O_n$ sampled i.i.d. where

$$O = (Y, X, C) \sim P \in \mathcal{M}$$

- $Y \in \mathbb{R}$: outcome (binary, count, or continuous)
- $X \in \mathbb{R}^p$: continuous exposure mixture
- $C \in \mathbb{R}^q$: measured baseline covariates

Standard setup of the observed data

Observed data: n-vectors $O_1, ..., O_n$ sampled i.i.d. where

$$O = (Y, X, C) \sim P \in \mathcal{M}$$

- $Y \in \mathbb{R}$: outcome (binary, count, or continuous)
- $X \in \mathbb{R}^p$: continuous exposure mixture
- $C \in \mathbb{R}^q$: measured baseline covariates

Question: What is the value of Y had we intervened and set $X = x$?

How are we currently answering this question?

Let Y^x be the potential outcome of Y after setting $X = x$.

We are interested in estimating $\mathbb{E}(Y^x)$, the **causal average dose-response**.

How are we currently answering this question?

Let Y^x be the potential outcome of Y after setting $X = x$.

We are interested in estimating $\mathbb{E}(Y^x)$, the **causal average dose-response**.

Current methods are:

- **BKMR**: Models $\mathbb{E}(Y \mid X, C) = h(X) + \gamma^\top C$ using Gaussian Process regression.

How are we currently answering this question?

Let Y^x be the potential outcome of Y after setting $X = x$.

We are interested in estimating $\mathbb{E}(Y^x)$, the **causal average dose-response**.

Current methods are:

- **BKMR**: Models $\mathbb{E}(Y \mid X, C) = h(X) + \gamma^\top C$ using Gaussian Process regression.
- **PCP**: Models $\mathbb{E}(Y \mid \beta^\top X, C) = h(\beta^\top X) + \gamma^\top C$, where β is obtained from PCA.

How are we currently answering this question?

Let Y^x be the potential outcome of Y after setting $X = x$.

We are interested in estimating $\mathbb{E}(Y^x)$, the **causal average dose-response**.

Current methods are:

- **BKMR**: Models $\mathbb{E}(Y \mid X, C) = h(X) + \gamma^\top C$ using Gaussian Process regression.
- **PCP**: Models $\mathbb{E}(Y \mid \beta^\top X, C) = h(\beta^\top X) + \gamma^\top C$, where β is obtained from PCA.
- **qgcomp**: Estimates $\mathbb{E}(Y^{a+1}) - \mathbb{E}(Y^a)$, where $A \in \mathbb{R}$ is a scalar joint quantile increase in all exposures in X.

How are we currently answering this question?

Let Y^x be the potential outcome of Y after setting $X = x$.

We are interested in estimating $\mathbb{E}(Y^x)$, the **causal average dose-response**.

Current methods are:

- **BKMR**: Models $\mathbb{E}(Y \mid X, C) = h(X) + \gamma^\top C$ using Gaussian Process regression.
- **PCP**: Models $\mathbb{E}(Y \mid \beta^\top X, C) = h(\beta^\top X) + \gamma^\top C$, where β is obtained from PCA.
- **qgcomp**: Estimates $\mathbb{E}(Y^{a+1}) - \mathbb{E}(Y^a)$, where $A \in \mathbb{R}$ is a scalar joint quantile increase in all exposures in X.

Problem: Each (Bobb et al. 2015; Gibson et al. 2022; Keil et al. 2020) has their pros and cons, but none of these are estimating $\mathbb{E}(Y^x)$.

Causal inference for continuous exposures

Flexible, nonparametric, doubly robust estimators of $\mu(x) := \mathbb{E}(Y^x)$ are available.

Let $m(x, c) := \mathbb{E}(Y \mid X = x, C = c)$ and $\pi(x \mid c) := f_{X|C}(x \mid c)$.

Causal inference for continuous exposures

Flexible, nonparametric, doubly robust estimators of $\mu(x) := \mathbb{E}(Y^x)$ are available.

Let $m(x, c) := \mathbb{E}(Y \mid X = x, C = c)$ and $\pi(x \mid c) := f_{X|C}(x \mid c)$.

A kernel-based doubly robust estimator (Colangelo and Lee 2026) is

$$\hat{\mu}_{DR}(x) = \frac{1}{n} \sum_{i=1}^{n} \left\{ \frac{K_h(X_i - x)}{\hat{\pi}(X_i \mid C_i)} [Y_i - \widehat{m}(X_i, C_i)] + \widehat{m}(x, C_i) \right\},$$

where $K_h(u) = h^{-p} K(u/h)$, and $\widehat{m}$ and $\hat{\pi}$ are estimated flexibly using machine learning (e.g., neural network, random forest, or ensembles).

What can go wrong with multiple exposures?

The doubly robust estimator only works well if we do a good job on $m(x, c)$ and $\pi(x \mid c)$.

$$\sqrt{nh^p} \cdot \|\widehat{m} - m\| \cdot \|\hat{\pi} - \pi\| \xrightarrow{p} 0 \implies \sqrt{nh^p}\,\{\hat{\mu}_{\mathrm{DR}}(x) - \mu(x)\} \rightsquigarrow \mathcal{N}(0, V_x),$$

What can go wrong with multiple exposures?

The doubly robust estimator only works well if we do a good job on $m(x, c)$ and $\pi(x \mid c)$.

$$\sqrt{nh^p} \cdot \|\widehat{m} - m\| \cdot \|\hat{\pi} - \pi\| \xrightarrow{p} 0 \implies \sqrt{nh^p}\,\{\hat{\mu}_{\mathrm{DR}}(x) - \mu(x)\} \rightsquigarrow \mathcal{N}(0, V_x),$$

For large p, fitting a $(p+q)$-regression and a p-variate density with q regressors is hard.

What can go wrong with multiple exposures?

The doubly robust estimator only works well if we do a good job on $m(x, c)$ and $\pi(x \mid c)$.

$$\sqrt{nh^p} \cdot \|\widehat{m} - m\| \cdot \|\hat{\pi} - \pi\| \xrightarrow{p} 0 \implies \sqrt{nh^p}\,\{\hat{\mu}_{\mathrm{DR}}(x) - \mu(x)\} \rightsquigarrow \mathcal{N}(0, V_x),$$

For large p, fitting a $(p+q)$-regression and a p-variate density with q regressors is hard.

Solution: Consider estimating the average dose-response in a lower dimensional space.

$$\boxed{\mathbb{E}(Y^x) = g(\beta^\top x), \qquad \beta \in \mathbb{R}^{p \times d}}$$

Sufficient dimension reduction (SDR)

There exists a rich literature on estimating β that preserves the *regression* relationship (Cook 2026):

$$\mathbb{E}(Y \mid X) = g(\beta^\top X).$$

Since β is not unique, the statistical estimand is the **central mean subspace**, $\mathcal{S}_{\mathbb{E}(Y|X)}$, the smallest subspace satisfying this identity.

Sufficient dimension reduction (SDR)

There exists a rich literature on estimating β that preserves the *regression* relationship (Cook 2026):

$$\mathbb{E}(Y \mid X) = g(\beta^\top X).$$

Since β is not unique, the statistical estimand is the **central mean subspace**, $\mathcal{S}_{\mathbb{E}(Y|X)}$, the smallest subspace satisfying this identity.

Problem: Because $\mathbb{E}(Y^x) \neq \mathbb{E}(Y \mid X = x)$ due to confounding, we cannot directly use regression SDR for our causal dose-response.

Ideal workflow of causal sufficient dimension reduction


Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).

Ideal workflow of causal sufficient dimension reduction

Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).

1. Exposures (p)

$X \in \mathbb{R}^p$




Ideal workflow of causal sufficient dimension reduction

Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).


1. Exposures (p)

$X \in \mathbb{R}^p$




2. Causal directions

$\beta \in \mathbb{R}^{p \times d}$



Ideal workflow of causal sufficient dimension reduction

Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).




Ideal workflow of causal sufficient dimension reduction

Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).


1. Exposures (p)

$X \in \mathbb{R}^p$




2. Causal directions

$\beta \in \mathbb{R}^{p \times d}$




3. Reduced (d)

$Z = \beta^\top X \in \mathbb{R}^d$



4. Flexible response

$\mathbb{E}(Y^x) = g(z)$




Ideal workflow of causal sufficient dimension reduction

Let $\mathbb{E}(Y^x) = g(\beta^\top x)$ and $\mathcal{S}_{\mathbb{E}(Y^x)}$ be the **causal central mean subspace** (CCMS).


1. Exposures (p)

$X \in \mathbb{R}^p$




2. Causal directions

$\beta \in \mathbb{R}^{p \times d}$




3. Reduced (d)

$Z = \beta^\top X \in \mathbb{R}^d$



4. Flexible response

$\mathbb{E}(Y^x) = g(z)$



Question: How can we estimate $\mathcal{S}_{\mathbb{E}(Y^x)}$ when $\mathbb{E}(Y^x) = g(\beta^\top x)$?

Estimation and Inference

Marginal structural model approach

Using the theory of marginal structural models (Robins 2000), Nabi et al. (2022) proposed estimating $\mathcal{S}_{\mathbb{E}(Y^x)}$ by solving the following efficient score:

$$
\begin{aligned}
0 = \frac{1}{n}\sum_{i=1}^{n}\Bigg[& \frac{\pi^*(X_i)}{\pi(X_i \mid C_i)}\Big\{Y_i - \mathbb{E}(Y \mid X_i, C_i)\Big\} \\
& \times \Big\{\alpha(X_i) - \mathbb{E}_q[\alpha(X) \mid \beta^\top X = \beta^\top X_i]\Big\} \\
& + \int_{\mathcal{X}} \Big\{\mathbb{E}(Y \mid X = x, C_i) - \mathbb{E}_q[Y \mid \beta^\top X = \beta^\top x]\Big\} \\
& \times \Big\{\alpha(x) - \mathbb{E}_q[\alpha(X) \mid \beta^\top X = \beta^\top x]\Big\}\pi^*(x)\,dx\Bigg].
\end{aligned}
$$

Marginal structural model approach difficulties

While the method is elegant, practical implementation is difficult.

Marginal structural model approach difficulties

While the method is elegant, practical implementation is difficult.

Nuisance functions to be estimated:

1. $m(x, c) := \mathbb{E}(Y \mid X, C)$
2. $\pi(x \mid c) := f_{X|C}(x \mid c)$
3. $\mathbb{E}_q(Y \mid \beta^\top X = z)$
4. $\mathbb{E}_q[\alpha(X) \mid \beta^\top X = z]$

Marginal structural model approach difficulties

While the method is elegant, practical implementation is difficult.

Nuisance functions to be estimated:

1. $m(x, c) := \mathbb{E}(Y \mid X, C)$
2. $\pi(x \mid c) := f_{X|C}(x \mid c)$
3. $\mathbb{E}_q(Y \mid \beta^\top X = z)$
4. $\mathbb{E}_q[\alpha(X) \mid \beta^\top X = z]$

Problem: Optimization is challenging when the parameter is matrix-valued and nuisance functions must be re-estimated at each iteration of β.

A simpler approach: recast the dose-response as a regression

Suppose we can construct a variable $\widetilde{Y}$ such that

$$\mathbb{E}(\widetilde{Y} \mid X = x) = \mathbb{E}(Y^x), \qquad \forall x \in \mathcal{X}.$$

A simpler approach: recast the dose-response as a regression

Suppose we can construct a variable $\widetilde{Y}$ such that

$$\mathbb{E}(\widetilde{Y} \mid X = x) = \mathbb{E}(Y^x), \qquad \forall x \in \mathcal{X}.$$

Then under our central assumption $\mathbb{E}(Y^x) = g(\beta^\top x)$, it follows that

$$\mathcal{S}_{\mathbb{E}(\widetilde{Y}|X)} = \mathcal{S}_{\mathbb{E}(Y^x)}.$$

A simpler approach: recast the dose-response as a regression

Suppose we can construct a variable $\widetilde{Y}$ such that

$$\mathbb{E}(\widetilde{Y} \mid X = x) = \mathbb{E}(Y^x), \qquad \forall x \in \mathcal{X}.$$

Then under our central assumption $\mathbb{E}(Y^x) = g(\beta^\top x)$, it follows that

$$\mathcal{S}_{\mathbb{E}(\widetilde{Y}|X)} = \mathcal{S}_{\mathbb{E}(Y^x)}.$$

Solution: Just run standard SDR for regression on $(\widetilde{Y}, X)$ to estimate the causal $\mathcal{S}_{\mathbb{E}(Y^x)}$!

Estimation of $\widetilde{Y}$

1. Causal dose-response:
$$\widetilde{Y} = \mu(X).$$
2. Doubly robust pseudo-outcome:
$$\widetilde{Y} = \xi = \frac{\pi(X)}{\pi(X \mid C)}\{Y - m(X, C)\} + \mathbb{E}_C[m(X, C)].$$

Both require estimation of the same nuisances

- $m(x, c) = \mathbb{E}(Y \mid X = x, C = c)$
- $\pi(x \mid c) = f_{X|C}(x \mid c)$.

Estimation of $\mathcal{S}_{\mathbb{E}(Y|X)}$ with MAVE

The minimum average variance estimator (MAVE) from Xia et al. (2002) estimates $\mathcal{S}_{\mathbb{E}(Y|X)}$:

$$\min_{\beta^\top\beta=I_d,\,\{a_j,b_j\}} \sum_{j=1}^{n}\sum_{i=1}^{n}\Big(Y_i - \{a_j + b_j^\top\beta^\top(X_i - X_j)\}\Big)^2 w_{ij}$$

where $w_{ij} = K_h(X_i - X_j)/\sum_{\ell=1}^{n} K_h(X_i - X_\ell)$.

Our CSDR Approach

CSDR Algorithm (Hsiao, Chang, et al. 2026):

1. Use $O = (Y, X, C)$ to estimate $\widetilde{Y}$.
2. Estimate $\mathcal{S}_{\mathbb{E}(Y^x)}$ by running MAVE on $(\widetilde{Y}, X)$.
3. Estimate $\mu(x)$ by replacing X with $Z := \widehat{\beta}^\top X$.

$$\widehat{\mu}_{DR,\widehat{\beta}}(z) = \frac{1}{n}\sum_{i=1}^{n}\left\{\frac{K_h(\widehat{Z}_i - z)}{\widehat{\pi}_{\widehat{\beta}}(\widehat{Z}_i \mid C_i)}\left[Y_i - \widehat{m}_{\widehat{\beta}}(\widehat{Z}_i, C_i)\right] + \widehat{m}_{\widehat{\beta}}(z, C_i)\right\},$$

4. Construct 95% confidence intervals for $\widehat{\mu}_{DR,\widehat{\beta}}(z)$ based on empirical variance of the influence function.

Alternative approach

The CSDR approach improves on the MSM approach by avoiding numerical optimization on β-dependent nuisances.

However, estimation of $\widetilde{Y}$ still involves nuisance estimation of $m(x, c)$ and $\pi(x \mid c)$.

We propose another alternative that can estimate $\mathcal{S}_{\mathbb{E}(Y^x)}$ without $m(x, c)$ and $\pi(x \mid c)$.

Partial dimension reduction on $\mathbb{E}(Y \mid X, C)$

For this approach, we make an additional assumption that

$$\mathbb{E}(Y^x \mid C = c) = g(\beta^\top x, c).$$

One can show that any β satisfying the above also satisfies our original central assumption:

$$\mathbb{E}(Y^x) = g(\beta^\top x).$$

It follows that the partial central mean subspace of Y given (X, C) is equal to $\mathcal{S}_{\mathbb{E}(Y^x)}$.

Partial dimension reduction on $\mathbb{E}(Y \mid X, C)$

For this approach, we make an additional assumption that

$$\mathbb{E}(Y^x \mid C = c) = g(\beta^\top x, c).$$

One can show that any β satisfying the above also satisfies our original central assumption:

$$\mathbb{E}(Y^x) = g(\beta^\top x).$$

It follows that the partial central mean subspace of Y given (X, C) is equal to $\mathcal{S}_{\mathbb{E}(Y^x)}$.

Solution: Run partial dimension reduction (Li et al. 2010) directly on $\mathbb{E}(Y \mid X, C)$.

Theoretical results: subspace estimation

Let $\delta_n = \sqrt{\log n/(nh^d)}$ and $R_n = \|\hat{\mu}(X) - \mu(X)\|_\infty$ (the error in estimating $\widetilde{Y}$).

Theoretical results: subspace estimation

Let $\delta_n = \sqrt{\log n/(nh^d)}$ and $R_n = \|\hat{\mu}(X) - \mu(X)\|_\infty$ (the error in estimating $\widetilde{Y}$).

1. **Subspace estimation**: We establish convergence rates for estimating $\mathcal{S}_{\mathbb{E}(Y^x)}$:

- Method 1: $\|\widehat{\beta} - \beta\| = O_p(h^3 + h\delta_n + h^{-1}R_n)$ (dependent on $\widetilde{Y}$ estimation)
- Method 2: $\|\widehat{\beta} - \beta\| = O_p(h^3 + h\delta_n + h^{-1}\delta_n^2)$ (nuisance estimation-free)

Theoretical results: subspace estimation

Let $\delta_n = \sqrt{\log n/(nh^d)}$ and $R_n = \|\hat{\mu}(X) - \mu(X)\|_\infty$ (the error in estimating $\widetilde{Y}$).

1. **Subspace estimation**: We establish convergence rates for estimating $\mathcal{S}_{\mathbb{E}(Y^x)}$:

- Method 1: $\|\widehat{\beta} - \beta\| = O_p(h^3 + h\delta_n + h^{-1}R_n)$ (dependent on $\widetilde{Y}$ estimation)
- Method 2: $\|\widehat{\beta} - \beta\| = O_p(h^3 + h\delta_n + h^{-1}\delta_n^2)$ (nuisance estimation-free)

2. **Structural dimension selection**: A cross-validation procedure consistently estimates the dimension of $\mathcal{S}_{\mathbb{E}(Y^x)}$.

$$\mathbb{P}(\hat{d} = d_0) \longrightarrow 1.$$

Theoretical results: asymptotic equivalence

One may wonder how estimating $\mu(x) = \mathbb{E}(Y^x)$ using $\widehat{\beta}_n$ compares to the true β.

Theoretical results: asymptotic equivalence

One may wonder how estimating $\mu(x) = \mathbb{E}(Y^x)$ using $\hat{\beta}_n$ compares to the true β.

Asymptotic equivalence: We show that error in estimating β can be *asymptotically negligible* for estimating $\mu(x)$ (Forzani et al. 2024):

$$\sqrt{nh_n^d}\{\hat{\mu}_{DR}(x;\hat{\beta}_n) - \hat{\mu}_{DR}(x;\beta)\} = o_p(1)$$

Insight: Estimation of β can achieve $O_p(n^{-1/2})$ convergence, faster than the $O_p\{(nh^d)^{-1/2}\}$ rate of $\hat{\mu}_{DR}(x)$.

Theoretical results: subspace importance scores

Problem: β is not uniquely identified, so direct interpretation of loadings is tricky.

Solution: Measure each exposure's contribution using the projection matrix, which is unique.

$$\lambda_j = e_j^\top \beta(\beta^\top \beta)^{-1}\beta^\top e_j = \left[\beta(\beta^\top \beta)^{-1}\beta^\top\right]_{jj}$$

- Basis invariant: unaffected by rotations in β.
- Bounded: $0 \leq \lambda_j \leq 1$
- Decomposition: $\sum_{j=1}^{p} \lambda_j = d$

Summary of our work

Existing approaches either

- Estimated outcome regression instead of causal dose-response
- Performed unsupervised dimension reduction that ignored causal structure
- Conduct causal inference but on a scalar, joint shift exposure

Summary of our work

Existing approaches either

- Estimated outcome regression instead of causal dose-response
- Performed unsupervised dimension reduction that ignored causal structure
- Conduct causal inference but on a scalar, joint shift exposure

Our approach builds on and combines the strengths of these previous approaches

- Flexibly estimates the nonparametric dose-response in a lower dimension
- Returns a causally and scientifically interpretable β
- Final dose-response estimation is asymptotically robust to estimation error of $\widehat{\beta}$

Simulations and Real Data

CSDR recovers the true $\mathcal{S}_{\mathbb{E}(Y^x)}$

Table 1: Frobenius norm error between estimated and true $\mathcal{S}_{\mathbb{E}(Y^x)}$ with estimated $\hat{d}$. Mean (SD).

Method	$n = 500$	$n = 1000$	$n = 5000$
CSDR	0.45 (0.39)	0.22 (0.29)	0.03 (0.04)
pCCA	1.20 (0.18)	1.10 (0.20)	0.74 (0.20)
MAVE	1.29 (0.17)	1.17 (0.10)	1.07 (0.02)
PCA	2.00 (0.00)	2.00 (0.00)	2.00 (0.00)

CSDR recovers the true structural dimension

Table 2: Proportion selecting each dimension ($d_0 = 2$).

n	Method	$d = 1$	$d = 2$	$d = 3$	$d = 4$
500	**CSDR**	0.03	**0.88**	0.07	0.01
	MAVE	0.00	**0.11**	0.75	0.14
5000	**CSDR**	0.00	**1.00**	0.00	0.00
	MAVE	0.00	**0.00**	1.00	0.00

CSDR improves inference of the causal dose-response




Figure 1: Pointwise 95% coverage for the exposure-response curve ($p = 10$, $d = 2$) at 100 test points; methods differ only in β estimation.

PFAS real data application

We use the **Atlanta African American Maternal-Child Cohort** ($n = 305$) to study the effect of a PFAS mixture ($p = 4$) on infant birthweight (Chang et al. 2022). Covariates include maternal age, BMI, education, tobacco use, marijuana use, and infant sex.

ESTIMATED REDUCTION ($\widehat{d} = 1$)

$$\hat{Z} = 0.77\,\text{PFOS} + 0.14\,\text{PFOA} + 0.26\,\text{PFNA} - 0.56\,\text{PFHxS}$$

SUBSPACE IMPORTANCE SCORE $\widehat{\lambda}_j$

PFOS	0.60
PFOA	0.02
PFNA	0.07
PFHxS	0.31





Figure 2: PFAS exposure-response curve from ATL-AA


BKMR and qgcomp comparison analysis

BKMR



All four PFAS share a quantile versus the median.

qgcomp



All four PFAS shift together across exposure quartiles.

Takeaway: Different estimands, but in general, increasing PFAS leads to decreasing birthweight.


Future Directions

Causal dimension reduction for mediation analysis


Colleagues in EH want to study biological mechanisms of exposures using omics.

Can CSDR ideas preserve mediation effects and identify important mediators in M?

Reduction between X and M



Reduction between M and Y




Confounders C omitted for clarity.

Problem: Preserving the mediation effect alone may not capture both the exposure–mediator and mediator–outcome relationships.

Causal representation learning

Can we extend CSDR from linear subspaces to **nonlinear, interpretable representations**?

Variational autoencoders offer a template.



Why I am excited: could help address many common problems in biomarker discovery including curse of dimensionality, positivity violations, and interpretability.

Data fusion for improved health surveillance

In previous work (Hsiao et al. 2025; Hsiao, Fede, et al. 2026; Varghese et al. 2026), we estimated county-level hypertension care cascade comparing

- NHANES (national, probability)
- BRFSS survey data (county, probability)
- Self-service kiosk data (county, nonprobability)
- Epic COSMOS EHR (county, nonprobability)

Problem: Can we combine NHANES and large nonprobability samples (e.g., kiosks, EHR) in an intelligent way? Can spatiotemporal dependence further improve estimates?

Final thoughts

What I bring

- Causal inference, dimension reduction, spatiotemporal statistics
- Applications in environmental health, EHR, disease surveillance for public health impact
- Engagement and leadership in scientific collaborations, teaching, and mentorship

I would love to build this agenda at Wake Forest and contribute to its growing research community. I am especially excited about opportunities across the School of Medicine, Advocate Health, and Statistical Sciences.

References i

Bobb, Jennifer F., Linda Valeri, Birgit Claus Henn, et al. 2015. "Bayesian Kernel Machine Regression for Estimating the Health Effects of Multi-Pollutant Mixtures." *Biostatistics* 16 (3): 493–508.

Chang, Che-Jung, Dana Boyd Barr, P.Barry Ryan, et al. 2022. "Per- and Polyfluoroalkyl Substance (PFAS) Exposure, Maternal Metabolomic Perturbation, and Fetal Growth in African American Women: A Meet-in-the-Middle Approach." *Environment International* 158 (January): 106964.

Colangelo, Kyle, and Ying-Ying Lee. 2026. "Double Debiased Machine Learning Nonparametric Inference with Continuous Treatments." *Journal of Business & Economic Statistics* 44 (1): 67–79.

Cook, R. Dennis. 2026. "On the Foundational Arguments of Sufficient Dimension Reduction." *WIREs Computational Statistics* 18 (2): e70064.

Forzani, Liliana, Daniela Rodriguez, and Mariela Sued. 2024. "Asymptotic Results for Nonparametric Regression Estimators After Sufficient Dimension Reduction Estimation." *TEST* 33 (4): 987–1013.

Gibson, Elizabeth A., Junhui Zhang, Jingkai Yan, et al. 2022. "Principal Component Pursuit for Pattern Identification in Environmental Mixtures." *Environmental Health Perspectives* 130 (11): 117008.

Hsiao, Thomas W., Howard H. Chang, and Razieh Nabi. 2026. *Causal Sufficient Dimension Reduction for Multiple Continuous Exposures with an Application to Environmental Mixtures*. arXiv:2606.14840. arXiv.

Hsiao, Thomas W., Lauren Fede, Cameron Gocke, Lance A. Waller, Mohammed K. Ali, and Jithin Sam Varghese. 2026. "County Prevalence, Awareness, and Control of High Blood Pressure From Health Kiosks in the United States, 2017–2024." *Hypertension* 83 (2): e25829.

Hsiao, Thomas W., Jithin Sam Varghese, Lauren Fede, Cameron Gocke, Lance A. Waller, and Mohammed K. Ali. 2025. "Blood Pressure Measurements From Self-Service Health Kiosks in US Retail Stores, 2017-2024." *JAMA Cardiology* 10 (10): 1044–48.

Keil, Alexander P., Jessie P. Buckley, Katie M. O'Brien, Kelly K. Ferguson, Shanshan Zhao, and Alexandra J. White. 2020. "A Quantile-Based g-Computation Approach to Addressing the Effects of Exposure Mixtures." *Environmental Health Perspectives* 128 (4): 047004.

Li, Lexin, Bing Li, and Li-Xing Zhu. 2010. "Groupwise Dimension Reduction." *Journal of the American Statistical Association* 105 (491): 1188–201.

References ii

Nabi, Razieh, Todd McNutt, and Ilya Shpitser. 2022. "Semiparametric Causal Sufficient Dimension Reduction of Multidimensional Treatments." *Proceedings of the Thirty-Eighth Conference on Uncertainty in Artificial Intelligence*, August, 1445–55.

Robins, James M. 2000. "Marginal Structural Models Versus Structural Nested Models as Tools for Causal Inference." In *Statistical Models in Epidemiology, the Environment, and Clinical Trials*, edited by Willard Miller, M. Elizabeth Halloran, and Donald Berry, vol. 116. Springer New York.

Varghese, Jithin Sam, Thomas W. Hsiao, Lauren Fede, et al. 2026. "National, State, and County Estimates of Adult Overweight and Obesity From Electronic Health Records and Kiosks in Retail Locations, 2024–2025." *Obesity* 34 (4): 908–16.

Xia, Yingcun, Howell Tong, W. K. Li, and Li-Xing Zhu. 2002. "An Adaptive Estimation of Dimension Reduction Space." *Journal of the Royal Statistical Society. Series B (Statistical Methodology)* 64 (3): 363–410.

Acknowledgments

Postdoctoral advisors

Howard Chang | **Razieh Nabi**

For their mentorship, guidance, and support.

Collaborators

Anne Dunlop | **Stephanie Eick** | **Donghai Liang** | **Youran Tan**

For sharing the ATL-AA cohort data and for their thoughtful discussions.

Questions?



CSDR preprint

arxiv.org/abs/2606.14840



csdr R package

github.com/tXiao95/csdr

Appendix

Setup outcome, exposures, and confounders

R code

```r
library(csdr)
data(csdr_example2)

Y <- csdr_example2$Y
A <- csdr_example2$A
C <- csdr_example2$C

head(Y, 5)
head(A[, c(1:3, 18:20)], 5)
head(C, 5)
```

Console

```
> head(Y, 5)
     obs1      obs2      obs3      obs4      obs5
  0.3805    0.1838   -2.0486    0.4977    5.4798

> head(A[, c(1:3, 18:20)], 5)
         A1     A2     A3    A18    A19    A20
obs1 -1.066  1.573  0.953  0.033 -0.480  0.500
obs2 -1.028  0.325 -0.364 -0.967 -1.543 -0.572
obs3 -0.800 -0.103 -0.433 -0.011  0.505  0.623
obs4  2.492  0.582 -1.134 -1.473 -1.138  1.551
obs5  1.733 -0.114  0.615  1.083  0.350  0.133

> head(C, 5)
         C1    C2     C3     C4     C5
obs1 -0.935 1.312 -0.294 -1.979  1.150
obs2 -0.694 0.531  0.367 -2.259 -1.293
obs3 -0.548 0.712 -0.088 -0.407  1.828
obs4  0.766 1.050 -0.498 -0.593 -0.604
obs5  1.852 -0.554  0.849  0.643 -0.312
```

Fit the causal subspace

R code

```r
sl_lib <- c(
  "SL.glm", "SL.earth", "SL.ranger"
)

learners <- csdr_learners(
  sl_library = sl_lib
)

set.seed(42)
fit <- csdr(
  Y = Y, A = A, C = C,
  L = 5, seed = 42,
  learners = learners,
  max_dim = 5,
  keep_nuisance = TRUE
)

summary(fit)
```

Console

```
> summary(fit)
Summary of causal sufficient dimension reduction fit

 variant d_hat  n p mave_method target_exposure
      DR     2 100 20   meanMAVE               A

Learners:
 outcome: SuperLearner regression
   family: gaussian
   SL.library: SL.glm, SL.earth, SL.ranger
 gps: MVN GPS with linear conditional mean
   method_gps: linear; density_floor: 1e-16

Target construction:
Effective folds: 5
PO marginalization: crossfit
```

Inspect the fitted subspace and scores

R code

```r
B_hat <- coef(fit)
Z_hat <- scores(fit)

mave <- mave_fits(fit)
dim_selection <- mave$dimension_selection

B_hat[c(1, 2, 20), , drop = FALSE]
sis(fit)
head(Z_hat, 2)
dim_selection
```

Console

```
> B_hat[c(1, 2, 20), , drop = FALSE]
       dir1    dir2
A1  -0.3401  0.5777
A2  -0.3524  0.4195
A20 -0.0521 0.0784

> sis(fit)
    A1    A2    A3    A4    A5    A6    A7    A8    A9   A10
 0.464 0.312 0.533 0.390 0.000 0.004 0.002 0.020 0.001 0.021
   A11   A12   A13   A14   A15   A16   A17   A18   A19   A20
 0.068 0.059 0.028 0.020 0.006 0.005 0.015 0.036 0.006 0.009

> head(Z_hat, 2)
       score1  score2
obs1  -0.1862  0.4515
obs2   1.5224 -0.3484

> dim_selection
Dimensions      0    1    2    3    4    5
CV-value     1.58 1.13 0.75 0.77 0.92 0.97
Selected dimension: 2
```

Estimate the response in the reduced exposure

R code

```r
z_grid <- matrix(
  rep(apply(Z_hat, 2, median), each = 50),
  nrow = 50,
  dimnames = list(NULL, colnames(Z_hat))
)
z_grid[, 1] <- seq(
  quantile(Z_hat[, 1], 0.05),
  quantile(Z_hat[, 1], 0.95),
  length.out = 50
)

ers <- estimate_ers(
  Y = Y, A = Z_hat, C = C,
  a_eval = z_grid, estimator = "DR",
  L = 5, seed = 42,
  verbose = FALSE
)

head(ers$results)
```

Console

```
> head(ers$results)
  score1 score2 estimate     se ci_lower ci_upper effective_n
1 -1.990 0.0015    2.676 0.5599    1.579    3.774       4.502
2 -1.912 0.0015    2.693 0.4762    1.760    3.627       4.559
3 -1.834 0.0015    2.703 0.3920    1.935    3.472       4.671
4 -1.756 0.0015    2.703 0.3136    2.088    3.317       4.851
5 -1.678 0.0015    2.687 0.2493    2.198    3.176       5.113
6 -1.601 0.0015    2.652 0.2100    2.241    3.064       5.476
    min_gps h_used_1 h_used_2
1 4.049e-05    0.578    0.596
2 4.049e-05    0.578    0.596
3 4.049e-05    0.578    0.596
4 4.049e-05    0.578    0.596
5 4.049e-05    0.578    0.596
6 4.049e-05    0.578    0.596
```

Dimension Selection

NW leave-one-out estimator

$$\hat{a}_{d0,j} = \frac{\sum_{i \neq j} K_{h_d}^{(i,j)} \tilde{Y}_i}{\sum_{i \neq j} K_{h_d}^{(i,j)}}, K_{h_d}^{(i,j)} = K_{h_d}\{\hat{\beta}_1^\top(\tilde{X}_i - \tilde{X}_j), \ldots, \hat{\beta}_d^\top(\tilde{X}_i - \tilde{X}_j)\}$$

Cross-validated error

$$\mathsf{CV}_n(d) = \frac{1}{n}\sum_{j=1}^{n}\left(\tilde{Y}_j - \hat{a}_{d0,j}\right)^2, \qquad \mathsf{CV}_n(0) = \frac{1}{n}\sum_{i=1}^{n}(\tilde{Y}_i - \bar{Y})^2$$

Estimated structural dimension

$$\hat{d} = \arg\min_{0 \leq d \leq p} \mathsf{CV}_n(d)$$