On the precision of dynamic gene regulation

Tarek Tohme

Under the supervision of

Aleksandra Walczak and Thierry Mora

November 9, 2026



Neuron



Kidney cell



Gene expression is regulated



Overview of gene regulation



Transcription

DNA
Promoter
Gene
RNA Polymerase

Overview of gene regulation



Transcription

DNA
Promoter
Gene
RNA Polymerase
mRNA

Overview of gene regulation



Transcription



Translation

DNA
Promoter
Gene
RNA Polymerase
mRNA

Overview of gene regulation



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Expression levels are continuous ...



Expression levels are continuous ...



Expression levels are continuous ...



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



Sensitive response to shallow concentration gradients

Early Drosophila embryo



Grimm et al., *Development,* 2010

Positional information of the bicoid gradient: Gregor et al., *Cell* 2007

Precise sensing on a tight schedule



Precise sensing on a tight schedule



Precise sensing on a tight schedule



Precise sensing on a tight schedule



Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

2. Which promoter architectures are optimal?

Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

2. Which promoter architectures are optimal?

3. How are these optimal architectures shaped by their constraints and by the objective function they maximize?

Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

2. Which promoter architectures are optimal?

3. How are these optimal architectures shaped by their constraints and by the objective function they maximize?





Lammers, Flamholz, Garcia *PNAS* 2023

Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

2. Which promoter architectures are optimal?

3. How are these optimal architectures shaped by their constraints and by the objective function they maximize?









Lammers, Flamholz, Garcia *PNAS* 2023

Understanding the concentration sensitivity of promoters

Questions:

1. How to quantify sensitivity, to evaluate/compare promoter models?

2. Which promoter architectures are optimal?

3. How are these optimal architectures shaped by their constraints and by the objective function they maximize?









Lammers, Flamholz, Garcia *PNAS* 2023

Issues:

1. No accounting for physics-imposed rate constraints

2. Measure sensitivity in bits/burst

Understanding the concentration sensitivity of promoters

Approach:

1. Defining a measure of precision

Understanding the concentration sensitivity of promoters

Approach:

1. Defining a measure of precision

2. Identifying physics-imposed constraints on promoter architecture

Understanding the concentration sensitivity of promoters

Approach:

1. Defining a measure of precision

2. Identifying physics-imposed constraints on promoter architecture

2. Constrained optimization of precision

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Concentration: high



A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Concentration: high



low

Sequential hypothesis testing

$P($ $| c_{\text{low}})$ $P($ $| c_{\text{high}})$

Sequential hypothesis testing

$P($ $| c_{\text{low}})$ $\qquad$ $P($ $| c_{\text{high}})$

$$\text{R(t)= log } \frac{P(\text{ } | c_{\text{low}})}{P(\text{ } | c_{\text{high}})}$$

Sequential hypothesis testing

$P(\text{ } \mid c_{\text{low}})$ $\qquad$ $P(\text{ } \mid c_{\text{high}})$

$$R(t) = \log \frac{P(\text{ } \mid c_{\text{low}})}{P(\text{ } \mid c_{\text{high}})} \begin{cases} > \text{confidence value K} \longrightarrow \boxed{\text{Decision: } c_{\text{low}}} \\ \\ \end{cases}$$

Sequential hypothesis testing

$P(\text{ } \mid c_{low})$ $P(\text{ } \mid c_{high})$

$$R(t) = \log \frac{P(\text{ } \mid c_{low})}{P(\text{ } \mid c_{high})} \begin{cases} > \text{confidence value K} \rightarrow \boxed{\text{Decision: } c_{low}} \\ < -K \rightarrow \boxed{\text{Decision: } c_{high}} \end{cases}$$

Sequential hypothesis testing

$P(\text{⎍⎍⎍⎍} \mid c_{\text{low}})$ $\qquad$ $P(\text{⎍⎍⎍⎍} \mid c_{\text{high}})$

$$R(t) = \log \frac{P(\text{⎍⎍⎍⎍} \mid c_{\text{low}})}{P(\text{⎍⎍⎍⎍} \mid c_{\text{high}})} \begin{cases} > \text{confidence value K} \rightarrow \boxed{\text{Decision: } c_{\text{low}}} \\ < -\text{K} \rightarrow \boxed{\text{Decision: } c_{\text{high}}} \end{cases}$$

Sequential Probability Ratio Test (SPRT)

Wald 1949 (conception)
Siggia, Vergassola 2012 (application to biological receptors)
Desponds, Vergassola, Walczak 2022 (optimization)
Lammers, Flamholz, Garcia, 2024 (nonequilibrium receptors)

Sequential hypothesis testing



Sequential hypothesis testing



Sequential hypothesis testing



Sequential hypothesis testing



Fisher information rate and decision time

$$\mathcal{R}(t) = \log \frac{P(\text{⎍⎍⎾⎍⎍⎿⎍⎍}|c)}{P(\text{⎍⎍⎾⎍⎍⎿⎍⎍}\,|c+\delta c)}$$

Fisher information rate and decision time

$$\mathcal{R}(t) = \log \frac{P(\text{⎍⎍⎍⎍⎍}|c)}{P(\text{⎍⎍⎍⎍⎍}|c+\delta c)}$$

Drift-diffusion approximation when $\delta c/c \ll 1$

$$\frac{d\mathcal{R}}{dt} = V + \sqrt{2V}\eta \leftarrow \text{Gaussian noise } \langle \eta(t)\eta(t') \rangle \sim \delta(t-t')$$

Drift = Diffusion

Fisher information rate and decision time

$$\mathcal{R}(t) = \log \frac{P(\text{⎍⎍⎍⎍⎍⎍⎍}|c)}{P(\text{⎍⎍⎍⎍⎍⎍⎍}|c+\delta c)}$$

Drift-diffusion approximation when $\delta c/c \ll 1$

$$\frac{d\mathcal{R}}{dt} = \underbrace{V}_{\text{Drift}} + \underbrace{\sqrt{2V}\eta}_{\text{Diffusion}} \leftarrow \begin{array}{c}\text{Gaussian noise} \\ \langle \eta(t)\eta(t') \rangle \sim \delta(t-t')\end{array}$$

SPRT mean decision time (Siggia & Vergassola 2012)

$$\langle T_{\text{dec}} \rangle = \frac{K}{V} \tanh\left(\frac{K}{2}\right)$$

Fisher information rate and decision time

$$\mathcal{R}(t) = \log \frac{P(\text{⎍⎍⎍⎍⎍⎍⎍}|c)}{P(\text{⎍⎍⎍⎍⎍⎍⎍}|c+\delta c)}$$

Drift-diffusion approximation when $\delta c/c \ll 1$

$$\frac{d\mathcal{R}}{dt} = V + \sqrt{2V}\eta \leftarrow \text{Gaussian noise } \langle \eta(t)\eta(t') \rangle \sim \delta(t-t')$$

Drift = Diffusion

SPRT mean decision time (Siggia & Vergassola 2012)

$$\langle T_{\text{dec}} \rangle = \frac{K}{V} \tanh\left(\frac{K}{2}\right)$$

Drift (our result)

$$V \approx \frac{1}{2}\left(\frac{\delta c}{c}\right)^2 \dot{\mathcal{F}}(c)$$

Holds for any model

Fisher information rate and decision time

$$\mathcal{R}(t) = \log \frac{P(\text{⎍⎍⎍⎍⎍}|c)}{P(\text{⎍⎍⎍⎍⎍}|c+\delta c)}$$

Drift-diffusion approximation when $\delta c/c \ll 1$

$$\frac{d\mathcal{R}}{dt} = V + \sqrt{2V}\eta \leftarrow \text{Gaussian noise } \langle \eta(t)\eta(t') \rangle \sim \delta(t-t')$$

Drift = Diffusion

SPRT mean decision time (Siggia & Vergassola 2012)

$$\langle T_{\text{dec}} \rangle = \frac{K}{V} \tanh\left(\frac{K}{2}\right)$$

Drift (our result)

$$V \approx \frac{1}{2}\left(\frac{\delta c}{c}\right)^2 \dot{\mathcal{F}}(c)$$

Holds for any model

Fisher information rate

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{⎍⎍⎍⎍⎍}|c)$$

The optimization problem

$$\text{Maximize } \dot{\mathcal{F}}(c)$$

subject to the constraints

$$k_b \leq 4aD \text{ and } k_{\text{u,on,off}} \leq k_{\max}$$

Diffusion Activation

The optimization problem

$$\text{Maximize } \dot{\mathcal{F}}(c)$$

subject to the constraints

$$k_b \leq 4aD \text{ and } k_{\text{u,on,off}} \leq k_{\max}$$

Diffusion Activation

In practice:

1. numerical first pass (dynamic programming)

All solutions:



The optimization problem

Maximize $\dot{\mathcal{F}}(c)$

subject to the constraints

$k_b \leq 4aD$ and $k_{\mathrm{u,on,off}} \leq k_{\max}$

Diffusion Activation

In practice:

1. numerical first pass (dynamic programming)
2. ansatz obtained from solutions (analytical)

All solutions:



Optimal promoters: two distinct regimes



Optimal promoters: two distinct regimes





Cooperative binding



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Cooperative binding



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Cooperative binding



$\downarrow$ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints $\downarrow$

Cooperative binding



- DNA
- Promoter
- Gene
- Transcription factor
- RNA Polymerase
- mRNA

↓ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints ↓



Cooperative binding



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

↓ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints ↓



Diffusion limited
Activation-limited
No constraint saturated



Cooperative binding



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Diffusion limited
Activation-limited
No constraint saturated

↓ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints ↓



Proofreading



Proofreading



Proofreading



ω : [decoy TF] / [correct TF]

Proofreading



ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Proofreading



ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Selectivity f = P(transcribing | correct TF)

Proofreading



ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Selectivity f = P(transcribing | correct TF)

$$\max f = \gamma^2 / (\omega + \gamma^2)$$

$$\min f = \gamma / (\omega(\gamma+1)/2 + \gamma)$$

Proofreading



↓ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints ↓

ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Selectivity f = P(transcribing | correct TF)

$$\max f = \gamma^2 / (\omega + \gamma^2)$$

$$\min f = \gamma / (\omega(\gamma+1)/2 + \gamma)$$



Proofreading



$\downarrow$ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints $\downarrow$

ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Selectivity f = P(transcribing | correct TF)

$$\max f = \gamma^2 / (\omega + \gamma^2)$$

$$\min f = \gamma / (\omega(\gamma+1)/2 + \gamma)$$



Takeaways

The limiting constraint shapes the optimal architecture



Takeaways

The limiting constraint shapes the optimal architecture



Solution set is disconnected: each cluster points to a different sensing strategy



count rare
binding events



leverage
cooperative binding

Takeaways

The limiting constraint shapes the optimal architecture



Solution set is disconnected: each cluster points to a different sensing strategy



count rare
binding events



leverage
cooperative binding

Sharpness, selectivity do not lie at extrema





Transcriptional control by distal elements



Transcriptional control by distal elements



Transcriptional control by distal elements





DNA
Promoter
Gene
RNA Polymerase
mRNA

Fukaya et al., *Cell,* 2016
Chen et al., *Nature Genetics,* 2018
Brückner et al., *Science,* 2023

Transcriptional control through liquid phase separation?



Liquid condensate
150-350 nm



Du et al.,*Cell* 2024

- DNA
- Promoter
- Gene
- Transcription factor

Hnisz et al., *Cell* 2017

Sabari et al., *Science* 2018

Munshi et al., *Science Advances* 2025

Proteins condense on the surface of DNA





Morin et al., *Nature Physics* 2022

DNA
Promoter
Gene
Transcription factor

Pancholi et al., *MolecularSystems Biology* 2021

Rouches & Machta, *Biophysical Journal* 2025

Proteins condense on the surface of DNA



Influence on precision of gene expression?

Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics







Coupling condensation and polymer dynamics





Polymer configuration depends sensitively on concentration





Polymer configuration depends sensitively on concentration









Continuous model of surface condensation



Flory-Huggins potential

$$f(\phi) = \frac{\phi}{\nu}\ln\phi + (1-\phi)\ln(1-\phi) - \chi\phi^2 - \mu\phi$$

Cahn-Hilliard total energy

$$F = \int_B \left[\frac{\kappa}{2}(\nabla\phi)^2 + f(\phi)\right] d^3x - \int_S h_0\phi(\mathbf{x})dS$$

ϕ TF volume fraction
ν TF / solvent volume ratio
χ TF-TF attraction
μ chemical potential
h_0 surface attraction
κ Interface stiffness

The prewetting transition: varying surface attraction





The prewetting transition: varying surface attraction



Varying surface attraction h_0



The prewetting transition: varying surface attraction



Varying surface attraction h_0



The prewetting transition: varying surface attraction



Varying surface attraction h_0



The prewetting transition: varying surface attraction



Varying surface attraction h_0



Quantifying sensitivity of local concentration



Free receptor

$$\dot{\mathcal{F}}_{\text{free}}(\phi) = \frac{1}{\tau_{\text{free}}(\phi)}$$

Receptor at the attractive surface

$$\dot{\mathcal{F}}(\phi) = \underbrace{\frac{1}{\tau_{\text{free}}(\phi_{\text{local}})}}_{\text{Probe}} \underbrace{\left(\frac{d\log\phi_{\text{local}}}{d\log\phi}\right)^2}_{\text{Surface}}$$

Quantifying sensitivity of local concentration





Rouse model of DNA



$$P_{\text{poly}}(d) \propto d^2 e^{-3d^2/2sa^2} \propto e^{-U_{\text{poly}}(d)/k_B T}$$



Promoter-enhancer attraction



Cahn-Hilliard total energy

$$F = \int_B \left[\frac{\kappa}{2}(\nabla\phi)^2 + f(\phi)\right] d^3x - \int_{S_1} h_0\phi(\mathbf{x})dS - \int_{S_2} h_0\phi(\mathbf{x})dS$$

$$U_{\text{attr}}(d) = \min_{\phi(\mathbf{x})}\{F\}$$

$$\text{Force} = \partial_d U_{\text{attr}}(d)$$





Polymer + capillary attraction

$$U_{\text{attr}}(d) + U_{\text{poly}}(d) = U_{\text{tot}}(d)$$

$$P_{eq}(d) \propto e^{-U_{tot}(d)/k_B T}$$



low concentration high concentration

Polymer dynamics

$$U_{\text{attr}}(d) + U_{\text{poly}}(d) = U_{\text{tot}}(d)$$

$$P_{eq}(d) \propto e^{-U_{tot}(d)/k_B T}$$



Sensitivity from polymer dynamics

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{⎍⎍⎍⎍⎍}\,|c)$$



$$\dot{\mathcal{F}}(c) \approx \dot{\mathcal{F}}_{local}(c) + \dot{\mathcal{F}}_{DNA}(c)$$



$$\dot{\mathcal{F}}_o(c) = \underbrace{\frac{1}{\tau_{\text{burst}}(c)}}_{\text{Probe}} \underbrace{\left(\frac{d\log c_o}{d\log c}\right)^2}_{\text{Surface}}$$





Total sensitivity

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{⎍⎍⎍⎍⎍} \,|c)$$

$$\dot{\mathcal{F}}(c) \approx \dot{\mathcal{F}}_{local}(c) + \dot{\mathcal{F}}_{\mathrm{DNA}}(c)$$





Takeaways

Surface condensation can boost concentration sensitivity

Polymer dynamics could contribute to gene expression sensitivity

Limitations of the model:

- Rouse model unrealistic for DNA
- Loop extrusion, TADs
- Membrane binding behaves differently than bulk

Conclusion: Cells are complicated



Conclusion: Cells are complicated



Conclusion: Transcriptional regulation is very precise

Conclusion

Q. What is the most concentration-sensitive promoter architecture?

sharp response?
cooperative?
proofreading?

A. It depends on the limiting constraint

1. Cooperativity only optimal above critical concentration

"Keep binding sites free" (diffusion-limited)



"Bind cooperatively" (activation-limited)



Proofreading only optimal above critical **decoy** concentration

2. Solutions sets are generically **discontinuous**

3. Optimal promoters neither maximize sharpness nor selectivity







Discussing optimality requires precise accounting of underlying constraints

Conclusion

Q. How does a promoter's environment influence concentration sensitivity?

Surface condensation of transcription factors can amplify gene expression sensitivity



Polymer dynamics itself can encode a signal about concentration in transcription activity





Thanks.

StatBioPhys@ENS

Supervisors

Aleksandra Walczak
Thierry Mora

Collaborators

Massimo Vergassola (1st project)
Jacqueline Janssen (2nd project)
Henry Alston (2nd project)

Group

Past...

Antoine Aragon
Maria-Francesca Abbate
Xiaowen Chen
Francesco Camaglia
Andrea Mazzolini
Maria Ruiz Ortega
Vadim Karnaukhov
Mirna Elizabeta Kramar
Paul Pereira
Suchismita Das

... and present

Henry Alston
Jacqueline Janssen
Gabriel Athènes
Clarisse Marchesin
Natalie Blot
Daniel Wong
Evgenii Egorov
Mattia Greco
Anthia Prapa
Monika Rasz
Max Zayas
Johnathan Schrohe
Lander Bogers