On the precision of dynamic gene regulation

Tarek Tohme

Under the supervision of

Aleksandra Walczak and Thierry Mora

November 6, 2026



Neuron



Kidney cell



Gene expression is regulated



Overview of gene regulation



Transcription

DNA
Promoter
Gene
RNA Polymerase

Overview of gene regulation



Transcription

DNA
Promoter
Gene
RNA Polymerase
mRNA

Overview of gene regulation



Transcription



Translation

DNA
Promoter
Gene
RNA Polymerase
mRNA

Overview of gene regulation



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Expression levels are continuous ...



Expression levels are continuous ...



Expression levels are continuous ...



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



... but transcriptional control is noisy



Sensitive response to shallow concentration gradients



Grimm et al., *Development,* 2010

Precise sensing on a tight schedule



Precise sensing on a tight schedule



Precise sensing on a tight schedule



Precise sensing on a tight schedule



Understanding the regulatory precision of promoters

Goals
How to quantify precision, to evaluate/compare promoter models?
Which promoter architectures are optimal?
How are these optimal architectures shaped by their constraints and by the objective function they maximize?

Understanding the regulatory precision of promoters

Approach
Defining a measure of precision
Optimizing precision under constraints

A simple model of transcriptional control



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Concentration: high



low

Sequential hypothesis testing

$P(\text{⎍⎍⎍⎍} \mid c_{\text{low}})$ $\qquad$ $P(\text{⎍⎍⎍⎍} \mid c_{\text{high}})$

$$R(t) = \log \frac{P(\text{⎍⎍⎍⎍} \mid c_{\text{low}})}{P(\text{⎍⎍⎍⎍} \mid c_{\text{high}})} \begin{cases} > \text{confidence value K} \rightarrow \boxed{\text{Decision: } c_{\text{low}}} \\ < -\text{K} \rightarrow \boxed{\text{Decision: } c_{\text{high}}} \end{cases}$$

Sequential Probability Ratio Test (SPRT)

Wald 1949 (conception)
Siggia, Vergassola 2012 (application to biological receptors)
Desponds, Vergassola, Walczak 2022 (optimization)
Lammers, Flamholz, Garcia, 2024 (nonequilibrium receptors)

Sequential hypothesis testing



Sequential hypothesis testing



Sequential hypothesis testing



Sequential hypothesis testing



Fisher information rate and decision time

Drift-diffusion approximation when $\delta c/c \ll 1$

$$\frac{d\mathcal{R}}{dt} = V + \sqrt{2V}\eta \quad \leftarrow \quad \text{Gaussian noise } \langle \eta(t)\eta(t') \rangle \sim \delta(t-t')$$

V = Drift, $\sqrt{2V}$ = Diffusion

SPRT mean decision time (Siggia & Vergassola 2012)

$$\langle T_{\text{dec}} \rangle = \frac{K}{V} \tanh\left(\frac{K}{2}\right)$$

$$V \approx \frac{1}{2}\left(\frac{\delta c}{c}\right)^2 \dot{\mathcal{F}}(c)$$

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{ } \,|c)$$

The optimization problem

$$\text{Maximize } \dot{\mathcal{F}}(c)$$

subject to the constraints

$$k_b \leq 4aD \text{ and } k_{\text{u,on,off}} \leq k_{\max}$$

Diffusion Activation

In practice:

1. numerical first pass (dynamic programming)
2. ansatz obtained from solutions (analytical)



Optimal promoters: two distinct regimes



Cooperative binding



DNA
Promoter
Gene
Transcription factor
RNA Polymerase
mRNA

Diffusion limited
Activation-limited
No constraint saturated

↓ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints ↓



Spurious binding



$\downarrow$ Maximize $\dot{\mathcal{F}}(c)$ subject to the constraints $\downarrow$

ω : [decoy TF] / [correct TF]

$$\gamma : \frac{\text{decoy unbinding rate}}{\text{correct unbinding rate}}$$

Selectivity f = P(transcribing | correct TF)

$$\max f = \gamma^2 / (\omega + \gamma^2)$$

$$\min f = \gamma / (\omega(\gamma+1)/2 + \gamma)$$



Takeaways

The limiting constraint shapes the optimal architecture



Solution set is disconnected: each cluster points to a different sensing strategy



count rare
binding events



leverage
cooperative binding

Sharpness, selectivity do not lie at extrema



Transcriptional control by distal elements



Transcriptional control by distal elements



Transcriptional control by distal elements





DNA
Promoter
Gene
RNA Polymerase
mRNA

Fukaya et al., *Cell,* 2016
Chen et al., *Nature Genetics,* 2018
Brückner et al., *Science,* 2023

Transcriptional control through liquid phase separation?



Liquid condensate
150-350 nm



Du et al.,*Cell* 2024

- DNA
- Promoter
- Gene
- Transcription factor

Hnisz et al., *Cell* 2017

Sabari et al., *Science* 2018

Munshi et al., *Science Advances* 2025

Proteins condense on the surface of DNA





Morin et al., *Nature Physics* 2022

DNA
Promoter
Gene
Transcription factor

Pancholi et al., *MolecularSystems Biology* 2021

Rouches & Machta, *Biophysical Journal* 2025

Proteins condense on the surface of DNA



Influence on precision of gene expression?

Coupling condensation and polymer dynamics



Coupling condensation and polymer dynamics





Polymer configuration depends sensitively on concentration





Continuous model of surface condensation



Flory-Huggins potential

$$f(\phi) = \frac{\phi}{\nu}\ln\phi + (1-\phi)\ln(1-\phi) - \chi\phi^2 - \mu\phi$$

Cahn-Hilliard total energy

$$F = \int_B \left[\frac{\kappa}{2}(\nabla\phi)^2 + f(\phi)\right] d^3x - \int_S h_0\phi(\mathbf{x})dS$$

ϕ TF volume fraction
ν TF / solvent volume ratio
χ TF-TF attraction
μ chemical potential
h_0 surface attraction
κ Interface stiffness

The prewetting transition: varying surface attraction





The prewetting transition: varying surface curvature





Quantifying sensitivity of local concentration



Free receptor

$$\dot{\mathcal{F}}_{\text{free}}(c) = \frac{1}{\tau_{\text{burst}}(c)}$$

Receptor at the attractive surface

$$\dot{\mathcal{F}}(c) = \underbrace{\frac{1}{\tau_{\text{burst}}(c)}}_{\text{Probe}} \underbrace{\left(\frac{d\log c_o}{d\log c}\right)^2}_{\text{Surface}}$$

$\tau_{\text{burst}}(c)$ mean cycle time of the probe

c_o TF volume fraction at the probe

Quantifying sensitivity of local concentration





Rouse model of DNA



$$P_{\text{poly}}(d) \propto d^2 e^{-3d^2/2sa^2} \propto e^{-U_{\text{poly}}(d)/k_B T}$$



Promoter-enhancer attraction



Cahn-Hilliard total energy

$$F = \int_B \left[\frac{\kappa}{2}(\nabla\phi)^2 + f(\phi)\right] d^3x - \int_{S_1} h_0\phi(\mathbf{x})dS - \int_{S_2} h_0\phi(\mathbf{x})dS$$

$$U_{\text{attr}}(d) = \min_{\phi(\mathbf{x})}\{F\}$$

$$\text{Force} = \partial_d U_{\text{attr}}(d)$$





Polymer + capillary attraction

$$U_{\text{attr}}(d) + U_{\text{poly}}(d) = U_{\text{tot}}(d)$$

$$P_{eq}(d) \propto e^{-U_{tot}(d)/k_B T}$$



low concentration high concentration

Polymer dynamics

$$U_{\text{attr}}(d) + U_{\text{poly}}(d) = U_{\text{tot}}(d)$$

$$P_{eq}(d) \propto e^{-U_{tot}(d)/k_B T}$$



Sensitivity from polymer dynamics

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{⎍⎍⎍⎍⎍}\,|c)$$



$$\dot{\mathcal{F}}(c) \approx \dot{\mathcal{F}}_{local}(c) + \dot{\mathcal{F}}_{DNA}(c)$$



$$\dot{\mathcal{F}}_o(c) = \underbrace{\frac{1}{\tau_{\text{burst}}(c)}}_{\text{Probe}} \underbrace{\left(\frac{d\log c_o}{d\log c}\right)^2}_{\text{Surface}}$$





Total sensitivity

$$\dot{\mathcal{F}}(c) = -\lim_{t\to\infty} \frac{1}{t} \left\langle \frac{\partial^2 \mathcal{L}(c)}{(\partial \ln c)^2} \right\rangle_c$$

$$\mathcal{L}(c;t) = \ln P(\text{⎍⎍⎍⎍⎍} \,|c)$$

$$\dot{\mathcal{F}}(c) \approx \dot{\mathcal{F}}_{local}(c) + \dot{\mathcal{F}}_{\mathrm{DNA}}(c)$$





Takeaways

Surface condensation can boost concentration sensitivity

Polymer dynamics could contribute to gene expression sensitivity

Limitations of the model:

- Rouse model unrealistic for DNA
- Loop extrusion, TADs
- Membrane binding behaves differently than bulk

A Eukaryotic cell



A Eukaryotic cell



A Eukaryotic cell