

Brief protocol for observing protein dimerization in bacteria via fluorescence microscopy

Ryan J. Butcher^a, Jeffrey J. Tabor^{a,b}

^aDepartment of Bioengineering, ^bDepartment of Biosciences, Rice University, Houston, TX 77005;

Protein-protein interactions are the driving force behind a vast and diverse set of cellular functions. Fluorescence resonance energy transfer (FRET), the proximity-dependent energy transfer between a donor and acceptor fluorophore, provides real-time measurements of protein-protein interactions in living cells¹. Alternatively, energy transfer can occur between identical fluorophores across the overlap between excitation and emission spectra (homoFRET). In the case of fluorescent proteins with relatively slow molecular rotation speed, this interaction decreases fluorescence polarization and has been used to measure GFP oligomerization², kinase activity³, and bacterial response regulator dimerization⁴ in living cells. This document is intended to be a practical guide to measure fluorescence polarization via microscopy, specifically the difference between monomeric and dimeric mNeonGreen⁵ (mNG) expressed in *E. coli*. For further reading on fluorescence polarization microscopy, see references 6-7.

Approach

To measure fluorescence polarization (FP), the sample must be excited by linearly polarized light. The subsequent emission light needs to be separated into the parallel and perpendicular components via an image splitter or similar device (**Fig. 1A**). A polarizing filter can easily be inserted into the excitation cup of the microscope, but determining the orientations of the two emission-side polarizing filters is not immediately obvious. Observed fluorescence will be maximal if the emission-side polarizing filter is aligned with the excitation-side polarizing filter (parallel), and minimal when the excitation- and emission-side polarizers are perpendicular. By measuring fluorescence while incrementally rotating the emission-side polarizers across 180 degrees, the parallel and perpendicular orientations will be revealed. (**Fig. 1B**).

Here we measure the fluorescence of *E. coli* expressing monomeric mNG or dimeric mNG (**Fig. 1C**). For simple analysis, the total pixel intensity of each image after background subtraction is summed, representing the total fluorescence of the cells in the frame (**Fig. 1D**). After plotting this total fluorescence quantity versus emission-side polarizer orientation, a sinusoidal relationship is apparent as the emission-side polarizer aligns and de-aligns with the excitation-side polarizer. Finally, one emission-side polarizer is set in the parallel position and the other in the perpendicular position before acquiring test images. Using **Eqn. 1**, we clearly observe that mNG dimerization decreases FP (**Fig. 1F**). If desired, this approach can be refined with segmentation to obtain single-cell measurements.

$$FP = \frac{I_{parallel} - I_{perpendicular}}{I_{parallel} + I_{perpendicular}}$$

Eqn. 1




Figure 1: Observing mNG dimerization in *E. coli* via fluorescence polarization. **A.** Light path schematic for measuring fluorescence polarization via microscopy **B.** Filter holder for image-splitting device. Overlaid white arrows indicate polarizer orientation, marked by the red stripe. **C.** Plasmid diagrams for strains expressing monomeric (high polarization) or dimeric (low polarization) mNG. **D.** Representative image field of cells expressing mNG. **E.** Modulation of total pixel intensity for either filter channel during manual rotation across 180 degrees. Minima occur at 51 and 46 degrees for filter 1 and 2, respectively. Maxima occur at 140 and 138 degrees for filter 1 and 2, respectively. Lines are fitted to the median of whole-image pixel intensity sums for 16 total frames (8 frames for either strain) containing roughly 50-100 cells. **F.** Final comparison of fluorescence polarization between monomeric and dimeric mNG. Data points represent single frames containing roughly 50-100 cells.

Detailed Protocol

Positive and negative control strains

E. coli K-12 strain BW28357 ($\Delta(araD-araB)567$, $\Delta lacZ4787(::rrnB-3)$, λ^- , $\Delta(rhaD-rhaB)568$, $hsdR514$) was transformed with either pRB035 (aTc-inducible mNeonGreen) or pRB121 (aTc-inducible dimeric mNeonGreen). Both plasmids are available on Addgene (<https://www.addgene.org/browse/article/28224612/>). Strains were stored in LB media + 15% glycerol at -80°C.

Data processing

Images were analyzed using MATLAB. Code and raw images are available on the Tabor lab Github (<https://github.com/taborlab/PolarizationMicroscopy>).

Equipment

- Inverted fluorescence microscope (Ti-E, Nikon)
- Excitation source (BDX module, X-LED1, Excelitas)
- Image splitting device (W-View Gemini, Hamamatsu)
- Filters (470/40 ex., 525/50 em., 59022 bs, from 89021 set, Chroma)
- 3x linear polarizing filters (DP – 100 – VIS, Meadowlark Optics)
- Non-polarizing beamsplitting cube (NB – 100 – VIS, Meadowlark Optics)
- 37°C shaking incubator

Protocol

1. Incubate cultures of positive and negative control strains in LB media at 37°C with 250 rpm shaking overnight (12-18hrs).
2. Dilute saturated cultures 1000x into M9 media (1× M9 salts (Teknova, M1902), 0.4% (wt/vol) glucose (Avantor, 4908-06), 0.2% (wt/vol) casamino acids (EMD Millipore, 2240-500GM), 2 mM MgSO₄ (VWR, BDH9246-500G) and 0.1 mM CaCl₂ (Alfa Aesar, L13191), plus 9 ng/mL anhydrotetracycline (aTc- Fisher Scientific, NC9451757) and antibiotics (chloramphenicol- Fisher Scientific, AAB2084114).
3. Incubate 3-4 hours at 37°C with 250 rpm shaking until cultures reach the middle of exponential phase.
4. Transfer 0.2 µL culture onto solidified 1 mm thickness, 1% agarose pads.
5. Collect fluorescence images of strains at a variety of polarizer orientations across 180°. In this case, we imaged eight fields per sample with nine filter orientations (22.5° filter rotation between images).
6. After identifying parallel and perpendicular filter orientations as in **Fig. 1E**, set one polarizer in the parallel orientation and the other in the perpendicular orientation.
7. Image samples and calculate the differences in fluorescence polarization.

References

1. Truong, Kevin, and Mitsuhiro Ikura. "The use of FRET imaging microscopy to detect protein–protein interactions and protein conformational changes in vivo." *Current opinion in structural biology* 11.5 (2001): 573-578.
2. Bader, Arjen N., et al. "Homo-FRET imaging enables quantification of protein cluster sizes with subcellular resolution." *Biophysical journal* 97.9 (2009): 2613-2622.
3. Ross, Brian L., et al. "Single-color, ratiometric biosensors for detecting signaling activities in live cells." *Elife* 7 (2018): e35458.
4. Butcher, Ryan J., and Jeffrey J. Tabor. "Real-time detection of response regulator phosphorylation dynamics in live bacteria." *Proceedings of the National Academy of Sciences* 119.35 (2022): e2201204119.
5. Shaner, Nathan C., et al. "A bright monomeric green fluorescent protein derived from *Branchiostoma lanceolatum*." *Nature methods* 10.5 (2013): 407-409.
6. Tramier, Marc, and Maite Coppey-Moisan. "Fluorescence anisotropy imaging microscopy for homo-FRET in living cells." *Methods in cell biology* 85 (2008): 395-414.
7. Piston, David W., and Mark A. Rizzo. "FRET by fluorescence polarization microscopy." *Methods in cell biology* 85 (2008): 415-430.