

Resolving Super Fine-Resolution SIF via Coarsely-Supervised U-Net Regression

Joshua Fan, Di Chen, Jiaming Wen, Ying Sun, Carla Gomes
Cornell University

NeurIPS 2021 Workshop, Tackling Climate Change with Machine Learning



Motivation: monitoring crop productivity

- Crop production is extremely sensitive to climate change
 - Anthropogenic climate change has slowed agricultural productivity growth by 21% since 1961 (Ortiz-Bobea et al, 2021)
- Heat and drought stress are particularly damaging
- Monitoring the effects of climate change on crop growth in real time is crucial, particularly for **food security** (He et al, 2020)




Solar-induced chlorophyll fluorescence (SIF)

- Satellites can inexpensively monitor crop growth from space
- A lot of work uses vegetation indices (VIs) such as NDVI
 - Simple combinations of a few spectral bands (“greenness”)
- By contrast: **Solar-Induced Chlorophyll Fluorescence (SIF)** has direct **mechanistic linkages to photosynthesis** (Magney et al, 2019)
 - Better at predicting crop yields than VIs (Peng et al, 2020)
 - Can detect crop response to heat stress (Song et al, 2018) and drought (Zhang et al, 2019)


Problem: coarse resolution of SIF

- SIF is difficult to measure from space
- Existing satellites (such as OCO-2) can capture at most 3x3km resolution; coverage is limited
- Within a large area (many square km), SIF can vary dramatically between farms due to **crop type**, **management practices**, etc.
- Want to predict SIF at finer resolution (e.g. 30x30m)



Task: coarsely supervised regression


- To predict SIF at a very fine resolution (30 meters), we can use 30m-resolution satellite imagery from Landsat, plus other features
- Goal is to learn mapping from (fine-resolution features) \rightarrow (fine-resolution SIF)
- However, during training, we only have **coarse-resolution SIF labels** (3km resolution) for supervision (no fine-resolution labels). So supervised super-resolution methods don't work
- We call this type of problem “**coarsely-supervised regression**”
 - Problem is highly under-determined; need regularization and prior knowledge




Existing baselines: averaging-based

- Existing baselines train by taking average of each feature across large tile.
- Spatial variation lost during training


Train: for each **large** tile, compute average of each variable. Train model to predict SIF from averages.



Test: for each **small region/pixel**, compute average of each band. Pass averages through model to predict SIF.




Our approach: Coarsely Supervised Regression U-Net (CSR-U-Net)



1. Apply augmentations to input tile
2. Pass transformed tile through U-Net to obtain predicted pixel SIFs
3. Take average SIF across valid pixels in tile
4. Compare predicted average SIF with true (coarse) average SIF; compute loss (error)
5. Backpropagate to adjust model parameters

Our approach: CSR-U-Net



NOTE - model does not look at fine-resolution SIF labels during training!

Regularization techniques: augmentations, multiplicative noise, early stopping

- Augmentations: flip/rotate, jigsaw, random subset
- **Multiplicative noise** - multiply all channels of entire image by a single random constant
 - Inspired by prior knowledge from vegetation indices, that the **ratio** between spectral bands can be more important than the absolute values
- Early stopping, based on fine-resolution validation set
 - Ensures similar farms have similar predictions. If we didn't do this, model can overfit (see below)




Figure 5: Example of overfitting. **Left:** ground-truth SIF map. **Right:** prediction by a U-Net that is over-trained. Note that the average tile SIFs are not too different, but the model tends to output extremely low and high values that do not reflect reality.



Datasets

Input features: Each input tile is of shape [24 features x 100 height x 100 width].

- Landsat reflectance bands
- FLDAS land data (temperature, rainfall, surface radiation)
- Land cover masks (corn, soybean, deciduous forest, grassland, etc)

Output SIF labels:

- During training, we use coarse SIF labels (gridded to 3km) from OCO-2 and CFIS
- During model selection and evaluation, we use a limited validation set of fine-resolution (30m) labels from CFIS to test accuracy of model's fine-resolution predictions

Total of 2,102 training tiles




Results

Results on predicting SIF for 30m pixels (in train tiles - coarse-resolution SIF was seen during training, but not fine-resolution SIF)

Method	NRMSE (lower better)	R^2 (higher better)
<i>Predict coarse</i>	<i>0.248</i>	<i>0.373</i>
Ridge Regression	0.213	0.537
Gradient Boosting	0.225	0.486
ANN	0.244 ± 0.006	0.396 ± 0.027
CSR-U-Net	0.196 ± 0.002	0.609 ± 0.007


Our method outperforms baselines (reducing NRMSE by 8% and increasing R^2 by 13%). Also true for completely unseen tiles, and across resolutions/crop types

Example output



Connection to “statistical downscaling” of climate models

- This task (inferring a fine-resolution map of a variable, given only coarse-resolution observations and other auxiliary data) is not exclusive to SIF
- Many related problems in climate science, including **downscaling climate models / climate variables**
 - Example: most global circulation models (GCMs) have a very coarse resolution.
 - Predicting at a finer resolution could enable more accurate understanding of climate impacts
- AI community should study this more!



Global climate model representation of California elevations (left) compared to LOCA

Image credit: USGS (public domain)



Conclusions

- We developed **Coarsely-Supervised Regression U-Net (CSR-U-Net)**, which can predict SIF at a super fine resolution (30m), even when only coarse-resolution (3km) SIF measurements are available
- Due to its localization properties, CSR-U-Net can figure out which farms within a large tile had higher and lower SIF.
- CSR-U-Net can avoid overfitting, thanks to techniques informed by prior knowledge, such as multiplicative noise and early stopping.
- Fine-resolution SIF estimates can facilitate improvements in crop yield prediction and monitoring.
- CSR-U-Net could be generalized to other related problems, such as predicting climate model variables (including precipitation, soil moisture, and evapotranspiration) at finer resolutions



References

Ariel Ortiz-Bobea, Toby R Ault, Carlos M Carrillo, Robert G Chambers, and David B Lobell. Anthropogenic climate change has slowed global agricultural productivity growth. *Nature Climate Change*, 11(4):306–312, 2021.

Liyin He, Troy Magney, Debsunder Dutta, Yi Yin, Philipp Köhler, Katja Grossmann, Jochen Stutz, Christian Dold, Jerry Hatfield, Kaiyu Guan, et al. From the ground to space: Using solar-induced chlorophyll fluorescence to estimate crop productivity. *Geophysical Research Letters*, 47(7):e2020GL087474, 2020

Bin Peng, Kaiyu Guan, Wang Zhou, Chongya Jiang, Christian Frankenberg, Ying Sun, Liyin He, and Philipp Köhler. Assessing the benefit of satellite-based solar-induced chlorophyll fluorescence in crop yield prediction. *International Journal of Applied Earth Observation and Geoinformation*, 90:102126, 2020.197

Magney, T. S., Bowling, D. R., Logan, B. A., Grossmann, K., Stutz, J., Blanken, P. D., ... Frankenberg, C. (2019). Mechanistic evidence for tracking the seasonality of photosynthesis with solar-induced fluorescence. *Proceedings of the National Academy of Sciences of the United States of America*, 116(24), 11640–11645. <https://doi.org/10.1073/pnas.1900278116>

Yang Song, Jing Wang, and Lixin Wang. Satellite solar-induced chlorophyll fluorescence reveals heat stress impacts on wheat yield in India. *Remote Sensing*, 12(20):3277, 2020.

Lifu Zhang, Na Qiao, Changping Huang, and Siheng Wang. Monitoring drought effects on vegetation productivity using satellite solar-induced chlorophyll fluorescence. *Remote Sensing*, 19911(4):378, 2019.200