

✔ Congratulations! You passed!

Grade received 100%

Latest Submission Grade 100%

To pass 80% or higher

Go to next item

1. Select the characteristic polynomial for the given matrix.

1 / 1 point

$$\begin{bmatrix} 2 & 1 \\ -3 & 6 \end{bmatrix}$$

☐

$$\lambda^2 - 8\lambda - 1$$

☒

$$\lambda^2 - 8\lambda + 15$$

☐

$$\lambda^2 + 8\lambda + 15$$

☐

$$\lambda^3 - 8\lambda + 15$$



Correct

Correct! $\lambda^2 - (2 + 6)\lambda + (2 * 6 - 1(-3)) = 0$

2. Select the eigenvectors for the previous matrix in Q1, as given below:

1 / 1 point

$$\begin{bmatrix} 2 & 1 \\ -3 & 6 \end{bmatrix}$$

☐ $\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

☐ $\begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}$

☒ $\begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

☐ $\begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

✓ **Correct**

Correct! You first find the eigenvalues for the given matrix: $\lambda = 5, \lambda = 3$.
Now you solve the equations using each of the eigenvalues.

For $\lambda = 5$, you have $\begin{cases} 2x + y = 5x \\ -3x + 6y = 5y \end{cases}$, which has solutions for
 $x = 1, y = 3$. Your eigenvector is $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$.

For $\lambda = 3$, you have $\begin{cases} 2x + y = 3x \\ -3x + 6y = 3y \end{cases}$, which has solutions for
 $x = 1, y = 1$. Your eigenvector is $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$.

3. Which of the following is an eigenvalue for the given identity matrix.

1 / 1 point

$$ID = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

☐

$\lambda = -1$

☐

$\lambda = 2$

☒

$\lambda = 1$

**Correct**

Correct! The eigenvalue for the identity matrix is always 1.

4. Find the eigenvalues of matrix $A \cdot B$ where:**1 / 1 point**

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Hint: What type of matrix is B ? Does it change the output when multiplied with A ? If not, focus only on one of the matrices to find the eigenvalues.

☐

$\lambda_1 = 4, \lambda_2 = 2$

☐

$\lambda_1 = 3, \lambda_2 = 1$



$$\lambda_1 = 4, \lambda_2 = 1$$



Eigenvalues cannot be determined.



Correct

Correct!

$$A \cdot B = \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 \cdot 1 + 2 \cdot 0 & 1 \cdot 0 + 2 \cdot 1 \\ 0 \cdot 1 + 4 \cdot 0 & 0 \cdot 0 + 4 \cdot 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix}$$

Since the second matrix is an identity matrix, you wouldn't need to solve the above multiplication since identity matrix does not change the result.

The eigenvalues of A are the roots of the characteristic equation $\det(A - \lambda I) = 0$.

By solving $\lambda^2 - 5\lambda + 4 = 0$, you get $\lambda_1 = 4, \lambda_2 = 1$.

5. Select the eigenvectors, using the eigenvalues you found for the above matrix A·B in Q2.

1 / 1 point



$$\vec{v}_1 = (2, 3); \vec{v}_2 = (1, 0)$$



$$\vec{v}_1 = (2, 3); \vec{v}_2 = (2, 3)$$



$$\vec{v}_1 = (2, 0); \vec{v}_2 = (1, 0)$$



$$\vec{v}_1 = (1, 3); \vec{v}_2 = (1, 0)$$



Correct

Correct!

For $\lambda = 4$, you have $\begin{cases} x + 2y = 4x \\ 0x + 4y = 4y \end{cases}$, which has solutions for $x = 2, y = 3$. Your eigenvector \vec{v}_1 is $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$.

For $\lambda = 1$, you have $\begin{cases} x + 2y = x \\ 0x + 4y = y \end{cases}$, which has solutions for $x = 1, y = 0$. Your eigenvector \vec{v}_2 is $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

6. For which value of a (in real numbers) does the matrix have real eigenvalues?

1 / 1 point

$$\begin{bmatrix} 2 & a \\ -1 & 1 \end{bmatrix}$$



$$a \leq 1/2$$



$$a \leq 1/4$$



$$a < 1/4$$



$$a \geq 1/4$$

**Correct**

Correct!

7.

Which of the vectors span the matrix $W = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 5 \\ 3 & -2 & -1 \end{bmatrix}$?

1 / 1 point

$$V_1 = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \quad V_2 = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix} \quad V_3 = \begin{bmatrix} 3 \\ -2 \\ -1 \end{bmatrix}$$



$$\vec{V}_1 = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} \quad \vec{V}_2 = \begin{bmatrix} 3 \\ 2 \\ -2 \end{bmatrix} \quad \vec{V}_3 = \begin{bmatrix} 0 \\ 5 \\ -1 \end{bmatrix}$$

**Correct**

Correct! There are linearly independent columns that span the matrix, which individually form three vectors $\vec{V}_1, \vec{V}_2, \vec{V}_3$. These vectors span the matrix W .

8. Given matrix **P** select the answer with the correct eigenbasis.

1 / 1 point

$$P = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$$

Hint: First compute the eigenvalues, eigenvectors and contrast the eigenbasis matrix with the spanning eigenvectors.



$$\text{Eigenbasis} = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$



$$\text{Eigenbasis} = \begin{bmatrix} 0 & -1 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$



$$\text{Eigenbasis} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

**Correct**

Correct! After solving the characteristic equations to find the eigenvalues, you should get $\lambda_1 = 1$ and $\lambda_2 = 2$.

The eigenvector for $\lambda_1 = 1$ is $\vec{V}_1 = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}$.

The eigenvectors for $\lambda_2 = 2$ are $\vec{V}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$, $\vec{V}_3 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$.

The eigenvectors form the eigenbasis: $\begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$

9. Select the characteristic polynomial for the given matrix.

1 / 1 point

$$\begin{bmatrix} 3 & 1 & -2 \\ 4 & 0 & 1 \\ 2 & 1 & -1 \end{bmatrix}$$

☐ $-\lambda^2 + 2\lambda^3 + 4\lambda - 5$

☐ $-\lambda^3 + 2\lambda^2 + 9$

☐ $\lambda^3 + 2\lambda^2 + 4\lambda - 5$

☒ $-\lambda^3 + 2\lambda^2 + 4\lambda - 5$

✓ **Correct**

Correct! The characteristic polynomial of a matrix A is given by

$$f(\lambda) = \det(A - \lambda I).$$

First, you find the following:

$$\begin{bmatrix} 3 & 1 & -2 \\ 4 & 0 & 1 \\ 2 & 1 & -1 \end{bmatrix} \cdot \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Now you compute the determinant of the result:

$$\det \begin{pmatrix} 3 - \lambda & 1 & -2 \\ 4 & -\lambda & 1 \\ 2 & 1 & -1 - \lambda \end{pmatrix} = -\lambda^3 + 2\lambda^2 + 4\lambda - 5$$

10. You are given a non-singular matrix A with real entries and eigenvalue i .

1 / 1 point

Which of the following statements is correct?

- ☒ $1/i$ is an eigenvalue of A^{-1} .
- ☐ i is an eigenvalue of $A^{-1} \cdot A \cdot I$.
- ☐ i is an eigenvalue of $A^{-1} + A$.

☒ **Correct**

Correct! You know that the eigenvalues of a matrix A are the solutions of its characteristic polynomial equation $\det(A - \lambda I) = 0$.