## Contents

1 Section 10.2															1	1							
	1.1	10.2.1																				1	_
	1 2	10.2.3																				1	

## 1 Section 10.2

Laplace transform of the second derivative of a function

$$\mathcal{L}(f'') = s^2 \mathcal{L}(f) - sf(0) - f'(0) \tag{1}$$

## 1.1 10.2.1

Solve the following differential equation by Laplace transforms. The function is subject to the given conditions.

$$y'' + 64y = 0, y(0) = 0, y'(0) = 1$$

$$\mathcal{L}(y'') + 64\mathcal{L}(y) = \mathcal{L}(0)$$

$$[s^{2}\mathcal{L}(y) - sy(0) - y'(0)] + 64\mathcal{L}(y) = \mathcal{L}(0)$$

$$s^{2}\mathcal{L}(y) - 0 - 1 + 64\mathcal{L}(y) = \mathcal{L}(0)$$

$$\mathcal{L}(y)(s^{2} + 64) = 1$$

$$\mathcal{L}(y) = \frac{1}{s^{2} + 64}$$

$$y = \mathcal{L}^{-1}\left(\frac{1}{s^{2} + 64}\right)$$

$$y = \frac{1}{8}\sin(8t)$$

## 1.2 10.2.3

Solve the following differential equation by Laplace transforms. The function is subject to the given conditions.

$$y'' + 9y = 9, y(0) = 1, y'(0) = 3$$

$$\mathcal{L}(y'') + 9\mathcal{L}(y) = \mathcal{L}(9)$$

$$s^{2}\mathcal{L}(y) + sy(0) + y'(0) + 9\mathcal{L}(y) = \frac{9}{s}$$

$$s^{2}\mathcal{L}(y) + s + 1 + 9\mathcal{L}(y) = \frac{9}{s}$$

$$\mathcal{L}(y)(s^{2} + 9) = \frac{9}{s} - s - 1$$

$$\mathcal{L}(y) = \frac{-s^{2} - s + 9}{s(s^{2} + 9)}$$

$$y = \mathcal{L}^{-1} \left( \frac{-2s - 1}{s^2 + 9} \right) + \mathcal{L}^{-1} \left( \frac{1}{s} \right)$$
$$y =$$

$$mx'' + kx = 0$$

$$m(s^{2}\mathcal{L}(x) - sx(0) - x'(0)) + k\mathcal{L}(x) = 0$$

$$ms^{2}\mathcal{L}(x) - smx(0) - mx'(0) + k\mathcal{L}(x) = 0$$

$$\mathcal{L}(x)(ms^{2} + k) = smx(0) + mx'(0)$$

$$\mathcal{L}(x) = \frac{smx(0) + mx'(0)}{ms^{2} + k}$$

$$\mathcal{L}(x) = \frac{smx(0)}{ms^{2} + k} + \frac{mx'(0)}{ms^{2} + k}$$

$$\mathcal{L}(x) = \frac{ms}{ms^{2} + k} + \frac{m}{ms^{2} + k}, \quad x(0) = 1, x'(0) = 1$$

$$\mathcal{L}(x) = \frac{s}{s^{2} + \frac{k}{m}} + \frac{1}{s^{2} + \frac{k}{m}}$$

$$x = \mathcal{L}^{-1}\left(\frac{s}{s^{2} + \frac{k}{m}}\right) + \mathcal{L}^{-1}\left(\frac{1}{s^{2} + \frac{k}{m}}\right)$$

$$x = \cos\left(\sqrt{\frac{k}{m}t}\right) + \sqrt{\frac{m}{k}}\sin\left(\sqrt{\frac{k}{m}t}\right)$$

$$x = \cos\left(\omega t\right) + \omega^{-1}\sin\left(\omega t\right)$$