

Contents

| | |
|--|----------|
| 1 Chapter 02 - Basic Structures | 1 |
| 1.1 Sets | 1 |
| 1.1.1 Definition 2 | 1 |
| 1.1.2 Definition 3 | 1 |
| 1.1.3 Definition 4 | 2 |
| 1.1.4 Countable and Uncountable Sets | 2 |
| 1.1.5 Example | 2 |
| 1.1.6 Definition 6 | 2 |
| 1.2 Set Operations | 2 |
| 1.2.1 Definition 1 | 2 |
| 1.2.2 Definition 2 | 3 |
| 1.2.3 Definition 3 | 3 |
| 1.2.4 Definition 4 | 3 |
| 1.2.5 Definition 5 | 3 |
| 1.2.6 Proof | 3 |
| 1.2.7 Proof: | 4 |
| 1.2.8 Union | 4 |
| 1.2.9 Intersection | 4 |
| 1.3 Functions | 4 |
| 1.3.1 Example | 4 |
| 1.3.2 Properties | 5 |
| 1.3.3 Example 2.3.24 | 5 |
| 1.3.4 Example 2.3.73.b | 5 |
| 1.4 Sequences and Summations | 6 |
| 1.5 Cardinality of Sets | 6 |
| 1.6 Matrices | 6 |

1 Chapter 02 - Basic Structures

1.1 Sets

\in : belong to, is in

1.1.1 Definition 2

Two sets are equal if and only if they have the same elements. Therefore, if A and B are sets, then A and B are equal if and only if $\forall x (x \in A \iff x \in B)$. We write $A = B$ if A and B are equal sets.

1.1.2 Definition 3

The set A is also a subset of B , and B is a superset of A , if and only if every element of A is also an element of B . We use the notation $A \subseteq B$ to indicate that A is a subset of the set B . If, instead, we want to stress that B is a

superset of A , we use the equivalent notation $B \supseteq A$. (So, $A \subseteq B$ and $B \supseteq A$ are equivalent statements.)

1.1.3 Definition 4

Let S be a set. If there are exactly n distinct elements in S where n is a nonnegative integer, we say that S is a finite set and that n is the cardinality of S . The cardinality of S is denoted by $|S|$.

1.1.4 Countable and Uncountable Sets

- Countable
 - \mathbb{N}
 - \mathbb{Z}
 - \mathbb{Q}
- Uncountable
 - \mathbb{R}
 - \mathbb{C}

Let $S_0 = \{x\}$, and $S_1 = \{\{x\}\}$.

$$S_0 \neq S_1 \tag{1}$$

1.1.5 Example

1. List the members of these sets.

a) $\{x \mid x \text{ is a real number such that } x^2 = 1\}$

$$S = \{x \in \mathbb{R} \mid x^2 = 1\}$$

b) $\{x \mid x \text{ is a positive integer less than } 12\}$

$$S = \{x \in \mathbb{R} \mid 0 \leq x < 12\}$$

1.1.6 Definition 6

Given a set S , the power set of S is the set of all subsets of the set S . The power set of S is denoted by $\mathcal{P}(S)$.

1.2 Set Operations

1.2.1 Definition 1

Let A and B be sets. The union of the sets A and B , denoted by $A \cup B$, is the set that contains those elements that are either in A or in B , or in both.

$$A \cup B = \{x \in U \mid (x \in A) \vee (x \in B)\} \tag{2}$$

1.2.2 Definition 2

Let A and B be sets. The intersection of the sets A and B , denoted by $A \cap B$, is the set containing those elements in both A and B .

$$A \cap B = \{x \in U \mid (x \in A) \wedge (x \in B)\} \quad (3)$$

1.2.3 Definition 3

Two sets are disjoint if their intersection is the empty set.

1.2.4 Definition 4

Let A and B be sets. The difference of A and B , denoted by $A - B$, is the set containing those elements that are in A but not in B . The difference of A and B is also called the complement of B with respect to A .

1.2.5 Definition 5

Let U be the universal set. The complement of the set A , denoted by \bar{A} , is the complement of A with respect to U . Therefore, the complement of the set A is $U - A$.

1.2.6 Proof

Let A, B be sets from U . Show that $A \subseteq B$ if and only if $\bar{B} \subseteq \bar{A}$.

Proof:

- For “ \implies ”

Given $A \subseteq B$, need to show $\bar{B} \subseteq \bar{A}$. Then $\forall x \in \bar{B}$, we have $x \in \bar{A}$.

By contrapositive we have

$$\begin{aligned} \neg(x \in B) &\implies \neg(x \in A) \\ x \notin B &\implies x \notin A, \quad \bar{B} \subseteq \bar{A} \end{aligned}$$

- For “ \impliedby ”

Given $\bar{B} \subseteq \bar{A}$, we have $\forall y \in \bar{B}, y \in \bar{A}$ then the contrapositive is

$$\begin{aligned} \neg(y \in \bar{A}) &\implies \neg(y \in \bar{B}) \\ y \in A &\implies y \in B \end{aligned}$$

1.2.7 Proof:

Use the identities to show that $\overline{(A \cup B)} \cap \overline{(B \cup C)} \cap \overline{(A \cup C)} = \overline{A} \cap \overline{B} \cap \overline{C}$

Proof:

$$\begin{aligned}
\overline{A} \cap \overline{B} \cap \overline{C} &= (\overline{A} \cap \overline{B}) \cap (\overline{B} \cap \overline{C}) \cap (\overline{A} \cap \overline{C}) \\
&= \overline{A} \cap (\overline{B} \cap \overline{B}) \cap (\overline{C} \cap \overline{C}) \cap \overline{A} \\
&= \overline{A} \cap \overline{B} \cap \overline{C}
\end{aligned}$$

1.2.8 Union

The union of a collection of sets is the set that contains those elements that are members of at least one set in the collection.

$$A_1 \cup A_2 \cup \cdots \cup A_n = \bigcup_{i=1}^n A_i$$

1.2.9 Intersection

The intersection of a collection of sets is the set that contains those elements that are members of all the sets in the collection.

$$A_1 \cap A_2 \cap \cdots \cap A_n = \bigcap_{i=1}^n A_i$$

1.3 Functions

- Sets: A , B , C , domain, codomain, range
- Relations: functions (f, g, h)
- Elements: image, preimage

1.3.1 Example

$$f_1 : \mathbb{R} \rightarrow \mathbb{R} \text{ as } y = f_1(x) = 3x - 2$$

$$f_2 : \mathbb{R} \rightarrow \mathbb{R} \text{ as } y = e^x$$

$$f_3 : \mathbb{R} \rightarrow \mathbb{R} \text{ as } y = \sqrt{x}$$

$$f : A \rightarrow B$$

$$\text{range}(f) = \{y \in B \mid \forall x \in S \subseteq A\}$$

1.3.2 Properties

1) f is injective – one-to-one

2) f is surjective –

3) f is bijective if 1) and 2)

Let $f : A \rightarrow B$ be a function.

We say f is injective if

$$(x_1 \neq x_2 \implies f(x_1) \neq f(x_2)) \iff (f(x_1) = f(x_2) \implies x_1 = x_2)$$

We say f is surjective if $\forall y \in B, \exists x \in A$ such that $y = f(x) \iff \text{range}(f) = B$

We say f is bijective if f is injective and surjective.

Monotonic Function: We say f is increasing (strictly increasing) if $x_1 > x_2 \implies f(x_1) > f(x_2)$

1.3.3 Example 2.3.24

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and let $f(x) > 0$ for all $x \in \mathbb{R}$. Show that $f(x)$ is strictly increasing if and only if the function $g(x) = \frac{1}{f(x)}$ is strictly decreasing.

$$\begin{aligned} &\rightarrow \frac{1}{f(x_1)} < \frac{1}{f(x_2)} \\ &\rightarrow f(x_1) \cdot f(x_2) \left(\frac{1}{f(x_1)} \right) < f(x_1) \cdot f(x_2) \left(\frac{1}{f(x_2)} \right) \\ &\rightarrow f(x_2) < f(x_1) \end{aligned}$$

where the first implication comes from the fact that $f(x) > 0, \forall x \in \mathbb{R}$.

1.3.4 Example 2.3.73.b

Prove or disprove each of these statements about the floor and ceiling functions.

$$\lfloor 2x \rfloor = 2 \lfloor x \rfloor$$

Consider $x = 1.6$

$$\begin{aligned} \lfloor 2x \rfloor &= \lfloor 2 \cdot 1.6 \rfloor = \lfloor 3.2 \rfloor = 3 \\ 2 \lfloor x \rfloor &= 2 \lfloor 1.6 \rfloor = 2(1) = 2 \end{aligned}$$

Hence $\lfloor 2x \rfloor = 2 \lfloor x \rfloor$ for all $x \in \mathbb{R}$ is false.

1.4 Sequences and Summations

1.5 Cardinality of Sets

1.6 Matrices