

Sage - *Salvia officinalis* - Oral

Source: [https://webprod.hc-sc.gc.ca/nhp/nd-bdipsn/atReq?atid=sage.or2\(=eng](https://webprod.hc-sc.gc.ca/nhp/nd-bdipsn/atReq?atid=sage.or2(=eng)

Extracted: 2025-08-26T06:35:54.915745

SAGE - SALVIA OFFICINALIS Oral Help on accessing alternative formats, such as Portable Document Format (PDF), Microsoft Word and PowerPoint (PPT) files, can be obtained in the alternate format help section. (PDF Version - 90 KB) This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredient. Notes Text in parentheses is additional optional information which can be included on the label at the applicant's discretion. The solidus (/) indicates that the terms and/or statements are synonymous. Either term or statement may be selected by the applicant on the label. Date January 10, 2025 Proper name(s), Common name(s), Source information Table 1. Proper name(s), Common name(s), Source information Proper name(s) Common name(s) Source information Source material(s) Part(s) Preparation(s) *Salvia officinalis* Common sage Dalmatian sage Garden sage Sage *Salvia officinalis* Leaf Dry References: Proper name: USDA 2024; Gardner and McGuffin 2013; Common names: USDA 2024; Gardner and McGuffin 2013; Source information: Blumenthal et al. 2000; BHP 1996; Cook 1869. Route of Administration Oral Dosage Form(s) This monograph excludes foods or food-like dosage forms as indicated in the Compendium of Monographs Guidance Document. Acceptable dosage forms for oral use are indicated in the dosage form drop-down list of the web-based Product Licence Application form for Compndial applications. Use(s) or Purpose(s) Source of antioxidants/Provides antioxidants (Bradley 2006; Lima et al. 2005). Source of antioxidants/Provides antioxidants that help fight/protect (cell) against/reduce (the oxidative effect of/the oxidative damage caused by/cell damage caused by) free radicals (Bradley 2006; Lima et al. 2005). (Traditionally) used in Herbal Medicine (as a carminative) to help provide relief of upset stomach and flatulence (flatulent dyspepsia) (EMA 2016; Godfrey et al. 2010; Bradley 2006; Mills and Bone 2005; BHP 1996; Culbreth 1927; Felter 1922). (Traditionally) used in Herbal Medicine (as a diaphoretic) to increase perspiration/sweating (PDR 2007; Culbreth 1927; Felter 1922; Cook 1869). (Traditionally) used in Herbal Medicine (as an antihidrotic) to reduce hyperhidrosis/excessive sweating or perspiration (EMA 2016; Bradley 2006; Mills and Bone 2005; BHP 1996; Culbreth 1927; Felter 1922). Used in Herbal Medicine to help reduce hot flushes/hot flashes/night sweats associated with menopause (Romm 2010; Bradley 2006; Mills and Bone 2000). (Traditionally) used in Herbal Medicine (as an antigalactagogue) to stop the production of breast milk (Godfrey et al. 2010; Wichtl 2004; Yarnell et al. 2003; Felter and Lloyd 1983; Culbreth 1927; Cook 1869). Notes The above uses can be combined on the product label if from the same traditional or non-traditional system of medicine (e.g., Traditionally used in Herbal Medicine to reduce excessive sweating or perspiration and to stop the production of breast milk). For multi-ingredient products: To prevent the product from being represented as a "traditional medicine", any indicated traditional use claim must refer to the specific medicinal ingredient(s) and recognized traditional system of medicine from which the claim originates when 1) both traditional and modern claims are present or 2) when claims originate from multiples systems of traditional medicine (e.g. Sage is traditionally used in Herbal Medicine to increase sweating). When ALL of the medicinal ingredients (MIs) in the product are used within the SAME identified system of traditional medicine AND the product makes ONLY traditional claims, listing of MIs in the traditional claim(s) is not required. Dose(s) Subpopulation(s) Adults 18 years and older Quantity(ies) Antioxidant Methods of preparation: Dry 1 , Non-Standardized Aqueous Extracts (Dry extract, Infusion, Infusion concentrate) Not to exceed 12 grams of dried leaf, per day (Bradley 2006; Mills et Bone 2005; Blumenthal et al. 2000; BHP 1996; Culbreth 1927). 1 Note: Dried leaf should be prepared as an infusion (see direction for use). Method of preparation: Non-Standardized Ethanolic Extracts (Dry extract, Tincture) Not to exceed 2.25 grams of dried leaf, per day (EMA 2016; Mills and Bone 2005; ESCOP 2003). Method of preparation: Fluid extract Not to exceed 6 grams of dried leaf, per day (EMA 2016; Bradley 2006; Mills and Bone 2005; ESCOP 2003; BHP 1996). Other uses Methods of preparation: Dry 1 , Non-Standardized Aqueous Extracts (Dry extract, Infusion, Infusion concentrate) 1 - 12 grams of dried leaf, per day (Mills and Bone 2005; Blumenthal et al. 2000; BHP 1996; Culbreth 1927). 1 Note: Dried leaf should be prepared as an infusion (see direction for use). Methods of preparation: Non-Standardized Ethanolic Extracts (Dry extract, Tincture) 0.3 - 2.25 grams dry leaf, per day (EMA 2016; Mills and Bone 2005; ESCOP 2003). Method of preparation: Fluid extract 1 - 6 grams of dried leaf, per day (EMA 2016; Bradley 2006; Mills and Bone 2005; ESCOP 2003; BHP

1996). Direction(s) for use Night sweats Take 1 hour before bedtime (EMA 2016). Dried leaves - All uses Pour hot boiled water on dried leaf and steep. Dried leaves - Antihidrotic (To reduce sweating,)* let infusion cool before drinking (Bradley 2006; Mills and Bone 2005; BHP 1996; Culbreth 1927; Felter 1922). Dried leaves - Diaphoretic (To increase sweating,)* drink infusion while still warm (PDR 2007; Culbreth 1927; Felter 1922; Cook 1869). *Note: Information in brackets is required on the label, when both uses to reduce and increase sweating are listed on the label. Duration(s) of Use Ask a health care practitioner/health care provider/health care professional/doctor/physician for use beyond 4 weeks (Gardner and McGuffin 2013; Wichtl 2004; ESCOP 2003). Risk Information Caution(s) and warning(s) Carminative, Diaphoretic, Antihidrotic, Antigalactagogue Ask a health care practitioner/health care provider/health care professional/doctor/physician if symptoms persist or worsen. Contraindication(s) All products Do not use if you are pregnant or breastfeeding (Gardner and McGuffin 2013; Bradley 2006; Mills and Bone 2005). Do not use if you have a seizure disorder such as epilepsy (Gardner and McGuffin 2013; Bradley 2006; Mills and Bone 2005). Known adverse reaction(s) No statement required. Non-medicinal ingredients Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database. Storage conditions Must be established in accordance with the requirements described in the Natural Health Products Regulations . Specifications The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal ingredient must comply with the requirements outlined in the NHPID. Sage contains thujone. For adults, the upper limit for total daily intake of thujone from health products is 6 mg. Product licence applications for oral products should include a copy of a certificate of analysis or any other equivalent document demonstrating that the thujone content of a daily dose of the product is acceptable. Because thujone content of the herbal materials can vary, the thujone content should be determined for each batch during production of the product. Example of Product Facts: Consult the Guidance Document, Labelling of Natural Health Products for more details. References Cited BHP 1996: British Herbal Medicine Association's Scientific Committee. British Herbal Pharmacopoeia. Bournemouth (GB): The British Herbal Medicine Association; 1996. Blumenthal M, Goldberg A, Brinckmann J. Herbal Medicine: Expanded Commission E Monographs. Boston (MA): American Botanical Council. 2000. Bradley PR, editor. 2006: Bradley PR, editor. British Herbal Compendium Volume 2: A Handbook of Scientific Information on Widely Used Plant Drug-Companion to the British Herbal Pharmacopoeia. Bournemouth (GB): British Herbal Medicine Association; 2006. Cook WMH. The Physio-Medical Dispensatory: A Treatise on Therapeutics, Materia Medica, and Pharmacy, in Accordance with the Principles of Physiological Medication. Cincinnati (OH): WM.H. Cook; 1869. Reprint version by Medical Herbalism and medherb.com, Boulder (CO). [Accessed 2024 March 12]. Available from: <http://www.henriettesherbal.com/eclectic/cook/index.html> Culbreth DMR. A Manual of Materia Medica and Pharmacology. 7 th edition Philadelphia (PA): Lea & Febiger; 1927. Abridged and alphabetized by Michael Moore, director. The Southwest School of Botanical Medicine, Bisbee (AZ). [Accessed 2024 March 12]. Available from: <http://www.swsbm.com/ManualsOther/Culbreth.html> ESCOP 2003: E/S/C/O/P Monographs: The Scientific Foundation for Herbal Medicinal Products. 2 nd edition. Completely revised and Expanded. Exeter (GB): ESCOP, the European Scientific Cooperative on Phytotherapy in collaboration with Georg Thieme Verlag and Thieme; 2003. EMA 2016: European Union herbal monograph on *Salvia officinalis* L., folium. Final. 20 September 2016. London (GB): European Medicines Agency: Committee on Herbal Medicinal Products (HMPC). [Accessed 2024 March 12]. Available from: https://www.ema.europa.eu/en/documents/herbal-monograph/final-european-union-herbal-monograph-salvia-officinalis-l-folium-revision-1_en.pdf Felter HW. The Eclectic Materia Medica, Pharmacology and Therapeutics. Cincinnati (OH): John K. Scudder; 1922. Reprint version by Michael Moore, The Southwest School of Botanical Medicine, Bisbee (AZ). [Accessed 2024 March 12]. Available from: <http://www.swsbm.com/FelterMM/Felters.html> Felter HW, Lloyd JU. King's American Dispensatory. Volume 2, 18 th edition. Sandy (OR): Eclectic Medical Publications; 1983 [Reprint of 1898 original]. Gardner Z, McGuffin M, editors. American Herbal Products Association's Botanical Safety Handbook. 2nd edition. Boca Raton (FL): Taylor and Francis Group; 2013. Godfrey A, Saunders PR, Barlow K, Gilbert C, Gowan M, Smith F. Principles and Practices of Naturopathic Botanical Medicine. Volume 1: Botanical Medicine Monographs. Toronto (ON): CCNM Press; 2010. Lima CF, Andrade PB, Seabra RM, Fernandes-Ferreira M, Pereira-Wilson C. The drinking of a *Salvia officinalis* infusion improves liver antioxidant status in mice and rats. *Journal of Ethnopharmacology* 2005;97(2):383-389. Mills S, Bone K. Principles and Practice of Phytotherapy. Toronto (ON): Churchill Livingstone; 2000. Mills S, Bone K. The Essential Guide to Herbal Safety. St. Louis (MO): Elsevier Churchill Livingstone; 2005. PDR 2007: Gruenwald J, Brendler T, Jaenicke C, editors. PDR for Herbal Medicines. 4 th edition. Montvale (NJ): Thomson Healthcare Inc; 2007. Romm A. Botanical Medicine for Women's Health. St. Louis (MO): Churchill Livingstone; 2010. USDA 2024: United States Department of Agriculture, Agricultural Research Service (USDA ARS), Germplasm Resources Information Network (GRIN) - Global. U.S. National Plant Germplasm System. [Accessed 2024 November 14]. Available from: <https://npgsweb.ars-grin.gov/gringlobal/taxon/taxonomysearch> Wichtl M, editor. Herbal Drugs and

Phytopharmaceuticals: A Handbook for Practice on a Scientific Basis. 3rd edition. Stuttgart (DE): Medpharm Scientific Publishers; 2004. Yarnell E, Abascal JD, Hooper CG. Clinical Botanical Medicine. Larchmont (NY): Mary Ann Leibert Inc.; 2003. References Reviewed ASHP 2005: American Society of Health-System Pharmacists. American Hospital Formulary Service (AHFS) Drug Information. Philadelphia (PA): Lippincott Williams and Wilkins; 2005. AU TGA 1999: Australian Therapeutic Goods Administration. TGA Approved Terminology for Medicines, Section 3 - Herbal Substances. July 1999. Symonston (AU): Australian Government Department of Health and Ageing, Therapeutic Goods Administration. [Accessed 2012 April 27]. Available from: <http://www.tga.gov.au/pdf/medicines-approved-terminologyherbal.pdf> Azevedo MF, Lima CF, Fernandes-Ferreira M, Almeida MJ, Wilson JM, Pereira-Wilson C. Rosmarinic acid, major phenolic constituent of Greek sage herbal tea, modulates rat intestinal SGLT1 levels with effects on blood glucose. *Molecular Nutrition & Food Research* 2011;55(S1): S15-S25. Barnes J, Anderson LA, Philipson JD. Herbal Medicines. 3rd edition. London (GB): The Pharmaceutical Press; 2007. Bartram T. Bartram's Encyclopaedia of Herbal Medicine: The definitive guide to the herbal treatment of diseases. New York: Marlowe & Company; 1998. Blumenthal M, Brusse WR, Goldberg A, Gruenwald J, Hall T, Riggins CW, Rister RS. The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines. Austin (TX): American Botanical Council in Cooperation with Integrative Medicine Communications; 1998. Bommer S, Klein P, Suter A. First time proof of sage's tolerability and efficacy in menopausal women with hot flushes. *Advances in Therapy* 2011;28(6):490-500. Brinker F. Herbal Contraindications and Drug Interactions: Plus Herbal Adjuncts With Medicines, expanded 4th edition. Sandy (OR): Eclectic Medical Publications; 2010. Canadian Nutrient File (CNF), 2023. Ottawa (ON): Food and Nutrition, Health Canada. [Accessed 2024 March 12]. Available from: <http://webprod3.hc-sc.gc.ca/cnf-fce/index-eng.jsp> Christensen KB, Jorgensen M, Kotowska D, Petersen RK, Kristiansen K, Christensen LP. Activation of the nuclear receptor PPAR γ by metabolites isolated from sage (*Salvia officinalis* L.). *Journal of Ethnopharmacology* 2010;132(1):127-133. derMarderosian A, Beutler JA, editors. The Review of Natural Products. "Sage: Date of Issue February 2010" St Louis (MO): Facts and Comparisons, Wolters Kluwer Health; Printed in 2008 and Updated to May 2012. Dos Santos-Neto LL, de Vilhena Toledo MA, Medeiros-Souza P, de Souza GA. The Use of Herbal Medicine in Alzheimer's Disease - A Systematic Review. *Evidence-Based Complementary and Alternative Medicine* 2006;3(4):441-445. Duke JA, Bogenschutz-Godwin MJ, duCellier J, Duke P-AK. Handbook of Medicinal Herbs. 2nd edition. Boca Raton (FL): CRC Press LLC; 2002. European Medicines Agency. Public statement on *Salvia officinalis* L., aetheroleum. London (GB): EMA Committee on Herbal Medicinal Products (HMPC); 2 February 2016. [Accessed 2024 March 12]. Available from: https://www.ema.europa.eu/en/documents/public-statement/public-statement-salvia-officinalis-l-aetheroleum_en.pdf European Scientific Cooperative of Phytotherapy. Monographs on the Medicinal Uses of Plant Drugs. Exeter (GB): European Scientific Cooperative on Phytotherapy; 1996. Facciola S. Cornucopia II: A source book of edible plants. Vista (CA): Kampong Publications; 1998. Felter HW, Lloyd JU. Kings American Dispensatory. 18th edition, 3rd revision. Cincinnati (OH): Ohio Valley Co.; 1898. Scanned version by Henriette Kress 1999-2011. [Accessed 2024 March 12]. Available from: <http://www.henriettesherbal.com/eclectic/kings/index.html> Grieve, M. A Modern Herbal, Volume 1-2. New York (NY): Dover Publications, 1971 [Reprint of 1931 Harcourt, Brace & Company publication]. Halicioglu O, Astarcioglu G, Yaprak I, Aydinlioglu H. Toxicity of *Salvia officinalis* in a newborn and a child: an alarming report. *Pediatric Neurology* 2011;45(4):259-260. Kennedy DO, Scholey AB. The psychopharmacology of European herbs with cognition enhancing properties. *Current Pharmaceutical Design* 2006;12(35):4613-4623. Kennedy DO, Pace S, Haskell C, Okello EJ, Milne A, Scholey AB. 2006. Effects of Cholinesterase Inhibiting Sage (*Salvia officinalis*) on Mood, Anxiety and Performance on a Psychological Stressor Battery. *Neuropsychopharmacology* 2006;31:845-852. Kennedy DO, Dodd FL, Robertson BC, Okello EJ, Reay JL, Scholey AB, Haskell CF. Monoterpenoid extract of sage (*Salvia lavandulaefolia*) with cholinesterase inhibiting properties improves cognitive performance and mood in healthy adults. *Journal of Psychopharmacology* 2011;25(8):1088-1100. Lachenmeier DW, Uebelacker M. Risk assessment of thujone in foods and medicines containing sage and wormwood - Evidence for a need of regulatory changes? *Regulatory Toxicology and Pharmacology* 2010;58:437-443. Lima CF, Azevedo MF, Araujo R, Fernandes-Ferreira M, Pereira-Wilson C. Metform-like effect of *Salvia officinalis* (common sage): is it useful in diabetes prevention? *British Journal of Nutrition* 2006; 92(2):326-333. Newall CA, Anderson LA, Phillipson JD. Herbal Medicines: A Guide for Health-care Professionals. London (GB): Pharmaceutical Press; 1996. Perry NS, Bollen C, Perry EK, Ballard C. *Salvia* for dementia therapy: review of pharmacological activity and pilot tolerability clinical trial. *Pharmacology, Biochemistry, and Behaviour* 2003;75(3):651-659. Remington JP, Woods HC, editors. 1918. The Dispensatory of the United States of America 20th > edition. Abridged; botanicals only. Scanned by Michael Moore, director, The Southwest School of Botanical Medicine, Bisbee (AZ). [Accessed 2024 March 12]. Available from: <http://www.swsbm.com/Dispensatory/USD-1918-complete.pdf> Sa CM, Ramos AA, Azevedo MF, Lima CF, Fernandes-Derreira M, Pereira-Wilson C. Sage tea drinking improves lipid profile and antioxidant defences in humans. *International Journal of Molecular Sciences* 2009;10(9):3937-3950. Swanston-Flatt SK,

Day C, Flatt PR, Gould BJ, Bailey CJ. Diabetes Research 1989;10(2):69-73. Tildesley NTJ, Kennedy DO, Perry EK, Ballard CG, Savelev S, Wesnes KA, Scholey AB. Salvia lavandulaefolia (Spanish Sage) enhances memory in healthy young volunteers. Pharmacology, Biochemistry and Behavior 2003;75:669-674. Tildesley NTJ, Kennedy DO, Perry EK, Ballard CG, Wesnes KA, Scholey AB. Positive modulation of mood and cognitive performance following administration of acute doses of Salvia lavandulaefolia essential oil to healthy young volunteers. Physiology & Behavior 2005;83:699-709. Umezu T. Evaluation of the effects of plant-derived essential oils on central nervous system function using discrete shuttle-type conditioned avoidance response in mice. Phytotherapy Research 2012;26(6):884-891. Wake G, Court J, Pickering A, Lewis R, Wilkins R, Perry E. CNS acetylcholine receptor activity in European medicinal plants traditionally used to improve failing memory. Journal of Ethnopharmacology 2000; 69:105-114. Walch SG, Kuballa T, Stühlinger W, Lachenmeier DW. Determination of the biologically active flavour substances thujone and camphor in foods and medicines containing sage (Salvia officinalis L.). Chemistry Central Journal 2011;5:44. Weiss RF. Fintelmann VF. Herbal Medicine. 2 nd edition, revised and expanded. Stuggart (NY): Thieme; 2000. World Health Organization. International Programme on Chemical Safety. Toxicological Evaluation of Certain Food Additives: WHO Food Additives Series 16. Geneva (CH): World Health Organization; 1981. [Accessed 2024 March 12]. Available from: <http://www.inchem.org/documents/jecfa/jecmono/v16je01.htm> World Health Organization. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives. Thujone. [Accessed 2024 March 12]. Available from: http://www.inchem.org/documents/jecfa/jecval/jec_2273.htm World Health Organization. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives. Isothujone. [Accessed 2024 March 12]. Available from: http://www.inchem.org/documents/jecfa/jecval/jec_2273.htm Report a problem on this page Date modified: 2019-03-01

DOSAGE FORM(S)

Acceptable dosage forms for oral use are indicated in the dosage form drop-down list of the web-based Product Licence Application form for Compendial applications.

DOSE(S)

Umezu T. Evaluation of the effects of plant-derived essential oils on central nervous system function using discrete shuttle-type conditioned avoidance response in mice. Phytotherapy Research 2012;26(6):884-891. Wake G, Court J, Pickering A, Lewis R, Wilkins R, Perry E. CNS acetylcholine receptor activity in European medicinal plants traditionally used to improve failing memory. Journal of Ethnopharmacology 2000; 69:105-114. Walch SG, Kuballa T, Stühlinger W, Lachenmeier DW. Determination of the biologically active flavour substances thujone and camphor in foods and medicines containing sage (Salvia officinalis L.). Chemistry Central Journal 2011;5:44. Weiss RF. Fintelmann VF. Herbal Medicine. 2nd edition, revised and expanded. Stuggart (NY): Thieme; 2000. World Health Organization. International Programme on Chemical Safety. Toxicological Evaluation of Certain Food Additives: WHO Food Additives Series 16. Geneva (CH): World Health Organization; 1981. [Accessed 2024 March 12]. Available from: <http://www.inchem.org/documents/jecfa/jecmono/v16je01.htm> World Health Organization. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives. Thujone. [Accessed 2024 March 12]. Available from: http://www.inchem.org/documents/jecfa/jecval/jec_2273.htm World Health Organization. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives. Isothujone. [Accessed 2024 March 12]. Available from: http://www.inchem.org/documents/jecfa/jecval/jec_2273.htm

RISK INFORMATION

Caution(s) and warning(s) Carminative, Diaphoretic, Antihidrotic, Antigalactagogue Ask a health care practitioner/health care provider/health care professional/doctor/physician if symptoms persist or worsen. Contraindication(s) All products Do not use if you are pregnant or breastfeeding (Gardner and McGuffin 2013; Bradley 2006; Mills and Bone 2005). Do not use if you have a seizure disorder such as epilepsy (Gardner and McGuffin 2013; Bradley 2006; Mills and Bone 2005). Known adverse reaction(s) No statement required.

NON-MEDICINAL INGREDIENTS

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database.

STORAGE CONDITION(S)

Must be established in accordance with the requirements described in the Natural Health Products Regulations.

SPECIFICATIONS

The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal ingredient must comply with the requirements outlined in the NHPID. Sage contains thujone. For adults, the upper limit for total daily intake of thujone from health products is 6 mg. Product licence applications for oral products should include a copy of a certificate of analysis or any other equivalent document demonstrating that the thujone content of a daily dose of the product is acceptable. Because thujone content of the herbal materials can vary, the thujone content should be determined for each batch during production of the product.

Proper name(s)	Common name(s)	Source information		
Source material(s)	Part(s)	Preparation(s)		
Salvia officinalis	Common sageDalmatian sageGarden sageSage	Salvia officinalis	Leaf	Dry