

# Wart Remover

Source: [https://webprod.hc-sc.gc.ca/nhp/nd-bdipsn/atReq?atid=wart\\_remover\\_anti\\_verrues&lang=eng](https://webprod.hc-sc.gc.ca/nhp/nd-bdipsn/atReq?atid=wart_remover_anti_verrues&lang=eng)

Extracted: 2025-08-26T06:38:24.323299

WART REMOVER Help on accessing alternative formats, such as Portable Document Format ( PDF ), Microsoft Word and PowerPoint ( PPT ) files, can be obtained in the alternate format help section. (PDF Version - 85 KB) This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredient. Notes Text in parentheses is additional optional information which can be included on the PLA and product label at the applicant's discretion. The solidus (/) indicates that the terms and/or statements are synonymous. Either term or statement may be selected by the applicant. Date July 1, 2019 Proper name(s), Common name(s), Source material(s) Table 1. Proper name(s), Common name(s), Source material(s) Proper name(s) Common name(s) Source ingredient(s) 1 Common name(s) 2-Hydroxybenzoic acid Salicylic acid Salicylic acid 1 The ingredient must be pharmacopoeial grade. References: Proper name: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Common name: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Source ingredient: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001. Table 2. Proper name(s), Common name(s), Source material(s) - Complementary ingredient (safety only) Proper name(s) Common name(s) Source ingredient(s) 1 Common name(s) 2-Hydroxy-2-methylacetic acid DL-Lactic acid DL-Lactic acid Lactic acid Lactic acid 1 The ingredient must be pharmacopoeial grade. References: Proper names: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Common names: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Source ingredient: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001. Route of Administration Topical (FDA 1990) Dosage Form(s) Collodion-like solution; Karaya gum, glycol plaster; Plaster (FDA 1990). Use(s) or Purpose(s) Wart remover (FDA 1990). For the removal of common warts (FDA 1990). For the removal of plantar warts on the bottom of the feet (FDA 1990). Common and/or plantar wart treatment (FDA 1990). Dose(s) Subpopulation(s) Children 2-11 years, Adolescents 12-17 years and Adults 18 years and older Quantity(ies) Table 3. Doses of Salicylic acid associated with dosage forms Dosage forms Doses Collodion-like solution 5 - 17% Karaya gum, glycol plaster 15% Other types of plasters 12 - 40% Reference: FDA 1990. Complementary Ingredient (Safety only) Not to exceed 30% of lactic acid (Fiume 1998) Direction(s) for use Products formulated in a collodion-like solution (FDA 1990) Wash affected area ( Optional: May soak wart in warm water for five minutes) and dry thoroughly. Apply one drop at a time to sufficiently cover each wart and let dry. Do not apply to surrounding skin. Repeat procedure once or twice daily as needed (until wart is removed). Products formulated in a karaya gum, glycol plaster (FDA 1990) Wash affected area ( Optional: May soak wart in warm water for five minutes) and dry thoroughly. Gently smooth wart surface with emery file provided (FDA 1994). Cut plaster to fit wart (if appropriate). Apply a drop of warm water to the wart, keeping the surrounding skin dry (FDA 1994). Apply medicated plaster at bedtime, leave in place for at least 8 hours. Remove and discard plaster in morning. Repeat every 24 hours as needed (until wart is removed). Products formulated with other types of plaster (FDA 1990) Wash affected area ( Optional: May soak wart in warm water for five minutes) and dry thoroughly. Cut plaster to fit wart (if appropriate). Apply medicated plaster. Repeat every 48 hours as needed (until wart is removed). Duration(s) of Use Do not use this product beyond 12 weeks (FDA 1990). Risk Information Caution(s) and warning(s) All products For external use only (FDA 1990). Consult a health care practitioner/health care provider/health care professional/doctor/ physician if discomfort persists (FDA 1990). Products formulated in a collodion-like solution only Extremely flammable or flammable or combustible (FDA 1990). Keep away from fire or flame (FDA 1990). Cap bottle tightly and store at room temperature, in an upright position, away from heat (FDA 1990). Flush with water for 15 minutes if product gets into the eye (FDA 1990). Avoid inhaling vapours (FDA 1990). Contraindication(s) All products Do not use this product on irritated or reddened skin or any area that is infected (FDA 1990). Do not use this product if you have diabetes or poor blood circulation (FDA 1990). Do not use this product on moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes (FDA 1990). Known adverse reaction(s) No statement required. Non-medicinal ingredients Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database. Storage conditions No statement required. Specifications The finished product specifications must be established in accordance with

the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal ingredient must comply with the requirements outlined in the NHPID. References Cited FDA 1994: The USA Department of Health and Human Services: Food and Drug Administration. 21 CFR Part 358. Wart remover drug products for over-the-counter human use, amendment of the final monograph; 1994. FR Citation: 59FR4015 [Accessed 2019 June 13]. Available at: <https://www.fda.gov/drugs/status-otc-rulemakings/rulemaking-history-otc-wart-remover-drug-products> FDA 1990: The USA Department of Health and Human Services: Food and Drug Administration. 21 CFR Part 358. Wart remover drug products for over-the-counter human use, final monograph, 1990. FR Citation: 55FR33246 [Accessed 2019 June 13]. Available at: <https://www.fda.gov/drugs/status-otc-rulemakings/rulemaking-history-otc-wart-remover-drug-products> Fiume MZ. Final report on the safety assessment of glycolic acid, ammonium, calcium, potassium, and sodium glycolates, methyl, ethyl, propyl, and butyl glycolates, and lactic acid, ammonium, calcium, potassium, sodium, and TEA-lactates, methyl, ethyl, isopropyl, and butyl lactates, and lauryl, myristyl, and cetyl lactates. International Journal of Toxicology, 1998;17 (Suppl. 1):1-3. Gottschalck TE and McEwen GN, editors. International Cosmetic Ingredient Dictionary and Handbook, 10 th ed. Washington (DC): Cosmetic, Toiletry and Fragrance Association; 2004. O'Neil MJ, Smith A, Heckelman PE, Budavari S, editors. Merck Index: An Encyclopedia of Chemicals, Drugs, & Biologicals, 13 th ed. Whitehouse Station (NJ): Merck & Co., Inc.; 2001. USP 29: The United States Pharmacopeia and the National Formulary (USP 29/NF 24). Rockville (MD): United States Pharmacopeial Convention, Inc.; 2006. Date Modified 2008-12-19 Top of page Important Notices

## MEDICINAL INGREDIENT(S)

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database. Storage conditions No statement required.

## DOSAGE FORM(S)

Dosage forms Doses Collodion-like solution 5 - 17% Karaya gum, glycol plaster 15% Other types of plasters 12 - 40%

## DOSE(S)

Dosage forms Doses Collodion-like solution 5 - 17% Karaya gum, glycol plaster 15% Other types of plasters 12 - 40%

## RISK INFORMATION

Caution(s) and warning(s) All products For external use only (FDA 1990). Consult a health care practitioner/health care provider/health care professional/doctor/ physician if discomfort persists (FDA 1990). Products formulated in a collodion-like solution only Extremely flammable or flammable or combustible (FDA 1990). Keep away from fire or flame (FDA 1990). Cap bottle tightly and store at room temperature, in an upright position, away from heat (FDA 1990). Flush with water for 15 minutes if product gets into the eye (FDA 1990). Avoid inhaling vapours (FDA 1990). Contraindication(s) All products Do not use this product on irritated or reddened skin or any area that is infected (FDA 1990). Do not use this product if you have diabetes or poor blood circulation (FDA 1990). Do not use this product on moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes (FDA 1990). Known adverse reaction(s) No statement required.

## NON-MEDICINAL INGREDIENTS

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database. Storage conditions No statement required.

## STORAGE CONDITION(S)

No statement required.

## SPECIFICATIONS

The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal ingredient must comply with the requirements outlined in the NHPID.

## REFERENCES

Table 2. Proper name(s), Common name(s), Source material(s) - Complementary ingredient (safety only) Proper name(s) Common name(s) Source ingredient(s) 1 Common name(s) 2-Hydroxy-2-methylacetic acid DL-Lactic acid DL-Lactic acid Lactic acid Lactic acid 1 The ingredient must be pharmacopoeial grade. References: Proper names: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Common names: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001; Source ingredient: USP 29 2006, Gottschalk & McEwen 2004, O'Neil et al. 2001. Route of Administration Topical (FDA 1990)

Proper name(s)	Common name(s)	Source ingredient(s)1
Common name(s)		
2-Hydroxybenzoic acid	Salicylic acid	Salicylic acid
1The ingredient must be pharmacopoeial grade.		

Proper name(s)	Common name(s)	Source ingredient(s)1
Common name(s)		
2-Hydroxy-2-methylacetic acidDL-Lactic acidDL-Lactic acidLactic acid		Lactic acid
1The ingredient must be pharmacopoeial grade.		

Dosage forms	Doses
--------------	-------

Collodion-like solution	5 - 17%
Karaya gum, glycol plaster	15%
Other types of plasters	12 - 40%