

Cassia - Cinnamomum aromaticum

Source: <https://webprod.hc-sc.gc.ca/nhp/ids/bdipsn/atReq?atid=cassia.cinnamomum.aromaticum2&=eng>

Extracted: 2025-08-26T06:31:30.345104

CASSIA - CINNAMOMUM AROMATICUM (PDF Version - 159 KB) This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredient. Notes Text in parentheses is additional optional information which can be included on the label at the applicant's discretion. The solidus (/) indicates that the terms and/or statements are synonymous. Either term or statement may be selected by the applicant on the label. Date May 30, 2025 Proper name(s), Common name(s), Source information Table 1. Proper name(s), Common name(s), Source information Proper name(s) Common name(s) Source information Source material(s) Part(s) Preparation(s) Cinnamomum aromaticum Cassia Cassia cinnamon Chinese cinnamon Chinese cinnamon tree Rou Gui Cinnamomum aromaticum Branch bark Stem bark Trunk bark Dry References: Proper name: USDA 2024; Gardner and McGuffin 2013; Common name: USDA 2024; Gardner and McGuffin 2013; Brinker 2010; Chen and Chen 2004; Blumenthal et al. 2000; Source information: PPRC 2020; Crawford 2009; BHC 2006; Mang et al. 2006; Bensky et al. 2004; Chen and Chen 2004; Khan et al. 2003; Blumenthal et al. 2000,1998. Route of administration Oral Dosage form(s) This monograph excludes foods or food-like dosage forms as indicated in the Compendium of Monographs Guidance Document. Acceptable dosage forms for oral use are indicated in the dosage form drop-down list of the web-based Product Licence Application form for Compendial applications. Use(s) or Purpose(s) Branch bark, Stem bark or Trunk bark Helps to support/maintain healthy blood glucose levels (Davis and Yokoyama 2011; Crawford 2009; Mang et al. 2006). Source of antioxidants/Provides antioxidants (Gruenwald et al. 2010; Roussel et al. 2009; Halvorsen et al. 2006; Shan et al. 2005). Source of antioxidants/Provides antioxidants that help fight/protect (cell) against/reduce (the oxidative effect of/the oxidative damage caused by/cell damage caused by) free radicals (Gruenwald et al. 2010; Roussel et al. 2009; Halvorsen et al. 2006; Shan et al. 2005). (Traditionally) used in Herbal Medicine for digestive disturbances/dyspeptic complaints/indigestion (such as mild spasms of the gastrointestinal tract, bloating and flatulence) (BHC 2006; Blumenthal et al. 2000, 1998). (Traditionally) used in Herbal Medicine for loss of appetite (BHC 2006; Blumenthal et al. 2000, 1998). Trunk bark only Used in Traditional Chinese Medicine (TCM) to dispel cold, relieve pain, open channels and collaterals (Bensky et al. 2004; Chen and Chen 2004). Used in Traditional Chinese Medicine (TCM) to dispel cold, warm the spleen and relieve pain (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Used in Traditional Chinese Medicine (TCM) to encourage production of qi and blood (Bensky et al. 2004; Chen and Chen 2004). Used in Traditional Chinese Medicine (TCM) to tonify kidney yang and augment ming men (life gate) fire (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Notes The above uses can be combined on the product label if from the same traditional or non-traditional system of medicine (e.g., Used in Traditional Chinese Medicine to encourage production of qi and blood, tonify kidney yang and augment ming men fire). For multi-ingredient products: To prevent the product from being represented as a "traditional medicine", any indicated traditional use claim must refer to the specific medicinal ingredient(s) and recognized traditional system of medicine from which the claim originates when 1) both traditional and modern claims are present or 2) when claims originate from multiple systems of traditional medicine (e.g., Cassia is traditionally used in Herbal Medicine for loss of appetite). When ALL of the medicinal ingredients (MIs) in the product are used within the SAME identified system of traditional medicine AND the product makes ONLY traditional claims, listing of MIs in the traditional claim(s) is not required. Dose(s) Subpopulation(s) Adults 18 years and older Quantity(ies) Branch bark, Stem bark or Trunk bark Methods of preparation: Dry, Powdered, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract) Antioxidant Not to exceed 6 grams of dried branch bark, stem bark and/or trunk bark, per day and 4 grams per single dose (Gruenwald et al. 2010; Roussel et al. 2009; Halvorsen et al. 2006; Shan et al. 2005). Appetite loss; Digestive disturbances/Indigestion 1 - 6 grams of dried branch bark, stem bark and/or trunk bark, per day; Not to exceed 4 grams per single dose (Gruenwald et al. 2010; Al Jamal et al. 2009; Crawford 2009; Mang et al. 2006; Safdar et al. 2004; Khan et al. 2003). Healthy glucose 3 - 6 grams of dried branch bark, stem bark and/or trunk bark, per day; Not to exceed 4 grams per single dose (Davis and Yokoyama 2011; Gruenwald et al. 2010; Crawford 2009; Mang et al. 2006). Trunk bark only Traditional Chinese Medicine claims Method of preparation: Decoction, Decoction concentrate 2 - 5 grams

of dried trunk bark, per day (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Methods of preparation: Dry, Powdered, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract) 1 - 2 grams of dried trunk bark, per day (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Direction(s) for use Appetite loss (For appetite loss)*, take 30 minutes before meals. Digestive disturbances/Indigestion (For digestive disturbances)*, take with a meal/food (Crawford 2009). *Note: Information in brackets is required on the label when both uses for appetite loss and digestive disturbances are listed on the label. Duration(s) of use Products providing 4 - 6 g of branch bark, stem bark and/or trunk bark, per day Ask a health care practitioner/health care provider/health care professional/doctor/physician for use beyond 6 weeks (Safdar et al. 2004; Khan et al. 2003). Risk information Caution(s) and warning(s) Appetite loss; Digestive disturbances/Indigestion; Traditional Chinese Medicine claims Ask a health care practitioner/health care provider/health care professional/doctor/physician if symptoms persist or worsen. Products providing 1 g or more of branch bark, stem bark and/or trunk bark, per day Ask a health care practitioner/health care provider/health care professional/doctor/physician before use if you are breastfeeding (Blumenthal et al. 2000; WHO 1999). Ask a health care practitioner/health care provider/health care professional/doctor/physician before use if you have diabetes (Gardner and McGuffin 2013; Brinker 2010). Contraindication(s) Products providing 1 g or more of branch bark, stem bark and/or trunk bark, per day Do not use if you are pregnant (PPRC 2020; Brinker 2010; BHC 2006; Chen and Chen 2004; Blumenthal 2000, 1998). Traditional Chinese Medicine claims Do not use if you have excess heat, blood heat or deficiency, or yin-deficient fire (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Known adverse reaction(s) Stop use if hypersensitivity/allergy occurs (Gardner and McGuffin 2013; Blumenthal 2000; WHO 1999). Non-medicinal ingredients Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database. Storage conditions Must be established in accordance with the requirements described in the Natural Health Products Regulations. Specifications The finished product specifications must be established with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal ingredient must comply with the requirements outlined in the NHPID. EXAMPLE OF PRODUCT FACTS: Consult the Guidance Document, Labelling of Natural Health Products for more details. References cited Al Jamal AR. Effects of cinnamon on blood glucose and lipid levels in diabetic patients (type 1). *African Journal of Biochemistry Research* 2009;3(5):181-184. Bensky D, Clavey, Stöger E, Gamble A. *Chinese Herbal Medicine: Materia Medica*. 3rd edition. Seattle (WA): Eastland Press, Incorporated; 2004. BHC 2006: Bradley PR, editor. *British Herbal Compendium Volume 2: A Handbook of Scientific Information on Widely Used Plant Drugs - Companion to the British Herbal Pharmacopoeia*. Bournemouth (GB): British Herbal Medicine Association; 2006. Blumenthal M, Goldberg A, Brinckmann J. *Herbal Medicine: Expanded Commission E Monographs*. Boston (MA): American Botanical Council; 2000. Blumenthal M, editor. *The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines*. Austin (TX): American Botanical Council in cooperation with Integrative Medicine Communications; 1998. Brinker F. *Herb Contraindications and Drug Interactions*, 4th edition. Sandy (OR): Eclectic Medical Publications; 2010. Chen JK, Chen TT. *Chinese Medical Herbology and Pharmacology*. Crampton L, editor. City of Industry (CA): Art of Medicine Press Inc.; 2004. Crawford P. Effectiveness of cinnamon for lowering hemoglobin A1C in patients with type 2 diabetes: a randomized, controlled trial. *Journal of the American Board of Family Medicine* 2009;22(5):507-512. Davis PA, Yokoyama W. Cinnamon intake lowers fasting blood glucose: meta-analysis. *Journal of Medicinal Food* 2011;14(9):884-889. Gardner Z, McGuffin M, editors. *American Herbal Products Association's Botanical Safety Handbook*. 2nd edition. Boca Raton (FL): Taylor and Francis Group; 2013. Gruenwald J, Freder J, Armbruester N. Cinnamon and health. *Critical Reviews in Food Science and Nutrition* 2010;50(9):822-834. Halvorsen BL, Carlsen MH, Phillips KM, Bohn SK, Holte K, Jacobs DR Jr, Blomhoff R. Content of redox-active compounds (ie, antioxidants) in foods consumed in the United States. *American Journal of Clinical Nutrition* 2006;84(1):95-135. Khan A, Safdar M, Ali Khan MM, Khattak KN, Anderson RA. Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabetes Care* 2003;26(12):3215-3218. Mang B, Wolters M, Schmitt B, Kelb K, Lichtinghagen R, Stichtenoth DO, Hahn A. Effects of a cinnamon extract on plasma glucose, HbA1C, and serum lipids in diabetes mellitus type 2. *European Journal of Clinical Investigation* 2006;36(5):340-344. PPRC 2020: *Pharmacopoeia of the People's Republic of China*. Volume 1, English edition. Beijing (CN): The State Pharmacopoeia Commission of the People's Republic of China; 2020. Roussel AM, Hininger I, Benaraba R, Ziegenfuss TN, Anderson RA. Antioxidant effects of a cinnamon extract in people with impaired fasting glucose that are overweight or obese. *Journal of the American College of Nutrition* 2009;28(1):16-21. Safdar M, Khan A., Khan MMA, Siddique M. Effect of various doses of cinnamon on blood glucose in diabetic individuals. *Pakistan Journal of Nutrition* 2004;3:268-272. Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *Journal of Agricultural and Food Chemistry* 2005;53(20):7749- 7759. USDA 2024: United States Department of Agriculture, Agricultural Research Service (USDA ARS), Germplasm Resources Information Network (GRIN) - Global. U.S. National Plant Germplasm System. [Accessed 2024 August 12]. Available from:

<https://npgsweb.ars-grin.gov/gringlobal/taxon/taxonomysearch> WHO 1999: World Health Organization. WHO Monographs on Selected Medicinal Plants, Volume 1. Geneva (CH): World Health Organization; 1999. References reviewed American Society of Health-System Pharmacists. American Hospital Formulary Service (AHFS) Drug Information. Philadelphia (PA): Lippincott Williams and Wilkins; 2005. Altschuler JA, Casella SJ, MacKenzie TA, Curtis KM. The effects of cinnamon on A1C among adolescence with type 1 diabetes. *Diabetes Care* 2007;30:813-816. Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, Graves DJ. Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin-like biological activity. *Journal of Agricultural and Food Chemistry* 2004;52(1):65- 70. Baker W, Gutierrez-Williams G, White CM, Kluger J, Coleman CI. Effect of cinnamon on glucose control and lipid parameters. *Diabetes Care* 2008;31:41-43. Bandara T, Uluwaduge I, Jansz ER. Bioactivity of cinnamon with special emphasis on diabetes mellitus: a review. *International Journal of Food Sciences and Nutrition* 2012;63(3):380-386. Blevins SM, Leyva MJ, Brown J, Wright J, Scofield RH, Aston CE. Effect of cinnamon on glucose and lipid levels in non-insulin dependent type 2 diabetes mellitus. *Diabetes Care* 2007;30:2236-2237. Brinker F. *The Toxicology of Botanical Medicines*. Sandy (OR): Eclectic Medical Publications; 2000. Broadhurst CL, Polansky MM, Anderson RA. Insulin like biological activity of culinary and medicinal plant aqueous extracts in vitro. *J Agric Food Chem* 2000;48:849-852. CNF 2023: Canadian Nutrient File, Food and Nutrition, Health Canada. [Accessed 2024 August 13]. Available from: <https://food-nutrition.canada.ca/cnf-fce> Carter JS, Pugh JA, Monterrosa A. Non-insulin-dependent diabetes mellitus in minorities in the United States. *Ann Intern Med* 1996;125(1):221-232. Clinical Practice Guidelines for the Prevention and Management of Diabetes in Canada. *Canadian Journal of Diabetes* 2008;32(suppl 1). Dugoua JJ, Seely D, Perri D, Cooley K, Forelli T, Mills E, Koren G. From type 2 diabetes to antioxidant activity: a systematic review of the safety and efficacy of common and cassia cinnamon bark. *Can. J. Physiol. Pharmacol* 2007;85:837-847. Facciola S. *Cornucopia II A Source Book of Edible Plants*. Vista (CA): Kampong Publications; 1998. Hlebowicz J, Darwiche G, Björgell O, Almé LO. Effect of cinnamon on postprandial blood glucose, gastric emptying, and satiety in healthy subjects. *American Journal of Clinical Nutrition* 2007;85:1552-1556. Hlebowicz J, Hlebowicz A, Lindstedt S, Björgell O, Höglund P, Holst JJ, et al. Effects of 1 and 3g cinnamon on gastric emptying, satiety, and postprandial blood glucose, insulin, glucose- dependent insulinotropic polypeptide, glucagon-like peptide 1, and ghrelin concentrations in healthy subjects. *American Journal of Clinical Nutrition* 2009;89:815-821. Imparl-Radosevich J, Deas S, Polansky MM et al. Regulation of PTP-1 and insulin receptor kinase by fractions from cinnamon: implications for cinnamon regulation of insulin signaling. *Horm Res* 1998;50:177-182. Jarvill-Taylor KJ, Anderson RA, Graves DJ. A hydroxychalcone derived from cinnamon functions as a mimetic for insulin in 3T3-L1 adipocytes. *J Am Coll Nutr* 2001;20(4):327-236. JEFCA Evaluation. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives: Cinnamaldehyde. [Accessed 2024 August 13]. Available from: http://www.inchem.org/documents/jecfa/jecval/jec_418.htm Joint Food and Agriculture Organization of the United Nations (FAO)/World Health Organization (WHO) Expert Committee on Food Additives. WHO Food Additives Series: 60. Safety evaluation of certain food additives. Geneva (CH): World Health Organization. 2009. [Accessed 2024 August 20] Available from: <http://www.inchem.org/documents/jecfa/jecmono/v60je01.pdf> Qin B, Nagasaki M, Ren M, Bajotto G, Oshida Y, Sato Y. Cinnamon extract (traditional herb) potentiates in vivo insulin-regulated glucose utilization via enhancing insulin signaling in rats. *Diabetes Res Clin Pract* 2003;62:139-148. Solomon TPJ, Blannin AK. Effects of short-term cinnamon ingestion on in vivo glucose tolerance. *Diabetes Obes Metab* 2007;8:895-901. Solomon TPJ, Blannin AK. Changes in glucose tolerance and insulin sensitivity following 2 weeks of daily cinnamon ingestion in healthy humans. *Eur J Appl Physiol* 2009;105:969-976. Soni R, Bhatnagar V. Effect of cinnamon (*Cinnamomum cassia*) intervention on blood glucose of middle aged adult male with non-insulin dependent diabetes mellitus (NIDDM). *Ethno-Med* 2009;3:141-144. United Kingdom Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837-853. Vanschoonbeek K, Thomassen BJW, Senden JM, Wodzig WKWH, van Loon LJC. Cinnamon supplementation does not improve glycemic control in postmenopausal type 2 diabetic patients. *J Nutr* 2006;136:977-980. WHO 2010a: World Health Organization. WHO Food Additives Series 46: Cinnamyl Alcohol and Related Substances. 2010. [Accessed 2024 August 13]. Available from: <http://www.inchem.org/documents/jecfa/jecmono/v46je07.htm> WHO 2010b: World Health Organization. WHO Food Additives Series 14: Cinnamaldehyde. 2010. [Accessed 2024 August 13]. Available from: <http://www.inchem.org/documents/jecfa/jecmono/v14je07.htm> Ziegenfuss TN, Hofheins JE, Mendel RW, Landis J., Anderson RA. Effects of a water-soluble cinnamon extract on body composition and features of the metabolic syndrome in pre-diabetic men and women. *J Int Soc Sports Nut* 2006;3:45-53. Report a problem on this page Date modified: 2019-03-01

DOSAGE FORM(S)

Acceptable dosage forms for oral use are indicated in the dosage form drop-down list of the web-based Product Licence Application form for Compendial applications.

DOSE(S)

Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *Journal of Agricultural and Food Chemistry* 2005;53(20):7749- 7759. USDA 2024: United States Department of Agriculture, Agricultural Research Service (USDA ARS), Germplasm Resources Information Network (GRIN) - Global. U.S. National Plant Germplasm System. [Accessed 2024 August 12]. Available from: <https://npgsweb.ars-grin.gov/gringlobal/taxon/taxonomysearch> WHO 1999: World Health Organization. WHO Monographs on Selected Medicinal Plants, Volume 1. Geneva (CH): World Health Organization; 1999.

RISK INFORMATION

Caution(s) and warning(s) Appetite loss; Digestive disturbances/Indigestion; Traditional Chinese Medicine claims Ask a health care practitioner/health care provider/health care professional/doctor/physician if symptoms persist or worsen. Products providing 1 g or more of branch bark, stem bark and/or trunk bark, per day Ask a health care practitioner/health care provider/health care professional/doctor/physician before use if you are breastfeeding (Blumenthal et al. 2000; WHO 1999).Ask a health care practitioner/health care provider/health care professional/doctor/physician before use if you have diabetes (Gardner and McGuffin 2013; Brinker 2010). Contraindication(s) Products providing 1 g or more of branch bark, stem bark and/or trunk bark, per day Do not use if you are pregnant (PPRC 2020; Brinker 2010; BHC 2006; Chen and Chen 2004; Blumenthal 2000, 1998). Traditional Chinese Medicine claims Do not use if you have excess heat, blood heat or deficiency, or yin-deficient fire (PPRC 2020; Bensky et al. 2004; Chen and Chen 2004). Known adverse reaction(s) Stop use if hypersensitivity/allergy occurs (Gardner and McGuffin 2013; Blumenthal 2000; WHO 1999).

NON-MEDICINAL INGREDIENTS

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database.

STORAGE CONDITION(S)

Must be established in accordance with the requirements described in the Natural Health Products Regulations.

SPECIFICATIONS

The finished product specifications must be established with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide. The medicinal

ingredient must comply with the requirements outlined in the NHPID.

Latin name(s)	Common name(s)	Source information	
Material(s)	Part(s)	Preparation(s)	
Cinnamomum aromaticum	CassiaCassia cinnamonChinese cinnamonCinnamomum aromaticum	Chinese cinnamon tree Bark	Branch barkStem barkTrunk bark