Addendum: Protocol for Paired Shoulder Abduction and Complementary Cervical Stabilization

Section 1: Biomechanical Framework for Shoulder Abduction

The act of raising the arm to the side, known as shoulder abduction, is a biomechanically sophisticated movement that requires the coordinated action of multiple joints, muscle groups, and neuromuscular control systems. A comprehensive understanding of this kinetic chain is fundamental to prescribing effective and safe exercises. This section deconstructs the phases of shoulder abduction and examines the critical force couples that govern its stability and efficiency.

1.1 The Kinetic Chain of Arm Elevation: A Multi-Phase Analysis

Shoulder abduction is not a monolithic action but a sequential process involving both the glenohumeral and scapulothoracic joints. The movement is classically divided into three distinct phases, each dominated by specific muscular contributors. Treating this motion as a simple "shoulder" exercise overlooks the intricate interplay of the entire shoulder girdle, a misstep that can lead to inefficient movement patterns and an increased risk of pathology. The movement must be understood as a coordinated event involving the humerus, scapula, and clavicle.

Phase 1 (0-15°): Initiation by the Supraspinatus

The initial phase of abduction, from to approximately , is primarily initiated by the supraspinatus muscle, one of the four rotator cuff muscles.[1] While the deltoid muscle is also active during this phase, the supraspinatus plays a uniquely critical role. Its line of pull allows it to act as a primary abductor at smaller angles of elevation.[3] More importantly, the supraspinatus generates a vital compressive force, pulling the humeral head medially into the glenoid fossa. This action stabilizes the glenohumeral joint and prevents the larger, more powerful deltoid muscle from causing superior migration of the humeral head upon its contraction.[3] This initial stabilization is a prerequisite for smooth, pain-free motion throughout the rest of the range. Failure of the supraspinatus to adequately perform this dual function of initiation and compression can lead to instability and is often implicated in subacromial impingement syndrome.[2]

Phase 2 (15-90°): Primary Abduction by the Deltoid

Once the arm has been raised to approximately 15˚ the middle fibers of the deltoid muscle become the prime mover for abduction up to 90˚. The deltoid's leverage and moment arm are more effective in this range, allowing it to generate significant abduction torque.[3] However, the deltoid's line of pull is predominantly superior. If left unopposed, its powerful contraction would translate the humeral head upward, potentially impinging the soft tissues in the subacromial space, such as the supraspinatus tendon and the subacromial bursa.[5] Therefore, throughout this phase,

the other rotator cuff muscles (infraspinatus, teres minor, and subscapularis) must remain active to provide a counteracting inferior and compressive force, ensuring the humeral head remains centered within the glenoid fossa.[3] This dynamic stabilization is crucial for the health and function of the glenohumeral joint.

Phase 3 (>90°): Scapulothoracic Contribution

Achieving a full range of abduction beyond is impossible without significant contribution from the scapulothoracic articulation.[1] To allow the humerus to continue its upward path, the scapula must rotate upwardly, tilting the glenoid fossa superiorly to maintain joint congruity. This upward rotation is not produced by a single muscle but by a synergistic force couple involving the upper trapezius, lower trapezius, and the serratus anterior.[7] The upper trapezius pulls the acromion and lateral clavicle upward and medially, the lower trapezius pulls the medial base of the scapular spine downward, and the serratus anterior pulls the inferior angle of the scapula forward and laterally around the rib cage. The synchronized contraction of these three muscles results in the smooth upward rotation of the scapula. This scapulohumeral rhythm, typically a 2:1 ratio of glenohumeral to scapulothoracic motion, is essential for achieving full, unimpeded arm elevation.[1] Any disruption in the function or coordination of these scapular stabilizers will limit the total range of abduction and force compensatory movements, such as excessive shrugging or lateral trunk bending.

1.2 Critical Force Couples in Shoulder Function

The stability and efficiency of the shoulder complex depend on the precise balance of forces generated by synergistic muscle groups. These "force couples" consist of two or more muscles pulling in different directions on a common attachment point to produce a pure rotation, minimizing unwanted translation. Two such force couples are paramount to shoulder abduction.

The Coronal Plane Force Couple (Deltoid-Rotator Cuff)

In the coronal plane, the primary force couple responsible for glenohumeral stability during abduction is formed by the deltoid and the rotator cuff.[5] As detailed previously, the deltoid muscle generates a strong superiorly directed shear force on the humerus. This is counteracted by the combined forces of the infraspinatus, teres minor, and subscapularis, which generate an inferiorly directed shear force.[6] The resultant vector of these opposing forces is a net compressive force that seats the humeral head firmly and centrally within the glenoid fossa throughout the arc of motion.[5] A tear or weakness in the rotator cuff disrupts this delicate balance, allowing the deltoid's upward pull to dominate. This can lead to superior humeral head migration, a hallmark of rotator cuff pathology that narrows the subacromial space and precipitates impingement and further tendon degradation.

The Scapular Force Couple (Trapezius-Serratus Anterior)

In the plane of the scapula, a second critical force couple produces the upward rotation necessary for full abduction. This couple is composed of the upper trapezius, lower trapezius, and serratus anterior.[7] The upper trapezius provides an upward and medial pull, the lower trapezius a downward and medial pull, and the serratus

anterior a forward and lateral pull. The coordinated action of these muscles rotates the scapula around an axis near the base of the scapular spine, elevating the acromion and orienting the glenoid fossa superiorly to support the elevating arm.[7] Dysfunction in any component of this triad, such as weakness in the serratus anterior or lower trapezius, leads to abnormal scapular motion (dyskinesis), which can compromise glenohumeral mechanics and contribute to shoulder pathology.
The interdependence of these two force couples cannot be overstated. They do not operate in isolation. A primary dysfunction in the glenohumeral joint, such as rotator cuff weakness, will invariably affect the function of the scapular stabilizers. In response to pain or instability at the glenohumeral joint, the central nervous system (CNS) often adopts a compensatory motor strategy.[9] A common maladaptation is the over-activation of the upper trapezius and levator scapulae muscles, which attempt to elevate the arm by shrugging the entire shoulder girdle. This action substitutes for true glenohumeral abduction and disrupts the balanced action of the scapular force couple, favoring excessive scapular elevation over the more efficient upward rotation. This creates a vicious cycle: glenohumeral instability leads to faulty scapular mechanics, which in turn further compromises the subacromial space and exacerbates the initial glenohumeral pathology. Therefore, any effective intervention must address both systems, recognizing that a stable glenohumeral joint requires a stable and mobile scapula, and vice versa.

Section 2: The Cervical-Scapular Nexus: Rationale for Integrated Exercise

The cervical spine and the shoulder girdle are not independent anatomical regions but form a single, integrated functional unit. Their intimate anatomical, neurological, and biomechanical relationships dictate that the status of one directly influences the function of the other. This section explores these connections and elucidates how cervical posture is a critical determinant of shoulder health and mechanics, providing the rationale for pairing a cervical stabilization exercise with shoulder abduction.

2.1 Anatomical and Functional Linkages

The direct physical and neurological connections between the neck and shoulder underscore the necessity of a holistic approach to assessment and treatment.

Key Musculature

Several key muscles span the cervicoscapular region, acting on both the neck and the shoulder girdle. The upper trapezius originates from the occiput and the cervical spinous processes and inserts onto the clavicle and acromion.[7] The levator scapulae originates from the transverse processes of the upper cervical vertebrae (C1-C4) and inserts on the superior medial border of the scapula.[11] Due to these attachments, these muscles are primary movers and stabilizers for both cervical and scapular motion. Consequently, tension, weakness, or dysfunction in these muscles will have a dual impact. For example, chronic tension in the upper trapezius can contribute to both neck pain and faulty scapular elevation during arm movements.

Neuromuscular Coupling

Beyond direct muscular connections, the neck and shoulder are linked through shared neurological pathways. The position of the head and neck can modulate the excitability of the motor neurons that supply the shoulder muscles. Research on the reticulospinal tract, a key pathway for postural control, has shown that it has an ipsilateral influence on shoulder abductors and elbow flexors.[12] This means that the neural drive to the muscles that lift the arm can be facilitated or inhibited based on signals originating from the brainstem, which are in turn influenced by head position. This provides a neurophysiological basis for the observation that correcting neck posture can improve the activation and strength of shoulder musculature. The re-emergence of primitive reflexes, such as the Asymmetrical Tonic Neck Reflex (ATNR) after cortical injury, further demonstrates this hard-wired link, where head rotation directly causes a pattern of limb extension and flexion.

2.2 Postural Influence on Scapulohumeral Rhythm

The most significant functional link between the neck and shoulder is the profound influence of head posture on scapular mechanics. Specifically, forward head posture (FHP) is a common postural fault that initiates a cascade of biomechanical impairments throughout the upper kinetic chain.

The Pathomechanics of Forward Head Posture (FHP)

FHP is characterized by an anterior translation of the head relative to the shoulders, typically involving flexion of the lower cervical spine and extension of the upper cervical spine.[11] This chronic malalignment leads to predictable patterns of muscle imbalance. The deep cervical flexors (DCFs), such as the longus colli and longus capitis, become lengthened and inhibited, losing their ability to provide segmental stability to the cervical vertebrae. Similarly, the scapular retractors and depressors, particularly the middle and lower trapezius, also become lengthened and weakened.[14] Conversely, a separate group of muscles becomes shortened, tight, and overactive. This group includes the suboccipital muscles at the base of the skull, the sternocleidomastoid (SCM), the upper trapezius, and the levator scapulae.

The Causal Chain from FHP to Impingement

This pattern of muscle imbalance directly leads to altered scapular positioning and faulty scapulohumeral rhythm, creating a high-risk environment for shoulder impingement. The causal sequence is as follows:

1. FHP is often accompanied by an increase in thoracic kyphosis (rounding of the upper back) and an adaptive shortening of the pectoralis minor muscle, which attaches to the coracoid process of the scapula.[11]
2. The combination of a tight pectoralis minor anteriorly and hypertonic upper trapezius/levator scapulae superiorly pulls the scapula into a resting position of anterior tilt, internal rotation, and protraction.[14]
3. An anteriorly tilted scapula causes the acromion to angle downward, physically narrowing the subacromial space through which the rotator cuff tendons pass.[2]
4. When shoulder abduction is attempted from this compromised position, the reduced subacromial space significantly increases the likelihood of the supraspinatus tendon and subacromial bursa being compressed between the

greater tuberosity of the humerus and the acromion.[3] This repetitive mechanical compression is the primary mechanism of subacromial impingement syndrome.

This analysis reveals that posture is not merely a static alignment but a dynamic reflection of the CNS's default motor control strategy. FHP represents a neuromuscular "set point" where the CNS has learned to upregulate the activity of postural muscles like the upper trapezius and pectorals while downregulating the activity of key stabilizers like the DCFs and lower trapezius.[9] Attempting to perform a dynamic movement like shoulder abduction from this dysfunctional set point will inevitably reinforce the faulty pattern, as the CNS will preferentially recruit the already facilitated, overactive muscles. Therefore, the most logical and effective therapeutic approach is not to simply strengthen the weak muscles in isolation. Instead, the primary goal should be to first reset this neuromuscular set point by correcting the foundational posture. By doing so, a neurological environment is created in which the correct muscles *can* be activated and integrated into the movement pattern.

Section 3: Analysis and Selection of the Optimal Complementary Neck Exercise

Given the established biomechanical link between cervical posture and shoulder function, the selection of a neck exercise to complement shoulder abduction must be deliberate and strategic. The goal is not merely to address neck symptoms but to optimize the foundational stability of the entire upper quarter. An evaluation of common neck exercises reveals that while many are beneficial, one stands out as uniquely suited for this preparatory role.

3.1 Evaluation of Candidate Exercises from Video Source

The exercises commonly recommended for neck health, as seen in the provided video and supporting materials, can be broadly categorized into dynamic mobility drills and isometric strengthening exercises.

Dynamic Mobility Drills

Exercises such as lateral neck flexion (side bending), neck rotation, and forward/backward tilting are excellent for improving the extensibility of superficial cervical muscles and enhancing joint range of motion.[13] A side-bending stretch, for instance, effectively targets the contralateral upper trapezius and scalene muscles, which are often tight in individuals with FHP.[14] While valuable for alleviating stiffness and restoring mobility, these stretches do not primarily address the underlying weakness of the deep stabilizing muscles or correct the segmental instability characteristic of poor cervical posture. They manage the symptoms of muscle tightness rather than the root cause of the postural fault.

Isometric Strengthening Drills

Isometric exercises, where the neck muscles contract against resistance without producing movement, are effective for increasing the strength and endurance of the global cervical musculature.[11] Pressing the head against the hand in flexion, extension, or side bending creates a co-contraction of agonist and antagonist

muscles, enhancing overall neck stability. However, these exercises tend to recruit the larger, more superficial muscles (e.g., SCM, upper trapezius) and may not specifically target the deep cervical flexors. While beneficial for general strengthening, they are less precise in correcting the specific muscular imbalance and postural deviation of FHP.

3.2 Selection Rationale: Cervical Retraction (Chin Tuck)

After a thorough analysis, **Cervical Retraction**, commonly known as the "chin tuck," is selected as the optimal complementary exercise to be paired with shoulder abduction. This selection is based on its unique ability to directly counteract the pathomechanics of FHP and to prime the neuromuscular system for correct movement. The justification is fourfold.

Justification 1: Direct Antagonist to Forward Head Posture

Cervical retraction is the precise kinematic opposite of FHP. Whereas FHP involves lower cervical flexion and upper cervical extension, the chin tuck maneuver promotes lower cervical extension and upper cervical flexion (a slight nod).[11] This movement pattern directly strengthens the muscles that are inhibited by FHP—namely, the deep cervical flexors (longus colli and longus capitis). Simultaneously, it produces a gentle stretch of the tight, overactive suboccipital extensor muscles at the base of the skull.[14] No other single exercise targets this specific dysfunctional pattern with such precision.

Justification 2: Promotes Proximal Stability

The principle of "proximal stability for distal mobility" is a cornerstone of biomechanics and rehabilitation. It posits that efficient movement of the limbs requires a stable core or base. In the context of the upper quarter, the cervical and thoracic spine serve as this proximal base. By activating and strengthening the DCFs, cervical retraction enhances the segmental stability of the individual cervical vertebrae. This creates a solid, reliable anchor for the origins of the key scapular muscles, including the levator scapulae and trapezius. A stable cervical platform allows these muscles to function more effectively and efficiently in controlling scapular motion during arm elevation.

Justification 3: Facilitates Optimal Scapular Positioning

By directly addressing FHP and its associated increase in thoracic kyphosis, the act of cervical retraction initiates a positive postural chain reaction. Pulling the head back into alignment over the shoulders encourages a more upright thoracic posture. This, in turn, allows the scapulae to rest in a more neutral position on the rib cage, moving them out of the pathological state of anterior tilt and protraction.[11] This repositioning of the scapula is critically important because it re-aligns the glenoid fossa and inherently widens the subacromial space. This effect occurs *before* the shoulder abduction movement even begins, creating a safer and more mechanically advantageous starting position for the glenohumeral joint.

Justification 4: Neuromuscular Re-education

Perhaps the most significant benefit of cervical retraction in this context is its role as a motor learning task. Performing a chin tuck is not simply a muscular contraction; it

is an act of neuromuscular re-education. It actively teaches the CNS to disengage from the faulty FHP pattern and find a neutral, stacked head-on-neck posture. It retrains the proprioceptive system to recognize this new, efficient alignment as the correct default "set point." By repeatedly practicing this movement, an individual can begin to override the ingrained motor program of FHP.

In this light, the role of cervical retraction extends far beyond being a simple "neck exercise." It functions as a foundational "pre-habilitation" or "postural priming" drill. While other exercises may stretch tight muscles or strengthen weak ones, cervical retraction fundamentally reorganizes the biomechanical and neurological foundation upon which all upper limb movements are built. Its primary purpose in this pairing is to prepare the entire upper quarter for the demands of shoulder abduction, addressing the root postural cause of mechanical inefficiency and impingement risk. This preparatory function makes it uniquely and ideally suited as a complementary exercise.

Section 4: Prescriptive Exercise Protocol

This section provides a detailed, evidence-based protocol for the performance of the selected exercises: Active Shoulder Abduction and Cervical Retraction. The protocol is designed for direct implementation, with an emphasis on precise execution, safety cues, and a logical programming sequence that maximizes therapeutic benefit.

4.1 Exercise A: Active Shoulder Abduction (Scaption)

Objective: To strengthen the primary shoulder abductors (supraspinatus, deltoid) and to train coordinated, efficient scapulohumeral rhythm involving the scapular stabilizing muscles (trapezius, serratus anterior).[1]
Execution:

1. **Starting Position:** Assume a standing position with feet positioned shoulder-width apart to provide a stable base. Engage the core musculature to stabilize the lumbar spine and pelvis. The cervical spine should be held in a neutral position, ideally after performing the cervical retraction exercise to establish proper alignment. Begin with light dumbbells in each hand or no weight, depending on strength and comfort level.
2. **Arm Position:** Position the arms at the sides but slightly in front of the body,

   within the scapular plane. This plane is approximately anterior to the coronal (frontal) plane and represents the natural orientation of the scapula on the thorax. This position is less likely to cause impingement than abduction in the pure coronal plane.[3] The hands should be oriented so the thumbs point towards the ceiling, which places the glenohumeral joint in a position of slight external rotation, further clearing the greater tuberosity from the acromion.[22]
3. **Movement:** Initiate the movement by slowly and smoothly raising the arms

   upward and outward to shoulder level (approximately of abduction). The motion should be deliberate and controlled, focusing on the feeling of the shoulder blades rotating on the rib cage to assist the movement. Avoid any momentum or swinging.

4. **Hold and Lower:** At the apex of the movement ( ), pause for one to two seconds. During this hold, consciously depress the shoulders, ensuring they are not shrugging up towards the ears. The descent should be even slower and more controlled than the ascent, taking approximately three to four seconds to lower the arms back to the starting position.

Dosage: Perform 2 to 3 sets of 10 to 15 repetitions. The resistance should be challenging enough that the final repetitions of each set are difficult to complete with perfect form, but not so heavy as to cause pain or compensation.

4.2 Exercise B: Cervical Retraction (Chin Tuck)

Objective: To activate and strengthen the deep cervical flexor muscles, correct forward head posture, and promote a stable cervical base to optimize scapular function.[11]

Execution (Seated Variation):

1. **Starting Position:** Sit upright in a chair with a firm back for support. Place feet flat on the floor. The gaze should be directed straight ahead, level with the horizon, and fixed on a distant point to prevent inadvertent head tilting.
2. **Movement:** Without tilting the head up or down, gently glide the head and chin horizontally backward. The motion should feel as if the entire head is sliding back on the neck. A useful cue is to imagine trying to make a "double chin." The primary sensation should be a mild activation or contraction in the muscles deep in the front of the neck and a gentle lengthening or stretch at the base of the skull (suboccipital region).[11]
3. **Hold and Release:** Maintain the fully retracted position for a duration of 5 to 10 seconds. Focus on breathing naturally during the hold. After the hold, slowly and smoothly release the contraction, allowing the head to return to the neutral starting position. Avoid letting the chin jut forward past neutral upon release.

Execution (Supine Variation):

This variation is often recommended for beginners or individuals with significant neck pain as it unloads the spine from gravity.

1. **Starting Position:** Lie on your back on a firm surface. A pillow should not be used, but a small, rolled-up towel may be placed in the curve of the neck for support if needed. The knees should be bent with feet flat on the surface.
2. **Movement:** Gently press the back of the head into the floor while simultaneously performing a slight chin tuck towards the chest. This is a very subtle movement focused on elongating the back of the neck and flattening its curve against the floor. It is crucial to avoid lifting the head off the surface.[16]
3. **Hold and Release:** Hold the gentle contraction for 5 to 10 seconds, then relax completely.

Dosage: Perform 2 sets of 10 repetitions, holding each repetition for 5 to 10 seconds. The focus should be on the quality of the contraction, not the force.

4.3 Protocol for Integrated Programming and Performance

Sequencing: For optimal results, Cervical Retraction (Exercise B) should always be performed **before** Active Shoulder Abduction (Exercise A).

Rationale for Sequencing: This specific sequence is critical. Performing the cervical retraction drill first serves as a neuromuscular primer. It actively corrects the

resting head and shoulder posture, activates the deep cervical stabilizers, and inhibits the overactive postural muscles. This process establishes an optimal mechanical and neurological state for the more complex, dynamic shoulder movement that follows. By "setting the foundation" first, the shoulder abduction exercise can be performed with better mechanics, reduced risk of impingement, and more effective recruitment of the target musculature. This sequence transforms the protocol from two isolated exercises into a synergistic preparatory and execution sequence.

Frequency: This paired protocol can be performed 3 to 5 times per week. It can be integrated as part of a warm-up routine before upper body resistance training or used as a standalone corrective exercise session on non-training days to address postural deficits.

Table 4.1: Paired Exercise Protocol Summary

Feature	**Exercise A: Active Shoulder Abduction (Scaption)**	**Exercise B: Cervical Retraction (Chin Tuck)**
Objective	To strengthen primary abductors (supraspinatus, deltoid) and train coordinated scapulohumeral rhythm.	To enhance cervical spine stability, activate deep cervical flexors, and correct forward head posture to create a stable base for shoulder movement.
Primary Muscles	Middle Deltoid, Supraspinatus.	Deep Cervical Flexors (Longus Colli, Longus Capitis).
Secondary Muscles	Upper Trapezius, Lower Trapezius, Serratus Anterior (as scapular rotators); Infraspinatus, Teres Minor, Subscapularis (as humeral head stabilizers).	Suboccipital muscles (as antagonists being stretched), Sternocleidomastoid (anterior fibers).
Key Cues	"Maintain a proud chest." "Raise arms in front of your body." "Lead with your thumbs." "Avoid shrugging your shoulders toward your ears."	"Imagine a string pulling the back of your head straight back." "Create a 'double chin'." "Keep your eyes level; do not look up or down."
Common Faults	Shrugging the shoulders (upper trapezius dominance); leaning the torso; arching the lower back; raising the arms in the direct coronal plane.	Tilting the head down (flexion) instead of gliding back (retraction); jutting the chin forward on release; pushing head forcefully into the floor (supine).

Feature	**Exercise A: Active Shoulder Abduction (Scaption)**	**Exercise B: Cervical Retraction (Chin Tuck)**
Rationale for Pairing	**Shoulder Abduction** is a complex movement highly dependent on scapular position. **Cervical Retraction** directly corrects the forward head posture that causes faulty scapular mechanics, thereby creating the necessary proximal stability for safe and effective distal mobility at the shoulder. Performing retraction first primes the system for optimal performance.	

Conclusion

The analysis confirms a profound biomechanical and neuromuscular interdependence between the cervical spine and the shoulder complex. Optimal shoulder function, particularly during abduction, is predicated on a stable and well-aligned cervical and thoracic base. The prevalence of forward head posture creates a cascade of muscular imbalances and postural faults that directly compromise scapular mechanics, narrow the subacromial space, and increase the risk of shoulder impingement syndrome.

Based on this understanding, the selection of Cervical Retraction (chin tuck) as a complementary exercise to Shoulder Abduction is clinically justified. Cervical Retraction is uniquely suited to this role as it directly counteracts the pathomechanics of FHP, activates the deep cervical stabilizing muscles, and serves as a powerful neuromuscular re-education tool to reset dysfunctional postural patterns.

The recommended protocol, which prioritizes the performance of cervical retraction *before* shoulder abduction, is not arbitrary. This sequence leverages the principle of proximal stability for distal mobility, using the retraction exercise as a preparatory drill to create a favorable biomechanical environment for the subsequent dynamic movement. This integrated approach, which addresses the root postural cause rather than just the symptomatic area, represents a more robust and effective strategy for improving shoulder function, reducing injury risk, and restoring optimal movement patterns across the entire upper quarter.

Works cited

1. Anatomy, Shoulder and Upper Limb, Glenohumeral Joint - StatPearls - NCBI Bookshelf, accessed on October 12, 2025, https://www.ncbi.nlm.nih.gov/books/NBK537018/
2. The Shoulder Joint - Structure - Movement - TeachMeAnatomy, accessed on October 12, 2025, https://teachmeanatomy.info/upper-limb/joints/shoulder/
3. Current Concepts in the Scientific and Clinical Rationale Behind Exercises for Glenohumeral and Scapulothoracic Musculature | Journal of Orthopaedic & Sports Physical Therapy, accessed on October 12, 2025, https://www.jospt.org/doi/10.2519/jospt.2009.2835
4. ANATOMY AND BIOMECHANICS OF THE SHOULDER, accessed on October 12, 2025, https://courses.washington.edu/bioen520/notes/Shoulder_Anatomy_&_Biomechanics_%28Halder%29.pdf
5. Force Couple Between Rotator Cuff and Deltoid - Physiopedia, accessed on October 12, 2025, https://www.physio-pedia.com/Force_Couple_Between_Rotator_Cuff_and_Deltoid
6. Shoulder biomechanics in normal and selected pathological conditions - PMC, accessed on October 12, 2025, https://pmc.ncbi.nlm.nih.gov/articles/PMC7484714/
7. Scapular Force Couple - Muscle collaboration in the shoulder - YouTube, accessed on October 12, 2025, https://www.youtube.com/watch?v=IN2-lqX_mFc
8. Muscle coordination and scapular kinematic patterns in individuals post rotator-cuff repair - Iowa Research Online, accessed on October 12, 2025, https://iro.uiowa.edu/view/pdfCoverPage?instCode=01IOWA_INST&filePid=13873772140002771&download=true
9. The Changes in Synergistic Patterns of Peri-Shoulder Muscles During Shoulder Abduction and Flexion in Patients With Rotator Cuff Tears - PubMed, accessed on October 12, 2025, https://pubmed.ncbi.nlm.nih.gov/40911448/
10. The Changes in Synergistic Patterns of Peri-shoulder Muscles during Shoulder Abduction and Flexion in Patients with Rotator Cuff Tears - ResearchGate, accessed on October 12, 2025, https://www.researchgate.net/publication/395304189_The_Changes_in_Synergistic_Patterns_of_Peri-shoulder_Muscles_during_Shoulder_Abduction_and_Flexion_in_Patients_with_Rotator_Cuff_Tears
11. 9 Exercises to Strengthen Your Neck & Shoulders - Athletico, accessed on October 12, 2025, https://www.athletico.com/2020/10/16/9-exercises-to-strengthen-your-neck-shoulders/
12. Neck rotation modulates flexion synergy torques, indicating an ipsilateral reticulospinal source for impairment in stroke, accessed on October 12, 2025, https://pmc.ncbi.nlm.nih.gov/articles/PMC3544866/
13. Neck Stretches and Exercises for Pain Relief - WebMD, accessed on October 12, 2025, https://www.webmd.com/fitness-exercise/fitness-neck-stretches
14. Exercises for Neck and Shoulder Pain - The BioMechanics Method, accessed on October 12, 2025, https://www.thebiomechanicsmethod.com/2023/07/19/exercises-for-neck-and-shoulder-pain/

15. Decrease Neck & Shoulder Pain And Improve Posture - JUST ONE MUSCLE! - YouTube, accessed on October 12, 2025, https://www.youtube.com/watch?v=VyVsjByS4uY
16. Neck Strain or Sprain: Rehab Exercises - MyHealth Alberta, accessed on October 12, 2025, https://myhealth.alberta.ca/Health/aftercareinformation/pages/conditions.aspx?hwid=zp4468
17. Deskercise - Neck and Shoulder Stretches - YouTube, accessed on October 12, 2025, https://www.youtube.com/watch?v=5lbe9oZbpDs
18. Exercises for neck muscle and joint problems - NHS inform, accessed on October 12, 2025, https://www.nhsinform.scot/illnesses-and-conditions/muscle-bone-and-joints/neck-and-back-problems-and-conditions/exercises-for-neck-problems
19. Neck: Exercises | Kaiser Permanente, accessed on October 12, 2025, https://healthy.kaiserpermanente.org/health-wellness/health-encyclopedia/he.neck-exercises.ad1500
20. Exercises to Relieve Neck Pain - Swift Institute, accessed on October 12, 2025, https://theswiftinstitute.com/neck-exercises/
21. 5 Back and Neck Stretches to Do Every Day - HSS, accessed on October 12, 2025, https://www.hss.edu/health-library/move-better/back-neck-stretches
22. Shoulder Surgery Exercise Guide - OrthoInfo - AAOS, accessed on October 12, 2025, https://orthoinfo.aaos.org/en/recovery/shoulder-surgery-exercise-guide/