

# Introduction to Quantum Computing

## Week 4: Bloch Sphere manipulation, quantum gates, and basic quantum circuits

Taha Selim

t.i.m.m.selim2@hva.nl

**Quantum Talent and Learning Center**  
Amsterdam University of Applied Sciences

Week 4  
April 12th, 2024

## Agenda for today:

- Recap of previous sessions.
- Quick review on Bra-Ket notation, wavefunction, and qubit states.
- Gate actions and Bloch Sphere manipulation.
- Constructing basic quantum circuits; get familiar with Qiskit.

Every 30 min or so we will have a feed from other QTLC groups.

Feel free to ask questions at any time!

It is an interactive workshop, we all learn from each other!

- Join Discord and display your name instead of the nickname or the username.
- Check the invite email for the Discord link.

- Check the list of active users from all universities.
- So far, we have the list from Saxion, Fontys, and HvA.
- We are missing the list from HHS.

List needs to be sent to the QTLC team/me by today or so.

- The session is being recorded.

However, the recording will not be shared unless with get the consent of each participant. For that, we will send a form via email for all participant to tell whether

they are ok for distributing the recordings or not.

each TLC can help by distributing a form on their participants

# Recap of the previous sessions

- Classical superposition vs. quantum superposition.
- Measurements in quantum mechanics.
- Quantum mechanics relies on linear algebra.
- Matrix operations and manipulations.
- Bra-Ket notation.
- Classical bits vs. qubits.
- Qubit states.
- Bloch Sphere representation.
- Unitary matrices and quantum gates.

**Quantum mechanics:**

**Wavefunctions** are used to describe the state of a quantum system.

## Quantum mechanics:

**Wavefunctions** are used to describe the state of a quantum system.

**Example: Free particle**

$$\Psi(x) = Ae^{ikx}$$

Wavefunction is a complex-valued function, for example, of position and time.



## Quantum mechanics:

**Wavefunctions** are used to describe the state of a quantum system.

**Example: Free particle**

$$\Psi(x) = Ae^{ikx}$$

Wavefunction is a complex-valued function, for example, of position and time.

**Probability density:**

$|\Psi(x, t)|^2$  gives the probability of finding the particle at position  $x$  at time  $t$ .

# Dirac Bra-Ket Notation

We can write the wavefunction as a ket  $|\psi\rangle$ :

$$|\psi\rangle = Ae^{ikx}$$

# Dirac Bra-Ket Notation

We can write the wavefunction as a ket  $|\psi\rangle$ :

$$|\psi\rangle = Ae^{ikx}$$

Bra-Ket notation is a standard notation in quantum mechanics.

It is used to describe quantum states and operations.


It is named after Paul Dirac.

Qubit states are represented as kets.

# Dirac Bra-Ket Notation

A vector  $\vec{A}$  in a two-dimensional space can be written as a column matrix:

$$\vec{A} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$$



Using the Dirac notation, we can write the vector  $\vec{A}$  as a ket:

$$|A\rangle = a_1|x\rangle + a_2|y\rangle$$

# Dirac Bra-Ket Notation

Similarly, a vector  $\vec{B}$  in a three-dimensional space can be written as a column matrix:


$$\vec{B} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

# Dirac Bra-Ket Notation

Similarly, a vector  $\vec{B}$  in a three-dimensional space can be written as a column matrix:

$$\vec{B} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$


Example



# Dirac Bra-Ket Notation

Similarly, a vector  $\vec{B}$  in a three-dimensional space can be written as a column matrix:

$$\vec{B} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$




Using the Dirac notation, we can write the vector  $\vec{B}$  as a ket:

$$|B\rangle = b_1|x\rangle + b_2|y\rangle + b_3|z\rangle$$

# Dirac Bra-Ket Notation

In analogy to vectors, we can write the wavefunction as a ket  $|\Psi\rangle$ :

$$|\Psi\rangle = \alpha|0\rangle + \beta|1\rangle$$



with  $\alpha$  and  $\beta$  called the amplitudes of the states and they are generally complex numbers.



The norm of a 2D vector is:

$$||A|| = \sqrt{a_1^2 + a_2^2}$$

Similarly, the norm of a 3D vector is:

$$||B|| = \sqrt{b_1^2 + b_2^2 + b_3^2}$$

# Dirac Bra-Ket Notation

**Amplitudes** give the probability of finding the system in a given state when performing a measurement.

# Dirac Bra-Ket Notation

**Amplitudes** give the probability of finding the system in a given state when performing a measurement.

The **probability** of finding the system in state  $|0\rangle$  is  $|\alpha|^2$ , and the probability of finding the system in state  $|1\rangle$  is  $|\beta|^2$ .

# Dirac Bra-Ket Notation

**Amplitudes** give the probability of finding the system in a given state when performing a measurement.

The **probability** of finding the system in state  $|0\rangle$  is  $|\alpha|^2$ , and the probability of finding

the system in state  $|1\rangle$  is  $|\beta|^2$ .

The **sum** of the **probabilities** of finding the system in the two states must be equal to

1.

# Dirac Bra-Ket Notation

**Amplitudes** give the probability of finding the system in a given state when performing a measurement.

The **probability** of finding the system in state  $|0\rangle$  is  $|\alpha|^2$ , and the probability of finding

the system in state  $|1\rangle$  is  $|\beta|^2$ .

The **sum** of the **probabilities** of finding the system in the two states must be equal to

1.

Hence,

$$|\alpha|^2 + |\beta|^2 = 1$$

# Dirac Bra-Ket Notation

**Amplitudes** give the probability of finding the system in a given state when performing a measurement.

The **probability** of finding the system in state  $|0\rangle$  is  $|\alpha|^2$ , and the probability of finding

the system in state  $|1\rangle$  is  $|\beta|^2$ .

The **sum** of the **probabilities** of finding the system in the two states must be equal to

1.

Hence,

$$|\alpha|^2 + |\beta|^2 = 1$$


The particle exists by itself in a superposition of states.

# Qubit vs. classical bit

Qubit is a quantum bit.

**Classical bit:** can be in one of two states: 0 or 1.

**Qubit:** can be in a superposition of 0 and 1.



# Matrix representation of qubit

We use matrix algebra to represent qubits.

State of a single qubit:

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle \quad (1)$$

In a vector form/representation:

$$|\psi\rangle = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \quad (2)$$

The states  $|0\rangle$  and  $|1\rangle$  are represented by the following column matrices:

$$|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad |1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad (3)$$

The coefficients  $\alpha$  and  $\beta$  are complex numbers, and they are generally called the amplitudes of the states.



# Matrix representation of qubit

Experimentally, we can manipulate qubits using lasers or passing them through optical devices.

# Matrix representation of qubit

Experimentally, we can manipulate qubits using lasers or passing them through optical devices.

Changing the qubit state is equivalent to changing the amplitudes  $\alpha$  and  $\beta$ .

This can be done using the action of an unitary matrix  $U$  on the qubit.

# Matrix representation of qubit

Experimentally, we can manipulate qubits using lasers or passing them through optical devices.

Changing the qubit state is equivalent to changing the amplitudes  $\alpha$  and  $\beta$ .

This can be done using the action of an unitary matrix  $U$  on the qubit.

Let the state of the qubit be  $|\psi\rangle$ . We can change the state of the qubit to  $|\psi'\rangle$  using the action of the unitary matrix  $U$  on the qubit:

$$|\psi'\rangle = \mathbf{U}|\psi\rangle \quad (4)$$

# Matrix representation of qubit

Experimentally, we can manipulate qubits using lasers or passing them through optical devices.

Changing the qubit state is equivalent to changing the amplitudes  $\alpha$  and  $\beta$ .

This can be done using the action of an unitary matrix  $U$  on the qubit.

Let the state of the qubit be  $|\psi\rangle$ . We can change the state of the qubit to  $|\psi'\rangle$  using the action of the unitary matrix  $U$  on the qubit:

$$|\psi'\rangle = \mathbf{U}|\psi\rangle \quad (4)$$

Unitary means that the matrix  $U$  acts on the qubit without changing the norm of the qubit, i.e.  $|\alpha|^2 + |\beta|^2 = 1$ .

The matrix  $U$  is unitary if its conjugate transpose is equal to its inverse:

$$U^\dagger U = UU^\dagger = I \quad (5)$$

# Matrix representation of qubit

## Example:

What is the conjugate transpose of the following matrix:

$$A = \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \quad (6)$$

# Matrix representation of qubit

## Example:

What is the conjugate transpose of the following matrix:

$$A = \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \quad (6)$$

The conjugate transpose of the matrix  $U$  is:

$$A^\dagger = \begin{bmatrix} 1 & 1 \\ -i & -i \end{bmatrix} \quad (7)$$

# Matrix representation of qubit

## Example:

What is the conjugate transpose of the following matrix:

$$A = \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \quad (6)$$

The conjugate transpose of the matrix  $U$  is:

$$A^\dagger = \begin{bmatrix} 1 & 1 \\ -i & -i \end{bmatrix} \quad (7)$$

Is the matrix  $A$  unitary?

# Matrix representation of qubit

## Example:

What is the conjugate transpose of the following matrix:

$$A = \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \quad (6)$$

The conjugate transpose of the matrix  $U$  is:

$$A^\dagger = \begin{bmatrix} 1 & 1 \\ -i & -i \end{bmatrix} \quad (7)$$

Is the matrix  $A$  unitary?

No, the matrix  $A$  is not unitary:

$$AA^\dagger = \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -i & -i \end{bmatrix} = 2 \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \neq \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



# Matrix representation of qubit

Given the state of a qubit in  $|0\rangle$ . What is the result of applying the unitary operator

$$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \text{ to the qubit?}$$

# Matrix representation of qubit

Given the state of a qubit in  $|0\rangle$ . What is the result of applying the unitary operator

$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  to the qubit?

The result of applying the unitary operator  $X$  to the qubit is:

$$X|0\rangle = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |1\rangle \quad (8)$$

# Matrix representation of qubit

Given the state of a qubit in  $|0\rangle$ . What is the result of applying the unitary operator

$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  to the qubit?

The result of applying the unitary operator  $X$  to the qubit is:

$$X|0\rangle = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |1\rangle \quad (8)$$

Hence the matrix  $X$  flips the state of the qubit from  $|0\rangle$  to  $|1\rangle$ .

# Matrix representation of qubit

Let's now perform two successive operations on the qubit in state  $|0\rangle$ . First, we apply the unitary operator  $X$  to the qubit, and then we apply the unitary operator

$Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$  to the qubit.

What is the result of applying the unitary operator  $Y$  to the qubit?

# Matrix representation of qubit

Let's now perform two successive operations on the qubit in state  $|0\rangle$ . First, we apply the unitary operator  $X$  to the qubit, and then we apply the unitary operator

$Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$  to the qubit.

What is the result of applying the unitary operator  $Y$  to the qubit?

The result of applying the unitary operator  $Y$  to the qubit is:

$$Y |1\rangle = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -i \\ 0 \end{bmatrix} = -i |0\rangle \quad (9)$$

# Matrix representation of qubit

Let's now perform two successive operations on the qubit in state  $|0\rangle$ . First, we apply the unitary operator  $X$  to the qubit, and then we apply the unitary operator

$Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$  to the qubit.

What is the result of applying the unitary operator  $Y$  to the qubit?

The result of applying the unitary operator  $Y$  to the qubit is:


$$Y |1\rangle = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -i \\ 0 \end{bmatrix} = -i |0\rangle \quad (9)$$

Hence the matrix  $Y$  flips the state of the qubit from  $|1\rangle$  to  $-i |0\rangle$ .

# Bloch Sphere

Bloch sphere is a geometrical representation of a qubit. It is a representation of the qubit state as a vector on unit sphere.


Before we go into the details of the Bloch sphere, let's first understand the spherical coordinates and the concept of a unit sphere.



# Spherical coordinates

Spherical coordinates are a system of three coordinates used to describe the position of a point in space.

Typical coordinate systems are Cartesian, cylindrical, and spherical.  
These are orthogonal coordinate systems.







# Bloch Sphere

Bloch sphere is a geometrical representation of a qubit. It is a representation of the qubit state as a vector on unit sphere.

Before we go into the details of the Bloch sphere, let's first understand the spherical coordinates and the concept of a unit sphere.



# Bloch Sphere



- The spherical coordinates are  $(r, \theta, \phi)$ .
- The unit sphere is defined by the equation  $x^2 + y^2 + z^2 = 1$ .
- Spherical coordinates are related to the Cartesian coordinates by the following equations:

$$x = r \sin(\theta) \cos(\phi)$$

$$y = r \sin(\theta) \sin(\phi)$$


$$z = r \cos(\theta)$$

# Computer Lab

Now, open your notebook and make a function that converts the Cartesian coordinates to the spherical coordinates using the previously formulas. Make sure you can make another function to plot the transformed point on a sphere.

Note: There is difference in defining the polar angles  $\theta$  and  $\phi$  between the physics and mathematics. In physics, the polar angle  $\theta$  is measured from the positive z-axis, while in mathematics, the polar angle  $\theta$  is measured from the positive x-axis.


$$\begin{aligned}x &= r \sin(\theta) \cos(\phi) \\y &= r \sin(\theta) \sin(\phi) \\z &= r \cos(\theta)\end{aligned}$$



# Schrödinger's cat on a Blochsphere

Schrödinger's cat is determined to be alive. What location on the Earth in the figure could the cat have been before the quantum measurement?

- ① Russia.
- ② North Pole.
- ③ Australia.
- ④ All the above.



Question source: Hughes et al. (2021), Springer. Quantum Computing for the Quantum Curious.

# Schrödinger's cat on a Blochsphere

The cat could have been anywhere on Earth except for the South Pole. Notice that in Australia the cat has a smaller probability of being alive since it is further away from the North Pole.


Question source: Hughes et al. (2021), Springer. Quantum Computing for the Quantum Curious.

# Blochsphere

The state of a qubit can be represented by a point on the Blochsphere.  
This point is defined by the spherical coordinates  $(\theta, \phi)$ .  
It also gives us an idea about the superposition of the qubit.


# Blochsphere

The state of a qubit can be represented by a point on the Blochsphere. This point is defined by the spherical coordinates  $(\theta, \phi)$ . It also gives us an idea about the superposition of the qubit.




# Blochsphere

The state of a qubit can be represented by a point on the Blochsphere. This point is defined by the spherical coordinates  $(\theta, \phi)$ . It also gives us an idea about the superposition of the qubit.



$$|\psi\rangle = |0\rangle$$




$$|\psi\rangle = |1\rangle$$




# Blochsphere


The state of a qubit can be represented by a point on the Blochsphere. This point is defined by the spherical coordinates  $(\theta, \phi)$ . It also gives us an idea about the superposition of the qubit.



$$|\psi\rangle = |0\rangle$$




$$|\psi\rangle = |1\rangle$$




$$|\psi\rangle = \frac{1}{\sqrt{2}} [|0\rangle + |1\rangle]$$

# Blochsphere


The state of a qubit can be represented by a point on the Blochsphere. This point is defined by the spherical coordinates  $(\theta, \phi)$ . It also gives us an idea about the superposition of the qubit.




$$|\psi\rangle = |0\rangle$$



$$|\psi\rangle = |1\rangle$$



$$|\psi\rangle = \frac{1}{\sqrt{2}} [|0\rangle + |1\rangle]$$



$$|\psi\rangle = \frac{1}{\sqrt{2}} [|0\rangle - |1\rangle]$$

Question source: Hughes et al. (2021), Springer. Quantum Computing for the Quantum Curious.

Let's have fun with the Blochsphere!

Go to the following link and play with the Blochsphere:

First, we check the action manually:

`https://attilakun.net/bloch/`

`https://bits-and-electrons.github.io/bloch-sphere-simulator/`

Then, we check the action using the IBM Quantum Experience:

`https://quantum.ibm.com/`

Now, open your IBM Quantum Experience account.

We are going to learn how to initialize qubits, create quantum circuits, manipulate qubits, and measure them.

First, let's understand the effect of gates on the qubit state using the Blochsphere.

# What is a qubit made of?

Classical bits are made of transistors.

It is easy to realize a classical bit in practice. For example, if the voltage is above a certain threshold, the bit is 1, otherwise it is 0.

Qubits are made of quantum systems in a sophisticated way.

We cannot simply create a superposition of 0 and 1 by using a transistor in which two currents are there, one flows and the other does not.

Quantum computers are still in their infancy. There are several technologies being developed to realize qubits. Still is a long way to go. It is unclear which candidate will provide more benefits than the other or the best qubits.

Answers will be there when quantum computers are more mature and more commercially available.

# What is a qubit made of?

A qubit can be made of different physical systems.

- **Superconducting qubits:**

- Made of superconducting circuits.
- They are cooled to very low temperatures.
- They are the most widely used qubits in quantum computers.

- **Trapped ions:**


- Made of individual atoms.
- They are trapped using electromagnetic fields.
- They are cooled to very low temperatures.

# What is a qubit made of?

- **Photonic qubits:**
  - Made of photons.
  - They are used in quantum communication.
- **Topological qubits:**
  - Made of exotic particles called anyons.
  - They are not yet realized in practice.

# Check your understanding!

If the qubit represented by the figure is measured, what are the possible outcomes?  
Numerical values for the amplitudes are not needed, only conceptual statements.





- Hughes, R. J., Nordholt, J. E., Mink, A., & Lanzagorta, M. (2021). Quantum Computing for the Quantum Curious. Springer.
- Nielsen, M. A., & Chuang, I. L. (2010). Quantum Computation and Quantum Information: 10th Anniversary Edition. Cambridge University Press.
- IBM Quantum Experience: <https://quantum-computing.ibm.com/>

Thank you

Thank you!