

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4}$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49}$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|}$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a}$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a} = \text{unit vector parallel to } \mathbf{a}.$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a} = \text{unit vector parallel to } \mathbf{a}.$$

- So,

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a} = \text{unit vector parallel to } \mathbf{a}.$$

- So,

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\mathbf{b} \cdot \mathbf{n})\mathbf{n} = \frac{\mathbf{b} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a}$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a} = \text{unit vector parallel to } \mathbf{a}.$$

- So,

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\mathbf{b} \cdot \mathbf{n})\mathbf{n} = \frac{\mathbf{b} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a} =$$

$$\frac{1}{49} \langle 1, 2, 3 \rangle \cdot \langle 3, 6, -2 \rangle \langle 3, 6, -2 \rangle$$

Problem 1(a) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write down the vector projection of \mathbf{b} along \mathbf{a} . (Hint: Use projections.)

Solution:

- We have $|\mathbf{a}| = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$.
- Then

$$\mathbf{n} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{7}\mathbf{a} = \text{unit vector parallel to } \mathbf{a}.$$

- So,

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\mathbf{b} \cdot \mathbf{n})\mathbf{n} = \frac{\mathbf{b} \cdot \mathbf{a}}{|\mathbf{a}|^2} \mathbf{a} =$$

$$\frac{1}{49} \langle 1, 2, 3 \rangle \cdot \langle 3, 6, -2 \rangle \langle 3, 6, -2 \rangle = \frac{9}{49} \langle 3, 6, -2 \rangle.$$

Problem 1(b) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Problem 1(b) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- We have

$$\mathbf{b} = \langle 1, 2, 3 \rangle$$

Problem 1(b) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- We have

$$\mathbf{b} = \langle 1, 2, 3 \rangle = \langle 1, 2, 3 \rangle - \frac{9}{49} \langle 3, 6, -2 \rangle + \frac{9}{49} \langle 3, 6, -2 \rangle$$

Problem 1(b) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- We have

$$\begin{aligned}\mathbf{b} = \langle 1, 2, 3 \rangle &= \langle 1, 2, 3 \rangle - \frac{9}{49} \langle 3, 6, -2 \rangle + \frac{9}{49} \langle 3, 6, -2 \rangle \\ &= \frac{1}{49} \langle 22, 44, 165 \rangle + \frac{9}{49} \langle 3, 6, -2 \rangle.\end{aligned}$$

Problem 1(b) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- We have

$$\begin{aligned}\mathbf{b} = \langle 1, 2, 3 \rangle &= \langle 1, 2, 3 \rangle - \frac{9}{49} \langle 3, 6, -2 \rangle + \frac{9}{49} \langle 3, 6, -2 \rangle \\ &= \frac{1}{49} \langle 22, 44, 165 \rangle + \frac{9}{49} \langle 3, 6, -2 \rangle.\end{aligned}$$

- Here

$$\frac{9}{49} \langle 3, 6, -2 \rangle \text{ parallel to } \mathbf{a} = \langle 3, 6, -2 \rangle$$

and

$$\frac{1}{49} \langle 22, 44, 165 \rangle \text{ orthogonal to } \mathbf{a} = \langle 3, 6, -2 \rangle.$$



Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

- Of course this is the same as

$$\mathbf{b} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\mathbf{b}) + \text{proj}_{\mathbf{a}}\mathbf{b}.$$

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

- Of course this is the same as

$$\mathbf{b} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\mathbf{b}) + \text{proj}_{\mathbf{a}}\mathbf{b}.$$

That is, we write \mathbf{b} as $\text{proj}_{\mathbf{a}}\mathbf{b}$ plus “the rest”.

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

- Of course this is the same as

$$\mathbf{b} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\mathbf{b}) + \text{proj}_{\mathbf{a}}\mathbf{b}.$$

That is, we write \mathbf{b} as $\text{proj}_{\mathbf{a}}\mathbf{b}$ plus “the rest”. But “the rest” is orthogonal to \mathbf{n} (and to \mathbf{a}), since

$$(\mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n}) \cdot \mathbf{n}$$

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

- Of course this is the same as

$$\mathbf{b} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\mathbf{b}) + \text{proj}_{\mathbf{a}}\mathbf{b}.$$

That is, we write \mathbf{b} as $\text{proj}_{\mathbf{a}}\mathbf{b}$ plus “the rest”. But “the rest” is orthogonal to \mathbf{n} (and to \mathbf{a}), since

$$(\mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n}) \cdot \mathbf{n} = \mathbf{b} \cdot \mathbf{n} - (\mathbf{b} \cdot \mathbf{n})(\mathbf{n} \cdot \mathbf{n})$$

Problem 1(b) Continuation - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Write \mathbf{b} as a sum of a vector parallel to \mathbf{a} and a vector orthogonal to \mathbf{a} . (Hint: Use projections.)

Solution:

- Why so? All we did was to write

$$\mathbf{b} = \mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n} + (\mathbf{b} \cdot \mathbf{n})\mathbf{n}$$

where $\mathbf{n} = \frac{\mathbf{a}}{7}$, $\mathbf{n}^2 = 1$.

- Of course this is the same as

$$\mathbf{b} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\mathbf{b}) + \text{proj}_{\mathbf{a}}\mathbf{b}.$$

That is, we write \mathbf{b} as $\text{proj}_{\mathbf{a}}\mathbf{b}$ plus “the rest”. But “the rest” is orthogonal to \mathbf{n} (and to \mathbf{a}), since

$$(\mathbf{b} - (\mathbf{b} \cdot \mathbf{n})\mathbf{n}) \cdot \mathbf{n} = \mathbf{b} \cdot \mathbf{n} - (\mathbf{b} \cdot \mathbf{n})(\mathbf{n} \cdot \mathbf{n}) = 0, \text{ as } \mathbf{n} \cdot \mathbf{n} = 1.$$



Problem 1(c) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Let θ be the angle between \mathbf{a} and \mathbf{b} . Find $\cos \theta$.

Problem 1(c) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Let θ be the angle between \mathbf{a} and \mathbf{b} . Find $\cos \theta$.

Solution:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$$

Problem 1(c) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Let θ be the angle between \mathbf{a} and \mathbf{b} . Find $\cos \theta$.

Solution:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \frac{\langle 3, 6, -2 \rangle \cdot \langle 1, 2, 3 \rangle}{|\langle 3, 6, -2 \rangle||\langle 1, 2, 3 \rangle|}$$

Problem 1(c) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Let θ be the angle between \mathbf{a} and \mathbf{b} . Find $\cos \theta$.

Solution:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \frac{\langle 3, 6, -2 \rangle \cdot \langle 1, 2, 3 \rangle}{|\langle 3, 6, -2 \rangle||\langle 1, 2, 3 \rangle|} = \frac{9}{\sqrt{49}\sqrt{14}}$$

Problem 1(c) - Spring 2009

Given $\mathbf{a} = \langle 3, 6, -2 \rangle$, $\mathbf{b} = \langle 1, 2, 3 \rangle$.

Let θ be the angle between \mathbf{a} and \mathbf{b} . Find $\cos \theta$.

Solution:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \frac{\langle 3, 6, -2 \rangle \cdot \langle 1, 2, 3 \rangle}{|\langle 3, 6, -2 \rangle||\langle 1, 2, 3 \rangle|} = \frac{9}{\sqrt{49}\sqrt{14}} = \frac{9}{7\sqrt{14}}.$$



Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line.

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line. For example, take the point to be A and the vector to be \overrightarrow{AB} .

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line. For example, take the point to be A and the vector to be \overrightarrow{AB} .
- The vector equation of L is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB}$$

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line. For example, take the point to be A and the vector to be \overrightarrow{AB} .
- The vector equation of L is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle -1, 7, 5 \rangle + t \langle 4, -5, -3 \rangle$$

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line. For example, take the point to be A and the vector to be \overrightarrow{AB} .
- The vector equation of L is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle -1, 7, 5 \rangle + t \langle 4, -5, -3 \rangle = \langle -1 + 4t, 7 - 5t, 5 - 3t \rangle,$$

where O is the origin.

Problem 2(a) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.

Let L be the line which passes through the points $A = (-1, 7, 5)$ and $B = (3, 2, 2)$. Find the parametric equations for L .

Solution:

- To get **parametric equations** for L you need a point through which the line passes and a vector parallel to the line. For example, take the point to be A and the vector to be \overrightarrow{AB} .
- The vector equation of L is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle -1, 7, 5 \rangle + t \langle 4, -5, -3 \rangle = \langle -1 + 4t, 7 - 5t, 5 - 3t \rangle,$$

where O is the origin.

- The **parametric equations** are:

$$\begin{cases} x = -1 + 4t \\ y = 7 - 5t, \\ z = 5 - 3t \end{cases} \quad t \in \mathbb{R}.$$



Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Solution:

Denote the fourth vertex by D .

Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Solution:

Denote the fourth vertex by D . Then

$$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{CB}$$

Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Solution:

Denote the fourth vertex by D . Then

$$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{CB} = \langle -1, 7, 5 \rangle + \langle 2, 0, -1 \rangle$$

Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Solution:

Denote the fourth vertex by D . Then

$$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{CB} = \langle -1, 7, 5 \rangle + \langle 2, 0, -1 \rangle = \langle 1, 7, 4 \rangle,$$

where O is the origin.

Problem 2(b) - Spring 2009

Given $A = (-1, 7, 5)$, $B = (3, 2, 2)$ and $C = (1, 2, 3)$.
 A , B and C are three of the four vertices of a parallelogram,
while CA and CB are two of the four edges. Find the fourth
vertex.

Solution:

Denote the fourth vertex by D . Then

$$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{CB} = \langle -1, 7, 5 \rangle + \langle 2, 0, -1 \rangle = \langle 1, 7, 4 \rangle,$$

where O is the origin. That is,

$$D = (1, 7, 4).$$



Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 .
Find an equation for the plane containing P , Q and R .

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 .
Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} .

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR}$$

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & -2 & -3 \\ 0 & -2 & -4 \end{vmatrix}$$

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & -2 & -3 \\ 0 & -2 & -4 \end{vmatrix} = \langle 2, -12, 6 \rangle$$

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & -2 & -3 \\ 0 & -2 & -4 \end{vmatrix} = \langle 2, -12, 6 \rangle = 2\langle 1, -6, 3 \rangle.$$

Problem 3(a) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find an equation for the plane containing P , Q and R .

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point P , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & -2 & -3 \\ 0 & -2 & -4 \end{vmatrix} = \langle 2, -12, 6 \rangle = 2\langle 1, -6, 3 \rangle.$$

So the **equation of the plane** is:

$$(x - 1) - 6(y - 3) + 3(z - 5) = 0.$$



Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 .
Find the area of the triangle with vertices P , Q , R .

Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find the area of the triangle with vertices P , Q , R .

Solution:

The area of the triangle \triangle with vertices P , Q , R can be found by taking the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} and dividing it by 2.

Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find the area of the triangle with vertices P , Q , R .

Solution:

The area of the triangle Δ with vertices P , Q , R can be found by taking the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} and dividing it by 2. Thus, using **a)**, we have:

$$\text{Area}(\Delta) = \frac{|\overrightarrow{PQ} \times \overrightarrow{PR}|}{2}$$

Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find the area of the triangle with vertices P , Q , R .

Solution:

The area of the triangle Δ with vertices P , Q , R can be found by taking the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} and dividing it by 2. Thus, using **a)**, we have:

$$\text{Area}(\Delta) = \frac{|\overrightarrow{PQ} \times \overrightarrow{PR}|}{2} = \frac{1}{2} |2\langle 1, -6, 3 \rangle|$$

Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find the area of the triangle with vertices P , Q , R .

Solution:

The area of the triangle Δ with vertices P , Q , R can be found by taking the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} and dividing it by 2. Thus, using **a)**, we have:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\overrightarrow{PQ} \times \overrightarrow{PR}|}{2} = \frac{1}{2} |2\langle 1, -6, 3 \rangle| \\ &= \sqrt{1 + 36 + 9}\end{aligned}$$

Problem 3(b) - Spring 2009

Consider the points $P(1, 3, 5)$, $Q(-2, 1, 2)$, $R(1, 1, 1)$ in \mathbb{R}^3 . Find the area of the triangle with vertices P , Q , R .

Solution:

The area of the triangle Δ with vertices P , Q , R can be found by taking the area of the parallelogram spanned by \vec{PQ} and \vec{PR} and dividing it by 2. Thus, using **a)**, we have:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{PQ} \times \vec{PR}|}{2} = \frac{1}{2} |2\langle 1, -6, 3 \rangle| \\ &= \sqrt{1 + 36 + 9} = \sqrt{46}.\end{aligned}$$



Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Solution:

- A vector \mathbf{v} parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, 1, 3 \rangle \times \langle 1, -1, 2 \rangle$$

Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Solution:

- A vector \mathbf{v} parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, 1, 3 \rangle \times \langle 1, -1, 2 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 3 \\ 1 & -1 & 2 \end{vmatrix}$$

Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Solution:

- A vector \mathbf{v} parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, 1, 3 \rangle \times \langle 1, -1, 2 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 3 \\ 1 & -1 & 2 \end{vmatrix} = \langle 5, 1, -2 \rangle.$$

- A point on \mathbf{L} is any (x_0, y_0, z_0) that satisfies the equations of **both** planes.

Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Solution:

- A vector \mathbf{v} parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, 1, 3 \rangle \times \langle 1, -1, 2 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 3 \\ 1 & -1 & 2 \end{vmatrix} = \langle 5, 1, -2 \rangle.$$

- A point on \mathbf{L} is any (x_0, y_0, z_0) that satisfies the equations of **both** planes.
- Setting $z = 0$, we obtain the equations $x + y = 1$ and $x - y = 1$ and find such a point $(\frac{1}{2}, \frac{1}{2}, 0)$.

Problem 4 - Spring 2009

Find parametric equations for the line of intersection of the planes $x + y + 3z = 1$ and $x - y + 2z = 0$.

Solution:

- A vector \mathbf{v} parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, 1, 3 \rangle \times \langle 1, -1, 2 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 3 \\ 1 & -1 & 2 \end{vmatrix} = \langle 5, 1, -2 \rangle.$$

- A point on \mathbf{L} is any (x_0, y_0, z_0) that satisfies the equations of **both** planes.
- Setting $z = 0$, we obtain the equations $x + y = 1$ and $x - y = 1$ and find such a point $(\frac{1}{2}, \frac{1}{2}, 0)$. Therefore **parametric equations** for \mathbf{L} are:

$$\begin{cases} x = \frac{1}{2} + 5t \\ y = \frac{1}{2} + t \\ z = -2t. \end{cases}$$



Problem 6(a) - Spring 2009

Consider the sphere **S** in \mathbb{R}^3 given by the equation

$$x^2 + y^2 + z^2 - 4x - 6z - 3 = 0.$$

Find its center **C** and its radius **R**.

Problem 6(a) - Spring 2009

Consider the sphere **S** in \mathbb{R}^3 given by the equation

$$x^2 + y^2 + z^2 - 4x - 6z - 3 = 0.$$

Find its center **C** and its radius **R**.

Solution:

- Completing the square we get

$$(x - 2)^2 - 4 + y^2 + (z - 3)^2 - 9 - 3 = 0$$

$$\iff$$

$$(x - 2)^2 + y^2 + (z - 3)^2 = 16.$$

Problem 6(a) - Spring 2009

Consider the sphere **S** in \mathbb{R}^3 given by the equation

$$x^2 + y^2 + z^2 - 4x - 6z - 3 = 0.$$

Find its center **C** and its radius **R**.

Solution:

- Completing the square we get

$$(x - 2)^2 - 4 + y^2 + (z - 3)^2 - 9 - 3 = 0$$

$$\iff$$

$$(x - 2)^2 + y^2 + (z - 3)^2 = 16.$$

- This gives:

$$\mathbf{C} = (2, 0, 3) \quad \mathbf{R} = 4$$



Problem 6(b) - Spring 2009

What does the equation $x^2 + z^2 = 4$ describe in \mathbb{R}^3 ? Make a sketch.

Problem 6(b) - Spring 2009

What does the equation $x^2 + z^2 = 4$ describe in \mathbb{R}^3 ? Make a sketch.

Solution:

- This is a (straight, circular) **cylinder** determined by the circle in the xz -plane of radius 2 and center $(0,0)$ and parallel to the y -axis.



Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.
- Take the point to be A and the vector to be the \overrightarrow{AB} .

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.
- Take the point to be A and the vector to be the \overrightarrow{AB} .
- The vector equation of **L** is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB}$$

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.
- Take the point to be A and the vector to be the \overrightarrow{AB} .
- The vector equation of **L** is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle 1, 2, 3 \rangle + t \langle 3, 4, 2 \rangle$$

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.
- Take the point to be A and the vector to be the \overrightarrow{AB} .
- The vector equation of **L** is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle 1, 2, 3 \rangle + t \langle 3, 4, 2 \rangle = \langle 1 + 3t, 2 + 4t, 3 + 2t \rangle,$$

where O is the origin.

Problem 1(a) - Fall 2008

Find **parametric equations** for the line **L** which contains $A(1, 2, 3)$ and $B(4, 6, 5)$.

Solution:

- To get the **parametric equations** of **L** you need a point through which the line passes and a vector parallel to the line.
- Take the point to be A and the vector to be the \overrightarrow{AB} .
- The vector equation of **L** is

$$\mathbf{r}(t) = \overrightarrow{OA} + t\overrightarrow{AB} = \langle 1, 2, 3 \rangle + t \langle 3, 4, 2 \rangle = \langle 1 + 3t, 2 + 4t, 3 + 2t \rangle,$$

where O is the origin.

- The **parametric equations** are:

$$\begin{aligned}x &= 1 + 3t \\y &= 2 + 4t, \quad t \in \mathbb{R}. \\z &= 3 + 2t\end{aligned}$$



Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$.

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle$$

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix}$$

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero.

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero. Setting $z = 0$, we obtain:

$$x - 2y = 10$$

$$2x + y = 0.$$

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero. Setting $z = 0$, we obtain:

$$x - 2y = 10$$

$$2x + y = 0.$$

Solving, we find that $x = 2$ and $y = -4$.

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero. Setting $z = 0$, we obtain:

$$x - 2y = 10$$

$$2x + y = 0.$$

Solving, we find that $x = 2$ and $y = -4$. Hence, $P = \langle 2, -4, 0 \rangle$ lies on the line **L**.

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero. Setting $z = 0$, we obtain:

$$x - 2y = 10$$

$$2x + y = 0.$$

Solving, we find that $x = 2$ and $y = -4$. Hence,

$P = \langle 2, -4, 0 \rangle$ lies on the line **L**.

- The **parametric equations** are:

Problem 1(b) - Fall 2008

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 10$ and $2x + y - z = 0$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, -1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}.$$

- Choose $P \in \mathbf{L}$ so the z -coordinate of P is zero. Setting $z = 0$, we obtain:

$$x - 2y = 10$$

$$2x + y = 0.$$

Solving, we find that $x = 2$ and $y = -4$. Hence, $P = \langle 2, -4, 0 \rangle$ lies on the line **L**.

- The **parametric equations** are:

$$x = 2 + t$$

$$y = -4 + 3t$$

$$z = 0 + 5t = 5t.$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 1

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -2, 0 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -2 \rangle$ which lie parallel to the plane.

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 1

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -2, 0 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -2 \rangle$ which lie parallel to the plane.
- Then consider the normal vector:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR}$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 1

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -2, 0 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -2 \rangle$ which lie parallel to the plane.
- Then consider the normal vector:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 0 \\ 3 & 0 & -2 \end{vmatrix}$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 1

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -2, 0 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -2 \rangle$ which lie parallel to the plane.
- Then consider the normal vector:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 0 \\ 3 & 0 & -2 \end{vmatrix} = 4\mathbf{i} + 4\mathbf{j} + 6\mathbf{k}.$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 1

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -2, 0 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -2 \rangle$ which lie parallel to the plane.
- Then consider the normal vector:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 0 \\ 3 & 0 & -2 \end{vmatrix} = 4\mathbf{i} + 4\mathbf{j} + 6\mathbf{k}.$$

- So the **equation of the plane** is given by:

$$\langle 4, 4, 6 \rangle \cdot \langle x + 1, y, z - 1 \rangle = 4(x + 1) + 4y + 6(z - 1) = 0.$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 2

- The plane consists of all the points $S(x, y, z) \in \mathbb{R}^3$, such that \overrightarrow{PS} , \overrightarrow{PQ} and \overrightarrow{PR} are in the same plane (coplanar).

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 2

- The plane consists of all the points $S(x, y, z) \in \mathbb{R}^3$, such that \overrightarrow{PS} , \overrightarrow{PQ} and \overrightarrow{PR} are in the same plane (coplanar).
- But this happens if and only if their box product is zero.

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 2

- The plane consists of all the points $S(x, y, z) \in \mathbb{R}^3$, such that \overrightarrow{PS} , \overrightarrow{PQ} and \overrightarrow{PR} are in the same plane (coplanar).
- But this happens if and only if their box product is zero.
- So the **equation of the plane** is:

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

Method 2

- The plane consists of all the points $S(x, y, z) \in \mathbb{R}^3$, such that \overrightarrow{PS} , \overrightarrow{PQ} and \overrightarrow{PR} are in the same plane (coplanar).
- But this happens if and only if their box product is zero.
- So the **equation of the plane** is:

$$\begin{vmatrix} x+1 & y & z-1 \\ 2 & -2 & 0 \\ 3 & 0 & -2 \end{vmatrix}$$

Problem 2(a) - Fall 2008

Find an **equation of the plane** which contains the points $P(-1, 0, 1)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.


Solution:

Method 2

- The plane consists of all the points $S(x, y, z) \in \mathbb{R}^3$, such that \overrightarrow{PS} , \overrightarrow{PQ} and \overrightarrow{PR} are in the same plane (coplanar).
- But this happens if and only if their box product is zero.
- So the **equation of the plane** is:


$$\begin{vmatrix} x+1 & y & z-1 \\ 2 & -2 & 0 \\ 3 & 0 & -2 \end{vmatrix} = 4(x+1) + 4y + 6(z-1) = 0.$$





Problem 2(b) - Fall 2008

Find the distance D from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.




Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.




Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $2x + y - 2z - 19 = 0$.




Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $2x + y - 2z - 19 = 0$.
- So, the distance from $(1, 2, -1)$ to the plane is:




Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $2x + y - 2z - 19 = 0$.
- So, the distance from $(1, 2, -1)$ to the plane is:

$$D = \frac{|(2 \cdot 1) + (1 \cdot 6) + (-2 \cdot -1) - 19|}{\sqrt{2^2 + 1^2 + (-2)^2}}$$




Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $2x + y - 2z - 19 = 0$.
- So, the distance from $(1, 2, -1)$ to the plane is:

$$D = \frac{|(2 \cdot 1) + (1 \cdot 6) + (-2 \cdot -1) - 19|}{\sqrt{2^2 + 1^2 + (-2)^2}} = \frac{|-9|}{\sqrt{9}}$$



Problem 2(b) - Fall 2008

Find the distance **D** from the point $(1, 6, -1)$ to the plane $2x + y - 2z = 19$.

Solution:

- Recall the distance formula $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $2x + y - 2z - 19 = 0$.
- So, the distance from $(1, 2, -1)$ to the plane is:

$$D = \frac{|(2 \cdot 1) + (1 \cdot 6) + (-2 \cdot -1) - 19|}{\sqrt{2^2 + 1^2 + (-2)^2}} = \frac{|-9|}{\sqrt{9}} = 3.$$

Problem 2(c) - Fall 2008

Find the point Q in the plane $2x + y - 2z = 19$ which is closest to the point $(1, 6, -1)$. (Hint: You can use part b) of this problem to help find Q or first find the equation of the line L passing through Q and the point $(1, 6, -1)$ and then solve for Q .)

Problem 2(c) - Fall 2008

Find the point Q in the plane $2x + y - 2z = 19$ which is closest to the point $(1, 6, -1)$. (Hint: You can use part b) of this problem to help find Q or first find the equation of the line L passing through Q and the point $(1, 6, -1)$ and then solve for Q .)

Solution:

- The line L in the Hint passes through $(1, 6, -1)$ and is parallel to $\mathbf{n} = \langle 2, 1, -2 \rangle$.

Problem 2(c) - Fall 2008

Find the point Q in the plane $2x + y - 2z = 19$ which is closest to the point $(1, 6, -1)$. (Hint: You can use part b) of this problem to help find Q or first find the equation of the line L passing through Q and the point $(1, 6, -1)$ and then solve for Q .)

Solution:

- The line L in the Hint passes through $(1, 6, -1)$ and is parallel to $\mathbf{n} = \langle 2, 1, -2 \rangle$.
- So, L has **parametric equations**:

$$\begin{aligned}x &= 1 + 2t \\y &= 6 + t, \\z &= -1 - 2t\end{aligned} \quad t \in \mathbb{R}.$$

Problem 2(c) - Fall 2008

Find the point Q in the plane $2x + y - 2z = 19$ which is closest to the point $(1, 6, -1)$. (Hint: You can use part b) of this problem to help find Q or first find the equation of the line L passing through Q and the point $(1, 6, -1)$ and then solve for Q .)

Solution:

- The line L in the Hint passes through $(1, 6, -1)$ and is parallel to $\mathbf{n} = \langle 2, 1, -2 \rangle$.
- So, L has **parametric equations**:

$$\begin{aligned}x &= 1 + 2t \\y &= 6 + t, \quad t \in \mathbb{R}. \\z &= -1 - 2t\end{aligned}$$

- L intersects the plane $2x + y - 2z = 19$ if and only if
$$2(1 + 2t) + (6 + t) - 2(-1 - 2t) = 19 \iff 9t = 9 \iff t = 1.$$

Problem 2(c) - Fall 2008

Find the point Q in the plane $2x + y - 2z = 19$ which is closest to the point $(1, 6, -1)$. (Hint: You can use part b) of this problem to help find Q or first find the equation of the line L passing through Q and the point $(1, 6, -1)$ and then solve for Q .)

Solution:

- The line L in the Hint passes through $(1, 6, -1)$ and is parallel to $\mathbf{n} = \langle 2, 1, -2 \rangle$.
- So, L has **parametric equations**:

$$\begin{aligned}x &= 1 + 2t \\y &= 6 + t, & t \in \mathbb{R}. \\z &= -1 - 2t\end{aligned}$$

- L intersects the plane $2x + y - 2z = 19$ if and only if
$$2(1 + 2t) + (6 + t) - 2(-1 - 2t) = 19 \iff 9t = 9 \iff t = 1.$$
- Substituting $t = 1$ in the **parametric equations** of L gives the point $Q = (3, 7, -3)$.



Problem 3(a) - Fall 2008

Find the volume V of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Problem 3(a) - Fall 2008

Find the volume **V** of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The **parallelepiped** is determined by its edges

$$\overrightarrow{AB} = \langle 0, -3, -2 \rangle, \quad \overrightarrow{AC} = \langle 1, 1, -3 \rangle, \quad \overrightarrow{AD} = \langle -2, -4, -1 \rangle.$$

Problem 3(a) - Fall 2008

Find the volume **V** of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The **parallelepiped** is determined by its edges

$$\overrightarrow{AB} = \langle 0, -3, -2 \rangle, \quad \overrightarrow{AC} = \langle 1, 1, -3 \rangle, \quad \overrightarrow{AD} = \langle -2, -4, -1 \rangle.$$

Its volume can be computed as the absolute value of the box product $\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})$,

Problem 3(a) - Fall 2008

Find the volume V of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The **parallelepiped** is determined by its edges

$$\overrightarrow{AB} = \langle 0, -3, -2 \rangle, \quad \overrightarrow{AC} = \langle 1, 1, -3 \rangle, \quad \overrightarrow{AD} = \langle -2, -4, -1 \rangle.$$

Its volume can be computed as the absolute value of the box product $\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})$, i.e.,

$$V = \left| \begin{vmatrix} 0 & -3 & -2 \\ 1 & 1 & -3 \\ -2 & -4 & -1 \end{vmatrix} \right|$$

Problem 3(a) - Fall 2008

Find the volume V of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The **parallelepiped** is determined by its edges

$$\overrightarrow{AB} = \langle 0, -3, -2 \rangle, \quad \overrightarrow{AC} = \langle 1, 1, -3 \rangle, \quad \overrightarrow{AD} = \langle -2, -4, -1 \rangle.$$

Its volume can be computed as the absolute value of the box product $\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})$, i.e.,

$$V = \left| \begin{vmatrix} 0 & -3 & -2 \\ 1 & 1 & -3 \\ -2 & -4 & -1 \end{vmatrix} \right| = |3(-1 - 6) - 2(-4 + 2)|$$

Problem 3(a) - Fall 2008

Find the volume **V** of the **parallelepiped** such that the following four points $A = (3, 4, 0)$, $B = (3, 1, -2)$, $C = (4, 5, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The **parallelepiped** is determined by its edges

$$\overrightarrow{AB} = \langle 0, -3, -2 \rangle, \quad \overrightarrow{AC} = \langle 1, 1, -3 \rangle, \quad \overrightarrow{AD} = \langle -2, -4, -1 \rangle.$$

Its volume can be computed as the absolute value of the box product $\overrightarrow{AB} \cdot (\overrightarrow{AC} \times \overrightarrow{AD})$, i.e.,

$$\mathbf{V} = \left| \begin{vmatrix} 0 & -3 & -2 \\ 1 & 1 & -3 \\ -2 & -4 & -1 \end{vmatrix} \right| = |3(-1 - 6) - 2(-4 + 2)| = |-17| = 17.$$



Problem 3(b) - Fall 2008

Find the **center** and **radius** of the sphere

$$x^2 - 4x + y^2 + 4y + z^2 = 8.$$

Problem 3(b) - Fall 2008

Find the **center** and **radius** of the sphere

$$x^2 - 4x + y^2 + 4y + z^2 = 8.$$

Solution:

- Completing the square we get

$$x^2 - 4x + y^2 + 4y + z^2 = (x^2 - 4x + 4) - 4 + (y^2 + 4y + 4) - 4 + (z^2)$$

Problem 3(b) - Fall 2008

Find the **center** and **radius** of the sphere

$$x^2 - 4x + y^2 + 4y + z^2 = 8.$$

Solution:

- Completing the square we get

$$x^2 - 4x + y^2 + 4y + z^2 = (x^2 - 4x + 4) - 4 + (y^2 + 4y + 4) - 4 + (z^2)$$

$$= (x - 2)^2 - 4 + (y + 2)^2 - 4 + z^2 = 8$$

$$\iff$$

$$(x - 2)^2 + (y + 2)^2 + z^2 = 16.$$

Problem 3(b) - Fall 2008

Find the **center** and **radius** of the sphere

$$x^2 - 4x + y^2 + 4y + z^2 = 8.$$

Solution:

- Completing the square we get

$$x^2 - 4x + y^2 + 4y + z^2 = (x^2 - 4x + 4) - 4 + (y^2 + 4y + 4) - 4 + (z^2)$$

$$= (x - 2)^2 - 4 + (y + 2)^2 - 4 + z^2 = 8$$

$$\iff$$

$$(x - 2)^2 + (y + 2)^2 + z^2 = 16.$$

- This gives:

$$\text{Center} = (2, -2, 0)$$

$$\text{Radius} = 4$$



Problem 5(a) - Fall 2008

Consider the points $A(2, 1, 0)$, $B(3, 0, 2)$ and $C(0, 2, 1)$. Find the area of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Problem 5(a) - Fall 2008

Consider the points $A(2, 1, 0)$, $B(3, 0, 2)$ and $C(0, 2, 1)$. Find the area of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- Recall that:

$$\text{Area} = \frac{1}{2} \left| \overrightarrow{AB} \times \overrightarrow{AC} \right|.$$

Problem 5(a) - Fall 2008

Consider the points $A(2, 1, 0)$, $B(3, 0, 2)$ and $C(0, 2, 1)$. Find the area of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- Recall that:

$$\text{Area} = \frac{1}{2} \left| \overrightarrow{AB} \times \overrightarrow{AC} \right|.$$

- Since $\overrightarrow{AB} = \langle 1, -1, 2 \rangle$ and $\overrightarrow{AC} = \langle -2, 1, 1 \rangle$,

Problem 5(a) - Fall 2008

Consider the points $A(2, 1, 0)$, $B(3, 0, 2)$ and $C(0, 2, 1)$. Find the area of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- Recall that:

$$\text{Area} = \frac{1}{2} \left| \overrightarrow{AB} \times \overrightarrow{AC} \right|.$$

- Since $\overrightarrow{AB} = \langle 1, -1, 2 \rangle$ and $\overrightarrow{AC} = \langle -2, 1, 1 \rangle$,

$$\text{Area} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\| = \frac{1}{2} |\langle -3, -5, -1 \rangle|$$

Problem 5(a) - Fall 2008

Consider the points $A(2, 1, 0)$, $B(3, 0, 2)$ and $C(0, 2, 1)$. Find the area of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- Recall that:

$$\text{Area} = \frac{1}{2} \left| \overrightarrow{AB} \times \overrightarrow{AC} \right|.$$

- Since $\overrightarrow{AB} = \langle 1, -1, 2 \rangle$ and $\overrightarrow{AC} = \langle -2, 1, 1 \rangle$,

$$\text{Area} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\| = \frac{1}{2} |\langle -3, -5, -1 \rangle| = \frac{1}{2} \sqrt{35}.$$



Problem 5(b) - Fall 2008

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(2, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Problem 5(b) - Fall 2008

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(2, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by **S**.

Problem 5(b) - Fall 2008

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(2, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by S . Then

$$\overrightarrow{OS} = \overrightarrow{OQ} + \overrightarrow{PR} = \langle 0, 1, 0 \rangle + \langle 2, 2, 0 \rangle = \langle 2, 3, 0 \rangle,$$

where O is the origin.

Problem 5(b) - Fall 2008

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(2, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by S . Then

$$\overrightarrow{OS} = \overrightarrow{OQ} + \overrightarrow{PR} = \langle 0, 1, 0 \rangle + \langle 2, 2, 0 \rangle = \langle 2, 3, 0 \rangle,$$

where O is the origin. That is,

$$S = (2, 3, 0).$$



Problem 6(a) - Spring 2008

Find an **equation of the plane** through the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Problem 6(a) - Spring 2008

Find an **equation of the plane** through the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point A , it suffices to find \mathbf{n} .

Problem 6(a) - Spring 2008

Find an **equation of the plane** through the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point A , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{vmatrix} = \langle -1, 1, 2 \rangle.$$

Problem 6(a) - Spring 2008

Find an **equation of the plane** through the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Solution:

Since a plane is determined by its normal vector \mathbf{n} and a point on it, say the point A , it suffices to find \mathbf{n} . Note that:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{vmatrix} = \langle -1, 1, 2 \rangle.$$

So the **equation of the plane** is:

$$-(x - 1) + (y - 2) + 2(z - 3) = 0.$$



Problem 6(b) - Spring 2008

Find the area of the triangle \triangle with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Problem 6(b) - Spring 2008

Find the area of the triangle \triangle with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$.

Problem 6(b) - Spring 2008

Find the area of the triangle \triangle with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle \triangle with these vertices can be found by taking the area of the parallelogram spanned by \overrightarrow{AB} and \overrightarrow{AC} and dividing by 2.

Problem 6(b) - Spring 2008

Find the area of the triangle Δ with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \overrightarrow{AB} and \overrightarrow{AC} and dividing by 2. Thus:

$$\text{Area}(\Delta) = \frac{|\overrightarrow{AB} \times \overrightarrow{AC}|}{2}$$

Problem 6(b) - Spring 2008

Find the area of the triangle Δ with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \overrightarrow{AB} and \overrightarrow{AC} and dividing by 2. Thus:

$$\text{Area}(\Delta) = \frac{|\overrightarrow{AB} \times \overrightarrow{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{array} \right\|$$

Problem 6(b) - Spring 2008

Find the area of the triangle Δ with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -1, 1, 2 \rangle| \end{aligned}$$

Problem 6(b) - Spring 2008

Find the area of the triangle Δ with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -1, 1, 2 \rangle| = \frac{1}{2} \sqrt{1 + 1 + 4}\end{aligned}$$

Problem 6(b) - Spring 2008

Find the area of the triangle Δ with vertices at the points $A = (1, 2, 3)$, $B = (0, 1, 3)$, and $C = (2, 1, 4)$.

Hint: the area of this triangle is related to the area of a certain parallelogram

Solution:

Consider the points $A = (1, 2, 3)$, $B = (0, 1, 3)$ and $C = (2, 1, 4)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 0 \\ 1 & -1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -1, 1, 2 \rangle| = \frac{1}{2} \sqrt{1 + 1 + 4} = \frac{1}{2} \sqrt{6}.\end{aligned}$$



Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Solution:

- The vector part of the line **L** is the normal vector **n** = $\langle 3, -1, 5 \rangle$ to the plane.

Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Solution:

- The vector part of the line **L** is the normal vector **n** = $\langle 3, -1, 5 \rangle$ to the plane.
- The **vector equation** of **L** is:

$$\mathbf{r}(t) = \langle 2, 4, 1 \rangle + t\mathbf{n}$$

Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Solution:

- The vector part of the line **L** is the normal vector **n** = $\langle 3, -1, 5 \rangle$ to the plane.
- The **vector equation** of **L** is:

$$\begin{aligned}\mathbf{r}(t) &= \langle 2, 4, 1 \rangle + t\mathbf{n} \\ &= \langle 2, 4, 1 \rangle + t\langle 3, -1, 5 \rangle\end{aligned}$$

Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Solution:

- The vector part of the line **L** is the normal vector **n** = $\langle 3, -1, 5 \rangle$ to the plane.
- The **vector equation** of **L** is:

$$\begin{aligned}\mathbf{r}(t) &= \langle 2, 4, 1 \rangle + t\mathbf{n} \\ &= \langle 2, 4, 1 \rangle + t\langle 3, -1, 5 \rangle = \langle 2 + 3t, 4 - t, 1 + 5t \rangle.\end{aligned}$$

Problem 7(a) - Spring 2008

Find the **parametric equations** of the line passing through the point $(2, 4, 1)$ that is perpendicular to the plane $3x - y + 5z = 77$.

Solution:

- The vector part of the line **L** is the normal vector **n** = $\langle 3, -1, 5 \rangle$ to the plane.
- The **vector equation** of **L** is:

$$\begin{aligned}\mathbf{r}(t) &= \langle 2, 4, 1 \rangle + t\mathbf{n} \\ &= \langle 2, 4, 1 \rangle + t\langle 3, -1, 5 \rangle = \langle 2 + 3t, 4 - t, 1 + 5t \rangle.\end{aligned}$$

- The **parametric equations** are:

$$\begin{aligned}x &= 2 + 3t \\ y &= 4 - t \\ z &= 1 + 5t.\end{aligned}$$



Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

- Plug these t -values into equation of plane and solve for t :

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

- Plug these t -values into equation of plane and solve for t :

$$3(2 + 3t) - (4 - t) + 5(1 + 5t) = 77,$$

$$6 + 9t - 4 + t + 5 + 25t = 77,$$

$$35t = 70;$$

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

- Plug these t -values into equation of plane and solve for t :

$$3(2 + 3t) - (4 - t) + 5(1 + 5t) = 77,$$

$$6 + 9t - 4 + t + 5 + 25t = 77,$$

$$35t = 70; \quad \implies t = 2.$$

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

- Plug these t -values into equation of plane and solve for t :

$$3(2 + 3t) - (4 - t) + 5(1 + 5t) = 77,$$

$$6 + 9t - 4 + t + 5 + 25t = 77,$$

$$35t = 70; \quad \implies t = 2.$$

So \mathbf{L} intersects the plane at time $t = 2$.

Problem 7(b) - Spring 2008

Find the intersection point of the line $\mathbf{L}(t) = \langle 2 + 3t, 4 - t, 1 + 5t \rangle$ in part (a) and the plane $3x - y + 5z = 77$.

Solution:

- By part (a), we have \mathbf{L} has **parametric equations**:

$$x = 2 + 3t$$

$$y = 4 - t$$

$$z = 1 + 5t.$$

- Plug these t -values into equation of plane and solve for t :

$$3(2 + 3t) - (4 - t) + 5(1 + 5t) = 77,$$

$$6 + 9t - 4 + t + 5 + 25t = 77,$$

$$35t = 70; \quad \implies t = 2.$$

So \mathbf{L} intersects the plane at time $t = 2$.

- At $t = 2$, the **parametric equations** give the point:

$$\langle 2 + 3 \cdot 2, 4 - 2, 1 + 5 \cdot 2 \rangle = \langle 8, 2, 11 \rangle.$$



Problem 11(a) - Spring 2007

Find **parametric equations** for the line **L** which contains $A(7, 6, 4)$ and $B(4, 6, 5)$.

Problem 11(a) - Spring 2007

Find **parametric equations** for the line **L** which contains $A(7, 6, 4)$ and $B(4, 6, 5)$.

Solution:

- A vector parallel to the line **L** is:

$$\mathbf{v} = \overrightarrow{AB} = \langle 4 - 7, 6 - 6, 5 - 4 \rangle = \langle -3, 0, 1 \rangle.$$

Problem 11(a) - Spring 2007

Find **parametric equations** for the line **L** which contains $A(7, 6, 4)$ and $B(4, 6, 5)$.

Solution:

- A vector parallel to the line **L** is:

$$\mathbf{v} = \overrightarrow{AB} = \langle 4 - 7, 6 - 6, 5 - 4, \rangle = \langle -3, 0, 1 \rangle.$$

- A point on the line is $A(7, 6, 4)$.

Problem 11(a) - Spring 2007

Find **parametric equations** for the line **L** which contains $A(7, 6, 4)$ and $B(4, 6, 5)$.

Solution:

- A vector parallel to the line **L** is:

$$\mathbf{v} = \overrightarrow{AB} = \langle 4 - 7, 6 - 6, 5 - 4, \rangle = \langle -3, 0, 1 \rangle.$$

- A point on the line is $A(7, 6, 4)$.
- Therefore **parametric equations** for the line **L** are:

$$x = 7 - 3t$$

$$y = 6$$

$$z = 4 + t.$$



Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle$$

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix}$$

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\begin{aligned}\mathbf{v} &= \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ &= \begin{vmatrix} -2 & 1 \\ 1 & -1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 1 \end{vmatrix} \mathbf{k}\end{aligned}$$

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\begin{aligned}\mathbf{v} &= \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ &= \begin{vmatrix} -2 & 1 \\ 1 & -1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 1 \end{vmatrix} \mathbf{k} = \langle 1, 3, 5 \rangle.\end{aligned}$$

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\begin{aligned}\mathbf{v} &= \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ &= \begin{vmatrix} -2 & 1 \\ 1 & -1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 1 \end{vmatrix} \mathbf{k} = \langle 1, 3, 5 \rangle.\end{aligned}$$

- A point on **L** is any (x_0, y_0, z_0) that satisfies **both** of the plane equations.

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\begin{aligned}\mathbf{v} &= \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ &= \begin{vmatrix} -2 & 1 \\ 1 & -1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 1 \end{vmatrix} \mathbf{k} = \langle 1, 3, 5 \rangle.\end{aligned}$$

- A point on **L** is any (x_0, y_0, z_0) that satisfies **both** of the plane equations. Setting $z = 0$, we obtain the equations $x - 2y = 5$ and $2x + y = 0$ and find such a point $(1, -2, 0)$.

Problem 11(b) - Spring 2007

Find the **parametric equations** for the **line L of intersection** of the planes $x - 2y + z = 5$ and $2x + y - z = 0$.

Solution:

- A vector **v** parallel to the line is the cross product of the normal vectors of the planes:

$$\begin{aligned}\mathbf{v} &= \langle 1, -2, 1 \rangle \times \langle 2, 1, -1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & -1 \end{vmatrix} \\ &= \begin{vmatrix} -2 & 1 \\ 1 & -1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 1 \end{vmatrix} \mathbf{k} = \langle 1, 3, 5 \rangle.\end{aligned}$$

- A point on **L** is any (x_0, y_0, z_0) that satisfies **both** of the plane equations. Setting $z = 0$, we obtain the equations $x - 2y = 5$ and $2x + y = 0$ and find such a point $(1, -2, 0)$.
- Therefore **parametric equations** for **L** are:

$$x = 1 + t$$

$$y = -2 + 3t$$

$$z = 5t.$$



Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of *any* two vectors that lie **in** the plane.

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of *any* two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of **any** two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

- So the normal vector is

$$\mathbf{n} = \langle 2, -2, -1 \rangle \times \langle 3, 0, -3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & -1 \\ 3 & 0 & -3 \end{vmatrix} =$$

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of **any** two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

- So the normal vector is

$$\mathbf{n} = \langle 2, -2, -1 \rangle \times \langle 3, 0, -3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & -1 \\ 3 & 0 & -3 \end{vmatrix} =$$
$$\begin{vmatrix} -2 & -1 \\ 0 & -3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 2 & -1 \\ 3 & -3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 2 & -2 \\ 3 & 0 \end{vmatrix} \mathbf{k} = \langle 6, 3, 6 \rangle.$$

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of **any** two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

- So the normal vector is

$$\mathbf{n} = \langle 2, -2, -1 \rangle \times \langle 3, 0, -3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & -1 \\ 3 & 0 & -3 \end{vmatrix} =$$
$$\begin{vmatrix} -2 & -1 \\ 0 & -3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 2 & -1 \\ 3 & -3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 2 & -2 \\ 3 & 0 \end{vmatrix} \mathbf{k} = \langle 6, 3, 6 \rangle.$$

- A point on the plane is $P(-1, 0, 2)$.

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of **any** two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

- So the normal vector is

$$\mathbf{n} = \langle 2, -2, -1 \rangle \times \langle 3, 0, -3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & -1 \\ 3 & 0 & -3 \end{vmatrix} =$$
$$\begin{vmatrix} -2 & -1 \\ 0 & -3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 2 & -1 \\ 3 & -3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 2 & -2 \\ 3 & 0 \end{vmatrix} \mathbf{k} = \langle 6, 3, 6 \rangle.$$

- A point on the plane is $P(-1, 0, 2)$. Therefore,

$$6(x - (-1)) + 3(y - 0) + 6(z - 2) = 0,$$

Problem 12(a) - Spring 2007

Find an **equation of the plane** which contains the points $P(-1, 0, 2)$, $Q(1, -2, 1)$ and $R(2, 0, -1)$.

Solution:

- A normal vector to the plane can be found by taking the cross product of **any** two vectors that lie **in** the plane. Two vectors that lie in the plane are $\overrightarrow{PQ} = \langle 2, -2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 0, -3 \rangle$.

- So the normal vector is


$$\mathbf{n} = \langle 2, -2, -1 \rangle \times \langle 3, 0, -3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & -1 \\ 3 & 0 & -3 \end{vmatrix} =$$
$$\begin{vmatrix} -2 & -1 \\ 0 & -3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 2 & -1 \\ 3 & -3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 2 & -2 \\ 3 & 0 \end{vmatrix} \mathbf{k} = \langle 6, 3, 6 \rangle.$$

- A point on the plane is $P(-1, 0, 2)$. Therefore,

$$6(x - (-1)) + 3(y - 0) + 6(z - 2) = 0,$$


or simplified, $6x + 3y + 6z - 6 = 0.$





Problem 12(b) - Spring 2007

Find the distance D from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.




Problem 12(b) - Spring 2007

Find the distance **D** from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane.




Problem 12(b) - Spring 2007

Find the distance **D** from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is **b** = $\langle 1, -1, -1 \rangle$.




Problem 12(b) - Spring 2007

Find the distance D from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is $\mathbf{b} = \langle 1, -1, -1 \rangle$. The distance D from $(1, 0, -1)$ to the plane is equal to:




Problem 12(b) - Spring 2007

Find the distance D from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is $\mathbf{b} = \langle 1, -1, -1 \rangle$. The distance D from $(1, 0, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| =$$




Problem 12(b) - Spring 2007

Find the distance D from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is $\mathbf{b} = \langle 1, -1, -1 \rangle$. The distance D from $(1, 0, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right|$$




Problem 12(b) - Spring 2007

Find the distance **D** from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is **b** = $\langle 1, -1, -1 \rangle$. The distance **D** from $(1, 0, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = |\langle 1, -1, -1 \rangle \cdot \frac{1}{3} \langle 2, 1, -2 \rangle|$$



Problem 12(b) - Spring 2007

Find the distance **D** from the point $P_1 = (1, 0, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 0, -1)$ which is $\mathbf{b} = \langle 1, -1, -1 \rangle$. The distance **D** from $(1, 0, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = |\langle 1, -1, -1 \rangle \cdot \frac{1}{3} \langle 2, 1, -2 \rangle| = 1.$$



Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$;

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.
- The point P in the plane closest to $(1, 0, -1)$ is the intersection of this line and the plane.

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.
- The point P in the plane closest to $(1, 0, -1)$ is the intersection of this line and the plane.
- Substitute the **parametric equations** of the line into the plane equation:
$$2(1 + 2t) + (t) - 2(-1 - 2t) = 1.$$

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.
- The point P in the plane closest to $(1, 0, -1)$ is the intersection of this line and the plane.
- Substitute the **parametric equations** of the line into the plane equation:
$$2(1 + 2t) + (t) - 2(-1 - 2t) = 1.$$

Simplifying and solving for t ,

$$9t + 4 = 1 \implies t = -\frac{1}{3}.$$

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.
- The point P in the plane closest to $(1, 0, -1)$ is the intersection of this line and the plane.
- Substitute the **parametric equations** of the line into the plane equation:
$$2(1 + 2t) + (t) - 2(-1 - 2t) = 1.$$

Simplifying and solving for t ,

$$9t + 4 = 1 \implies t = -\frac{1}{3}.$$

- Plugging this t -value into the **parametric equations**, we get the coordinates of the point of intersection: $x = 1 + 2(-\frac{1}{3}) = \frac{1}{3}$, $y = -\frac{1}{3}$, $z = -1 - 2(-\frac{1}{3}) = -\frac{1}{3}$.

Problem 12(c) - Spring 2007

Find the point P in the plane $2x + y - 2z = 1$ which is closest to the point $(1, 0, -1)$. (Hint: You can use part (b) of this problem to help find P or first find the equation of the line passing through P and the point $(1, 0, -1)$ and then solve for P .)

Solution:

- First find the **parametric equations** of the line that goes through the point $(1, 0, -1)$ that is normal to the plane: $x = 1 + 2t$, $y = t$, $z = -1 - 2t$; here $\mathbf{n} = \langle 2, 1, -2 \rangle$ is a normal to the plane.
- The point P in the plane closest to $(1, 0, -1)$ is the intersection of this line and the plane.
- Substitute the **parametric equations** of the line into the plane equation:
$$2(1 + 2t) + (t) - 2(-1 - 2t) = 1.$$

Simplifying and solving for t ,

$$9t + 4 = 1 \implies t = -\frac{1}{3}.$$

- Plugging this t -value into the **parametric equations**, we get the coordinates of the point of intersection: $x = 1 + 2(-\frac{1}{3}) = \frac{1}{3}$, $y = -\frac{1}{3}$, $z = -1 - 2(-\frac{1}{3}) = -\frac{1}{3}$.
- So the point on the plane closest to $(1, 0, -1)$ is $P = (\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3})$.

Problem 13(b) - Spring 2007

Find the **center** and **radius** of the sphere

$$x^2 + y^2 + 2y + z^2 + 4z = 20.$$

Problem 13(b) - Spring 2007

Find the **center** and **radius** of the sphere

$$x^2 + y^2 + 2y + z^2 + 4z = 20.$$

Solution:

- Completing the square in the y and z variables, we get

$$x^2 + (y^2 + 2y + 1) + (z^2 + 4z + 4) = 20 + 1 + 4.$$

Problem 13(b) - Spring 2007

Find the **center** and **radius** of the sphere

$$x^2 + y^2 + 2y + z^2 + 4z = 20.$$

Solution:

- Completing the square in the y and z variables, we get

$$x^2 + (y^2 + 2y + 1) + (z^2 + 4z + 4) = 20 + 1 + 4.$$

- Rewriting, we have

$$x^2 + (y + 1)^2 + (z + 2)^2 = 25 = 5^2.$$

Problem 13(b) - Spring 2007

Find the **center** and **radius** of the sphere

$$x^2 + y^2 + 2y + z^2 + 4z = 20.$$

Solution:

- Completing the square in the y and z variables, we get

$$x^2 + (y^2 + 2y + 1) + (z^2 + 4z + 4) = 20 + 1 + 4.$$

- Rewriting, we have

$$x^2 + (y + 1)^2 + (z + 2)^2 = 25 = 5^2.$$

- Hence, the **center** is **C** = $(0, -1, -2)$ and the **radius** is $r = 5$.



Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- The area of the parallelogram is

$$|\vec{AB} \times \vec{AC}| = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix}$$

Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- The area of the parallelogram is

$$\begin{aligned} |\vec{AB} \times \vec{AC}| &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix} \\ &= \left| \begin{vmatrix} -1 & 2 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} -1 & -1 \\ -2 & 1 \end{vmatrix} \mathbf{k} \right| \end{aligned}$$

Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- The area of the parallelogram is

$$\begin{aligned} |\vec{AB} \times \vec{AC}| &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix} \\ &= \left| \begin{vmatrix} -1 & 2 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} -1 & -1 \\ -2 & 1 \end{vmatrix} \mathbf{k} \right| \\ &= |\langle -3, -3, -3 \rangle| \end{aligned}$$

Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- The area of the parallelogram is

$$\begin{aligned} |\vec{AB} \times \vec{AC}| &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix} \\ &= \left| \begin{vmatrix} -1 & 2 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} -1 & -1 \\ -2 & 1 \end{vmatrix} \mathbf{k} \right| \\ &= |\langle -3, -3, -3 \rangle| = \sqrt{27}. \end{aligned}$$

Problem 15(a) - Spring 2008

Consider the points $A(2, 1, 0)$, $B(1, 0, 2)$ and $C(0, 2, 1)$. Find the area **A** of the triangle ABC . (Hint: If you know how to find the area of a parallelogram spanned by 2 vectors, then you should be able to solve this problem.)

Solution:

- The area of the parallelogram is

$$\begin{aligned} |\vec{AB} \times \vec{AC}| &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix} \\ &= \left| \begin{vmatrix} -1 & 2 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} -1 & -1 \\ -2 & 1 \end{vmatrix} \mathbf{k} \right| \\ &= |\langle -3, -3, -3 \rangle| = \sqrt{27}. \end{aligned}$$

- So the area of the triangle ABC is

$$\mathbf{A} = \frac{\sqrt{27}}{2}.$$



Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal **n** and a point on it.

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal **n** and a point on it. Setting $t = 0$, we find the point $(4, 3, 1)$ on the first line.

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal \mathbf{n} and a point on it. Setting $t = 0$, we find the point $(4, 3, 1)$ on the first line.
- The part vector \mathbf{v}_1 of the first line is $\langle -4, -1, 5 \rangle$ and the vector part \mathbf{v}_2 of the second line is $\langle -1, 2, 0 \rangle$.

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal \mathbf{n} and a point on it. Setting $t = 0$, we find the point $(4, 3, 1)$ on the first line.
- The part vector \mathbf{v}_1 of the first line is $\langle -4, -1, 5 \rangle$ and the vector part \mathbf{v}_2 of the second line is $\langle -1, 2, 0 \rangle$.
- Since the vector

$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & -1 & 5 \\ -1 & 2 & 0 \end{vmatrix} = \langle -10, -5, -9 \rangle,$$

is orthogonal to both \mathbf{v}_1 and \mathbf{v}_2 , it is the normal to the plane.

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal \mathbf{n} and a point on it. Setting $t = 0$, we find the point $(4, 3, 1)$ on the first line.
- The part vector \mathbf{v}_1 of the first line is $\langle -4, -1, 5 \rangle$ and the vector part \mathbf{v}_2 of the second line is $\langle -1, 2, 0 \rangle$.
- Since the vector

$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & -1 & 5 \\ -1 & 2 & 0 \end{vmatrix} = \langle -10, -5, -9 \rangle,$$

is orthogonal to both \mathbf{v}_1 and \mathbf{v}_2 , it is the normal to the plane.

- The **equation of the plane** is:

$$\langle -10, -5, -9 \rangle \cdot \langle x - 4, y - 3, z - 1 \rangle$$

Problem 16 - Fall 2007

Find the **equation of the plane** containing the lines

$$x = 4 - 4t, \quad y = 3 - t, \quad z = 1 + 5t \quad \text{and}$$

$$x = 4 - t, \quad y = 3 + 2t, \quad z = 1$$

Solution:

- To find the equation of a plane, we need to find its normal \mathbf{n} and a point on it. Setting $t = 0$, we find the point $(4, 3, 1)$ on the first line.
- The part vector \mathbf{v}_1 of the first line is $\langle -4, -1, 5 \rangle$ and the vector part \mathbf{v}_2 of the second line is $\langle -1, 2, 0 \rangle$.
- Since the vector


$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & -1 & 5 \\ -1 & 2 & 0 \end{vmatrix} = \langle -10, -5, -9 \rangle,$$

is orthogonal to both \mathbf{v}_1 and \mathbf{v}_2 , it is the normal to the plane.

- The **equation of the plane** is:


$$\begin{aligned} &\langle -10, -5, -9 \rangle \cdot \langle x - 4, y - 3, z - 1 \rangle \\ &= -10(x - 4) - 5(y - 3) - 9(z - 1) = 0. \end{aligned}$$





Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.




Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.




Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $4x - 6y - z - 5 = 0$.




Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.

Solution:

- Recall the distance formula $D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $4x - 6y - z - 5 = 0$.
- So, the distance from $(3, -2, 7)$ to the plane is:




Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $4x - 6y - z - 5 = 0$.
- So, the distance from $(3, -2, 7)$ to the plane is:

$$\mathbf{D} = \frac{|(4 \cdot 3) + (-6 \cdot -2) + (-1 \cdot 7) - 5|}{\sqrt{4^2 + (-6)^2 + (-1)^2}}$$



Problem 17 - Fall 2007

Find the distance **D** from the point $P_1 = (3, -2, 7)$ and the plane $4x - 6y - z = 5$.

Solution:

- Recall the distance formula $\mathbf{D} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$ from a point $P = (x_1, y_1, z_1)$ to a plane $ax + by + cz + d = 0$.
- In order to apply the formula, rewrite the equation of the plane in standard form: $4x - 6y - z - 5 = 0$.
- So, the distance from $(3, -2, 7)$ to the plane is:

$$\mathbf{D} = \frac{|(4 \cdot 3) + (-6 \cdot -2) + (-1 \cdot 7) - 5|}{\sqrt{4^2 + (-6)^2 + (-1)^2}} = \frac{12}{\sqrt{53}}.$$

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

- Equating x and y-coordinates:

$$\begin{aligned}x &= t = -4s + 3 \\ y &= 2t + 1 = -3s + 2.\end{aligned}$$

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

- Equating x and y -coordinates:

$$\begin{aligned}x &= t = -4s + 3 \\ y &= 2t + 1 = -3s + 2.\end{aligned}$$

- Solving gives $s = 1$ and $t = -1$.

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

- Equating x and y -coordinates:

$$\begin{aligned}x &= t = -4s + 3 \\ y &= 2t + 1 = -3s + 2.\end{aligned}$$

- Solving gives $s = 1$ and $t = -1$.
- $L_1(-1) = \langle -1, -1, -1 \rangle \neq \langle -1, -1, 3 \rangle = L_2(1)$.

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

- Equating x and y-coordinates:

$$\begin{aligned}x &= t = -4s + 3 \\ y &= 2t + 1 = -3s + 2.\end{aligned}$$

- Solving gives $s = 1$ and $t = -1$.
- $L_1(-1) = \langle -1, -1, -1 \rangle \neq \langle -1, -1, 3 \rangle = L_2(1)$. So these lines do **not intersect**.

Problem 18 - Fall 2007

Determine whether the lines L_1 and L_2 given below are **parallel**, **skew** or **intersecting**. If they intersect, find the point of intersection.

$$L_1 : \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

$$L_2 : \frac{x-3}{-4} = \frac{y-2}{-3} = \frac{z-1}{2}$$

Solution:

- Rewrite these lines as vector equations:

$$L_1(t) = \langle t, 2t + 1, 3t + 2 \rangle$$

$$L_2(s) = \langle -4s + 3, -3s + 2, 2s + 1 \rangle$$

- Equating x and y -coordinates:

$$\begin{aligned}x &= t = -4s + 3 \\ y &= 2t + 1 = -3s + 2.\end{aligned}$$

- Solving gives $s = 1$ and $t = -1$.
- $L_1(-1) = \langle -1, -1, -1 \rangle \neq \langle -1, -1, 3 \rangle = L_2(1)$. So these lines do **not intersect**.
- Since the lines are clearly **not parallel** (the direction vectors $\langle 1, 2, 3 \rangle$ and $\langle -4, -3, 2 \rangle$ are **not parallel**), the lines are **skew**.



Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$.

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$. Then the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} is:

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$. Then the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} is:

$$\text{Area}(\Delta) = |\overrightarrow{PQ} \times \overrightarrow{PR}|$$

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$. Then the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} is:

$$\text{Area}(\Delta) = |\overrightarrow{PQ} \times \overrightarrow{PR}| = \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & -1 \\ 3 & 2 & 0 \end{array} \right\|$$

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$. Then the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} is:

$$\begin{aligned}\text{Area}(\Delta) &= |\overrightarrow{PQ} \times \overrightarrow{PR}| = \left| \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & -1 \\ 3 & 2 & 0 \end{vmatrix} \right| \\ &= |\langle 2, -3, -6 \rangle|\end{aligned}$$

Problem 20(a) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the area of the parallelogram.

Solution:

Consider the vectors $\overrightarrow{PQ} = \langle 0, 2, -1 \rangle$ and $\overrightarrow{PR} = \langle 3, 2, 0 \rangle$. Then the area of the parallelogram spanned by \overrightarrow{PQ} and \overrightarrow{PR} is:

$$\begin{aligned}\text{Area}(\Delta) &= |\overrightarrow{PQ} \times \overrightarrow{PR}| = \left\| \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & -1 \\ 3 & 2 & 0 \end{vmatrix} \right\| \\ &= |\langle 2, -3, -6 \rangle| = \sqrt{4 + 9 + 36} = 7\end{aligned}$$



Problem 20(b) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the cosine of the angle between the vector PQ and PR .

Problem 20(b) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the cosine of the angle between the vector PQ and PR .

Solution:

- Note that:

$$\overrightarrow{PQ} = \langle 0, 2, -1 \rangle \quad \overrightarrow{PR} = \langle 3, 2, 0 \rangle.$$

Problem 20(b) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the cosine of the angle between the vector PQ and PR .

Solution:

- Note that:

$$\overrightarrow{PQ} = \langle 0, 2, -1 \rangle \quad \overrightarrow{PR} = \langle 3, 2, 0 \rangle.$$

- By our formula for dot products:

$$\cos \theta = \frac{\overrightarrow{PQ} \cdot \overrightarrow{PR}}{|\overrightarrow{PQ}| |\overrightarrow{PR}|}$$

Problem 20(b) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the cosine of the angle between the vector PQ and PR .

Solution:

- Note that:

$$\overrightarrow{PQ} = \langle 0, 2, -1 \rangle \quad \overrightarrow{PR} = \langle 3, 2, 0 \rangle.$$

- By our formula for dot products:

$$\cos \theta = \frac{\overrightarrow{PQ} \cdot \overrightarrow{PR}}{|\overrightarrow{PQ}| |\overrightarrow{PR}|} = \frac{\langle 0, 2, -1 \rangle \cdot \langle 3, 2, 0 \rangle}{\sqrt{5} \sqrt{13}}$$

Problem 20(b) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the cosine of the angle between the vector PQ and PR .

Solution:

- Note that:

$$\overrightarrow{PQ} = \langle 0, 2, -1 \rangle \quad \overrightarrow{PR} = \langle 3, 2, 0 \rangle.$$

- By our formula for dot products:

$$\cos \theta = \frac{\overrightarrow{PQ} \cdot \overrightarrow{PR}}{|\overrightarrow{PQ}| |\overrightarrow{PR}|} = \frac{\langle 0, 2, -1 \rangle \cdot \langle 3, 2, 0 \rangle}{\sqrt{5} \sqrt{13}} = \frac{4}{\sqrt{5} \sqrt{13}}.$$



Problem 20(c) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Problem 20(c) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by **S**.

Problem 20(c) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by S . Then

$$\overrightarrow{OS} = \overrightarrow{OQ} + \overrightarrow{PR} = \langle 0, 1, 0 \rangle + \langle 3, 2, 0 \rangle = \langle 3, 3, 0 \rangle,$$

where O is the origin.

Problem 20(c) - Fall 2007

Three of the four vertices of a parallelogram are $P(0, -1, 1)$, $Q(0, 1, 0)$ and $R(3, 1, 1)$. Two of the sides are PQ and PR . Find the coordinates of the fourth vertex.

Solution:

Denote the fourth vertex by S . Then

$$\overrightarrow{OS} = \overrightarrow{OQ} + \overrightarrow{PR} = \langle 0, 1, 0 \rangle + \langle 3, 2, 0 \rangle = \langle 3, 3, 0 \rangle,$$

where O is the origin. That is,

$$S = (3, 3, 0).$$



Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.
- When $y = 0$, then $0 = 2t - 1$ or $t = \frac{1}{2}$.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.
- When $y = 0$, then $0 = 2t - 1$ or $t = \frac{1}{2}$.
- Hence,

$$\left\langle 2 - \left(\frac{1}{2}\right)^2, 2 \cdot \frac{1}{2} - 1, \ln \frac{1}{2} \right\rangle = \left\langle 1\frac{3}{4}, 0, -\ln 2 \right\rangle$$

is the unique point of the **intersection** of **C** with xz -plane.



Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.
- When $y = 0$, then $0 = 2t - 1$ or $t = \frac{1}{2}$.

Problem 21(a) - Fall 2007

Let **C** be the parametric curve

$$x = 2 - t^2, \quad y = 2t - 1, \quad z = \ln t.$$

Determine the point(s) of **intersection** of **C** with the xz -plane.

Solution:

- The points of intersection of **C** with the xz -plane correspond to the points where the y -coordinate of **C** is 0.
- When $y = 0$, then $0 = 2t - 1$ or $t = \frac{1}{2}$.
- Hence,

$$\left\langle 2 - \left(\frac{1}{2}\right)^2, 2 \cdot \frac{1}{2} - 1, \ln \frac{1}{2} \right\rangle = \left\langle 1\frac{3}{4}, 0, -\ln 2 \right\rangle$$

is the unique point of the **intersection** of **C** with xz -plane.



Problem 22(a) - Fall 2006

Find **parametric equations** for the line r which contains $A(2, 0, 1)$ and $B(-1, 1, -1)$.

Problem 22(a) - Fall 2006

Find **parametric equations** for the line **r** which contains $A(2, 0, 1)$ and $B(-1, 1, -1)$.

Solution:

- Note that $\overrightarrow{AB} = \langle -3, 1, -2 \rangle$ and the **vector equation** is:

$$\mathbf{r}(t) = \vec{A} + t\overrightarrow{AB}$$

Problem 22(a) - Fall 2006

Find **parametric equations** for the line **r** which contains $A(2, 0, 1)$ and $B(-1, 1, -1)$.

Solution:

- Note that $\overrightarrow{AB} = \langle -3, 1, -2 \rangle$ and the **vector equation** is:

$$\mathbf{r}(t) = \vec{A} + t\overrightarrow{AB} = \langle 2, 0, 1 \rangle + t\langle -3, 1, -2 \rangle$$

Problem 22(a) - Fall 2006

Find **parametric equations** for the line **r** which contains $A(2, 0, 1)$ and $B(-1, 1, -1)$.

Solution:

- Note that $\overrightarrow{AB} = \langle -3, 1, -2 \rangle$ and the **vector equation** is:

$$\mathbf{r}(t) = \vec{A} + t\overrightarrow{AB} = \langle 2, 0, 1 \rangle + t\langle -3, 1, -2 \rangle = \langle 2 - 3t, t, 1 - 2t \rangle.$$

Problem 22(a) - Fall 2006

Find **parametric equations** for the line **r** which contains $A(2, 0, 1)$ and $B(-1, 1, -1)$.

Solution:

- Note that $\overrightarrow{AB} = \langle -3, 1, -2 \rangle$ and the **vector equation** is:

$$\mathbf{r}(t) = \vec{A} + t\overrightarrow{AB} = \langle 2, 0, 1 \rangle + t\langle -3, 1, -2 \rangle = \langle 2 - 3t, t, 1 - 2t \rangle.$$

- The **parametric equations** are:

$$x = 2 - 3t$$

$$y = t$$

$$z = 1 - 2t.$$



Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$.

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s$$

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

Problem 22(b) - Fall 2006

Determine whether the lines $\mathbf{L}_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $\mathbf{L}_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line \mathbf{L}_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line \mathbf{L}_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

$$y = 3t = 4 + s$$

Problem 22(b) - Fall 2006

Determine whether the lines $\mathbf{L}_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $\mathbf{L}_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line \mathbf{L}_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line \mathbf{L}_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

$$y = 3t = 4 + s \implies 3t = 4 + s.$$

Problem 22(b) - Fall 2006

Determine whether the lines $\mathbf{L}_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $\mathbf{L}_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line \mathbf{L}_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line \mathbf{L}_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

$$y = 3t = 4 + s \implies 3t = 4 + s.$$

Solving yields:

$$t = 6 \text{ and } s = 14.$$

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.
- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

$$y = 3t = 4 + s \implies 3t = 4 + s.$$

Solving yields:

$$t = 6 \text{ and } s = 14.$$

Plugging these values into $z = 2 - t = 1 + 3s$ yields the inequality $-4 \neq 43$, which means the z -coordinates are never equal and the lines do **not intersect**.

Problem 22(b) - Fall 2006

Determine whether the lines $L_1 : x = 1 + 2t, y = 3t, z = 2 - t$ and $L_2 : x = -1 + s, y = 4 + s, z = 1 + 3s$ are **parallel**, **skew** or **intersecting**.

Solution:

- Vector part of line L_1 is $\mathbf{v}_1 = \langle 2, 3, -1 \rangle$ and for line L_2 is $\mathbf{v}_2 = \langle 1, 1, 3 \rangle$. Clearly, \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 and so these lines are **not parallel**.

- If these lines intersect, then for some values of t and s :

$$x = 1 + 2t = -1 + s \implies 2t = -2 + s,$$

$$y = 3t = 4 + s \implies 3t = 4 + s.$$

Solving yields:

$$t = 6 \text{ and } s = 14.$$

Plugging these values into $z = 2 - t = 1 + 3s$ yields the inequality $-4 \neq 43$, which means the z -coordinates are never equal and the lines do **not intersect**.

- Thus, the lines are **skew**.



Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.
- The normal vector to the plane is:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR}$$

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.
- The normal vector to the plane is:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 0 \\ 2 & -1 & -2 \end{vmatrix}$$

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.
- The normal vector to the plane is:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 0 \\ 2 & -1 & -2 \end{vmatrix} = 8\mathbf{i} + 4\mathbf{j} + 6\mathbf{k}.$$

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:

- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.
- The normal vector to the plane is:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 0 \\ 2 & -1 & -2 \end{vmatrix} = 8\mathbf{i} + 4\mathbf{j} + 6\mathbf{k}.$$

- Since $P(-1, 2, 1)$ lies on the plane, the **equation of the plane** is:

Problem 23(a) - Fall 2006

Find an **equation of the plane** which contains the points $P(-1, 2, 1)$, $Q(1, -2, 1)$ and $R(1, 1, -1)$.

Solution:


- Consider the vectors $\overrightarrow{PQ} = \langle 2, -4, 0 \rangle$ and $\overrightarrow{PR} = \langle 2, -1, -2 \rangle$ which are parallel to the plane.
- The normal vector to the plane is:

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 0 \\ 2 & -1 & -2 \end{vmatrix} = 8\mathbf{i} + 4\mathbf{j} + 6\mathbf{k}.$$

- Since $P(-1, 2, 1)$ lies on the plane, the **equation of the plane** is:


$$\langle 8, 4, 6 \rangle \cdot \langle x+1, y-2, z-1 \rangle = 8(x+1) + 4(y-2) + 6(z-1) = 0.$$





Problem 23(b) - Fall 2006

Find the distance **D** from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.




Problem 23(b) - Fall 2006

Find the distance **D** from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane.




Problem 23(b) - Fall 2006

Find the distance **D** from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is **b** = $\langle 1, 1, -1 \rangle$.




Problem 23(b) - Fall 2006

Find the distance D from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is $\mathbf{b} = \langle 1, 1, -1 \rangle$. The distance D from $(1, 2, -1)$ to the plane is equal to:




Problem 23(b) - Fall 2006

Find the distance D from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is $\mathbf{b} = \langle 1, 1, -1 \rangle$. The distance D from $(1, 2, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| =$$




Problem 23(b) - Fall 2006

Find the distance D from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is $\mathbf{n} = \langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is $\mathbf{b} = \langle 1, 1, -1 \rangle$. The distance D from $(1, 2, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right|$$




Problem 23(b) - Fall 2006

Find the distance **D** from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is **b** = $\langle 1, 1, -1 \rangle$. The distance **D** from $(1, 2, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = |\langle 1, 1, -1 \rangle \cdot \frac{1}{3} \langle 2, 1, -2 \rangle|$$



Problem 23(b) - Fall 2006

Find the distance **D** from the point $(1, 2, -1)$ to the plane $2x + y - 2z = 1$.

Solution:

The normal to the plane is **n** = $\langle 2, 1, -2 \rangle$ and the point $P_0 = (0, 1, 0)$ lies on this plane. Consider the vector from P_0 to $P_1 = (1, 2, -1)$ which is **b** = $\langle 1, 1, -1 \rangle$. The distance **D** from $(1, 2, -1)$ to the plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = |\langle 1, 1, -1 \rangle \cdot \frac{1}{3} \langle 2, 1, -2 \rangle| = \frac{5}{3}.$$



Problem 26(b) - Fall 2006

Find the **center** and **radius** of the sphere $x^2 + y^2 + z^2 + 6z = 16$.

Problem 26(b) - Fall 2006

Find the **center** and **radius** of the sphere $x^2 + y^2 + z^2 + 6z = 16$.

Solution:

- Complete the square in order to put the equation in the form:

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2.$$

Problem 26(b) - Fall 2006

Find the **center** and **radius** of the sphere $x^2 + y^2 + z^2 + 6z = 16$.

Solution:

- Complete the square in order to put the equation in the form:

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2.$$

- We get:

$$x^2 + y^2 + (z^2 + 6z) = x^2 + y^2 + (z^2 + 6z + 9) - 9 = 16.$$

Problem 26(b) - Fall 2006

Find the **center** and **radius** of the sphere $x^2 + y^2 + z^2 + 6z = 16$.

Solution:

- Complete the square in order to put the equation in the form:

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2.$$

- We get:

$$x^2 + y^2 + (z^2 + 6z) = x^2 + y^2 + (z^2 + 6z + 9) - 9 = 16.$$

- This gives the equation

$$(x - 0)^2 + (y - 0)^2 + (z + 3)^2 = 25 = 5^2.$$

Problem 26(b) - Fall 2006

Find the **center** and **radius** of the sphere $x^2 + y^2 + z^2 + 6z = 16$.

Solution:

- Complete the square in order to put the equation in the form:

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2.$$

- We get:

$$x^2 + y^2 + (z^2 + 6z) = x^2 + y^2 + (z^2 + 6z + 9) - 9 = 16.$$

- This gives the equation

$$(x - 0)^2 + (y - 0)^2 + (z + 3)^2 = 25 = 5^2.$$

Hence, the **center** is **C** = (0, 0, -3) and the **radius** is $r = 5$.



Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB}$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:
$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$
- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

$$2(2 + 3t) - 3(1 + 2t) + 4(-1 - t) = -4t - 3 = 13$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

$$2(2 + 3t) - 3(1 + 2t) + 4(-1 - t) = -4t - 3 = 13$$

$$\implies -4t = 16$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

$$2(2 + 3t) - 3(1 + 2t) + 4(-1 - t) = -4t - 3 = 13$$

$$\implies -4t = 16 \implies t = -4.$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

$$2(2 + 3t) - 3(1 + 2t) + 4(-1 - t) = -4t - 3 = 13$$

$$\implies -4t = 16 \implies t = -4.$$

- So, the **point of intersection** is:

$$\mathbf{L}(-4) = \langle 2 - 12, 1 - 8, -1 - (-4) \rangle$$

Problem 27

Consider the line **L** through points $A = (2, 1, -1)$ and $B = (5, 3, -2)$. Find the **intersection** of the line **L** and the plane given by $2x - 3y + 4z = 13$.

Solution:

- The vector part of **L** is $\overrightarrow{AB} = \langle 3, 2, -1 \rangle$ and the point A is on the line.
- The **vector equation** of **L** is:

$$\mathbf{L} = \vec{A} + t\overrightarrow{AB} = \langle 2, 1, -1 \rangle + t\langle 3, 2, -1 \rangle = \langle 2+3t, 1+2t, -1-t \rangle.$$

- Plugging $x = 2 + 3t$, $y = 1 + 2t$ and $z = -1 - t$ into the **equation of the plane** gives:

$$2(2 + 3t) - 3(1 + 2t) + 4(-1 - t) = -4t - 3 = 13$$

$$\implies -4t = 16 \implies t = -4.$$

- So, the **point of intersection** is:

$$\mathbf{L}(-4) = \langle 2 - 12, 1 - 8, -1 - (-4) \rangle = \langle -10, -7, 3 \rangle.$$



Problem 29

Consider the parallelogram with vertices A, B, C, D such that B and C are adjacent to A . If $A = (2, 5, 1)$, $B = (3, 1, 4)$, $D = (5, 2, -3)$, find the point C .

Problem 29

Consider the parallelogram with vertices A, B, C, D such that B and C are adjacent to A . If $A = (2, 5, 1)$, $B = (3, 1, 4)$, $D = (5, 2, -3)$, find the point C .

Solution:

After drawing a picture, the point C is easily seen to be:

$$\overrightarrow{OA} + \overrightarrow{BD}$$

Problem 29

Consider the parallelogram with vertices A, B, C, D such that B and C are adjacent to A . If $A = (2, 5, 1)$, $B = (3, 1, 4)$, $D = (5, 2, -3)$, find the point C .

Solution:

After drawing a picture, the point C is easily seen to be:

$$\overrightarrow{OA} + \overrightarrow{BD} = \langle 2, 5, 1 \rangle + \langle 2, 1, -7 \rangle$$

Problem 29

Consider the parallelogram with vertices A, B, C, D such that B and C are adjacent to A . If $A = (2, 5, 1)$, $B = (3, 1, 4)$, $D = (5, 2, -3)$, find the point C .

Solution:

After drawing a picture, the point C is easily seen to be:

$$\overrightarrow{OA} + \overrightarrow{BD} = \langle 2, 5, 1 \rangle + \langle 2, 1, -7 \rangle = \langle 4, 6, -6 \rangle,$$

where O is the origin.



Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Solution:

- We just plug in the vectors $\vec{a} = \vec{AB} = \langle -1, -1, 2 \rangle$ and $\vec{b} = \vec{AC} = \langle -2, 1, 1 \rangle$ into the formula:

Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Solution:

- We just plug in the vectors $\mathbf{a} = \vec{AB} = \langle -1, -1, 2 \rangle$ and $\mathbf{b} = \vec{AC} = \langle -2, 1, 1 \rangle$ into the formula:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}} \mathbf{a}.$$

Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Solution:

- We just plug in the vectors $\mathbf{a} = \vec{AB} = \langle -1, -1, 2 \rangle$ and $\mathbf{b} = \vec{AC} = \langle -2, 1, 1 \rangle$ into the formula:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}} \mathbf{a}.$$

- Plugging in, we get:

$$\text{proj}_{\vec{AB}}(\vec{AC})$$

Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Solution:

- We just plug in the vectors $\mathbf{a} = \vec{AB} = \langle -1, -1, 2 \rangle$ and $\mathbf{b} = \vec{AC} = \langle -2, 1, 1 \rangle$ into the formula:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}} \mathbf{a}.$$

- Plugging in, we get:

$$\text{proj}_{\vec{AB}}(\vec{AC}) = \frac{\langle -1, -1, 2 \rangle \cdot \langle -2, 1, 1 \rangle}{\langle -1, -1, 2 \rangle \cdot \langle -1, -1, 2 \rangle} \langle -1, -1, 2 \rangle$$

Problem 30(a)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Find the **orthogonal projection** $\text{proj}_{\vec{AB}}(\vec{AC})$ of the vector \vec{AC} onto the vector \vec{AB} .

Solution:

- We just plug in the vectors $\mathbf{a} = \vec{AB} = \langle -1, -1, 2 \rangle$ and $\mathbf{b} = \vec{AC} = \langle -2, 1, 1 \rangle$ into the formula:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}} \mathbf{a}.$$

- Plugging in, we get:

$$\text{proj}_{\vec{AB}}(\vec{AC}) = \frac{\langle -1, -1, 2 \rangle \cdot \langle -2, 1, 1 \rangle}{\langle -1, -1, 2 \rangle \cdot \langle -1, -1, 2 \rangle} \langle -1, -1, 2 \rangle = \frac{1}{2} \langle -1, -1, 2 \rangle.$$



Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.
Find the area of triangle ABC .

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.
Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$.

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle \triangle with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2.

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\text{Area}(\Delta) = \frac{|\vec{AB} \times \vec{AC}|}{2}$$

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\text{Area}(\Delta) = \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\|$$

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -3, -3, -3 \rangle|\end{aligned}$$

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -3, -3, -3 \rangle| = \frac{1}{2} \sqrt{9 + 9 + 9}\end{aligned}$$

Problem 30(b)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the area of triangle ABC .

Solution:

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Then the area of the triangle Δ with these vertices can be found by taking the area of the parallelogram spanned by \vec{AB} and \vec{AC} and dividing by 2. Thus:

$$\begin{aligned}\text{Area}(\Delta) &= \frac{|\vec{AB} \times \vec{AC}|}{2} = \frac{1}{2} \left\| \begin{array}{ccc} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & -1 & 2 \\ -2 & 1 & 1 \end{array} \right\| \\ &= \frac{1}{2} |\langle -3, -3, -3 \rangle| = \frac{1}{2} \sqrt{9 + 9 + 9} = \frac{1}{2} \sqrt{27}.\end{aligned}$$



Problem 30(c)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the distance d from the point C to the line L that contains points A and B .

Problem 30(c)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the distance **d** from the point C to the line **L** that contains points A and B .

Solution:

- From the figure drawn on the blackboard, we see that the distance **d** from C to **L** is the absolute value of the scalar projection of \overrightarrow{AC} in the direction

$$\mathbf{v} = \overrightarrow{AC} - \text{proj}_{\overrightarrow{AB}} \overrightarrow{AC}.$$

Problem 30(c)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the distance d from the point C to the line L that contains points A and B .

Solution:

- From the figure drawn on the blackboard, we see that the distance d from C to L is the absolute value of the scalar projection of \overrightarrow{AC} in the direction

$$\mathbf{v} = \overrightarrow{AC} - \text{proj}_{\overrightarrow{AB}} \overrightarrow{AC}.$$

- The vector \mathbf{v} lies in the plane containing A, B, C and is perpendicular to \overrightarrow{AB} .

Problem 30(c)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the distance d from the point C to the line L that contains points A and B .

Solution:

- From the figure drawn on the blackboard, we see that the distance d from C to L is the absolute value of the scalar projection of \overrightarrow{AC} in the direction

$$\mathbf{v} = \overrightarrow{AC} - \text{proj}_{\overrightarrow{AB}} \overrightarrow{AC}.$$

- The vector \mathbf{v} lies in the plane containing A, B, C and is perpendicular to \overrightarrow{AB} .
- Hence,

$$d = |\text{comp}_{\mathbf{v}} \overrightarrow{AC}|.$$

Problem 30(c)

Consider the points $A = (2, 1, 0)$, $B = (1, 0, 2)$ and $C = (0, 2, 1)$. Find the distance d from the point C to the line L that contains points A and B .

Solution:

- From the figure drawn on the blackboard, we see that the distance d from C to L is the absolute value of the scalar projection of \overrightarrow{AC} in the direction

$$\mathbf{v} = \overrightarrow{AC} - \text{proj}_{\overrightarrow{AB}} \overrightarrow{AC}.$$

- The vector \mathbf{v} lies in the plane containing A, B, C and is perpendicular to \overrightarrow{AB} .
- Hence,

$$d = |\text{comp}_{\mathbf{v}} \overrightarrow{AC}|.$$

- Next, you the student, do the algebraic calculation of d .



Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$.

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle$$

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix}$$

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero.

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero. Setting $z = 0$, we get:

$$\begin{aligned} x - 2y &= 1 \\ 2x + y &= 1. \end{aligned}$$

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero. Setting $z = 0$, we get:

$$\begin{aligned} x - 2y &= 1 \\ 2x + y &= 1. \end{aligned}$$

Solving, we find that $x = \frac{3}{5}$ and $y = -\frac{1}{5}$.

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero. Setting $z = 0$, we get:

$$\begin{aligned} x - 2y &= 1 \\ 2x + y &= 1. \end{aligned}$$

Solving, we find that $x = \frac{3}{5}$ and $y = -\frac{1}{5}$. Hence, **P** = $\langle \frac{3}{5}, -\frac{1}{5}, 0 \rangle$ lies on the line **L**.

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero. Setting $z = 0$, we get:

$$\begin{aligned} x - 2y &= 1 \\ 2x + y &= 1. \end{aligned}$$

Solving, we find that $x = \frac{3}{5}$ and $y = -\frac{1}{5}$. Hence, **P** = $\langle \frac{3}{5}, -\frac{1}{5}, 0 \rangle$ lies on the line **L**.

- The **parametric equations** are:

Problem 31

Find **parametric equations** for the line **L** of intersection of the planes $x - 2y + z = 1$ and $2x + y + z = 1$.

Solution:

- The vector part **v** of the line **L** of intersection is orthogonal to the normal vectors $\langle 1, -2, 1 \rangle$ and $\langle 2, 1, 1 \rangle$. Hence **v** can be taken to be:

$$\mathbf{v} = \langle 1, -2, 1 \rangle \times \langle 2, 1, 1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 1 & 1 \end{vmatrix} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}.$$

- Choose **P** \in **L** so the z-coordinate of **P** is zero. Setting $z = 0$, we get:

$$\begin{aligned} x - 2y &= 1 \\ 2x + y &= 1. \end{aligned}$$

Solving, we find that $x = \frac{3}{5}$ and $y = -\frac{1}{5}$. Hence, **P** = $\langle \frac{3}{5}, -\frac{1}{5}, 0 \rangle$ lies on the line **L**.

- The **parametric equations** are:

$$\begin{aligned} x &= \frac{3}{5} - 3t \\ y &= -\frac{1}{5} + t \\ z &= 5t. \end{aligned}$$



Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t \langle -2, 4, 0 \rangle$$

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

$$L_2(s) = \langle 2, 3, -1 \rangle + s\langle 2, 1, -2 \rangle = \langle 2 + 2s, 3 + s, -1 - 2s \rangle$$

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

$$L_2(s) = \langle 2, 3, -1 \rangle + s\langle 2, 1, -2 \rangle = \langle 2 + 2s, 3 + s, -1 - 2s \rangle$$

- Equating z -coordinates, we find $1 = -1 - 2s \implies s = -1$.

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 intersect? If not, are they skew or parallel?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

$$L_2(s) = \langle 2, 3, -1 \rangle + s\langle 2, 1, -2 \rangle = \langle 2 + 2s, 3 + s, -1 - 2s \rangle$$

- Equating z -coordinates, we find $1 = -1 - 2s \implies s = -1$.
- Equating y -coordinates with $s = -1$, we find $4t = 3 - 1 \implies t = \frac{1}{2}$.

Problem 32

Let \mathbf{L}_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let \mathbf{L}_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines \mathbf{L}_1 and \mathbf{L}_2 **intersect**? If not, are they **skew** or **parallel**?

Solution:

- The vector equations of the lines are:

$$\mathbf{L}_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

$$\mathbf{L}_2(s) = \langle 2, 3, -1 \rangle + s\langle 2, 1, -2 \rangle = \langle 2 + 2s, 3 + s, -1 - 2s \rangle$$

- Equating z -coordinates, we find $1 = -1 - 2s \implies s = -1$.
- Equating y -coordinates with $s = -1$, we find $4t = 3 - 1 \implies t = \frac{1}{2}$.
- Equating x -coordinates with $s = -1$ and $t = \frac{1}{2}$, we find:

$$\mathbf{L}_1\left(\frac{1}{2}\right) = \langle 0, 2, 1 \rangle = \mathbf{L}_2(-1).$$

Problem 32

Let L_1 denote the line through the points $(1, 0, 1)$ and $(-1, 4, 1)$ and let L_2 denote the line through the points $(2, 3, -1)$ and $(4, 4, -3)$. Do the lines L_1 and L_2 **intersect**? If not, are they **skew** or **parallel**?

Solution:

- The vector equations of the lines are:

$$L_1(t) = \langle 1, 0, 1 \rangle + t\langle -2, 4, 0 \rangle = \langle 1 - 2t, 4t, 1 \rangle$$

$$L_2(s) = \langle 2, 3, -1 \rangle + s\langle 2, 1, -2 \rangle = \langle 2 + 2s, 3 + s, -1 - 2s \rangle$$

- Equating z -coordinates, we find $1 = -1 - 2s \implies s = -1$.
- Equating y -coordinates with $s = -1$, we find $4t = 3 - 1 \implies t = \frac{1}{2}$.
- Equating x -coordinates with $s = -1$ and $t = \frac{1}{2}$, we find:

$$L_1\left(\frac{1}{2}\right) = \langle 0, 2, 1 \rangle = L_2(-1).$$

- Hence, the lines **intersect**.



Problem 33(a)

Find the volume V of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Problem 33(a)

Find the volume **V** of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The volume **V** is equal to the absolute value of the determinant of the matrix with rows $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$, $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$, $\overrightarrow{AD} = \langle 0, -4, -3 \rangle$.

Problem 33(a)

Find the volume **V** of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The volume **V** is equal to the absolute value of the determinant of the matrix with rows $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$, $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$, $\overrightarrow{AD} = \langle 0, -4, -3 \rangle$.

$$\mathbf{V} = \left\| \begin{vmatrix} 2 & -3 & -4 \\ 3 & -1 & -5 \\ 0 & -4 & -3 \end{vmatrix} \right\|$$

Problem 33(a)

Find the volume **V** of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The volume **V** is equal to the absolute value of the determinant of the matrix with rows $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$, $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$, $\overrightarrow{AD} = \langle 0, -4, -3 \rangle$.

$$\mathbf{V} = \begin{vmatrix} 2 & -3 & -4 \\ 3 & -1 & -5 \\ 0 & -4 & -3 \end{vmatrix}$$

$$= |2 \cdot (-17) + -(-3) \cdot (-9) + (-4) \cdot (-12)|$$

Problem 33(a)

Find the volume **V** of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The volume **V** is equal to the absolute value of the determinant of the matrix with rows $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$, $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$, $\overrightarrow{AD} = \langle 0, -4, -3 \rangle$.

$$\mathbf{V} = \begin{vmatrix} 2 & -3 & -4 \\ 3 & -1 & -5 \\ 0 & -4 & -3 \end{vmatrix}$$

$$= |2 \cdot (-17) + -(-3) \cdot (-9) + (-4) \cdot (-12)| = |-13|$$

Problem 33(a)

Find the volume **V** of the **parallelepiped** such that the following four points $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$, $D = (1, 0, -1)$ are vertices and the vertices B, C, D are all adjacent to the vertex A .

Solution:

The volume **V** is equal to the absolute value of the determinant of the matrix with rows $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$, $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$, $\overrightarrow{AD} = \langle 0, -4, -3 \rangle$.

$$\mathbf{V} = \begin{vmatrix} 2 & -3 & -4 \\ 3 & -1 & -5 \\ 0 & -4 & -3 \end{vmatrix}$$

$$= |2 \cdot (-17) + -(-3) \cdot (-9) + (-4) \cdot (-12)| = |-13| = 13.$$



Problem 33(b)

Find an **equation of the plane** through

$$A = (1, 4, 2), B = (3, 1, -2), C = (4, 3, -3).$$

Problem 33(b)

Find an **equation of the plane** through

$$A = (1, 4, 2), B = (3, 1, -2), C = (4, 3, -3).$$

Solution:

- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.

Problem 33(b)

Find an **equation of the plane** through

$A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$.

Solution:

- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.
- The normal vector is:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC}$$

Problem 33(b)

Find an **equation of the plane** through

$$A = (1, 4, 2), B = (3, 1, -2), C = (4, 3, -3).$$

Solution:

- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.
- The normal vector is:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -3 & -4 \\ 3 & -1 & -5 \end{vmatrix}$$

Problem 33(b)

Find an **equation of the plane** through

$$A = (1, 4, 2), B = (3, 1, -2), C = (4, 3, -3).$$

Solution:

- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.
- The normal vector is:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -3 & -4 \\ 3 & -1 & -5 \end{vmatrix} = 11\mathbf{i} - 2\mathbf{j} + 7\mathbf{k}.$$

Problem 33(b)

Find an **equation of the plane** through

$$A = (1, 4, 2), B = (3, 1, -2), C = (4, 3, -3).$$

Solution:

- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.
- The normal vector is:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -3 & -4 \\ 3 & -1 & -5 \end{vmatrix} = 11\mathbf{i} - 2\mathbf{j} + 7\mathbf{k}.$$

- Since $A = (1, 4, 2)$, is on the plane, then the **equation of the plane** is given by:

Problem 33(b)

Find an **equation of the plane** through

$A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$.

Solution:


- Consider the vectors $\overrightarrow{AB} = \langle 2, -3, -4 \rangle$ and $\overrightarrow{AC} = \langle 3, -1, -5 \rangle$ which lie parallel to the plane.
- The normal vector is:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -3 & -4 \\ 3 & -1 & -5 \end{vmatrix} = 11\mathbf{i} - 2\mathbf{j} + 7\mathbf{k}.$$

- Since $A = (1, 4, 2)$, is on the plane, then the **equation of the plane** is given by:


$$11(x - 1) - 2(y - 4) + 7(z - 2) = 0.$$





Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$ and the xy -plane.




Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3 - 3)$ and the xy -plane.

Solution:

- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$,
 $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.




Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$ and the xy -plane.

Solution:

- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$, $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.
- If θ is the angle between the planes, then:




Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3 - 3)$ and the xy -plane.

Solution:

- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$, $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.
- If θ is the angle between the planes, then:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1||\mathbf{n}_2|}$$




Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$ and the xy -plane.

Solution:

- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$, $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.
- If θ is the angle between the planes, then:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1||\mathbf{n}_2|} = \frac{7}{\sqrt{11^2 + (-2)^2 + 7^2}}$$




Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$ and the xy -plane.

Solution:

- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$, $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.
- If θ is the angle between the planes, then:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|} = \frac{7}{\sqrt{11^2 + (-2)^2 + 7^2}} = \frac{7}{\sqrt{174}}.$$



Problem 33(c)

Find the angle between the plane through $A = (1, 4, 2)$, $B = (3, 1, -2)$, $C = (4, 3, -3)$ and the xy -plane.

Solution:


- The normal vectors of these planes are $\mathbf{n}_1 = \langle 0, 0, 1 \rangle$, $\mathbf{n}_2 = \langle 11, -2, 7 \rangle$.
- If θ is the angle between the planes, then:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|} = \frac{7}{\sqrt{11^2 + (-2)^2 + 7^2}} = \frac{7}{\sqrt{174}}.$$

•

$$\theta = \cos^{-1} \left(\frac{1}{\sqrt{174}} \right).$$






Problem 37

Find the distance D between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$




Problem 37

Find the distance D between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane.




Problem 37

Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$.




Problem 37

Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$. The distance **D** from P_1 to the first plane is equal to:



Problem 37


Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$. The distance **D** from P_1 to the first plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| =$$



Problem 37


Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$. The distance **D** from P_1 to the first plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right|$$



Problem 37


Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$. The distance **D** from P_1 to the first plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = \left| \langle 0, 0, \frac{5}{2} \rangle \cdot \frac{1}{\sqrt{6}} \langle 2, 1, -1 \rangle \right|$$



Problem 37

Find the distance **D** between the given parallel planes

$$z = 2x + y - 1, \quad -4x - 2y + 2z = 3.$$

Solution:

The normal to the first plane is $\mathbf{n} = \langle 2, 1, -1 \rangle$ and the point $P_0 = (0, 0, -1)$ lies on this plane. The point $P_1 = \langle 0, 0, \frac{3}{2} \rangle$ lies on the second plane. Consider the vector from P_0 to P_1 which is $\mathbf{b} = \langle 0, 0, \frac{5}{2} \rangle$. The distance **D** from P_1 to the first plane is equal to:

$$|\text{comp}_{\mathbf{n}} \mathbf{b}| = \left| \mathbf{b} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} \right| = \left| \langle 0, 0, \frac{5}{2} \rangle \cdot \frac{1}{\sqrt{6}} \langle 2, 1, -1 \rangle \right| = \frac{5}{2\sqrt{6}}.$$



Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Solution:

- The direction vector $\mathbf{a} = \langle 3, -1, 2 \rangle$ of the line is parallel to the plane.

Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Solution:

- The direction vector $\mathbf{a} = \langle 3, -1, 2 \rangle$ of the line is parallel to the plane.
- For $t = 0$, the point $Q = \langle 2, 1, 2 \rangle$ on the line and the plane.

Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Solution:

- The direction vector $\mathbf{a} = \langle 3, -1, 2 \rangle$ of the line is parallel to the plane.
- For $t = 0$, the point $Q = \langle 2, 1, 2 \rangle$ on the line and the plane.
- So $\mathbf{b} = \overrightarrow{PQ} = \langle 1, 0, 2 \rangle$ is also parallel to the plane.

Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Solution:

- The direction vector $\mathbf{a} = \langle 3, -1, 2 \rangle$ of the line is parallel to the plane.
- For $t = 0$, the point $Q = \langle 2, 1, 2 \rangle$ on the line and the plane.
- So $\mathbf{b} = \overrightarrow{PQ} = \langle 1, 0, 2 \rangle$ is also parallel to the plane.
- To find a normal vector to the plane, take cross products:

$$\mathbf{n} = \mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -1 & 2 \\ 1 & 0 & 2 \end{vmatrix} = \langle -2, -4, 1 \rangle.$$

Problem 41

Find an **equation of the plane** that passes through the point $P(1, 1, 0)$ and contains the line given by **parametric equations** $x = 2 + 3t$, $y = 1 - t$, $z = 2 + 2t$.

Solution:

- The direction vector $\mathbf{a} = \langle 3, -1, 2 \rangle$ of the line is parallel to the plane.
- For $t = 0$, the point $Q = \langle 2, 1, 2 \rangle$ on the line and the plane.
- So $\mathbf{b} = \overrightarrow{PQ} = \langle 1, 0, 2 \rangle$ is also parallel to the plane.
- To find a normal vector to the plane, take cross products:

$$\mathbf{n} = \mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -1 & 2 \\ 1 & 0 & 2 \end{vmatrix} = \langle -2, -4, 1 \rangle.$$

- Since $(1, 1, 0)$ is on the plane, the **equation of the plane** is:
 $\langle -2, -4, 1 \rangle \cdot \langle x - 1, y - 1, z \rangle = -2(x - 1) - 4(y - 1) + z = 0.$

