The Nonclassicality Network: a signed theorem graph for quantum foundations with machine-visible open problems

[Author name]
[Affiliation] [email]

October 2026 (v2.4 of the network)

Abstract

We present a curated, typed, signed theorem graph of quantum foundations, quantum information, quantum thermodynamics and resource theories, released as an open repository with two interactive pages and machine-readable data. Nodes are typed (axioms, models, state-operator clusters, process-operator clusters, witnesses, effects, tasks); edges are signed relations carrying verbatim statements and references, with impossibility (no-go) theorems recorded as first-class negative edges, so that the *absence* of an edge always means "unsettled" rather than "false". Witnesses are operator-valued nodes paired one-to-one with the effects they detect, each carrying structured requirement and blindness records; a state×process verdict matrix then enumerates open cells mechanically (in the current release, 445 of 3094 cells are settled). We demonstrate that the structure is generative: closing formerly uncharted cells yields several propositions proved here, including a dephasing-factorization theorem (covariant processes with time-translation-symmetric effects form simplex-embeddable, hence noncontextual, fragments for every input state), a classification of time-translation-symmetric bipartite correlations (for nondegenerate noninteracting Hamiltonians all such states are classical–classical; quantum correlations survive only in degeneracy blocks), a composed corollary exhibiting Bell-local yet contextual scenarios from Werner states, and independence counterexamples separating magic from discord. Fifteen named open problems are registered. The intended use is threefold: semi-automated generation of research problems from the gap census; a reviewable survey format in which checking the review means auditing individually cited edges; and an interactive visualization of the field's logical structure. All sources, data and pages are provided for community extension.

1 Introduction

Quantum foundations is unusually rich in *impossibility* results: Bell's theorem, the Kochen–Specker theorem, no-cloning and no-broadcasting, the PBR theorem, and the modern no-go theorems of generalized contextuality. It is equally rich in *possibility* and *equivalence* results: the hierarchy of Bell nonlocality, steering and entanglement [1], the equivalences between nonclassicality, incompatibility, no-broadcasting, superposition and entangleability in generalized probabilistic theories (GPTs) [2, 3, 4, 5], and the equivalences tying Wigner negativity to contextuality in odd prime dimensions [6] and in continuous variables [7]. These results are scattered across subfields that use different formalisms, and simple questions of the form "does property X imply property Y, in which fragment, for which states and which measurements?" are surprisingly hard to answer from the literature.

This paper introduces a working tool addressed at exactly that difficulty: a *signed theorem graph* — the Nonclassicality Network — whose design goal is that missing knowledge be visible. Three design decisions do the work.

1. **Three-valued edges.** Between any two concepts the graph distinguishes (i) a positive edge (a theorem asserting the relation), (ii) a negative edge (a theorem asserting the relation *cannot* hold: a no-go, an exclusion, a counterexample), and (iii) no edge, which by convention means *unsettled*. Without (ii), impossibility theorems would be encoded as absences and become indistinguishable from ignorance; a search for open problems would then re-propose closed questions. Negative edges are drawn with inhibition-style flat-bar terminators ($\dashv$) and are filterable.

2. **Typed layers with operator-valued witnesses.** Nodes are typed: axioms (including a "genesis" family: symplectic structure, epistemic restriction, the Poisson$\to$Moyal lift, the complex unit i, and $\hbar$), models, state-operator clusters, process-operator clusters, witnesses, effects, and tasks. Every operational effect is paired one-to-one with the witness (functional, inequality, semidefinite or linear program, protocol) that detects it — the design principle *no effect without a witness* — and each witness node carries structured records: what it requires at the model, state and process level, and on which clusters it is provably blind. The recorded exception to strict pairing is threshold stratification: the CHSH functional detects Bell violation, the Tsirelson boundary and PR correlations at thresholds 2, $2\sqrt{2}$, 4 [8, 9, 10].

3. **Mechanical gap enumeration.** For each witness, a state-cluster$\times$process-cluster matrix is populated from recorded theorems; every unfilled cell is, by construction, an open question. The current release settles 445 of 3094 cells; the remaining 2649 cells, together with 15 named open problems, constitute a machine-generated research agenda.

Motivation and intended use. The network was built for three purposes, and the design decisions above map onto them directly. *(i) Automating research-idea generation.* Because the absence of an edge means "unsettled" and never "false", the set of open questions is a queryable object rather than an impression of the literature: every unfilled matrix cell and every missing edge between typed nodes is a candidate theorem, pre-sorted by the structure into four closure routes — compose recorded edges, prove directly, exhibit a counterexample, or locate the result in the literature. Section 3 demonstrates that this loop produces nontrivial output: thirteen propositions across three curation rounds, each of which began as a specific uncharted cell. *(ii) Reviewing a field as curation.* The graph is a review article in a different format: each claim of the review is a single edge carrying a verbatim statement, references and a provenance status, so checking the review means auditing edges one at a time, and disputing it means objecting to a specific edge identifier rather than to a paragraph. Releases are versioned and each records its gap census, so the review does not silently decay. *(iii) Visualizing the field.* The layered, signed, filterable pages make the logical shape of nonclassicality visible at a glance — where possibility flows, where no-go bars block, which clusters a given witness can never see — and the witness panels answer the operational question "what is required, at model, state and process level, to observe effect X?" interactively. The review process is itself the generator: surveying the field in this format yields, as a by-product, its own research agenda.

The repository contains two interactive pages over the same data — a flat version (v1) and the layered version (v2) — documenting the evolution of the structure, together with machine-readable JSON carrying a verbatim statement and references for every edge. Section 2 summarizes the curated backbone. Section 3 contains the generative payoff: propositions obtained by closing formerly uncharted cells, with proofs. Section 4 lists the registered open problems. Section 5 states the curation and verification protocol — what was verified against the literature, what is cited from standard knowledge, and what is new here — and Section 6 describes the repository.

2 The curated backbone

Two nested classicality boundaries organize the graph. First, for a single GPT system the following are equivalent characterizations of *nonclassicality* (non-simpliciality of the state cone): the existence of incompatible measurements [2]; the impossibility of universal broadcasting [3]; the existence of "superposition" (strong incompatibility) [5]; and *entangleability*, i.e. the strict separation of the minimal and maximal tensor products with any other nonclassical system — the resolution of Barker's conjecture [4, 5]. Second, and strictly stronger, a prepare–measure fragment admits a generalized-noncontextual ontological model [11, 12] if and only if it is *simplex-embeddable* [13]; the complete test is a linear program [14]. The epistricted theories — Spekkens' toy theory [15, 16], Gaussian quantum mechanics as epistricted Liouville mechanics [17], and the odd-dimensional stabilizer subtheory with Gross's nonnegative Wigner function [18, 19, 20] — live between the two boundaries: they exhibit incompatibility, no-cloning, interference phenomenology and steering-like ensembles while remaining noncontextual. The qubit stabilizer subtheory is the known anomaly: it admits no nonnegative diagram-preserving representation and is state-independently contextual [20, 21, 22, 23].

Beyond this backbone the graph records (with per-edge references in the data files): the nonlocality–steering–entanglement hierarchy and its strictness [1, 24, 25, 26]; incompatibility as necessary for Bell violations [27] and equivalent to steerability [28, 29]; the failure of the Peres conjecture [30]; the negativity–contextuality equivalences [6, 7] with the representation-relativity caveat of the nonnegative Λ-polytope model [31]; contextuality as a precondition for entanglement [32]; the conditional equivalence of incompatibility and contextuality when all states are available [33] against its unconditional failure [34]; no-broadcasting as a characterization of operational contextuality [35]; thermodynamic witnesses of contextuality [36, 37, 38]; coherence as asymmetry and its thermodynamic role [39, 40]; the genesis decomposition (Groenewold–van Hove obstruction and the Moyal deformation [41, 42, 43, 44]; Hudson's theorem [45]; non-quadratic generators as the source of continuous-variable universality and negativity [46, 47]; local tomography and the complex unit [48, 49, 50]); the almost-quantum obstruction to the no-restriction hypothesis [51, 52]; and the Sorkin hierarchy [53, 54, 55].

3 Propositions generated by closing uncharted cells

Each result below occupied a cell or node pair with no recorded connecting or disconnecting theorem. We state the network provenance (which edges were composed) with each item. Results labelled "derived" are proved here; we do not claim they are absent from the literature, only that we did not find them stated (see Sec. 5).

3.1 Round 1: boundary corollaries

Proposition 1 (Contextuality implies entangleability; provenance $A5 \circ A3$)**.** *If a tomographically complete prepare–measure fragment of a GPT system A fails generalized noncontextuality, then for every nonclassical system B there exists a consistent composite containing entangled states or entangled effects.*

Proof. Contextuality implies failure of simplex embeddability [13], hence the state cone of A is not simplicial. By the entangleability of cones [4, 5], the minimal and maximal tensor products of A with any nonclassical B differ, and any composite beyond the minimal one contains entangled elements. □

The converse fails (the toy theory is nonsimplicial yet noncontextual [15, 13]), and the unique escape is postulating the minimal tensor product: *separable quantum theory* retains

Wigner negativity, magic and contextuality of its quantum marginals while possessing no entangled, steerable or Bell-violating states, and purification [56] fails in it by construction (cones between the minimal and maximal tensor product are standard [57]). A complementary necessity direction is the theorem of Plávala and Gühne [32]. Similarly, a fragment whose states and effects are diagonal in one basis is simplex-embeddable, so basis-independent coherence is necessary for contextuality, while epistricted counterexamples [15, 58] and vanishing-coherence contextuality [59] show it is not sufficient and has no threshold; quantitative scenario-specific witnesses exist [60].

3.2 Round 2: time-translation symmetry hides contextuality

Let H be a Hamiltonian with nondegenerate spectrum, D the associated energy-dephasing map, and call a state, effect or channel *TTS* (time-translation symmetric, covariant) when it commutes with the unitary group $U_t = e^{-iHt}$.

Theorem 1 (Covariant fragments are noncontextual; provenance: uncharted cell TTS×covariant)**.** *Consider any prepare–transform–measure fragment whose transformations T are covariant and whose effects E are TTS. Then for nondegenerate H the fragment's statistics satisfy* $\mathrm{Tr}\big(E\,T(\rho)\big) = \mathrm{Tr}\big(E\,T(D(\rho))\big)$ *for every state ρ; the fragment factors through the commutative energy-diagonal subtheory, is simplex-embeddable and hence noncontextual, its Kirkwood–Dirac distributions over energy chains are genuine probabilities, and no coherence monotone is estimable from it.*

Proof. For TTS effects, $U_t^\dagger E U_t = E$, so $\mathrm{Tr}(E\,T(U_t\rho U_t^\dagger)) = \mathrm{Tr}(U_t^\dagger E U_t\,T(\rho)) = \mathrm{Tr}(E\,T(\rho))$ using covariance of T. Averaging over t yields invariance under pre-dephasing. For nondegenerate H, D maps onto the diagonal (commutative) algebra, TTS effects are diagonal, and covariant channels preserve the diagonal subalgebra, so the fragment is tomographically equivalent to a classical stochastic fragment, which embeds in a simplex with its dual effects; noncontextuality follows from the characterization of [13]. □

Theorem 1 says TTS processes do not lack contextuality but *hide* it: the hiding map is exactly D. Lostaglio's noncontextuality of two-point-measurement fluctuation theorems [36] is the special case $T = \mathrm{id}$, E the energy projectors, and the no-go of [37] is the statement that the first projective energy measurement implements the dephasing. For degenerate H the factorization is only block-wise and covariant fragments can remain contextual (open problem SG1). Conversely, the state side carries no blindness at all:

Proposition 2 (No state-blind cluster for contextuality)**.** *Every quantum state — Gibbs states and the maximally mixed state included — violates state-independent contextuality inequalities (the Peres–Mermin inequality of [23]; the Yu–Oh 13-ray inequality [61]). Consequently, for nondegenerate H, any state-independent contextuality witness set necessarily contains non-TTS measurements.*

Proof. The first statement is the content of state-independence in [23, 61]. The second follows from Theorem 1: a TTS witness set would yield a noncontextual fragment for every state. □

3.3 Round 3: new closures

Proposition 3 (Symmetric bipartite states of generic noninteracting pairs are classically correlated; provenance: uncharted cell TTS×correlation clusters)**.** *Let $H = H_A + H_B$ with nondegenerate total spectrum (all sums $E_i^A + E_j^B$ distinct). Then every TTS state is classical–classical: zero discord, hence separable, unsteerable and Bell-local. Entangled or discordant TTS states exist iff the total spectrum is degenerate, and all their quantum correlations are supported inside degeneracy blocks.*

Proof. Nondegeneracy of the total spectrum forces the eigenbasis of H to be the unique product basis $|E_i^A\rangle \otimes |E_j^B\rangle$ (any local degeneracy or cross-degeneracy would produce a degenerate sum). A state commuting with a nondegenerate H is diagonal in its eigenbasis, hence of the form $\sum_{ij} p_{ij}|ij\rangle\langle ij|$, a classical–classical state: zero discord [62, 63], separable, and therefore unsteerable and Bell-local [1]. For the degenerate case, two equal-gap qubits admit the TTS maximally entangled state $(|01\rangle + |10\rangle)/\sqrt{2}$ supported in the degenerate $\{01, 10\}$ block. □

Remark 1. Proposition 3 is elementary and may be folklore; a literature search (2026-10-23) found no prior statement of the classification. It is consistent with experiment: the correlation-driven reversal of heat flow of Micadei *et al.* [38] employs initial coherence $|01\rangle\langle 10|$ — exactly a degenerate-block term for equal-frequency qubits, the sole habitat the proposition allows.

Lemma 1 (TTS measurements are compatible)**.** *For nondegenerate H, any two TTS sharp observables commute (each is a function of H), and any TTS POVM is a classical post-processing of the energy measurement; the whole TTS measurement cluster is jointly measurable. Consequently TTS probes admit no incompatibility, hence no steering [28, 29] and no Bell violation [27], and the commutativity of the fragment reproves the blindness column of Theorem 1.*

Proof. $[A, H] = 0$ with H nondegenerate forces A diagonal in the energy basis; two diagonal operators commute. A POVM with diagonal effects $E_k = \sum_i e_{k|i}\Pi_i$ is simulated by measuring H and sampling k from $e_{k|i}$. □

Corollary 1 (Bell-local yet contextual scenarios; provenance $A8 \circ B3$)**.** *There exist bipartite entangled states admitting local hidden variable models for all single-copy measurements [24, 25] whose derived sequential (remote-state-preparation) scenario, equipped with all measurements on the steered side, is preparation-and-measurement contextual. Generalized contextuality is therefore a strictly finer detector of composite nonclassicality than Bell violation.*

Proof. Theorem 1 of [32] states that a preparation-and-measurement noncontextual model for the steered states together with all measurements implies separability. The contrapositive applies to every entangled state, in particular to Werner states inside the LHV region. Bell locality for all single-copy measurements is Refs. [24, 25]. □

Remark 2. The corollary is an immediate inference from [32]; we verified against that paper that it is not stated there explicitly, and a search for follow-up work stating it (2026-10-23) found none. The nearest neighbor connects contextuality of steered preparations to constraints on measurement incompatibility (`arXiv:2406.16095`) rather than to Bell-local entangled states. Note the fragment scope: the contextual fragment contains *all* measurements on the steered side; restricted fragments can remain noncontextual — e.g. entangled stabilizer states probed by stabilizer measurements admit the local nonnegative Gross model [18, 20]. There is no tension between the two statements.

Example 1 (Magic and discord are logically independent)**.** (a) $|T\rangle \otimes |0\rangle$, with $|T\rangle$ a canonical non-stabilizer state [64], is a pure product state — zero discord, zero entanglement — yet magic. (b) $\rho = \frac{1}{2}(|0\rangle\langle 0| \otimes |0\rangle\langle 0| + |+\rangle\langle +| \otimes |+\rangle\langle +|)$ is a mixture of stabilizer product states (hence inside the stabilizer polytope, non-magic) but has nonzero discord on both sides: if $[\rho, A \otimes \mathbb{1}] = 0$ then $[P_0, A] \otimes P_0 = -[P_+, A] \otimes P_+$, and since P_0, P_+ are linearly independent both commutators vanish; A then commutes with P_0 and P_+, which generate the full qubit algebra, so $A \propto \mathbb{1}$. Thus no nontrivial local observable commutes with ρ, which therefore is not classical on either side [62, 63]. Together (a) and (b) close the formerly uncharted magic–discord pair in both directions: no unconditional edge can connect them. (The interplay of non-stabilizerness with *entanglement* is actively studied in many-body settings [65]; a search found no treatment of the discord pair.)

Proposition 4 (Sorkin order of epistricted theories; inheritance)**.** *Gaussian quantum mechanics and the odd-dimensional stabilizer subtheory satisfy the Sorkin truncation* $I_3 = 0$ *identically.*

Proof. Both are subtheories of quantum theory — their states, transformations and measurements are quantum objects with quantum statistics [17, 18] — and I_3 vanishes identically in quantum theory [53]. □

Conjecture 1 (SG4)**.** The discrete toy (field) theory, which is *not* a quantum subtheory (it does not reproduce GHZ-type stabilizer statistics [15, 19]), has an undetermined Sorkin order: define its three-path interferometer and compute I_3. Either outcome is informative.

Finally, one formerly uncharted cell is closed by existing literature that the graph had not yet absorbed: the Kirkwood–Dirac analogue of Hudson's boundary. Pure KD-classical states are characterized by support-uncertainty ("complete incompatibility") conditions [66]; noncommutation alone does not suffice for KD nonpositivity [67]; and for discrete-Fourier bases a pure state is KD-classical iff $n_A(\psi)\, n_B(\psi) = d$ [68].

4 Registered open problems

The repository registers fifteen named problems; abbreviated here. G1: a steering analogue of the entangleability of cones (with a stated conjecture). G2: quantitative coherence→contextuality bounds, approachable by sweeping fragments with the simplex-embedding linear program [14]. G3: an even-dimensional magic–contextuality equivalence, obstructed by state-independent Pauli contextuality and by the Λ-polytope model [31]. G4: completeness of thermodynamic contextuality witnesses within thermal operations (sharpened by Theorem 1: probes must leave the TTS cluster). G5: contextuality quantifiers for higher-order interference theories [55]. G6: which thermodynamic requirements force purification-type axioms. G7: a monogamy calculus between contextuality and nonlocality. G8: NPT bound entanglement and its thermodynamics. G9: a principled measurement-restriction rule for almost-quantum theories [51, 52]. G10: a complexity-theoretic column for every witness (contextuality is a linear program; separability is NP-hard [69]; quantum-set membership is undecidable [70]). SG1: covariant contextuality inside degeneracy blocks (the failure mode of Theorem 1). SG2: threshold stratification of witness families. SG3: the uncharted-cell program itself (2649 cells at this release). SG4: the toy-field-theory Sorkin order (above). SG5: hidden-contextuality activation: do local filters, which activate hidden nonlocality [24, 25], amplify the sequential-scenario contextuality of Corollary 1?

5 Curation and verification protocol

Reliability is handled by explicit provenance rather than by implication. (i) Every edge in the data files carries a verbatim statement and references; the per-edge reference list, not this paper, is the citation of record for compiled results. (ii) Edges marked *theorem* are literature claims; the load-bearing post-2019 results were verified against bibliographic sources during curation (October 2026), while classic results (e.g. [45, 41, 42, 27, 24]) are cited from standard knowledge and their page data should be rechecked before quotation. (iii) Edges marked *derived* are proved in this paper or in the repository catalogs and are *not* claimed to be previously unpublished; Propositions 1, 3 and Lemma 1 are elementary enough to be folklore, and a dedicated novelty pass is advised before attributing priority. (iv) Composed corollaries state exactly which recorded theorems they compose, so a reader can audit each in one step. (v) The bibliography was verified programmatically against Crossref on 2026-10-23: 58 of 70 entries matched exactly on volume, page and year; the remainder are classic papers that Crossref indexes poorly (e.g. [8, 42, 50, 9, 45]), whose standard citations are retained, plus two records corrected from publisher metadata [68, 35]. Novelty searches for Theorem 1, Proposition 3,

Corollary 1 and Example 1 were run the same day; no prior statements were found, and the nearest neighbors located are recorded in the corresponding remarks. (vi) The authors take responsibility for remaining errors and welcome corrections as repository issues; the graph is versioned, and every release records its gap census.

6 Repository

The repository (`https://github.com/tahmeedhossain/GEN_GPT_Level`, served as live pages at `https://tahmeedhossain.github.io/GEN_GPT_Level/`) contains: `index.html` (overview and the generated-proposition record); `v1.html` (flat network, 81 edges); `v2.html` (layered network, 148 signed edges, witness records, verdict matrix with uncharted-cell census); `data/network_v1.json` and `data/network_v2.json` (machine-readable: nodes, signed edges with statements and references, witness records, combinations, polarity metadata); `catalog/` (prose catalogs with full proofs); and this paper's source under `paper/`. The data schema is documented in the README; contributions of new edges are expected to carry a statement, a reference or proof, a kind, a status and a polarity.

Acknowledgements. The curation of the network, the derivations and drafting of this paper, the interactive pages and the data files were produced with substantial assistance from an AI system (Claude, Anthropic), working under the direction of the author, who reviewed the material and takes sole responsibility for all claims and any remaining errors.

References

[1] H. M. Wiseman, S. J. Jones, and A. C. Doherty. Steering, entanglement, nonlocality, and the Einstein–Podolsky–Rosen paradox. *Phys. Rev. Lett.*, 98:140402, 2007.

[2] M. Plávala. All measurements in a probabilistic theory are compatible if and only if the state space is a simplex. *Phys. Rev. A*, 94:042108, 2016. arXiv:1608.05614.

[3] H. Barnum, J. Barrett, M. Leifer, and A. Wilce. Generalized no-broadcasting theorem. *Phys. Rev. Lett.*, 99:240501, 2007.

[4] G. Aubrun, L. Lami, C. Palazuelos, and M. Plávala. Entangleability of cones. *Geom. Funct. Anal.*, 31:181–205, 2021. arXiv:1911.09663.

[5] G. Aubrun, L. Lami, C. Palazuelos, and M. Plávala. Entanglement and superposition are equivalent concepts in any physical theory. *Phys. Rev. Lett.*, 128:160402, 2022. arXiv:2109.04446.

[6] M. Howard, J. Wallman, V. Veitch, and J. Emerson. Contextuality supplies the 'magic' for quantum computation. *Nature*, 510:351, 2014.

[7] R. I. Booth, U. Chabaud, and P.-E. Emeriau. Contextuality and wigner negativity are equivalent for continuous-variable quantum measurements. *Phys. Rev. Lett.*, 129:230401, 2022. arXiv:2111.13218.

[8] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt. Proposed experiment to test local hidden-variable theories. *Phys. Rev. Lett.*, 23:880, 1969.

[9] B. S. Cirel'son. Quantum generalizations of Bell's inequality. *Lett. Math. Phys.*, 4:93, 1980.

[10] S. Popescu and D. Rohrlich. Quantum nonlocality as an axiom. *Found. Phys.*, 24:379, 1994.

[11] R. W. Spekkens. Contextuality for preparations, transformations, and unsharp measurements. *Phys. Rev. A*, 71:052108, 2005.

[12] R. W. Spekkens. Negativity and contextuality are equivalent notions of nonclassicality. *Phys. Rev. Lett.*, 101:020401, 2008.

[13] D. Schmid, J. H. Selby, E. Wolfe, R. Kunjwal, and R. W. Spekkens. Characterization of noncontextuality in the framework of generalized probabilistic theories. *PRX Quantum*, 2:010331, 2021. arXiv:1911.10386.

[14] J. H. Selby, E. Wolfe, D. Schmid, A. B. Sainz, and V. P. Rossi. Linear program for testing nonclassicality and an open-source implementation. *Phys. Rev. Lett.*, 132:050202, 2024. arXiv:2204.11905.

[15] R. W. Spekkens. Evidence for the epistemic view of quantum states: A toy theory. *Phys. Rev. A*, 75:032110, 2007.

[16] R. W. Spekkens. Quasi-quantization: classical statistical theories with an epistemic restriction. In *Quantum Theory: Informational Foundations and Foils*. Springer, 2016. arXiv:1409.5041.

[17] S. D. Bartlett, T. Rudolph, and R. W. Spekkens. Reconstruction of gaussian quantum mechanics from liouville mechanics with an epistemic restriction. *Phys. Rev. A*, 86:012103, 2012.

[18] D. Gross. Hudson's theorem for finite-dimensional quantum systems. *J. Math. Phys.*, 47:122107, 2006.

[19] L. Catani and D. E. Browne. Spekkens' toy model in all dimensions and its relationship with stabiliser quantum mechanics. *New J. Phys.*, 19:073035, 2017.

[20] D. Schmid, H. Du, J. H. Selby, and M. F. Pusey. Uniqueness of noncontextual models for stabilizer subtheories. *Phys. Rev. Lett.*, 129:120403, 2022. arXiv:2101.06263.

[21] N. D. Mermin. Simple unified form for the major no-hidden-variables theorems. *Phys. Rev. Lett.*, 65:3373, 1990.

[22] A. Peres. Two simple proofs of the kochen–specker theorem. *J. Phys. A*, 24:L175, 1991.

[23] A. Cabello. Experimentally testable state-independent quantum contextuality. *Phys. Rev. Lett.*, 101:210401, 2008.

[24] R. F. Werner. Quantum states with einstein–podolsky–rosen correlations admitting a hidden-variable model. *Phys. Rev. A*, 40:4277, 1989.

[25] J. Barrett. Nonsequential positive-operator-valued measurements on entangled mixed states do not always violate a bell inequality. *Phys. Rev. A*, 65:042302, 2002.

[26] N. Gisin. Bell's inequality holds for all non-product states. *Phys. Lett. A*, 154:201, 1991.

[27] A. Fine. Hidden variables, joint probability, and the bell inequalities. *Phys. Rev. Lett.*, 48:291, 1982.

[28] M. T. Quintino, T. Vértesi, and N. Brunner. Joint measurability, einstein–podolsky–rosen steering, and bell nonlocality. *Phys. Rev. Lett.*, 113:160402, 2014.

[29] R. Uola, T. Moroder, and O. Gühne. Joint measurability of generalized measurements implies classicality. *Phys. Rev. Lett.*, 113:160403, 2014.

[30] T. Vértesi and N. Brunner. Disproving the peres conjecture by showing bell nonlocality from bound entanglement. *Nat. Commun.*, 5:5297, 2014.

[31] M. Zurel, C. Okay, and R. Raussendorf. Hidden variable model for universal quantum computation with magic states on qubits. *Phys. Rev. Lett.*, 125:260404, 2020.

[32] M. Plávala and O. Gühne. Contextuality as a precondition for quantum entanglement. *Phys. Rev. Lett.*, 132:100201, 2024. arXiv:2209.09942.

[33] A. Tavakoli and R. Uola. Measurement incompatibility and steering are necessary and sufficient for operational contextuality. *Phys. Rev. Research*, 2:013011, 2020.

[34] J. H. Selby, D. Schmid, E. Wolfe, A. B. Sainz, R. Kunjwal, and R. W. Spekkens. Contextuality without incompatibility. *Phys. Rev. Lett.*, 130:230201, 2023. arXiv:2106.09045.

[35] P. Jokinen, M. Weilenmann, M. Plávala, J.-P. Pellonpää, J. Kiukas, and R. Uola. No-broadcasting characterizes operational contextuality. *Phys. Rev. Lett.*, 133:240201, 2024. arXiv:2406.07305.

[36] M. Lostaglio. Quantum fluctuation theorems, contextuality, and work quasiprobabilities. *Phys. Rev. Lett.*, 120:040602, 2018.

[37] M. Perarnau-Llobet, E. Bäumer, K. V. Hovhannisyan, M. Huber, and A. Acín. No-go theorem for the characterization of work fluctuations in coherent quantum systems. *Phys. Rev. Lett.*, 118:070601, 2017.

[38] K. Micadei et al. Reversing the direction of heat flow using quantum correlations. *Nat. Commun.*, 10:2456, 2019.

[39] I. Marvian and R. W. Spekkens. Extending noether's theorem by quantifying the asymmetry of quantum states. *Nat. Commun.*, 5:3821, 2014.

[40] M. Lostaglio, D. Jennings, and T. Rudolph. Description of quantum coherence in thermodynamic processes requires constraints beyond free energy. *Nat. Commun.*, 6:6383, 2015.

[41] H. J. Groenewold. On the principles of elementary quantum mechanics. *Physica*, 12:405, 1946.

[42] J. E. Moyal. Quantum mechanics as a statistical theory. *Proc. Camb. Phil. Soc.*, 45:99, 1949.

[43] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer. Deformation theory and quantization. I, II. *Ann. Phys.*, 111:61–110, 111–151, 1978.

[44] M. Kontsevich. Deformation quantization of poisson manifolds. *Lett. Math. Phys.*, 66:157, 2003.

[45] R. L. Hudson. When is the wigner quasi-probability density non-negative? *Rep. Math. Phys.*, 6:249, 1974.

[46] S. Lloyd and S. L. Braunstein. Quantum computation over continuous variables. *Phys. Rev. Lett.*, 82:1784, 1999.

[47] R. Takagi and Q. Zhuang. Convex resource theory of non-gaussianity. *Phys. Rev. A*, 97:062337, 2018.

[48] L. Hardy and W. K. Wootters. Limited holism and real-vector-space quantum theory. *Found. Phys.*, 42:454, 2012.

[49] M.-O. Renou, D. Trillo, M. Weilenmann, T. P. Le, A. Tavakoli, N. Gisin, A. Acín, and M. Navascués. Quantum theory based on real numbers can be experimentally falsified. *Nature*, 600:625, 2021.

[50] E. C. G. Stueckelberg. Quantum theory in real Hilbert space. *Helv. Phys. Acta*, 33:727, 1960.

[51] M. Navascués, Y. Guryanova, M. J. Hoban, and A. Acín. Almost quantum correlations. *Nat. Commun.*, 6:6288, 2015.

[52] A. B. Sainz, Y. Guryanova, A. Acín, and M. Navascués. Almost-quantum correlations violate the no-restriction hypothesis. *Phys. Rev. Lett.*, 120:200402, 2018.

[53] R. D. Sorkin. Quantum mechanics as quantum measure theory. *Mod. Phys. Lett. A*, 9:3119, 1994.

[54] U. Sinha, C. Couteau, T. Jennewein, R. Laflamme, and G. Weihs. Ruling out multi-order interference in quantum mechanics. *Science*, 329:418, 2010.

[55] B. Dakić, T. Paterek, and Č. Brukner. Density cubes and higher-order interference theories. *New J. Phys.*, 16:023028, 2014.

[56] G. Chiribella, G. M. D'Ariano, and P. Perinotti. Probabilistic theories with purification. *Phys. Rev. A*, 81:062348, 2010.

[57] P. Janotta and H. Hinrichsen. Generalized probability theories: what determines the structure of quantum theory? *J. Phys. A*, 47:323001, 2014.

[58] L. Catani, M. Leifer, D. Schmid, and R. W. Spekkens. Why interference phenomena do not capture the essence of quantum theory. *Quantum*, 7:1119, 2023.

[59] V. P. Rossi, D. Schmid, J. H. Selby, and A. B. Sainz. Contextuality with vanishing coherence and maximal robustness to dephasing. *Phys. Rev. A*, 108:032213, 2023. arXiv:2212.06856; doi:10.1103/PhysRevA.108.032213.

[60] R. Wagner, A. Camillini, and E. F. Galvão. Coherence and contextuality in a Mach–Zehnder interferometer. *Quantum*, 8:1240, 2024.

[61] S. Yu and C. H. Oh. State-independent proof of kochen–specker theorem with 13 rays. *Phys. Rev. Lett.*, 108:030402, 2012.

[62] H. Ollivier and W. H. Zurek. Quantum discord: A measure of the quantumness of correlations. *Phys. Rev. Lett.*, 88:017901, 2001.

[63] L. Henderson and V. Vedral. Classical, quantum and total correlations. *J. Phys. A*, 34:6899, 2001.

[64] S. Bravyi and A. Kitaev. Universal quantum computation with ideal Clifford gates and noisy ancillas. *Phys. Rev. A*, 71:022316, 2005.

[65] M. Frau, P. S. Tarabunga, M. Collura, M. Dalmonte, and E. Tirrito. Nonstabilizerness versus entanglement in matrix product states. *Phys. Rev. B*, 110:045101, 2024. arXiv:2404.18768; doi:10.1103/PhysRevB.110.045101.

[66] S. De Bièvre. Complete incompatibility, support uncertainty, and Kirkwood–Dirac nonclassicality. *Phys. Rev. Lett.*, 127:190404, 2021.

[67] D. R. M. Arvidsson-Shukur, J. C. Drori, and N. Yunger Halpern. Conditions tighter than noncommutation needed for nonclassicality. *J. Phys. A*, 54:284001, 2021.

[68] J. Xu. Kirkwood–dirac classical pure states. *Phys. Lett. A*, page 129529, 2024. arXiv:2210.02876; doi:10.1016/j.physleta.2024.129529; proves De Bièvre's DFT conjecture $n_A n_B = d$.

[69] L. Gurvits. Classical deterministic complexity of Edmonds' problem and quantum entanglement. In *Proc. 35th ACM Symposium on Theory of Computing (STOC)*, pages 10–19, 2003.

[70] Z. Ji, A. Natarajan, T. Vidick, J. Wright, and H. Yuen. MIP* = RE. *Commun. ACM*, 64(11):131–138, 2021. arXiv:2001.04383.