

When should I suspect hypoglycaemia in a person with type 1 diabetes?

Last revised in December 2024

- [Hypoglycaemia \(/topics/diabetes-type-1/background-information/complications/\)](/topics/diabetes-type-1/background-information/complications/) presents with a wide variety of symptoms (autonomic and neuroglycopenic).
 - Suspect mild hypoglycaemia in a person with type 1 diabetes who presents with the following:
 - Hunger.
 - Anxiety or irritability.
 - Sweating.
 - Tingling lips.
 - Irritability.
 - Palpitations.
 - Tremor.
 - As blood glucose levels fall lower, the person may experience:
 - Weakness and lethargy.
 - Impaired vision.
 - Incoordination.
 - Reduced orientation.
 - Confusion.
 - Irrational behaviour.
 - Emotional lability.
 - Deterioration of cognitive function (when blood glucose levels fall lower than 3.0 mmol/L).
 - Severe hypoglycaemia may result in:
 - Convulsions.
 - Inability to swallow.
 - Loss of consciousness.
 - Coma.

Basis for recommendation

▼ [Show](#)

The content on the NICE Clinical Knowledge Summaries site (CKS) is the copyright of [Clarity Informatics Limited \(trading as Agilio Software Primary Care\)](#) (<https://agiliosoftware.com/primary-care/>). By using CKS, you agree to the licence set out in the [CKS End User Licence Agreement](https://www.nice.org.uk/terms-and-conditions/cks-end-user-licence-agreement) (<https://www.nice.org.uk/terms-and-conditions/cks-end-user-licence-agreement>).

