

When should I suspect hypoglycaemia in a person with type 2 diabetes?

Last revised in March 2025

- Hypoglycaemia (low blood glucose) is generally defined as blood glucose levels less than 3.5 mmol/L.
 - It is the main adverse effect of insulin treatment and may occur in some people taking [sulfonylureas \(/topics/diabetes-type-2/prescribing-information/sulfonylureas/\)](/topics/diabetes-type-2/prescribing-information/sulfonylureas/), and should be avoided wherever possible.
 - See the section on [Hypoglycaemia \(/topics/insulin-therapy-in-type-2-diabetes/management/insulin-therapy-type-2-diabetes/#hypoglycaemia\)](/topics/insulin-therapy-in-type-2-diabetes/management/insulin-therapy-type-2-diabetes/#hypoglycaemia) in the CKS topic on [Insulin therapy in type 2 diabetes \(/topics/insulin-therapy-in-type-2-diabetes/\)](/topics/insulin-therapy-in-type-2-diabetes/) for more information on the prevention, diagnosis, and management of hypoglycaemia.

Basis for recommendation

▼ [Show](#)

The content on the NICE Clinical Knowledge Summaries site (CKS) is the copyright of [Clarity Informatics Limited \(trading as Agilio Software Primary Care\) \(https://agiliosoftware.com/primary-care/\)](#). By using CKS, you agree to the licence set out in the [CKS End User Licence Agreement \(https://www.nice.org.uk/terms-and-conditions/cks-end-user-licence-agreement\)](https://www.nice.org.uk/terms-and-conditions/cks-end-user-licence-agreement).