

No-hair theorem and Spontaneous scalarization of BHs

Taishi Ikeda (Sapienza University di Rome)




SAPIENZA
UNIVERSITÀ DI ROMA



DarkGRA 

My collaborators



Tomohiro Nakamura Masato Minamitsuji



Carlos Herdeiro



Eugen Radu

Outline of my talk

- 1. Introduction**
- 2. No-hair theorem for ST theory**
- 3. Spontaneous scalarization of BH**
 - 1. Scalar-Gauss-Bonnet theory**
 - 2. Other models**
 - 3. Dynamical behavior of the scalarizaion**
- 4. Summary**

Outline of my talk

1. Introduction

2. No-hair theorem for ST theory

3. Spontaneous scalarization of BH

1. Scalar-Gauss-Bonnet theory

2. Other models

3. Dynamical behavior of the scalarizaion

4. Summary

Modification of gravity

- General Relativity (GR)

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{16\pi} R - 2\Lambda + \mathcal{L}_m \right)$$

- GR is quite successful gravitational theory.
- It can explain cosmology, gravitational waves, and so on ...


- Why modify gravity ?

- Dark Energy ?, Dark Matter ?, Quantum correction ?
- Test of GR
- How special is GR ?
- How universal are several GR's properties and theorems ?

➡ Let us modify gravity, and compare with GR.

No-hair theorem of BH

- BH is the simplest compact object in GR.
 - vacuum solution of GR
 - No-hair theorem of BH
 - parameters : M , a , (Q)
- How universal is no-hair theorem ?



no-hair

- Real scalar field
(Bekenstein 1972)
- Non-canonical real scalar
(A.Graham et al 2014)
- Horndeski theory with some conditions (L.Hui et al 2013)

....

solution with scalar hair

- Shift symmetric Scalar Gauss Bonnet gravity
(T.P.Sotiriou et al 2014)
- GR with a conformally coupled scalar field
(N.M.Bocharova et al.1970)

....

Outline of my talk

1. Introduction

2. No-hair theorem for ST theory

3. Spontaneous scalarization of BH

1. Scalar-Gauss-Bonnet theory

2. Other models

**3. Dynamical behavior of the
scalarizaion**

4. Summary

Horndeski theory

$$X := -\frac{1}{2}g^{\mu\nu}\nabla_\mu\Phi\nabla_\nu\Phi$$

$$\Phi_\mu := \nabla_\mu\Phi$$

- Scalar-Tensor theory (with 2nd order EoM, 2+1 DoF)

- Physical degrees of freedom : $(g_{\mu\nu}, \Phi)$

- action

$$S[g, \Phi, A] = \int d^4x \sqrt{-g} \sum_{i=2}^5 \mathcal{L}_{\text{ST}}^i$$

$$\mathcal{L}_{\text{ST}}^2 = G_2(\Phi, X)$$

$$\mathcal{L}_{\text{ST}}^3 = G_3(\Phi, X)\Phi_\mu^\mu$$

* $G_i(\Phi, X)$ ($i = 2, 3, 4, 5$) are arbitrary functions.

$$\mathcal{L}_{\text{ST}}^4 = G_4(\Phi, X)R + G_{4,X}(\Phi, X)\left((\Phi_\mu^\mu)^2 - \Phi_\mu^\nu\Phi_\nu^\mu\right)$$

$$\mathcal{L}_{\text{ST}}^5 = G_5(\Phi, X)G_{\mu\nu}\Phi^{\mu\nu} - \frac{G_{5,X}(\Phi, X)}{6}\left((\Phi_\mu^\mu)^3 - 3(\Phi_\mu^\mu)(\Phi_\nu^\rho\Phi_\rho^\nu) + 2\Phi_\mu^\nu\Phi_\nu^\rho\Phi_\rho^\mu\right)$$

- If $G_i = G_i(X)$, the theory has shift symmetry.

$$\Phi \rightarrow \Phi + c \quad \longrightarrow \quad J_\Phi : \text{Noether current}$$

No-hair theorem

- No- hair theorem for Shift-symmetric Horndeski theory

(L.Hui et. al. (2013))

- Assumptions

1. Static, spherically symmetric, asymptotically flat spacetime
2. Scalar field is also static and spherical symmetry.
3. $\nabla_{\mu}\Phi \rightarrow 0$ at $r \rightarrow \infty$
4. $J_{\Phi\mu}J^{\mu}_{\Phi} < \infty$ on and outside the BH horizon
5. Action has canonical kinetic term of scalar field. : $X \subset \mathcal{L}$
6. $G_{2,3,4,5}(X)$ are analytic at $X = 0$.

- Statement

- Under these assumptions, **BHs can not have a nontrivial scalar hair.**

No-hair theorem

- Comment 1 : Extension


- We can naturally extend no-hair theorem to Scalar-Vector-Tensor theory with additional conditions.

(T.Ikeda et al (2019))

- Comment 2 : How wide theories are included ?

- Scalar-Gauss-Bonnet gravity is “not” included.

$$f(\Phi)\mathcal{G}_{\text{GB}}$$



$$G_2 = 8f^{(4)}(\Phi)X^2 (3 - \ln X)$$

$$G_3 = 4f^{(3)}(\Phi)X (7 - 3 \ln X)$$

$$G_4 = 4f^{(2)}(\Phi)X (2 - \ln X)$$

$$G_5 = -4f'(\Phi)\ln X$$

No-hair BH

- Horndeki theory with some conditions
- Non-canonical real scalar


.....

Hairy BH

- Scalar-Gauss-Bonnet with Shift-symmetry
- GR with a conformally coupled scalar field

.....

?



Spontaneous scalarization

- Weak gravity region : No-hair BH
- Strong gravity region : Hairy BH

Outline of my talk

1. Introduction

2. No-hair theorem for ST theory

3. Spontaneous scalarization of BH

1. Scalar-Gauss-Bonnet theory

2. Other models

**3. Dynamical behavior of the
scalarizaion**

4. Summary

Spontaneous scalarization

- Spontaneous scalarization is one of the mechanism to produce a scalar hair in extreme situation.

$$\square \Phi = \mu_{\text{eff}}^2(g_{\mu\nu}, T)\Phi$$

$\mu_{\text{eff}}^2 < 0$ around $\left\{ \begin{array}{l} \text{dense matter region} \\ \text{strong gravity region} \end{array} \right.$

\rightarrow Neutron star scalarization

\rightarrow Black hole scalarization

cf: T. Damour et al (1993)
T. Harada (1997)

\rightarrow The trivial scalar profile becomes unstable, and scalar hair appears.

- The theory can be distinguish from GR around extreme situation.

Outline of my talk

- 1. Introduction**
- 2. No-hair theorem for ST theory**
- 3. Spontaneous scalarization of BH**

1. Scalar-Gauss-Bonnet theory

2. Other models

3. Dynamical behavior of the scalarizaion

4. Summary

Scalar-Gauss-Bonnet theory

- Action

$$S = \int d^4x \sqrt{-g} \left(R - \frac{1}{2} \nabla_\mu \Phi \nabla^\mu \Phi + f(\Phi) \mathcal{G}_{\text{GB}} \right)$$
$$\mathcal{G}_{\text{GB}} := R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2$$

- EOM

$$\square \Phi = -f'(\Phi) \mathcal{G}$$

- We assume $f'(\Phi_0) = 0$, then Schwarzschild BH is solution with $\Phi = \Phi_0$.
- Around the Schwarzschild BH, $\Phi = \Phi_0 + \delta\Phi$

$$\square \delta\Phi = -\cancel{f''(\Phi_0) \mathcal{G}_{\text{GB}}} \delta\Phi$$

effective mass

Scalar-Gauss-Bonnet theory

- We can proof following theorem. cf: $\square \delta\Phi = -f''(\Phi_0)\mathcal{G}_{\text{GB}}\delta\Phi$
 - Assumption :
 - stationary (Killing vector : $\vec{\xi}$) and asymptotic flat BH spacetime
 - Event horizon is a Killing horizon associated with $\vec{\xi}$
 - $\mathcal{L}_{\vec{\xi}}\Phi = 0$
 - $-f''(\Phi)\mathcal{G}_{\text{GB}} > 0$ in the whole region
 - Then :
 - Scalar field is constant
- This theorem means the scalar field may be able to have nontrivial configuration when $-f''(\Phi)\mathcal{G}_{\text{GB}} < 0$ at some regions.
negative mass square

Spontaneous scalarization

- Spontaneous scalarization of BH in quadratic Scalar-Gauss-Bonnet theory

(H.O.Silva et al.(2018), G.Antoniou et al. (2018), D.D.Doneva et al. (2018))

$$S = \int d^4x \sqrt{-g} \left(R - \frac{1}{2} \nabla_\mu \Phi \nabla^\mu \Phi + f(\Phi) \mathcal{G}_{\text{GB}} \right)$$

with $f(\Phi) = \frac{\eta \Phi^2}{8}$

- EOM : $\square \Phi = \mu_{\text{eff}}^2 \Phi$
- effective mass : $\mu_{\text{eff}}^2(r) = -(\eta/4) \mathcal{G}_{\text{GB}}$
- Around Schwarzschild BH : $\mathcal{G}_{\text{GB}}(r) \Big|_{\text{Sch}} = 48M^2/r^6 > 0$

$$\mu_{\text{eff}}^2(r) < 0 \quad (\eta > 0)$$

➡ Scalar field can have non-trivial profile.


Spontaneous scalarization

$$\Phi(t, r, \theta, \phi) = \sum_{l,m} \frac{e^{i\omega t} \tilde{\sigma}_{lm}(r)}{r} Y_{lm}(\theta, \phi) + \Phi_0$$

- Perturbation around Schwarzschild BH

$$-\partial_{r_*}^2 \tilde{\sigma}_{lm\omega} + V_{\text{eff}}(r) \tilde{\sigma}_{lm\omega} = \omega^2 \tilde{\sigma}_{lm\omega}$$

$$V_{\text{eff}}(r) = (r - 2M) \left(\frac{2M}{r^7} (r^3 - 6M\eta) + \frac{l(l+1)}{r^3} \right)$$





- ▶ Schwarzschild BH is unstable when $\frac{10}{3}M^2 < \eta$
- ▶ New scalarized BH appears.

Spontaneous scalarization

- Scalarized BH

- Boundary condition : regular on r_H , $\Phi \rightarrow 0$ ($r \rightarrow \infty$)
- BHs are labeled by nodes and $\Phi(r_H)$.




Spontaneous scalarization

- Test field analysis

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{16\pi G} R - \frac{1}{2} \nabla_\mu \Phi \nabla^\mu \Phi + f(\Phi) \mathcal{G}_{\text{GB}} \right)$$

- Let us consider $G \rightarrow 0$ with fixed $GM = \text{const.}$
 - ▶ $\mathcal{O}(G^{-1})$: Schwarzschild BH
 - ▶ $\mathcal{O}(G^0)$: $\square \Phi = -f'(\Phi) \mathcal{G}$
- In this limit,

Scalarized BH  Schwarzschild BH
+ (test) non-trivial scalar profile

Spontaneous scalarization

- Stability analysis of test field

$$f(\Phi) = \frac{\eta\Phi^2}{8}$$

$$\Phi(t, r, \theta, \phi) = \sum_{l,m} \frac{e^{i\omega t} \tilde{\sigma}_{lm}(r)}{r} Y_{lm}(\theta, \phi) + \Phi_0(r)$$

↑
non-trivial profile around
Schwarzschild BH

- In quadratic coupling, the scalar field equation is linear.

➡ Effective potential is same as one of constant scalar field.

$$V_{\text{eff}}(r) = (r - 2M) \left(\frac{2M}{r^7} (r^3 - 6M\eta) + \frac{l(l+1)}{r^3} \right)$$

➡ Test non-trivial scalar field is unstable.

- All scalarized BHs of quadric coupling are unstable.

(J.L.Blazquez-Salcedo et al (2018))

Spontaneous scalarization



- Higher order coupling can stabilize the scalarized BH.

- Quartic coupling

M.Minamitsuji, T.Ikeda (2019), H.O.Silva (2019)

$$S = \int d^4x \sqrt{-g} \left(R - \frac{1}{2} \nabla_\mu \Phi \nabla^\mu \Phi + f(\Phi) \mathcal{E}_{\text{GB}} \right)$$

$$f(\Phi) = \frac{\eta}{8} (\Phi^2 + \alpha \Phi^4)$$



$$(\eta/r_H^2, \alpha) = \underline{(0.725, 0)}, \underline{(0.338, 10000)}, \underline{(7.31, -4990)}$$

$$\underline{(4.87, 0)}, \underline{(3.09, 10000)}, \underline{(20.2, -4990)}$$



Spontaneous scalarization

- Stability analysis of test field $G \rightarrow 0$, GM : fix $f(\Phi) = \frac{\eta}{8} (\Phi^2 + \alpha\Phi^4)$

$$\square \Phi = -\frac{\eta}{4} (\Phi + 2\alpha\Phi^3) \mathcal{G}_{\text{GB}}$$

$$V_{\text{eff}}(r) = (r - 2M) \left(\frac{2M}{r^7} (r^3 - 6M\eta(1 + 6\alpha\Phi_0(r)^2)) + \frac{l(l+1)}{r^3} \right)$$

if $\alpha < 0$, this term contribute plus.





$$(\eta/M^2, \alpha) = \underline{(2.14, 0.3)}, \underline{(2.9, 0)}, \underline{(3.32, -0.1)}, \underline{(3.4, -0.115)}, \underline{(3.59, -0.15)}$$

Spontaneous scalarization

- Stability of the scalarized BH

$$f(\Phi) = \frac{\eta}{8} (\Phi^2 + \alpha\Phi^4)$$



- Nodeless scalarized BH solution can stabilize by non-linear term.
- Scalarized BH with one or more nodes are unstable.

Outline of my talk

- 1. Introduction**
- 2. No-hair theorem for ST theory**
- 3. Spontaneous scalarization of BH**
 - 1. Scalar-Gauss-Bonnet theory**
 - 2. Other models**
 - 3. Dynamical behavior of the scalarizaion**
- 4. Summary**

Spontaneous scalarization

- Other model of the spontaneous scalarization

- Einstein-Maxwell-Scalar model C.A.R.Herdeiro et al (2018)


$$S = \int d^4x \sqrt{-g} \left(R + \frac{1}{2} (\nabla \Phi)^2 - \frac{1}{4} f(\Phi) F^{\mu\nu} F_{\mu\nu} \right)$$

- Scalar-Vector-Tensor with double-dual Riemann T.I et al (2019)

$$S = \int d^4x \sqrt{-g} \left(R + \frac{1}{2} (\nabla \Phi)^2 - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} + H(\Phi) L^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta} \right)$$

$$L^{\mu\nu\alpha\beta} = \frac{1}{4} \epsilon^{\mu\nu\rho\sigma} \epsilon^{\alpha\beta\gamma\delta} R_{\rho\sigma\gamma\delta}$$

Reissner-Nordström BH
+ constant scalar field



Charged BH
+ non-trivial scalar field

- Which class of ST theory can realize scalarization ?

➡ Only theory close to SGB gravity M.Minamitsuji, T.I, (2019)

Outline of my talk

- 1. Introduction**
- 2. No-hair theorem for ST theory**
- 3. Spontaneous scalarization of BH**
 - 1. Scalar-Gauss-Bonnet theory**
 - 2. Other models**
 - 3. Dynamical behavior of the scalarizaion**
- 4. Summary**




Scalarizaion on charged sphere

- Spontaneous scalarization is dynamical process.
 - To understand the process, let us consider simple toy model.
- Scalar-Maxwell theory on Minkowski spacetime with charged sphere.

C.A.R.Herdeiro, TI, et al (2018)

$$S = - \int d^4x \left(2\partial^\mu \Phi \partial_\mu \Phi + f(\Phi) F^{\mu\nu} F_{\mu\nu} \right)$$

charged sphere



Scalarizaion on charged sphere

- Equation of motion

$$\begin{cases} \square \Phi - \frac{1}{4} f'(\Phi) F^{\mu\nu} F_{\mu\nu} = 0 \\ D_\mu (f(\Phi) F^{\mu\nu}) = 0 \end{cases}$$

- First, we construct spherically symmetric scalarized solution.

$$\begin{aligned} \Phi &= \Phi_0(r) & A_\mu &= (A_0(r), 0, 0, 0) \\ \left\{ \begin{aligned} \Phi_0''(r) + \frac{2}{r} \Phi_0'(r) + \frac{1}{2} f'(\Phi_0) A_0'(r)^2 &= 0 \\ A_0''(r) + \frac{2}{r} A_0'(r) + \frac{f'(\Phi_0)}{f(\Phi_0)} \Phi_0'(r) A_0'(r) &= 0 \end{aligned} \right. &\Rightarrow A_0'(r) &= \frac{Q}{r^2 f(\Phi_0)} \\ &\Rightarrow \Phi_0''(r) + \frac{2}{r} \Phi_0'(r) + \frac{Q^2}{2r^4} \frac{f'(\Phi_0)}{f(\Phi)^2} &= 0 \end{aligned}$$

Scalarizaion on charged sphere

- Boundary conditions

- Asymptotic region : $\Phi_0 \rightarrow 0 \quad (r \rightarrow \infty)$
- On the charged sphere
 - Dirichlet B.C. (D): $\Phi_0|_{r_s} = 0$
 - Neumann B.C.(N) : $\partial_r (r^{-1}\Phi_0)|_{r_s} = 0$
 - Radiative B.C.(R) : $(\partial_r \Phi_\omega + i\omega \Phi_\omega)|_{r_s} = 0$

$$\longrightarrow \Phi_0|_{r_s} = \Phi_{\min}$$

- Coupling function

- inverse quartic model

$$f(\Phi)^{-1} = 1 - a\Phi^2 + \frac{k^2 a^2}{4}\Phi^4$$

- inverse cosine model

$$f(\Phi)^{-1} = \cos(\sqrt{2a}\Phi)$$

We can solve these models analytically.

Scalarizaion on charged sphere

- Scalarized solution with $\Phi(r \rightarrow \infty) \rightarrow 0$

- inverse quartic model

$$\bar{\Phi}(r) = \sqrt{\frac{ak^2}{2}} \Phi_0 = \sqrt{1 - \sqrt{1 - C_0}} \text{sn} \left(\frac{Q}{\sqrt{2}r} \sqrt{1 + \sqrt{1 - C_0}}, \frac{1 - \sqrt{1 - C_0}}{1 + \sqrt{1 - C_0}} \right)$$

Jacobi elliptic sine function

- inverse cosine model


$$\bar{\Phi} = \sqrt{\frac{a}{2}} \Phi_0 = \text{am} \left(\sqrt{\frac{C_1}{2}} \frac{Q}{r}, \frac{2}{C_1} \right)$$

Jacobi amplitude

- Constant C_0, C_1 are determined from boundary condition on the sphere.
- We can check the stability of each solutions with each B.C.


Scalarizaion on charged sphere

- (Un)stable region of the scalarized solutions.
 - Radiative boundary condition



Inverse quartic

Scalarized solution is stable.




Inverse cosine

Scalarized solution with $\Phi(r_s) = \frac{2n+1}{2}\pi$ is stable.

Scalarizaion on charged sphere

- (Un)stable region of the scalarized solutions.
 - Dirichlet Boundary condition




Inverse quartic

Inverse cosine

Nodeless scalarized solution is stable.

Scalarizaion on charged sphere

- (Un)stable region of the scalarized solutions.
 - Neumann Boundary condition



Inverse polynomial

Inverse cosine

Nodeless scalarized solution is stable.

Time evolution

- We can solve evolution equation of this system.
- Time evolution

$$-\ddot{\Phi} + \Phi'' + \frac{2}{r}\Phi' = -\frac{Q^2}{r^4} \frac{f'(\Phi)}{f(\Phi)^2}$$

- Initial data


$$\Phi = Ae^{-\left(\frac{r-r_0}{w}\right)^2} \quad \dot{\Phi} = 0 \quad (t = 0)$$

- Boundary condition

- Infinity : out-going boundary condition
- On the sphere
 - Dirichlet B.C. : $\Phi|_{r_s} = 0$
 - Neumann B.C. : $\partial_r (r^{-1}\Phi)|_{r_s} = 0$
 - Radiative B.C. : $(\partial_r \Phi_\omega + i\omega \Phi_\omega)|_{r_s} = 0$


Time evolution

- Inverse quartic, Dirichlet B.C.



Time evolution



- Inverse quartic, Dirichlet B.C.
 - Initial parameters: $(A, w, r_0) = (0.1\Phi_{\min}, 4r_s, 8r_s)$
 - charge : $\bar{Q} = Q\sqrt{a}/r_s = \sqrt{2}$



- In this parameter, the charged sphere does not scalarize.

Time evolution


- Inverse quartic, Dirichlet B.C.
 - Initial parameters: $(A, w, r_0) = (0.1\Phi_{\min}, 4r_s, 8r_s)$
 - charge : $\bar{Q} = Q\sqrt{a}/r_s = 4\sqrt{2}$



- The scalar field propagates speed of light, and the charged sphere scalarize as expected.



Time evolution

- Inverse quartic, Radiative B.C.



Time evolution



- Inverse quartic, Radiative B.C.
 - Initial parameters: $(A, w) = (0.01\Phi_{\min}, 4r_s)$
 - charge : $\bar{Q} = Q\sqrt{a}/r_s = 0.1\sqrt{2}$



- Scalar field propagates with speed of light, and reaches the expected scalarized solution, directly.

Time evolution


- Inverse quartic, Radiative B.C.
 - Initial parameters: $(A, w) = (0.07\Phi_{\min}, 4r_s)$
 - charge : $\bar{Q} = Q\sqrt{a}/r_s = 0.1\sqrt{2}$



- Scalar field reaches the expected scalarized solution after many oscillations.

Time evolution

- Inverse cosine, Radiative B.C.




Time evolution

- Inverse cosine, Radiative B.C.

- charge : $\bar{Q} = Q\sqrt{a}/r_s = 4\sqrt{2}$


$$(A\sqrt{a/2}, w/r_s) = (3.0, 8.0)$$

$$r\Phi(t, r)$$



$$(A\sqrt{a/2}, w/r_s) = (3.5, 8.0)$$

$$r\Phi(t, r)$$



- The final scalarized solution depends on the initial data.

Outline of my talk

- 1. Introduction**
- 2. No-hair theorem for ST theory**
- 3. Spontaneous scalarization of BH**
 - 1. Scalar-Gauss-Bonnet theory**
 - 2. Other models**
 - 3. Dynamical behavior of the scalarizaion**
- 4. Summary**

Summary

- Spontaneous scalarization is one of the mechanism to produce a scalar hair in extreme situation.
- Spontaneous scalarization in Scalar-Gauss-Bonnet gravity

$$S = \int d^4x \sqrt{-g} \left(R - \frac{1}{2} \nabla_\mu \Phi \nabla^\mu \Phi + f(\Phi) \mathcal{G}_{\text{GB}} \right)$$

$$\text{quadratic : } f(\Phi) = \frac{\eta \Phi^2}{8} \quad \text{quartic : } f(\Phi) = \frac{\eta}{8} (\Phi^2 + \alpha \Phi^4)$$

- Scalarized BHs with quadratic coupling are unstable.
- Non-linear term in quartic coupling can stabilize the scalarized BHs.
- We showed time evolution of the scalarization using the toy model.