

Confluent hypergeometric kernel determinant on multiple large intervals

Taiyang Xu, Lun Zhang and Zhengyang Zhao

School of Mathematical Sciences, Fudan University
arXiv:2508.10463

15/10/2025
©BMS RMT Seminar

1. Introduction
2. Main results
3. About the proofs

Introduction

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2.$$

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues with FH singularity for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} |\lambda_k|^{2\alpha} \chi_\beta(\lambda_k) \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2,$$

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues with FH singularity for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} |\lambda_k|^{2\alpha} \chi_\beta(\lambda_k) \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2,$$

where

- \mathcal{Z}_n : normalization constant.

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues with FH singularity for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} |\lambda_k|^{2\alpha} \chi_\beta(\lambda_k) \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2,$$

where

- \mathcal{Z}_n : normalization constant.
- $\chi_\beta(\lambda)$ has the jump-type singularity with $\beta \in \mathbb{R}$:

$$\chi_\beta(\lambda) := \begin{cases} e^{-i\pi\beta}, & \lambda \geq 0, \\ e^{i\pi\beta}, & \lambda < 0. \end{cases}$$

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues with FH singularity for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} |\lambda_k|^{2\alpha} \chi_\beta(\lambda_k) \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2,$$

where

- \mathcal{Z}_n : normalization constant.
- $\chi_\beta(\lambda)$ has the jump-type singularity with $\beta \in \mathbb{R}$:

$$\chi_\beta(\lambda) := \begin{cases} e^{-i\pi\beta}, & \lambda \geq 0, \\ e^{i\pi\beta}, & \lambda < 0. \end{cases}$$

- $|\lambda|^{2\alpha}$: root-type singularity, $\alpha > -1/2$.

Unitary ensembles with Fisher–Hartwig (FH) singularity

The joint probability density function (jpdf) of eigenvalues with FH singularity for GUE:

$$f_n(\lambda_1, \dots, \lambda_n) = \frac{1}{\mathcal{Z}_n} \prod_{k=1}^n e^{-\lambda_k^2} |\lambda_k|^{2\alpha} \chi_\beta(\lambda_k) \prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)^2,$$

where

- \mathcal{Z}_n : normalization constant.
- $\chi_\beta(\lambda)$ has the jump-type singularity with $\beta \in \mathbb{R}$:

$$\chi_\beta(\lambda) := \begin{cases} e^{-i\pi\beta}, & \lambda \geq 0, \\ e^{i\pi\beta}, & \lambda < 0. \end{cases}$$

- $|\lambda|^{2\alpha}$: root-type singularity, $\alpha > -1/2$.
- jump-type singularity & root-type singularity \rightsquigarrow **Fisher–Hartwig** singularity (at $z = 0$).

Unitary ensembles with Fisher–Hartwig (FH) singularity

The jpdf of Dyson circular unitary ensemble with FH singularity:

$$f_n(\theta_1, \dots, \theta_n) = \frac{1}{\mathcal{Z}'_n} \prod_{k=1}^n |1 - e^{i\theta_k}|^{2\alpha} e^{i\beta(\theta_k - \pi)} \prod_{1 \leq k < j \leq n} |e^{i\theta_k} - e^{i\theta_j}|^2,$$

where

- \mathcal{Z}'_n : normalization constant.
- $\theta_k \in [0, 2\pi)$, $k = 1, \dots, n$.
- FH singularity at $z = 1$, (i.e., $\theta = 0$).

The correlation kernel

The above eigenvalues form a **determinant point process** (DPP), which is characterized by a correlation kernel $K_n(x, y; \alpha, \beta)$, that is,

$$\rho(\lambda_1, \dots, \lambda_n) = \det [K_n(\lambda_i, \lambda_j; \alpha, \beta)]_{i,j=1}^n.$$

The correlation kernel

The above eigenvalues form a **determinant point process** (DPP), which is characterized by a correlation kernel $K_n(x, y; \alpha, \beta)$, that is,

$$\rho(\lambda_1, \dots, \lambda_n) = \det [K_n(\lambda_i, \lambda_j; \alpha, \beta)]_{i,j=1}^n.$$

- GUE: $K_n(x, y; \alpha, \beta) = \sqrt{w(x)}\sqrt{w(y)} \sum_{j=0}^{n-1} p_j(x)p_j(y)$.
 $p_j(x)$ are orthonormal polynomials with the weight function $w(x)$ over \mathbb{R} , i.e.,

$$\int_{\mathbb{R}} p_i(x)p_j(x)w(x) \, dx = \delta_{ij}, \quad w(x) = e^{-x^2}|x|^{2\alpha}\chi_{\beta}(x).$$

The correlation kernel

The above eigenvalues form a **determinant point process** (DPP), which is characterized by a correlation kernel $K_n(x, y; \alpha, \beta)$, that is,

$$\rho(\lambda_1, \dots, \lambda_n) = \det [K_n(\lambda_i, \lambda_j; \alpha, \beta)]_{i,j=1}^n.$$

- GUE: $K_n(x, y; \alpha, \beta) = \sqrt{w(x)}\sqrt{w(y)} \sum_{j=0}^{n-1} p_j(x)p_j(y)$.
 $p_j(x)$ are orthonormal polynomials with the weight function $w(x)$ over \mathbb{R} , i.e.,

$$\int_{\mathbb{R}} p_i(x)p_j(x)w(x) dx = \delta_{ij}, \quad w(x) = e^{-x^2}|x|^{2\alpha}\chi_{\beta}(x).$$

- CUE: $K_n(e^{i\theta}, e^{i\phi}; \alpha, \beta) = \sqrt{w(e^{i\theta})}\sqrt{w(e^{i\phi})} \sum_{j=0}^{n-1} p_j(e^{i\theta})\overline{p_j(e^{i\phi})}$.

$$\int_0^{2\pi} p_i(e^{i\theta})\overline{p_j(e^{i\theta})}w(e^{i\theta}) d\theta = \delta_{jk}, \quad w(z) = z^n|z-1|^{2\alpha}z^{\beta}e^{-i\pi\beta}, \quad z = e^{i\theta}.$$

Large- n limiting kernel

In the bulk (near the FH singular point), the correlation kernel $K_n(x, y; \alpha, \beta)$ converges to **confluent hypergeometric kernel** $K^{(\alpha, \beta)}(x, y)$ under a suitable scaling.

Large- n limiting kernel

In the bulk (near the FH singular point), the correlation kernel $K_n(x, y; \alpha, \beta)$ converges to **confluent hypergeometric kernel** $K^{(\alpha, \beta)}(x, y)$ under a suitable scaling.

- Unitary random matrix ensembles generated by a concrete weight function on the unit circle.

[Deift-Krasovsky-Vasilevska, '11]

Large- n limiting kernel

In the bulk (near the FH singular point), the correlation kernel $K_n(x, y; \alpha, \beta)$ converges to **confluent hypergeometric kernel** $K^{(\alpha, \beta)}(x, y)$ under a suitable scaling.

- Unitary random matrix ensembles generated by a concrete weight function on the unit circle.

[Deift-Krasovsky-Vasilevska, '11]

- it describes the local statistics of eigenvalues in the bulk of the spectrum near a FH singular point for a broad class of unitary ensemble of random matrices.

The confluent hypergeometric kernel

The confluent hypergeometric (CH) kernel with two parameters $\alpha > -1/2$ and $\beta \in i\mathbb{R}$ is defined by

$$K^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi i} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} \frac{A(x)B(y) - A(y)B(x)}{x - y},$$

The confluent hypergeometric kernel

The confluent hypergeometric (CH) kernel with two parameters $\alpha > -1/2$ and $\beta \in \mathbb{R}$ is defined by

$$K^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi i} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} \frac{A(x)B(y) - A(y)B(x)}{x - y},$$

- $\Gamma(z)$ denotes the usual Gamma function. $A(x)$ and $B(x)$ are defined by

$$A(x) := \chi_\beta(x)^{\frac{1}{2}} |2x|^\alpha e^{-ix} \phi(1 + \alpha + \beta, 1 + 2\alpha; 2ix),$$

$$B(x) := \chi_\beta(x)^{\frac{1}{2}} |2x|^\alpha e^{ix} \phi(1 + \alpha - \beta, 1 + 2\alpha; -2ix).$$

The confluent hypergeometric kernel

The confluent hypergeometric (CH) kernel with two parameters $\alpha > -1/2$ and $\beta \in \mathbb{R}$ is defined by

$$K^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi i} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} \frac{A(x)B(y) - A(y)B(x)}{x - y},$$

- $\Gamma(z)$ denotes the usual Gamma function. $A(x)$ and $B(x)$ are defined by

$$A(x) := \chi_\beta(x)^{\frac{1}{2}} |2x|^\alpha e^{-ix} \phi(1 + \alpha + \beta, 1 + 2\alpha; 2ix),$$

$$B(x) := \chi_\beta(x)^{\frac{1}{2}} |2x|^\alpha e^{ix} \phi(1 + \alpha - \beta, 1 + 2\alpha; -2ix).$$

- The confluent hypergeometric function is defined by

$$\phi(a, b; z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} \frac{z^n}{n!}, \quad b \neq 0, -1, -2, \dots,$$

where $(z)_n := z(z+1) \cdots (z+n-1) = \frac{\Gamma(z+n)}{\Gamma(z)}$ is the Pochhammer symbol.

Universal features of the confluent hypergeometric kernel

The confluent hypergeometric kernel arises in several different, but related areas

Universal features of the confluent hypergeometric kernel

The confluent hypergeometric kernel arises in several different, but related areas

- Infinite random matrices and Hua-Pickrell measure.

[Borodin-Olshanski, '01]

- Representation theory.

[Borodin-Deift, '01]

- Circular unitary ensemble.

[Deift-Krasovsky-Vasilevska, '11]

Relations to other kernels

Confluent hypergeometric kernel can reduce to other kernels

- If $\beta = 0$, $\alpha \neq 0$, following from the relation

$$\phi(\alpha, 2\alpha; 2ix) = \Gamma\left(\alpha + \frac{1}{2}\right) e^{ix} \left(\frac{x}{2}\right)^{-\alpha + \frac{1}{2}} J_{\alpha - \frac{1}{2}}(x),$$

Relations to other kernels

Confluent hypergeometric kernel can reduce to other kernels

- If $\beta = 0$, $\alpha \neq 0$, following from the relation

$$\phi(\alpha, 2\alpha; 2ix) = \Gamma\left(\alpha + \frac{1}{2}\right) e^{ix} \left(\frac{x}{2}\right)^{-\alpha + \frac{1}{2}} J_{\alpha - \frac{1}{2}}(x),$$

then we have type-I Bessel kernel

$$K^{(\alpha, 0)}(x, y) \equiv K^{(\text{Bessel1})}(x, y) = \left(\frac{|x|}{x}\right)^{\alpha} \left(\frac{|y|}{y}\right)^{\alpha} \frac{\sqrt{xy}}{2} \frac{J_{\alpha + \frac{1}{2}}(x) J_{\alpha - \frac{1}{2}}(y) - J_{\alpha + \frac{1}{2}}(y) J_{\alpha - \frac{1}{2}}(x)}{x - y}.$$

[Akemann-Damgaard-Magnea-Nishigaki, '97] [Kuijlaars-Vanlessen, '03]

Relations to other kernels

Confluent hypergeometric kernel can reduce to other kernels

- If $\beta = 0$, $\alpha \neq 0$, following from the relation

$$\phi(\alpha, 2\alpha; 2ix) = \Gamma\left(\alpha + \frac{1}{2}\right) e^{ix} \left(\frac{x}{2}\right)^{-\alpha + \frac{1}{2}} J_{\alpha - \frac{1}{2}}(x),$$

then we have type-I Bessel kernel

$$K^{(\alpha, 0)}(x, y) \equiv K^{(\text{Bessel1})}(x, y) = \left(\frac{|x|}{x}\right)^{\alpha} \left(\frac{|y|}{y}\right)^{\alpha} \frac{\sqrt{xy} J_{\alpha + \frac{1}{2}}(x) J_{\alpha - \frac{1}{2}}(y) - J_{\alpha + \frac{1}{2}}(y) J_{\alpha - \frac{1}{2}}(x)}{2(x - y)}.$$

[Akemann-Damgaard-Magnea-Nishigaki, '97] [Kuijlaars-Vanlessen, '03]

An episode: Type-II Bessel kernel:

$$K^{(\text{Bessel2})}(x, y) := \frac{J_{\alpha}(\sqrt{x})\sqrt{y}J'_{\alpha}(\sqrt{y}) - \sqrt{x}J'_{\alpha}(\sqrt{x})J_{\alpha}(\sqrt{y})}{2(x - y)}.$$

[Forrester, '93]

- If $\alpha = 0$, $\beta \neq 0$, we have a degenerated confluent hypergeometric kernel.

[Tibboel, '10] [Moreno- Martínez-Finkelshtein -Sousa, '11]

- If $\alpha = 0$, $\beta \neq 0$, we have a degenerated confluent hypergeometric kernel.

[Tibboel, '10] [Moreno- Martínez-Finkelshtein -Sousa, '11]

- If $\alpha = \beta = 0$, we obtain the sine kernel

$$K^{(0,0)}(x, y) \equiv K^{(\text{sine})}(x, y) = \frac{\sin(x - y)}{\pi(x - y)}.$$

- The **Gap probability**: finding no eigenvalues on a specific interval Σ .

- The **Gap probability**: finding no eigenvalues on a specific interval Σ .
- **Fredholm determinant representation**:

$$\mathbb{P}(\text{no points lie on } \Sigma) = \mathcal{F}(\Sigma) := \det(1 - \mathcal{K}|_{\Sigma}),$$

where \mathcal{K} is an integration operator acting on $L^2(\Sigma)$ with integrable kernel $K(x, y)$.

Gap probability for DPP

- The **Gap probability**: finding no eigenvalues on a specific interval Σ .
- **Fredholm determinant representation**:

$$\mathbb{P}(\text{no points lie on } \Sigma) = \mathcal{F}(\Sigma) := \det(1 - \mathcal{K}|_{\Sigma}),$$

where \mathcal{K} is an integration operator acting on $L^2(\Sigma)$ with integrable kernel $K(x, y)$.

- Our interest: **large gap asymptotics**

$$\mathcal{F}(s\Sigma) := \det(1 - \mathcal{K}|_{s\Sigma}) = ?, \quad \text{as } s \rightarrow +\infty,$$

especially the case that Σ is a **union of disjoint intervals**.

History: sine kernel determinant

$$\log \det (1 - \mathcal{K}^{(\text{sine})}|_{s\Sigma}) = C_1 s^2 + C_2 \log s + C_3 \log \theta(V(s)) + C_4 + \mathcal{O}(s^{-1}), \quad s \rightarrow +\infty.$$

¹Combined the earlier work [\[Widom '71\]](#) on Toeplitz determinant.

History: sine kernel determinant

$$\log \det (1 - \mathcal{K}^{(\text{sine})}|_{s\Sigma}) = C_1 s^2 + C_2 \log s + C_3 \log \theta(V(s)) + C_4 + \mathcal{O}(s^{-1}), \quad s \rightarrow +\infty.$$

Σ	C_1	C_2	C_3	C_4
$(-1, 1)$	<div>[Dyson, '62] [Cloizeaux-Mehta, '73] [Widom, '94] $C_1 = -1/2$</div>	<div>[Deift-Its-Zhou, '97] $C_2 = -1/4, C_3 = 0$</div>		<div>[Dyson, '76]¹ [Krasovsky, '04] [Ehrhardt, '04] [Deift-Its-Krasovsky, '07] $C_4 = (\log 2)/12 + 3\zeta'(-1)$</div>
$(-1, v_1) \cup (v_2, 1)$	<div>[Fahs-Krasovsky, '22]</div>			
$\cup_{j=0}^n(a_j, b_j)$	<div>[Deift-Its-Zhou, '97] C_2: integral expression</div>			<div>?</div>

Table: Large gap asymptotics for the sine kernel determinant.

¹Combined the earlier work [\[Widom '71\]](#) on Toeplitz determinant.

History: Airy kernel determinant

$$\log \det (1 - \mathcal{K}^{(\text{Ai})}|_{s\Sigma}) = C_1 s^3 + C_2 \log s + C_3 \log \theta(V(s)) + C_4 + e(s), \quad s \rightarrow +\infty.$$

History: Airy kernel determinant

$$\log \det (1 - \mathcal{K}^{(\text{Ai})}|_{s\Sigma}) = C_1 s^3 + C_2 \log s + C_3 \log \theta(V(s)) + C_4 + e(s), \quad s \rightarrow +\infty.$$

Σ	C_1	C_2	C_3	$e(s)$	C_4
$(-1, +\infty)$	[Tracy-Widom, '94] $C_1 = -1/12, C_2 = -1/8,$ $C_3 = 0, e(s) = \mathcal{O}(s^{-3/2})$				[Deift-Its-Krasovsky, '08] [Baik-Buckingham-DiFranco, '08] $C_4 = (\log 2)/24 + \zeta'(-1)$
$(x_2, x_1) \cup (x_0, +\infty)$ $x_2 < x_1 < x_0 < 0$	[Blackstone-Charlier-Lenells, '22] $\rightsquigarrow e(s) = \mathcal{O}(s^{-1})$ [Krasovsky-Maroudas, '24]				[Krasovsky-Maroudas, '24]
$\bigcup_{j=0}^g (x_{2j}, x_{2j-1})$ $x_{-1} = +\infty$?				
(x_2, x_1)	[Blackstone-Charlier-Lenells, '21] $e(s) = \mathcal{O}(s^{-3/2})$?
$\bigcup_{j=1}^g (x_{2j}, x_{2j-1})$?				

Table: Large gap asymptotics for the Airy kernel determinant.

History: type-II Bessel kernel determinant

$$\log \det(1 - \mathcal{K}^{(\text{Bes2})})|_{s\Sigma} = C_1 s + C_2 s^{1/2} + C_3 \log s + C_4 \log \theta(V(s)) + C_5 + \mathcal{O}(s^{-1/2}), \quad s \rightarrow +\infty.$$

History: type-II Bessel kernel determinant

$$\log \det(1 - \mathcal{K}^{(\text{Bes2})})|_{s\Sigma} = C_1 s + C_2 s^{1/2} + C_3 \log s + C_4 \log \theta(V(s)) + C_5 + \mathcal{O}(s^{-1/2}), \quad s \rightarrow +\infty.$$

Σ	C_1	C_2	C_3	C_4	C_5
$(0, x_1)$	[Tracy-Widom, '94] $C_1 = -x_1/4, C_2 = \alpha\sqrt{x_1},$ $C_3 = -\alpha^2/4, C_4 = 0$				$C_5 = G(1 + \alpha)(2\pi)^{-\alpha/2} - (\alpha^2 \log x_1)/4$ [Ehrhardt, '10] $\rightsquigarrow \alpha \in (-1, 1)$ [Deift-Krasovsky-Vasilevska, '11] $\rightsquigarrow \alpha \in (-1, +\infty)$
$\bigcup_{j=0}^{2g} (x_j, x_{j+1})$ $x_0 = 0$	[Blackstone-Charlier-Lenells, 21'] C_3 : exact value (under the ergodic condition)				?

Table: Large gap asymptotics for the type-II Bessel kernel determinant.

History: Confluent hypergeometric kernel determinant

As $s \rightarrow +\infty$,

$$\log \det(1 - \mathcal{K}^{(\alpha, \beta)}|_{s\Sigma}) = C_1 s^2 + C_2 s + (\beta^2 - \alpha^2 + C_3) \log s + C_4 \log \theta(V(s)) + C_5 + \mathcal{O}(s^{-1}).$$

History: Confluent hypergeometric kernel determinant

As $s \rightarrow +\infty$,

$$\log \det(1 - \mathcal{K}^{(\alpha, \beta)}|_{s\Sigma}) = C_1 s^2 + C_2 s + (\beta^2 - \alpha^2 + C_3) \log s + C_4 \log \theta(V(s)) + C_5 + \mathcal{O}(s^{-1}).$$

Σ	C_1	C_2	C_3	C_4	C_5
$0 \in (-1, 1)$	[Deift-Krasovsky-Vasilevska, '11] [Xu-Zhao, '20] $C_1 = -1/2, C_2 = 2\alpha, C_3 = -1/4, C_4 = 0$ $C_5 = \log\left(\frac{\sqrt{\pi} G^2(1/2) G(1+2\alpha)}{2^{2\alpha^2} G(1+\alpha+\beta) G(1+\alpha-\beta)}\right)$				
$\cup_{j=0}^n (a_j, b_j)$ $0 \in (a_m, b_m)$ for some $m: 1 \leq m \leq n$?				

Table: Large gap asymptotics for the confluent hypergeometric kernel determinant.

Today's topic

Aim: establish large gap asymptotics for the confluent hypergeometric kernel determinant $\det(1 - \mathcal{K}^{(\alpha, \beta)})$ on multiple large intervals.

$$\overline{a_0} \quad \overline{b_0} \quad \overline{a_1} \quad \overline{b_1} \quad \cdots \quad \overline{a_m} \quad \color{red}{0} \quad \overline{b_m} \quad \cdots \quad \overline{a_n} \quad \overline{b_n}$$

Today's topic

Aim: establish large gap asymptotics for the confluent hypergeometric kernel determinant $\det(1 - \mathcal{K}^{(\alpha, \beta)})$ on multiple large intervals.

$$\overline{a_0 \quad b_0} \quad \overline{a_1 \quad b_1} \quad \cdots \quad \overline{a_m \quad 0 \quad b_m} \quad \cdots \quad \overline{a_n \quad b_n}$$

Σ	C_1	C_2	C_3	C_4	C_5
$(-1, 1)$ $0 \in (-1, 1)$	[Deift-Krasovsky-Vasilevska, '11] [Xu-Zhao, '20] $C_1 = -1/2, C_2 = 2\alpha, C_3 = -1/4, C_4 = 0$ $C_5 = \log\left(\frac{\sqrt{\pi} G^2(1/2) G(1+2\alpha)}{2^{2\alpha^2} G(1+\alpha+\beta) G(1+\alpha-\beta)}\right)$				
$\cup_{j=0}^n (a_j, b_j)$ $0 \in (a_m, b_m)$ for some $m: 1 \leq m \leq n$	[Xu-Zhang-Zhao, '25]				?

Table: Large gap asymptotics for the confluent hypergeometric kernel determinant.

Main results

The Riemann surface


We will encounter a hyperelliptic Riemann surface \mathcal{W} associated to the algebraic equation

$$\sqrt{\mathcal{R}(z)} := \sqrt{\prod_{j=0}^n (z - a_j)(z - b_j)}.$$

The Riemann surface

We will encounter a hyperelliptic Riemann surface \mathcal{W} associated to the algebraic equation

$$\sqrt{\mathcal{R}(z)} := \sqrt{\prod_{j=0}^n (z - a_j)(z - b_j)}.$$



- $\sqrt{\mathcal{R}(z)} \sim \pm z^{n+1}$, as $z \rightarrow \infty$ on the first (second) sheet.
- Canonical homology basis $\{A_j, B_j\}$.

Figure: The canonical homology basis $\{A_j, B_j\}_{j=1}^n$ for the Riemann surface \mathcal{W} .

Preliminaries – The \mathbb{A} -matrix

The \mathbb{A} -matrix:

$$\mathbb{A} := (a_{k,l})_{0 \leq k \leq n, 0 \leq l \leq n} = \begin{pmatrix} a_{0,0} & a_{0,1} & \cdots & a_{0,n} \\ a_{1,0} & a_{1,1} & \cdots & a_{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,0} & a_{n,1} & \cdots & a_{n,n} \end{pmatrix},$$

$$\tilde{\mathbb{A}} := (a_{k,l})_{\substack{l=0,\dots,n-1 \\ k=1,\dots,n}} = \begin{pmatrix} a_{1,0} & a_{1,1} & \cdots & a_{1,n-1} \\ a_{2,0} & a_{2,1} & \cdots & a_{2,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,0} & a_{n,1} & \cdots & a_{n,n-1} \end{pmatrix},$$

$$\vec{a} = (a_{0,n+1}, a_{1,n+1}, \dots, a_{n,n+1})^T,$$

$$\text{where } a_{k,l} := \oint_{A_k} \frac{z^l}{\sqrt{\mathcal{R}(z)}} dz = 2i(-1)^{n-k+1} \int_{a_k}^{b_k} \frac{z^l}{|\mathcal{R}(z)|^{\frac{1}{2}}} dz.$$

Preliminaries – The \mathbb{A} -matrix

The \mathbb{A} -matrix:

$$\mathbb{A} := (a_{k,l})_{0 \leq k \leq n, 0 \leq l \leq n} = \begin{pmatrix} a_{0,0} & a_{0,1} & \cdots & a_{0,n} \\ a_{1,0} & a_{1,1} & \cdots & a_{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,0} & a_{n,1} & \cdots & a_{n,n} \end{pmatrix},$$

$$\tilde{\mathbb{A}} := (a_{k,l})_{\substack{l=0,\dots,n-1 \\ k=1,\dots,n}} = \begin{pmatrix} a_{1,0} & a_{1,1} & \cdots & a_{1,n-1} \\ a_{2,0} & a_{2,1} & \cdots & a_{2,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,0} & a_{n,1} & \cdots & a_{n,n-1} \end{pmatrix},$$

$$\vec{a} = (a_{0,n+1}, a_{1,n+1}, \dots, a_{n,n+1})^T,$$

where $a_{k,l} := \oint_{A_k} \frac{z^l}{\sqrt{\mathcal{R}(z)}} dz = 2i(-1)^{n-k+1} \int_{a_k}^{b_k} \frac{z^l}{|\mathcal{R}(z)|^{\frac{1}{2}}} dz$.

$\rightsquigarrow \mathbb{A}$ and $\tilde{\mathbb{A}}$ are invertible.

Preliminaries – The basis of one-form

Introduce the basis of holomorphic one-forms:

$$\vec{\omega} := (\omega_1, \omega_2, \dots, \omega_n) = \frac{dz}{\sqrt{\mathcal{R}(z)}} (1, z, \dots, z^{n-1}) \tilde{\mathbb{A}}^{-1},$$

such that

$$\oint_{A_k} \omega_j = \delta_{jk}, \quad j, k = 1, \dots, n.$$

Meanwhile, we have the Riemann matrix of B_j periods:

$$\tau := (\tau_{ij})_{i,j=1}^n = \left(\oint_{B_j} \omega_i \right)_{i,j=1}^n.$$

Preliminaries – The basis of one-form

Introduce the basis of holomorphic one-forms:

$$\vec{\omega} := (\omega_1, \omega_2, \dots, \omega_n) = \frac{dz}{\sqrt{\mathcal{R}(z)}} (1, z, \dots, z^{n-1}) \tilde{\mathbb{A}}^{-1},$$

such that

$$\oint_{A_k} \omega_j = \delta_{jk}, \quad j, k = 1, \dots, n.$$

Meanwhile, we have the Riemann matrix of B_j periods:

$$\tau := (\tau_{ij})_{i,j=1}^n = \left(\oint_{B_j} \omega_i \right)_{i,j=1}^n.$$

$\rightsquigarrow \tau$ is symmetric and has a positively definite imaginary part, i.e, $-i\tau$ is positive definite.

[Farkas-Kra, '92, *Riemann Surfaces* 2nd ed]

Preliminaries – Multi-dimensional Riemann- θ function and Abel's map

The multi-dimensional Riemann- θ function is defined by

$$\theta(\vec{z}) = \sum_{\vec{m} \in \mathbb{Z}^n} e^{2\pi i \vec{m}^T \vec{z} + i\pi \vec{m}^T \tau \vec{m}}, \quad \vec{z} = (z_1, \dots, z_n)^T \in \mathbb{C}^n \text{ modulo } \mathbb{Z}^n.$$

- Converging absolutely and uniformly on compact sets of the \mathbb{C}^n .
- Even ($\theta(\vec{z}) = \theta(-\vec{z})$), entire function for $\vec{z} \in \mathbb{C}^n$.
- Periodic properties: $\theta(\vec{z} + \vec{e}_j) = \theta(\vec{z})$ and $\theta(\vec{z} \pm \vec{\tau}_j) = e^{\mp 2\pi i z_j - \pi i \tau_{jj}} \theta(\vec{z})$, where $\vec{e}_j = (0, \dots, 1, \dots, 0)^T$ with 1 in the j -th position and $\vec{\tau} := \tau \vec{e}_j$.
- Vanishing at each odd half-period: $\theta(\vec{z}) = 0$ if $\vec{z} = \frac{\vec{m}}{2} + \frac{\tau \vec{n}}{2}$ with $\vec{m}, \vec{n} \in \mathbb{Z}^n$ and $\vec{m}^T \vec{n}$ is odd.

Preliminaries – Multi-dimensional Riemann- θ function and Abel's map

The multi-dimensional Riemann- θ function is defined by

$$\theta(\vec{z}) = \sum_{\vec{m} \in \mathbb{Z}^n} e^{2\pi i \vec{m}^T \vec{z} + i\pi \vec{m}^T \tau \vec{m}}, \quad \vec{z} = (z_1, \dots, z_n)^T \in \mathbb{C}^n \text{ modulo } \mathbb{Z}^n.$$

- Converging absolutely and uniformly on compact sets of the \mathbb{C}^n .
- Even ($\theta(\vec{z}) = \theta(-\vec{z})$), entire function for $\vec{z} \in \mathbb{C}^n$.
- Periodic properties: $\theta(\vec{z} + \vec{e}_j) = \theta(\vec{z})$ and $\theta(\vec{z} \pm \vec{\tau}_j) = e^{\mp 2\pi i z_j - \pi i \tau_{jj}} \theta(\vec{z})$, where $\vec{e}_j = (0, \dots, 1, \dots, 0)^T$ with 1 in the j -th position and $\vec{\tau} := \tau \vec{e}_j$.
- Vanishing at each odd half-period: $\theta(\vec{z}) = 0$ if $\vec{z} = \frac{\vec{m}}{2} + \frac{\tau \vec{n}}{2}$ with $\vec{m}, \vec{n} \in \mathbb{Z}^n$ and $\vec{m}^T \vec{n}$ is odd.

Abel's map:

$$\vec{\mathcal{A}}(z) := \int_{a_0}^z \vec{\omega}^T.$$

Preliminaries – The polynomial, linear vector and frequencies

The following polynomial of degree $n + 1$ plays an important role in our analysis:

$$p(z) = z^{n+1} + \sum_{j=0}^n p_j z^j.$$

$$\oint_{A_k} \frac{p(s)}{\sqrt{\mathcal{R}(s)}} ds = 0, \quad k = 0, 1, \dots, n. \implies (p_0, \dots, p_n)^T = -\mathbb{A}^{-1} \vec{a}.$$

Preliminaries – The polynomial, linear vector and frequencies

The following polynomial of degree $n + 1$ plays an important role in our analysis:

$$p(z) = z^{n+1} + \sum_{j=0}^n p_j z^j.$$

$$\oint_{A_k} \frac{p(s)}{\sqrt{\mathcal{R}(s)}} ds = 0, \quad k = 0, 1, \dots, n. \implies (p_0, \dots, p_n)^T = -\mathbb{A}^{-1} \vec{a}.$$

A column vector with linear components:

$$\vec{V}(s) := (V_1(s), \dots, V_n(s))^T, \quad V_j(s) := \frac{s}{2\pi} \Omega_j + \frac{1}{2\pi} \operatorname{Im}(\zeta_j) \in \mathbb{R}.$$

- Frequencies $\Omega_j := 2 \sum_{k=0}^{j-1} (-1)^{n-k} \int_{b_k}^{a_{k+1}} \frac{p(s)}{|\mathcal{R}(s)|^{\frac{1}{2}}} ds > 0$.
- $\operatorname{Im} \zeta_j \in \mathbb{R}$ is dependent of α and β .

Preliminaries – The linear flow on torus

A function $\mathcal{L} : \mathbb{C} \times \mathbb{R}^n / \mathbb{Z}^n \rightarrow \mathbb{C}$ defined by

$$\mathcal{L}(z, \vec{\mu}) = \frac{h(z)}{p(z)} \eta(z, \vec{\mu}).$$

where $h(z) = \prod_{k=0}^n (z - a_k) + \prod_{k=0}^n (z - b_k)$, and $\eta(z, \vec{\mu}) = \frac{\theta(\vec{0})^2 \theta(\vec{\mathcal{A}}(z) + \vec{\mu} + \vec{d}) \theta(\vec{\mathcal{A}}(z) - \vec{\mu} + \vec{d})}{\theta(\vec{\mu})^2 \theta(\vec{\mathcal{A}}(z) + \vec{d})^2}$.

Preliminaries – The linear flow on torus

A function $\mathcal{L} : \mathbb{C} \times \mathbb{R}^n / \mathbb{Z}^n \rightarrow \mathbb{C}$ defined by

$$\mathcal{L}(z, \vec{\mu}) = \frac{h(z)}{p(z)} \eta(z, \vec{\mu}).$$

where $h(z) = \prod_{k=0}^n (z - a_k) + \prod_{k=0}^n (z - b_k)$, and $\eta(z, \vec{\mu}) = \frac{\theta(\vec{0})^2 \theta(\vec{\mathcal{A}}(z) + \vec{\mu} + \vec{d}) \theta(\vec{\mathcal{A}}(z) - \vec{\mu} + \vec{d})}{\theta(\vec{\mu})^2 \theta(\vec{\mathcal{A}}(z) + \vec{d})^2}$.

Consider $\mathcal{L}(p, \vec{V}(t))$ for $p \in \mathcal{I}_e := \{a_j, b_j\}_{j=0}^n$

Preliminaries – The linear flow on torus

A function $\mathcal{L} : \mathbb{C} \times \mathbb{R}^n / \mathbb{Z}^n \rightarrow \mathbb{C}$ defined by

$$\mathcal{L}(z, \vec{\mu}) = \frac{h(z)}{p(z)} \eta(z, \vec{\mu}).$$

where $h(z) = \prod_{k=0}^n (z - a_k) + \prod_{k=0}^n (z - b_k)$, and $\eta(z, \vec{\mu}) = \frac{\theta(\vec{0})^2 \theta(\vec{\mathcal{A}}(z) + \vec{\mu} + \vec{d}) \theta(\vec{\mathcal{A}}(z) - \vec{\mu} + \vec{d})}{\theta(\vec{\mu})^2 \theta(\vec{\mathcal{A}}(z) + \vec{d})^2}$.

Consider $\mathcal{L}(p, \vec{V}(t))$ for $p \in \mathcal{I}_e := \{a_j, b_j\}_{j=0}^n$

- $\mathcal{L}(p, \vec{V}(t))$ is real-valued and analytic.

Preliminaries – The linear flow on torus

A function $\mathcal{L} : \mathbb{C} \times \mathbb{R}^n / \mathbb{Z}^n \rightarrow \mathbb{C}$ defined by

$$\mathcal{L}(z, \vec{\mu}) = \frac{h(z)}{p(z)} \eta(z, \vec{\mu}).$$

where $h(z) = \prod_{k=0}^n (z - a_k) + \prod_{k=0}^n (z - b_k)$, and $\eta(z, \vec{\mu}) = \frac{\theta(\vec{0})^2 \theta(\vec{\mathcal{A}}(z) + \vec{\mu} + \vec{d}) \theta(\vec{\mathcal{A}}(z) - \vec{\mu} + \vec{d})}{\theta(\vec{\mu})^2 \theta(\vec{\mathcal{A}}(z) + \vec{d})^2}$.

Consider $\mathcal{L}(p, \vec{V}(t))$ for $p \in \mathcal{I}_e := \{a_j, b_j\}_{j=0}^n$

- $\mathcal{L}(p, \vec{V}(t))$ is real-valued and analytic.
- $\mathcal{L}(p, \cdot)$ is periodic on $\mathbb{R}^n / \mathbb{Z}^n$.

Preliminaries – The linear flow on torus

A function $\mathcal{L} : \mathbb{C} \times \mathbb{R}^n / \mathbb{Z}^n \rightarrow \mathbb{C}$ defined by


$$\mathcal{L}(z, \vec{\mu}) = \frac{h(z)}{p(z)} \eta(z, \vec{\mu}).$$

where $h(z) = \prod_{k=0}^n (z - a_k) + \prod_{k=0}^n (z - b_k)$, and $\eta(z, \vec{\mu}) = \frac{\theta(\vec{0})^2 \theta(\vec{\mathcal{A}}(z) + \vec{\mu} + \vec{d}) \theta(\vec{\mathcal{A}}(z) - \vec{\mu} + \vec{d})}{\theta(\vec{\mu})^2 \theta(\vec{\mathcal{A}}(z) + \vec{d})^2}$.

Consider $\mathcal{L}(p, \vec{V}(t))$ for $p \in \mathcal{I}_e := \{a_j, b_j\}_{j=0}^n$

- $\mathcal{L}(p, \vec{V}(t))$ is real-valued and analytic.
- $\mathcal{L}(p, \cdot)$ is periodic on $\mathbb{R}^n / \mathbb{Z}^n$.

$\implies \vec{V}(t)$ is a **linear flow (orbit)** on the torus $\mathbb{R}^n / \mathbb{Z}^n$.



Large gap asymptotics: general case

Theorem (X.-Zhang-Zhao, '25)

Let $\mathcal{F}(s\Sigma) := \det(1 - \mathcal{K}^{(\alpha, \beta)}|_{s\Sigma})$ and $\Sigma := \cup_{j=0}^n (a_j, b_j)$ be such that $a_1 < b_1 < \dots < a_m < 0 < b_m < \dots < a_n < b_n$ for some $0 \leq m \leq n$. For $\alpha > -1/2$ and $\beta \in \mathbb{R}$, we have, as $s \rightarrow +\infty$,

$$\begin{aligned} \log \mathcal{F}(s\Sigma) = & -\gamma_0 s^2 - 2i\mathcal{D}_{\infty,1}s + \log \theta \left(\vec{V}(s) \right) + (\beta^2 - \alpha^2) \log s \\ & - \frac{1}{16} \sum_{j=0}^n \int_{\hat{s}}^s \left(\mathcal{L} \left(a_j, \vec{V}(t) \right) + \mathcal{L} \left(b_j, \vec{V}(t) \right) \right) \frac{dt}{t} + \check{C}_1 + \mathcal{O}(s^{-1}), \end{aligned}$$

where

$$\gamma_0 = -\frac{1}{\pi i} \sum_{j=0}^n \int_{a_j}^{b_j} \frac{z p(z)}{\sqrt{\mathcal{R}(z)}_+} dz \in \mathbb{R},$$

$\mathcal{D}_{\infty,1}$ is purely imaginary and depends on the parameters α and β , $\vec{V}(s) \in \mathbb{R}^n$ is defined above, $\hat{s} > 0$ is a sufficiently large number independent of s , $\mathcal{L}(p, \vec{V}(t))$ is real for $p \in \mathcal{I}_e := \{a_j, b_j\}_{j=0}^n$ and \check{C}_1 is an undetermined constant independent of s . Moreover, for $p \in \mathcal{I}_e$, as $s \rightarrow +\infty$, we have

$$\int_{\hat{s}}^s \mathcal{L} \left(p, \vec{V}(t) \right) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + o(\log s), \quad \hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L} \left(p, \vec{V}(t) \right) dt.$$

Remarks on the general case

- If $n = 0$, $\Sigma = (-1, 1)$, we have

$$p(z) = z, \quad \gamma_0 = \frac{1}{2}, \quad \mathcal{D}_{\infty,1} = i\alpha, \quad \mathcal{L}(p, \vec{V}(t)) = \mathcal{L}(z, 0) = 2.$$

Remarks on the general case

- If $n = 0$, $\Sigma = (-1, 1)$, we have

$$p(z) = z, \quad \gamma_0 = \frac{1}{2}, \quad \mathcal{D}_{\infty,1} = i\alpha, \quad \mathcal{L}(p, \vec{V}(t)) = \mathcal{L}(z, 0) = 2.$$

\implies Reduce to the results in [\[Deift-Krasovsky-Vasilevska, '11\]](#) (except the constant term).

Remarks on the general case

- If $n = 0$, $\Sigma = (-1, 1)$, we have

$$p(z) = z, \quad \gamma_0 = \frac{1}{2}, \quad \mathcal{D}_{\infty,1} = i\alpha, \quad \mathcal{L}(p, \vec{V}(t)) = \mathcal{L}(z, 0) = 2.$$

\implies Reduce to the results in [\[Deift-Krasovsky-Vasilevska, '11\]](#) (except the constant term).

- For general $n > 1$, $\beta = 0 \implies$ The large gap asymptotics for the type-I Bessel kernel determinant on multiple intervals.

Remarks on the general case

- If $n = 0$, $\Sigma = (-1, 1)$, we have

$$p(z) = z, \quad \gamma_0 = \frac{1}{2}, \quad \mathcal{D}_{\infty,1} = i\alpha, \quad \mathcal{L}(p, \vec{V}(t)) = \mathcal{L}(z, 0) = 2.$$

\implies Reduce to the results in [\[Deift-Krasovsky-Vasilevska, '11\]](#) (except the constant term).

- For general $n > 1$, $\beta = 0 \implies$ The large gap asymptotics for the type-I Bessel kernel determinant on multiple intervals.
- For general $n > 1$, if $\alpha = \beta = 0 \implies$ The results is consistent with the Eq. (1.34) in [\[Deift-Its-Zhou, '97\]](#).

Large gap asymptotics: “good Diophantine property” case

Question: Could we improve the asymptotics

$$\int_{\hat{s}}^s \mathcal{L} \left(p, \vec{V}(t) \right) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + o(\log s), \quad \hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L} \left(p, \vec{V}(t) \right) dt.$$

as $s \rightarrow +\infty$?

Large gap asymptotics: “good Diophantine property” case

Question: Could we improve the asymptotics

$$\int_{\hat{s}}^s \mathcal{L} \left(p, \vec{V}(t) \right) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + o(\log s), \quad \hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L} \left(p, \vec{V}(t) \right) dt.$$

as $s \rightarrow +\infty$?

Aim 1: $o(\log s) \rightsquigarrow \mathcal{O}(s^{-1})$

Large gap asymptotics: “good Diophantine property” case

Question: Could we improve the asymptotics

$$\int_{\hat{s}}^s \mathcal{L}(p, \vec{V}(t)) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + o(\log s), \quad \hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) dt.$$

as $s \rightarrow +\infty$?

Aim 1: $o(\log s) \rightsquigarrow \mathcal{O}(s^{-1})$

Definition (good Diophantine property)

The linear flow

$$(0, +\infty) \ni s \mapsto (V_1(s) \bmod 1, V_2(s) \bmod 1, \dots, V_n(s) \bmod 1), \quad V_j(s) = \frac{s}{2\pi} \Omega_j + \frac{1}{2\pi} \operatorname{Im} \zeta_j,$$

has “**good Diophantine properties**” if there exist $\delta_1, \delta_2 > 0$ such that

$$|\vec{m}^T \vec{\Omega}| \geq \delta_1 \|\vec{m}\|_2^{-\delta_2} \quad \text{for all } \vec{m} \in \mathbb{Z}^{n \times 1} \text{ with } \vec{m}^T \vec{\Omega} \neq 0,$$

where $\vec{\Omega} := (\Omega_1, \dots, \Omega_n)^T$ and $\|\vec{m}\|_2 = |\vec{m}^T \vec{m}|^{\frac{1}{2}}$.

Large gap asymptotics: “good Diophantine property” case

Theorem (X.-Zhang-Zhao, '25)

Let $\Sigma = \cup_{j=0}^n (a_j, b_j)$ be fixed such that $a_0 < b_0 < \dots < a_m < 0 < b_m < \dots < a_n < b_n$ for some $0 \leq m \leq n$ and assume that the good diophantine properties holds. As $s \rightarrow +\infty$, one has

$$\int_{\hat{s}}^s \mathcal{L} \left(p, \vec{V}(t) \right) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + C_p + \mathcal{O}(s^{-1}),$$

where $p \in \mathcal{I}_e$ and C_p is independent of s . Thus, we have, as $s \rightarrow +\infty$,

$$\log \mathcal{F}(s\Sigma) = -\gamma_0 s^2 - 2i\mathcal{D}_{\infty,1}s + \log \theta \left(\vec{V}(s) \right) + \left[\beta^2 - \alpha^2 - \frac{1}{16} \sum_{j=0}^n (\hat{\mathcal{L}}_{a_j} + \hat{\mathcal{L}}_{b_j}) \right] \log s + \check{\mathcal{C}}_2 + \mathcal{O}(s^{-1}),$$

where $\check{\mathcal{C}}_2 = \check{\mathcal{C}}_1 - \frac{1}{16} \sum_{j=0}^n (C_{a_j} + C_{b_j})$ is a constant independent of s with $\check{\mathcal{C}}_1$ as in the asymptotics of the general case.

Large gap asymptotics: ergodic case

Aim 2: Simplify the $\hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) dt$.

Large gap asymptotics: ergodic case

Aim 2: Simplify the $\hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) dt$.

Definition (ergodic property)

The linear flow

$$(0, +\infty) \ni s \mapsto (V_1(s) \bmod 1, V_2(s) \bmod 1, \dots, V_n(s) \bmod 1), \quad V_j(s) = \frac{s}{2\pi} \Omega_j + \frac{1}{2\pi} \operatorname{Im} \zeta_j,$$

is ergodic in the n -dimensional torus $\mathbb{R}^n/\mathbb{Z}^n$ if $\{\vec{V}(s) \bmod \mathbb{Z}^n\}_{s>0}$ is dense in $\mathbb{R}^n/\mathbb{Z}^n$. Equivalently, the linear flow is ergodic in $\mathbb{R}^n/\mathbb{Z}^n$ if $\{\Omega_j\}_{j=1}^n$ are rationally independent, that is, if there exist $(c_1, c_2, \dots, c_n) \in \mathbb{Z}^n$ such that

$$c_1 \Omega_1 + c_2 \Omega_2 + \dots + c_n \Omega_n = 0,$$

then $c_1 = c_2 = \dots = c_n = 0$.

Large gap asymptotics: ergodic case

Aim 2: Simplify the $\hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) dt$.

Definition (ergodic property)

The linear flow

$$(0, +\infty) \ni s \mapsto (V_1(s) \bmod 1, V_2(s) \bmod 1, \dots, V_n(s) \bmod 1), \quad V_j(s) = \frac{s}{2\pi} \Omega_j + \frac{1}{2\pi} \operatorname{Im} \zeta_j,$$

is ergodic in the n -dimensional torus $\mathbb{R}^n/\mathbb{Z}^n$ if $\{\vec{V}(s) \bmod \mathbb{Z}^n\}_{s>0}$ is dense in $\mathbb{R}^n/\mathbb{Z}^n$. Equivalently, the linear flow is ergodic in $\mathbb{R}^n/\mathbb{Z}^n$ if $\{\Omega_j\}_{j=1}^n$ are rationally independent, that is, if there exist $(c_1, c_2, \dots, c_n) \in \mathbb{Z}^n$ such that

$$c_1 \Omega_1 + c_2 \Omega_2 + \dots + c_n \Omega_n = 0,$$

then $c_1 = c_2 = \dots = c_n = 0$.

Theorem (Birkhoff's ergodic theorem)

The time average exists everywhere, and coincides with the space average if f is continuous (or merely Riemann integrable) and the frequencies Ω_j are independent.

Large gap asymptotics: ergodic case

Applying Birkhoff's ergodic theorem to $\hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) \, dt$

Large gap asymptotics: ergodic case

Applying Birkhoff's ergodic theorem to $\hat{\mathcal{L}}_p := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \mathcal{L}(p, \vec{V}(t)) dt$

Theorem (X.-Zhang-Zhao, '25)

Let $\Sigma = \cup_{j=0}^n (a_j, b_j)$ be fixed such that $a_0 < b_0 < \dots < a_m < 0 < b_m < \dots < a_n < b_n$ for some $0 \leq m \leq n$ and assume that $\vec{\Omega}$ satisfies ergodic properties. As $s \rightarrow +\infty$, one has

$$\int_{\hat{s}}^s \mathcal{L}(p, \vec{V}(t)) \frac{dt}{t} = \hat{\mathcal{L}}_p \log s + o(\log s), \quad \hat{\mathcal{L}}_p = \frac{h(p)}{p(p)} \int_{[0,1]^n} \eta(p; u_1, u_2, \dots, u_n) du_1 \cdots du_n.$$

Thus, we have, as $s \rightarrow +\infty$,

$$\log \mathcal{F}(s\Sigma) = -\gamma_0 s^2 - 2i\mathcal{D}_{\infty,1}s + \left[\beta^2 - \alpha^2 - \frac{1}{16} \sum_{j=0}^n (\hat{\mathcal{L}}_{a_j} + \hat{\mathcal{L}}_{b_j}) \right] \log s + o(\log s),$$

where the constant $\hat{\mathcal{L}}_p$, $p \in \mathcal{I}_e$, is explicitly given by the n -fold integral.

Large gap asymptotics: genus $n = 1$ case

In the case of $n = 1$, the linear flow satisfies both good Diophantine properties and ergodic properties.

Large gap asymptotics: genus $n = 1$ case

In the case of $n = 1$, the linear flow satisfies both good Diophantine properties and ergodic properties.

It could be calculated that $\hat{\mathcal{L}}_p = \frac{h(p)}{p(p)} \int_{[0,1)} \eta(p; u_1) \mathrm{d}u_1 = 2$ for $p \in \{a_0, b_0, a_1, b_1\}$.

Large gap asymptotics: genus $n = 1$ case

In the case of $n = 1$, the linear flow satisfies both good Diophantine properties and ergodic properties.

It could be calculated that $\hat{\mathcal{L}}_p = \frac{h(p)}{p(p)} \int_{[0,1)} \eta(p; u_1) du_1 = 2$ for $p \in \{a_0, b_0, a_1, b_1\}$.

Theorem (X.-Zhang-Zhao, '25)

Let $\Sigma := (a_0, b_0) \cup (a_1, b_1)$ be fixed such that $a_0 < b_0 < a_1 < 0 < b_1$ (or $a_0 < 0 < b_0 < a_1 < b_1$). For $\alpha > -1/2$ and $\beta \in i\mathbb{R}$, we have, as $s \rightarrow +\infty$,

$$\log \mathcal{F}(s\Sigma) = -\gamma_0 s^2 - 2iD_{\infty,1}s + \log \theta(V_1(s)) + \left(\beta^2 - \alpha^2 - \frac{1}{2}\right) \log s + C + \mathcal{O}(s^{-1}),$$

where C is an undetermined constant independent of s .

Remarks

Let

$$\mathcal{S}_D := \left\{ \vec{\Omega} : \vec{\Omega} \text{ has "good Diophantine property"} \right\}, \quad \mathcal{S}_E := \left\{ \vec{\Omega} : \vec{\Omega} \text{ is rationally independent} \right\}.$$

Let

$$\mathcal{S}_D := \left\{ \vec{\Omega} : \vec{\Omega} \text{ has "good Diophantine property"} \right\}, \quad \mathcal{S}_E := \left\{ \vec{\Omega} : \vec{\Omega} \text{ is rationally independent} \right\}.$$

We have some examples:

- Genus $n = 1$ case: $\mathcal{S}_D = \mathcal{S}_E = (0, +\infty)$;
- Genus $n = 2$ case: (a) $\vec{\Omega} := (1, \sqrt{2}) \in \mathcal{S}_D \cap \mathcal{S}_E$; (b) $\vec{\Omega} := (1, 1) \in \mathcal{S}_D \setminus \mathcal{S}_E$;
(c) $\vec{\Omega} := (1, C_L) \in \mathcal{S}_E \setminus \mathcal{S}_D$ with $C_L = \sum_{n=1}^{\infty} 10^{-n!}$ be Liouville's constant.
- Genus $n = 3$ case: $\vec{\Omega} = (1, C_L, 1) \notin \mathcal{S}_D \cup \mathcal{S}_E$.
-

Remarks

Let

$$\mathcal{S}_D := \left\{ \vec{\Omega} : \vec{\Omega} \text{ has "good Diophantine property"} \right\}, \quad \mathcal{S}_E := \left\{ \vec{\Omega} : \vec{\Omega} \text{ is rationally independent} \right\}.$$

We have some examples:

- Genus $n = 1$ case: $\mathcal{S}_D = \mathcal{S}_E = (0, +\infty)$;
- Genus $n = 2$ case: (a) $\vec{\Omega} := (1, \sqrt{2}) \in \mathcal{S}_D \cap \mathcal{S}_E$; (b) $\vec{\Omega} := (1, 1) \in \mathcal{S}_D \setminus \mathcal{S}_E$;
(c) $\vec{\Omega} := (1, C_L) \in \mathcal{S}_E \setminus \mathcal{S}_D$ with $C_L = \sum_{n=1}^{\infty} 10^{-n!}$ be Liouville's constant.
- Genus $n = 3$ case: $\vec{\Omega} = (1, C_L, 1) \notin \mathcal{S}_D \cup \mathcal{S}_E$.
-

For $n \geq 2$, all of the following cases can and do occur for certain choices of the edges points $\{a_j, b_j\}_{j=0}^n$.

$$\vec{\Omega} \notin \mathcal{S}_D \cup \mathcal{S}_E, \quad \vec{\Omega} \in \mathcal{S}_D \setminus \mathcal{S}_E, \quad \vec{\Omega} \in \mathcal{S}_E \setminus \mathcal{S}_D, \quad \vec{\Omega} \in \mathcal{S}_D \cap \mathcal{S}_E.$$

[Deift-Its-Zhou, '97]

- For the n -fold integral obtained in the ergodic case

$$\hat{\mathcal{L}}_p = \frac{h(p)}{p(p)} \int_{[0,1]^n} \eta(p; u_1, u_2, \dots, u_n) du_1 \cdots du_n,$$

- (a) It's been proved that $\hat{\mathcal{L}}_p \equiv 2$ for $n = 1$.
- (b) Numerically, $\hat{\mathcal{L}}_p \equiv 2$ for all finite $n > 1$.

- For the n -fold integral obtained in the ergodic case

$$\hat{\mathcal{L}}_p = \frac{h(p)}{p(p)} \int_{[0,1]^n} \eta(p; u_1, u_2, \dots, u_n) du_1 \cdots du_n,$$

- (a) It's been proved that $\hat{\mathcal{L}}_p \equiv 2$ for $n = 1$.
 - (b) Numerically, $\hat{\mathcal{L}}_p \equiv 2$ for all finite $n > 1$.
- The multiplicative constant \rightsquigarrow challenge.

About the proofs

An RH characterization of the $\mathcal{F}(s\Sigma)$

Rely on an integrable structure of the confluent hypergeometric kernel $K^{(\alpha,\beta)}(x,y)$

$$K^{(\alpha,\beta)}(x,y) = \frac{\vec{f}(x)^\top \vec{h}(y)}{x-y} = \frac{\sum_{k=1}^2 f_k(x) h_k(y)}{x-y}.$$

An RH characterization of the $\mathcal{F}(s\Sigma)$

Rely on an integrable structure of the confluent hypergeometric kernel $K^{(\alpha,\beta)}(x,y)$

$$K^{(\alpha,\beta)}(x,y) = \frac{\vec{f}(x)^\top \vec{h}(y)}{x-y} = \frac{\sum_{k=1}^2 f_k(x) h_k(y)}{x-y}.$$

- $Y(z) := I - \int_{s\Sigma} \frac{\vec{F}(x)\vec{h}(x)^\top}{x-z} dx$ satisfies an 2×2 RH problem, where $\vec{F}(z) = (1 - \mathcal{K}^{(\alpha,\beta)}|_{s\Sigma})^{-1} \vec{f}(z)$.

[Its-Izergin-Korepin-Slavnov, '90] [Deift-Its-Zhou, '97]

An RH characterization of the $\mathcal{F}(s\Sigma)$

Rely on an integrable structure of the confluent hypergeometric kernel $K^{(\alpha,\beta)}(x,y)$

$$K^{(\alpha,\beta)}(x,y) = \frac{\vec{f}(x)^\top \vec{h}(y)}{x-y} = \frac{\sum_{k=1}^2 f_k(x) h_k(y)}{x-y}.$$

- $Y(z) := I - \int_{s\Sigma} \frac{\vec{F}(x)\vec{h}(x)^\top}{x-z} dx$ satisfies an 2×2 RH problem, where $\vec{F}(z) = (1 - \mathcal{K}^{(\alpha,\beta)}|_{s\Sigma})^{-1} \vec{f}(z)$.

[Its-Izergin-Korepin-Slavnov, '90] [Deift-Its-Zhou, '97]

- $\vec{f}(z)$ and $\vec{h}(z)$ are expressed in terms of a confluent hypergeometric parametrix Φ_{CH} .

[Claeys-Its-Krasovsky, '11] [Xu-Zhao, '20]

An RH characterization of the $\mathcal{F}(s\Sigma)$

Rely on an integrable structure of the confluent hypergeometric kernel $K^{(\alpha,\beta)}(x,y)$

$$K^{(\alpha,\beta)}(x,y) = \frac{\vec{f}(x)^\top \vec{h}(y)}{x-y} = \frac{\sum_{k=1}^2 f_k(x) h_k(y)}{x-y}.$$

- $Y(z) := I - \int_{s\Sigma} \frac{\vec{F}(x)\vec{h}(x)^\top}{x-z} dx$ satisfies an 2×2 RH problem, where $\vec{F}(z) = (1 - \mathcal{K}^{(\alpha,\beta)}|_{s\Sigma})^{-1} \vec{f}(z)$.

[Its-Izergin-Korepin-Slavnov, '90] [Deift-Its-Zhou, '97]


- $\vec{f}(z)$ and $\vec{h}(z)$ are expressed in terms of a confluent hypergeometric parametrix Φ_{CH} .

[Claeys-Its-Krasovsky, '11] [Xu-Zhao, '20]

Undressing transform \rightsquigarrow an RH problem for X with constant jumps.


RH problem for X

(a) $X(z)$ is holomorphic for $z \in \mathbb{C} \setminus \Gamma_X$



RH problem for X

(b) For $z \in \Gamma_X$, we have $X_+(z) = X_-(z)J_X(z)$.



RH problem for X

(c) As $z \rightarrow \infty$, we have

$$X(z) = \left(I + \frac{X_1(s)}{z} + \mathcal{O}(z^{-2}) \right) z^{-\beta\sigma_3} e^{-isz\sigma_3},$$

where

$$X_1(s) = (2s)^{\beta\sigma_3} \begin{pmatrix} \frac{1}{2s}(\Phi_{\text{CH},1})_{11} + \frac{1}{s}(Y_1(s))_{11} & \frac{1}{2s}(\Phi_{\text{CH},1})_{12} + \frac{1}{s}(Y_1(s))_{12}e^{-\pi i\beta} \\ \frac{1}{2s}(\Phi_{\text{CH},1})_{21} + \frac{1}{s}(Y_1(s))_{21}e^{\pi i\beta} & \frac{1}{2s}(\Phi_{\text{CH},1})_{22} + \frac{1}{s}(Y_1(s))_{22} \end{pmatrix} (2s)^{-\beta\sigma_3}$$

with $\Phi_{\text{CH},1} := \lim_{z \rightarrow \infty} z(\Phi_{\text{CH}}(z)z^{\beta\sigma_3}e^{\frac{i}{2}z\sigma_3} - I)$.

(d) As $z \rightarrow p$ from $\Omega_{X,1}$, $p \in \mathcal{I}_e$, we have $X(z) = \mathcal{O}(\log(z-p))$.

(e) As $z \rightarrow 0$ from $\text{Im } z > 0$, we have

$$X(z) = X_0(z)z^{\alpha\sigma_3},$$

where $X_0(z)$ is holomorphic in the neighborhood of 0. The behavior of $X(z)$ as $z \rightarrow 0$ is determined by jump conditions.

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

For genus n case ($n \geq 2$) \rightsquigarrow create difficulties in further simplification (All quantities that we encounter are vector-valued).

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

For genus n case ($n \geq 2$) \rightsquigarrow create difficulties in further simplification (All quantities that we encounter are vector-valued). Particularly, one should expand a vector-valued function $f(\vec{z})$, $\vec{z} \in \mathbb{C}^n$ at a certain point \vec{z}_0 :

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

For genus n case ($n \geq 2$) \rightsquigarrow create difficulties in further simplification (All quantities that we encounter are vector-valued). Particularly, one should expand a vector-valued function $f(\vec{z})$, $\vec{z} \in \mathbb{C}^n$ at a certain point \vec{z}_0 :

$$f(\vec{z}) = f(\vec{z}_0) + \langle \nabla f(\vec{z}_0), \vec{z} - \vec{z}_0 \rangle + \frac{1}{2}(\vec{z} - \vec{z}_0)^T \nabla^2 f(\vec{z}_0)(\vec{z} - \vec{z}_0) + \text{tensors}(\vec{z} - \vec{z}_0)^3 + \dots$$

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

For genus n case ($n \geq 2$) \rightsquigarrow create difficulties in further simplification (All quantities that we encounter are vector-valued). Particularly, one should expand a vector-valued function $f(\vec{z})$, $\vec{z} \in \mathbb{C}^n$ at a certain point \vec{z}_0 :

$$f(\vec{z}) = f(\vec{z}_0) + \langle \nabla f(\vec{z}_0), \vec{z} - \vec{z}_0 \rangle + \frac{1}{2}(\vec{z} - \vec{z}_0)^T \nabla^2 f(\vec{z}_0)(\vec{z} - \vec{z}_0) + \text{tensors}(\vec{z} - \vec{z}_0)^3 + \dots$$

Difficulties: (a) several complex variables; (b) hard to evaluate the coefficients of higher order derivatives.

A “failed” differential identity

Usually, one could always relate $\partial_s \log \mathcal{F}(s\Sigma)$ to the local behavior of X at $z = p$ for $p \in \mathcal{I}_e$

Proposition

$$\partial_s \log \mathcal{F}(s\Sigma) = \sum_{p \in \mathcal{I}_e} f_p(\alpha, \beta) \lim_{z \rightarrow p} (X^{-1}(z)X'(z))_{21}.$$

For genus 1 case, It's okay! (All quantities that we encounter are scalar-valued)

For genus n case ($n \geq 2$) \rightsquigarrow create difficulties in further simplification (All quantities that we encounter are vector-valued). Particularly, one should expand a vector-valued function $f(\vec{z})$, $\vec{z} \in \mathbb{C}^n$ at a certain point \vec{z}_0 :

$$f(\vec{z}) = f(\vec{z}_0) + \langle \nabla f(\vec{z}_0), \vec{z} - \vec{z}_0 \rangle + \frac{1}{2}(\vec{z} - \vec{z}_0)^T \nabla^2 f(\vec{z}_0)(\vec{z} - \vec{z}_0) + \text{tensors}(\vec{z} - \vec{z}_0)^3 + \dots$$

Difficulties: (a) several complex variables; (b) hard to evaluate the coefficients of higher order derivatives. \rightsquigarrow **We need a new one!**

A new differential identity

Proposition (X.-Zhang-Zhao, '25)

We have

$$\partial_s \log \mathcal{F}(s\Sigma) = i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s},$$

where X_1 is the residue term of X at infinity.

A new differential identity

Proposition (X.-Zhang-Zhao, '25)

We have

$$\partial_s \log \mathcal{F}(s\Sigma) = i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s},$$

where X_1 is the residue term of X at infinity.

Only the 1-st order expansion of some vector-valued functions is needed (Not general!).

A new differential identity

Proposition (X.-Zhang-Zhao, '25)

We have

$$\partial_s \log \mathcal{F}(s\Sigma) = i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s},$$

where X_1 is the residue term of X at infinity.

Only the 1-st order expansion of some vector-valued functions is needed (Not general!).
The proof relies on the following lemma.

Lemma (X.-Zhang-Zhao, '25)

$$\partial_s K_s^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} (A(sx)B(sy) + A(sy)B(sx)).$$

A new differential identity

Proposition (X.-Zhang-Zhao, '25)

We have

$$\partial_s \log \mathcal{F}(s\Sigma) = i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s},$$

where X_1 is the residue term of X at infinity.

Only the 1-st order expansion of some vector-valued functions is needed (Not general!).
The proof relies on the following lemma.

Lemma (X.-Zhang-Zhao, '25)

$$\partial_s K_s^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} (A(sx)B(sy) + A(sy)B(sx)).$$

- Ideas of proof: play several recurrence relations of the CH functions.

A new differential identity

Proposition (X.-Zhang-Zhao, '25)

We have

$$\partial_s \log \mathcal{F}(s\Sigma) = i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s},$$

where X_1 is the residue term of X at infinity.

Only the 1-st order expansion of some vector-valued functions is needed (Not general!).
The proof relies on the following lemma.

Lemma (X.-Zhang-Zhao, '25)

$$\partial_s K_s^{(\alpha, \beta)}(x, y) = \frac{1}{2\pi} \frac{\Gamma(1 + \alpha + \beta)\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)^2} (A(sx)B(sy) + A(sy)B(sx)).$$

- Ideas of proof: play several recurrence relations of the CH functions.
- sine kernel: $\partial_s K_s^{(0,0)}(x, y) = \frac{\cos(s(x-y))}{\pi}.$


Deift-Zhou nonlinear steepest analysis for X

g -function mechanism is taken to normalize the RH problem for X .

Deift-Zhou nonlinear steepest analysis for X

g -function mechanism is taken to normalize the RH problem for X .

Opening lenses to obtain a solvable RH problem for P



Deift-Zhou nonlinear steepest analysis for X

Global parametrix for $P^{(\infty)}$

- A Szegő function \mathcal{D} to deal with the Fisher-Hartwig singularity.

$$\mathcal{D}(z) := \exp \left\{ \frac{\sqrt{\mathcal{R}(z)}}{2\pi i} \int_{\Sigma} \frac{\mathcal{H}(\xi)}{\xi - z} d\xi \right\},$$

where

$$\mathcal{H}(z) := \frac{\log z^{-2\beta} + \operatorname{sgn}(z)(\alpha - \beta)\pi i + \zeta_j}{\sqrt{\mathcal{R}(z)}_+}, \quad z \in (a_j, b_j), \quad j = 0, 1, \dots, n.$$

Deift-Zhou nonlinear steepest analysis for X

Global parametrix for $P^{(\infty)}$

- A Szegő function \mathcal{D} to deal with the Fisher-Hartwig singularity.

$$\mathcal{D}(z) := \exp \left\{ \frac{\sqrt{\mathcal{R}(z)}}{2\pi i} \int_{\Sigma} \frac{\mathcal{H}(\xi)}{\xi - z} d\xi \right\},$$

where

$$\mathcal{H}(z) := \frac{\log z^{-2\beta} + \operatorname{sgn}(z)(\alpha - \beta)\pi i + \zeta_j}{\sqrt{\mathcal{R}(z)}_+}, \quad z \in (a_j, b_j), \quad j = 0, 1, \dots, n.$$

- Construction of $P^{(\infty)}$: a model RH problem solvable in terms of Riemann- θ function.

[Deift-Its-Zhou, '97]

Deift-Zhou nonlinear steepest analysis for X

Global parametrix for $P^{(\infty)}$

- A Szegő function \mathcal{D} to deal with the Fisher-Hartwig singularity.

$$\mathcal{D}(z) := \exp \left\{ \frac{\sqrt{\mathcal{R}(z)}}{2\pi i} \int_{\Sigma} \frac{\mathcal{H}(\xi)}{\xi - z} d\xi \right\},$$

where

$$\mathcal{H}(z) := \frac{\log z^{-2\beta} + \operatorname{sgn}(z)(\alpha - \beta)\pi i + \zeta_j}{\sqrt{\mathcal{R}(z)}_+}, \quad z \in (a_j, b_j), \quad j = 0, 1, \dots, n.$$

- Construction of $P^{(\infty)}$: a model RH problem solvable in terms of Riemann- θ function.
[Deift-Its-Zhou, '97]
- Some exact formulas related to $P^{(\infty)}$ are required.

Deift-Zhou nonlinear steepest analysis for X

Local parametrix at the edge points: Bessel model RH problem.

Deift-Zhou nonlinear steepest analysis for X

Local parametrix at the edge points: Bessel model RH problem.

Final transformation: Small norm RH problem for R .

Deift-Zhou nonlinear steepest analysis for X

Local parametrix at the edge points: Bessel model RH problem.

Final transformation: Small norm RH problem for R .

Proposition

$$\begin{aligned}\partial_s \log \mathcal{F}(s\Sigma) &= i((X_1(s))_{11} - (X_1(s))_{22}) - \frac{\alpha^2 - \beta^2}{s} \\ &= -2s\gamma_0 + i\left((P_1^{(\infty)})_{11} - (P_1^{(\infty)})_{22}\right) + \frac{i}{s}\left((R_1^{(1)})_{11} - (R_1^{(1)})_{22}\right) - \frac{\alpha^2 - \beta^2}{s} + \mathcal{O}(s^{-2}),\end{aligned}$$

where $*_1$ is the residue term of $*$ at infinity.

How the two properties help?

Go back to the time average integral

$$I = \frac{1}{T} \int_{\hat{s}}^T \mathcal{Y}(\vec{V}(t)) \, dt, \quad \vec{V}(t) = t\vec{\Omega} + \vec{\zeta} \in \mathbb{R}^n / \mathbb{Z}^n.$$

How the two properties help?

Go back to the time average integral

$$I = \frac{1}{T} \int_{\hat{s}}^T \mathcal{Y}(\vec{V}(t)) dt, \quad \vec{V}(t) = t\vec{\Omega} + \vec{\zeta} \in \mathbb{R}^n/\mathbb{Z}^n.$$

Analytical function \mathcal{Y} on $\mathbb{R}^n/\mathbb{Z}^n$ admits a Fourier expansion

$$\mathcal{Y}(\vec{\mu}) = \sum_{\vec{m} \in \mathbb{Z}^n} \ell_{\vec{m}} e^{2\pi i \vec{m}^T \vec{\mu}}, \quad \vec{\mu} \in \mathbb{R}^n/\mathbb{Z}^n.$$

How the two properties help?

Go back to the time average integral

$$I = \frac{1}{T} \int_{\hat{s}}^T \mathcal{Y}(\vec{V}(t)) dt, \quad \vec{V}(t) = t\vec{\Omega} + \vec{\zeta} \in \mathbb{R}^n/\mathbb{Z}^n.$$

Analytical function \mathcal{Y} on $\mathbb{R}^n/\mathbb{Z}^n$ admits a Fourier expansion

$$\mathcal{Y}(\vec{\mu}) = \sum_{\vec{m} \in \mathbb{Z}^n} \ell_{\vec{m}} e^{2\pi i \vec{m}^T \vec{\mu}}, \quad \vec{\mu} \in \mathbb{R}^n/\mathbb{Z}^n.$$

Evaluate

$$I = \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} \neq 0}} A + \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} = 0}} \ell_{\vec{m}} e^{i \vec{m}^T \vec{\zeta}}. \quad (\text{Oscillatory term} + \text{Non-oscillatory term})$$

How the two properties help?

Go back to the time average integral

$$I = \frac{1}{T} \int_{\hat{s}}^T \mathcal{Y}(\vec{V}(t)) dt, \quad \vec{V}(t) = t\vec{\Omega} + \vec{\zeta} \in \mathbb{R}^n / \mathbb{Z}^n.$$

Analytical function \mathcal{Y} on $\mathbb{R}^n / \mathbb{Z}^n$ admits a Fourier expansion

$$\mathcal{Y}(\vec{\mu}) = \sum_{\vec{m} \in \mathbb{Z}^n} \ell_{\vec{m}} e^{2\pi i \vec{m}^T \vec{\mu}}, \quad \vec{\mu} \in \mathbb{R}^n / \mathbb{Z}^n.$$

Evaluate

$$I = \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} \neq 0}} \textcolor{red}{A} + \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} = 0}} \textcolor{blue}{\ell_{\vec{m}}} e^{i \vec{m}^T \vec{\zeta}}. \quad (\text{Oscillatory term} + \text{Non-oscillatory term})$$

- Good diophantine property + bound arguments (for $\ell_{\vec{m}}$) $\implies \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} \neq 0}} \textcolor{red}{A} = \mathcal{O}(t^{-1})$.

How the two properties help?

Go back to the time average integral

$$I = \frac{1}{T} \int_{\hat{s}}^T \mathcal{Y}(\vec{V}(t)) dt, \quad \vec{V}(t) = t\vec{\Omega} + \vec{\zeta} \in \mathbb{R}^n / \mathbb{Z}^n.$$

Analytical function \mathcal{Y} on $\mathbb{R}^n / \mathbb{Z}^n$ admits a Fourier expansion

$$\mathcal{Y}(\vec{\mu}) = \sum_{\vec{m} \in \mathbb{Z}^n} \ell_{\vec{m}} e^{2\pi i \vec{m}^T \vec{\mu}}, \quad \vec{\mu} \in \mathbb{R}^n / \mathbb{Z}^n.$$


Evaluate

$$I = \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} \neq 0}} \textcolor{red}{A} + \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} = 0}} \textcolor{blue}{\ell_{\vec{m}} e^{i \vec{m}^T \vec{\zeta}}}. \quad (\text{Oscillatory term} + \text{Non-oscillatory term})$$

- Good diophantine property + bound arguments (for $\ell_{\vec{m}}$) $\implies \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} \neq 0}} \textcolor{red}{A} = \mathcal{O}(t^{-1})$.
- Ergodic property + Birkhoff's ergodic theorem $\implies \sum_{\substack{\vec{m} \in \mathbb{Z}^n \\ \vec{m}^T \vec{\Omega} = 0}} \textcolor{blue}{\ell_{\vec{m}} e^{i \vec{m}^T \vec{\zeta}}} = \ell_{\vec{0}}$.


Future work

- Confluent hypergeometric kernel determinant with a FH singularity at the gap and the merging case at the FH singular point.



Future work

- Confluent hypergeometric kernel determinant with a FH singularity at the gap and the merging case at the FH singular point.



- New differential identity \rightsquigarrow Other integrable kernel determinants for general genus $g > 1$ case?
(Progress on the Airy kernel determinant)

Thanks for your attention!