

Contents

I Basic Set Theory	1
I.1 Club Sets	1
II Forcing Notions	3
II.1 Definition for the Forcing	3
II.2 General Properties for a Forcing	4
II.3 Iterated Forcing	7
III Type of Forcings	10
III.1 ccc Forcing	10
III.2 σ -centerd Forcing	11
III.3 σ -closed Forcing	11
III.4 Axiom A Forcing	12
III.5 Proper Forcing	18
IV List of Forcing Notions	25
IV.1 Cohen Forcing	25
IV.2 Hechler Forcing	26
IV.3 Mathias Forcing	27
IV.4 Sacks Forcing (Perfect Set Forcing)	32
IV.5 Shooting A Club by Finite Conditions	35
V The Concrete Forcing Examples	36
V.1 Axiom of Choice	36
V.2 Continuum Hypothesis	36
V.3 Martin Axiom	36
V.4 Borel Conjecture	36

I Basic Set Theory

I.1 Club Sets

Definition I.1.1 ([Jec03]). For a regular cardinal κ and a subset $C \subset \kappa$, C is *unbounded* provided that $\cup C = \kappa$ and C is *closed* provided that for any limit $\gamma \in \kappa$ if $\cup(C \cap \gamma) = \gamma$ then $\gamma \in C$. A subset C is a *club subset of κ* provided that C is closed and unbounded subset of κ . A subset $S \subset \kappa$ is a *stationary subset of κ* provided that S meets every club subset of κ .

Note that a subset C is closed is equivalent to that for any subset $S \subset \kappa$ we have $\cup S \in C$ when $S \subset C$ and the size of stationary subset is κ since for a strictly increasing convergence sequence $\langle \alpha_\xi \in \kappa ; \xi \in \kappa \rangle \nearrow \kappa$, $\langle \alpha_\xi ; \eta \in \xi \wedge \xi \in \kappa \rangle$ is club for any $\eta \in \kappa$.

Theorem I.1.2 ([Kun11]). For a regular uncountable cardinal κ , we have the following:

1. The intersection of less than κ club subsets of κ is club,
2. The *diagonal intersection* of a κ -sequence of club subsets of κ

$$\Delta\langle C_\alpha; \alpha \in \kappa \rangle = \{\xi \in \kappa \setminus 1; \xi \in \cap\{C_\alpha; \alpha \in \xi\}\}$$

is club.

Proof. 1. We shall prove the following by induction on $\beta_0 \in \kappa \setminus 1$:

For κ -club sets $\langle C_\xi; \xi \in \beta_0 \rangle$, the intersection $C = \cap\{C_\xi; \xi \in \beta_0\}$ is club.

The case of the leading stage and the successor stages are clear. Let β_0 be limit. Define $D_\xi = \cap\{C_\eta; \eta \in \xi\}$ for $\xi \in \beta_0$ and note that by induction hypothesis every D_ξ is club and we obtain that the decreasing club sets $\langle D_\xi; \xi \in \beta_0 \rangle$ such that the intersection, D , equals $\cap\{C_\xi; \xi \in \beta_0\}$. Note that manifestly D is closed. We shall show that D is unbounded in κ . Fix a $\gamma \in \kappa$ and define $\gamma_\xi \in D_\xi$, recursively such that:

- $\gamma_0 \in D_0$ such that $\gamma \in \gamma_0$,
- $\gamma_\xi \in D_\xi$ such that $\gamma_\eta \in \gamma_\xi$ for each $\eta \in \xi$.

Then we obtain an increasing sequence $x = \langle \gamma_\xi; \xi \in \beta_0 \rangle$ in κ with limit $\beta \in \kappa$. Since for every $\xi \in \beta_0$ x is eventually in D_ξ so does β . Therefore, β witnesses that D is unbounded in κ .

2. To see that $\Delta\langle C_\alpha; \alpha \in \kappa \rangle$ is closed. Let $\langle \beta_\xi; \xi \in \beta_0 \rangle \nearrow \beta$ be a sequence with limit point $\beta \in \kappa$. We shall show that $\forall \eta \in \beta$ ($\beta \in C_\eta$). For any $\eta \in \beta$ there is a $\xi_0 \in \beta_0$ such that $\eta \in \beta_\xi$ for any $\xi \supset \xi_0$. This shows that the sequence eventually contained in C_η so does the limit point β .

To see that $\Delta\langle C_\alpha; \alpha \in \kappa \rangle$ is unbounded. Fix $\delta_0 \in \kappa \setminus 1$ and define δ_n , $n \in \omega$, recursively such that:

$$\delta_{n+1} \supset \delta_n \text{ and } \delta_{n+1} \in \cap\{C_\alpha; \alpha \in \delta_n\} \text{ for any } n \in \omega.$$

Then for the limit point δ , since we have $\delta \in \cap\{C_\alpha; \alpha \in \delta\}$, $\delta \in \Delta\langle C_\alpha; \alpha \in \kappa \rangle$. This shows that $\Delta\langle C_\alpha; \alpha \in \kappa \rangle$ is unbounded. \square

Definition I.1.3 ([Jec03]). For a set of ordinals A , a function $f: A \rightarrow A$ is a *regressive function* provided that $f(\alpha) \in \alpha$ for any $\alpha \in A$.

Theorem I.1.4 (Hodor Theorem, [Jec03]). Let f be a regressive function on a stationary set $S \subset \kappa$. Then there is a stationary set $T \subset S$ and $\gamma \in \kappa$ such that $f(\alpha) = \gamma$ for any $\alpha \in T$.

Proof. Assume that “ $\{\alpha \in S; f(\alpha) = \gamma\}$ is not stationary.” for all $\gamma \in \kappa$. For each $\gamma \in \kappa$ choose a club set C_γ which does not meet $\{\alpha \in S; f(\alpha) = \gamma\}$. Since $C = \Delta\{C_\gamma; \gamma \in \kappa\}$ is a club set, there is $\alpha \in S \cap C$ and for $\gamma \in \alpha$ we obtain that $\alpha \in C_\gamma$. This shows that $f(\alpha) \neq \gamma$ for any $\gamma \in \alpha$, this contrary to that $f(\alpha) \in \alpha$. \square

Theorem I.1.5 (Delta System Lemma, [Kun11]). Let λ and κ be regular cardinals with $\omega \leq \lambda < \kappa$ and assume that $\forall \theta \in \kappa (\theta^{<\lambda} < \kappa)$. Then for a κ size family \mathcal{A} with $|A| < \lambda$ for each $A \in \mathcal{A}$, there is a delta system $\mathcal{B} \in [\mathcal{A}]^\kappa$. Moreover,

Proof. III.6.15 □

II Forcing Notions

II.1 Definition for the Forcing

Definition II.1.1. A *forcing notion* $(\mathbb{P}, \leq_{\mathbb{P}}, \mathbb{1})$ is a poset $(\mathbb{P}, \leq_{\mathbb{P}})$ with the maximal element $\mathbb{1}$. For $p \in \mathbb{P}$ an extension of p is $q \in \mathbb{P}$ satisfying $q \leq_{\mathbb{P}} p$.

1. A set $D \subset \mathbb{P}$ is a *dense subset below* $p \in \mathbb{P}$ provided that for every extension $q \leq_{\mathbb{P}} p$ there is an extension for p in D and we simply said D is *dense* in \mathbb{P} if D is dense below $\mathbb{1}$.
2. A set $G \subset \mathbb{P}$ is a (V, \mathbb{P}) -*generic filter* provided that:
 - (a) $G \subset \mathbb{P}$, $G \in V$,
 - (b) $\forall p, q \in G \exists r \in G r \leq_{\mathbb{P}} p, q$,
 - (c) $\forall p \in G \forall q \in \mathbb{P} (p \leq_{\mathbb{P}} q \implies q \in G)$,
 - (d) Every dense subset $D \subset \mathbb{P}$, D meets G if $D \in V$.

We will usually say that G is \mathbb{P} -generic and write \leq for the $\leq_{\mathbb{P}}$ when there is no danger of confusing.

Definition II.1.2. Let \mathbb{P} be a forcing notion and E be a subset of \mathbb{P} .

1. p and q are *compatible*, $p \not\perp q$, provided that there is a common extension $r \leq p, q$, for any conditions $p, q \in \mathbb{P}$,
2. p and q are *incompatible*, $p \perp q$, provided that $\neg(p \not\perp q)$,
3. E is *predense* provided that for every condition $p \in \mathbb{P}$ there is $q \in E$ such that $p \not\perp q$,
4. E is *open* provided that for $p \leq q$ if $p \in E$ then $q \in E$,
5. E is *dense open* provided that E is dense and open,
6. E is an *antichain* provided that there are no distinct $p, q \in E$ such that $p \not\perp q$,
7. E is a *maximal antichain* provided that E is an antichain and maximum among the subset relation,
8. For a condition $p \in \mathbb{P}$, $p \perp E$ is an abbreviation for $\forall q \in E p \perp q$.

Theorem II.1.3 ([Kun11]). III.3.60

Definition II.1.4. Let \mathbb{P} be a forcing notion.

1. For names τ, σ, θ in $V^{\mathbb{P}}$, define, recursively,:
 - (1-a) $p \Vdash \tau = \sigma$ provided that

$$\forall \theta \in \text{dom}(\tau) \cup \text{dom}(\sigma) \forall q \leq p (q \Vdash \theta \in \tau \Leftrightarrow q \Vdash \theta \in \sigma),$$
 - (1-b) $p \Vdash \tau \in \sigma$ provided that

$$\{q \leq p; \exists \langle \theta, r \rangle \in \sigma (q \leq r \wedge q \Vdash \tau = \theta)\} \text{ is dense below } p.$$
2. For formulas $\varphi, \psi \in \mathcal{FL}_{\mathbb{P}}$,
 - (2-a) $p \Vdash \varphi \wedge \psi$ provided that $p \Vdash \varphi$ and $p \Vdash \psi$,
 - (2-b) $p \Vdash \neg \varphi$ provided that $\neg \exists q \leq p (q \Vdash \varphi)$,
 - (2-c) $p \Vdash \forall x \varphi(x)$ provided that $p \Vdash \varphi(\tau)$ for any $\tau \in V^{\mathbb{P}}$,
 - (2-d) $p \Vdash \exists x \varphi(x)$ provided that $\{q \leq p; \exists \tau \in V^{\mathbb{P}} q \Vdash \varphi(\tau)\}$ is dense below p .

Definition II.1.5. Let \mathbb{P} be a forcing notion. A \mathbb{P} -name, τ , is a relation such that:

$$\forall \langle \sigma, p \rangle \in \tau \text{ ("}\sigma \text{ is a } \mathbb{P}\text{-name."} \wedge p \in \mathbb{P})$$

Let $V^{\mathbb{P}}$ be the class of all \mathbb{P} -names and for a transitive model M for ZF with $\mathbb{P} \in M$, define:

$$M^{\mathbb{P}} = M \cap V^{\mathbb{P}} = \{\tau \in M; M \models \text{"}\tau \text{ is } \mathbb{P}\text{-name."}\}$$

Theorem II.1.6 ([Kun11]). For a transitive model M for ZF–P (resp. ZF, ZFC), \mathbb{P} a forcing notion with $\mathbb{P} \in M$ and G a generic, $M[G]$ is a transitive model for ZF–P (resp. ZF, ZFC) with $M \subset M[G]$ and $G \in M[G]$. Conversely, if there is a transitive model N for ZF–P with $M \subset N$ and $G \in N$, we obtain that $M[G] \subset M$.

Proof. See IV.2.15, 2.19, 2.26. □

II.2 General Properties for a Forcing

Definition II.2.1. Let \mathbb{P} and \mathbb{Q} be forcing posets and $i: \mathbb{P} \rightarrow \mathbb{Q}$ a function. A function i is a *complete embedding* provided that

1. $i(1_{\mathbb{P}}) = 1_{\mathbb{Q}}$,
2. $q \leq_{\mathbb{P}} p$ implies $i(q) \leq_{\mathbb{Q}} i(p)$ for any $p, q \in \mathbb{P}$,
3. $p \not\leq_{\mathbb{P}} q$ iff $i(p) \not\leq_{\mathbb{Q}} i(q)$ for any $p, q \in \mathbb{P}$, and
4. If A is a maximal antichain in \mathbb{P} , $i(A)$ is a maximal antichain in \mathbb{Q} , for any subset $A \subset \mathbb{P}$.

And a function i is a *dense embedding* provided that it is a complete embedding and

- 5. $i(\mathbb{P})$ is a dense subset of \mathbb{Q} .

Theorem II.2.2 ([Kun11]). Let M be a transitive model for ZFC, \mathbb{P}, \mathbb{Q} forcing notions and a function $i: \mathbb{P} \rightarrow \mathbb{Q}$ with $\mathbb{Q}, \mathbb{P}, i \in M$ and i is a dense embedding. Define:

- $i_*: M^{\mathbb{P}} \rightarrow M^{\mathbb{Q}}$ be the function which assign to τ the

$$i_*(\tau) = \{ \langle i_*(\sigma), i(p) \rangle ; \langle \sigma, p \rangle \in \tau \}.$$

- $\tilde{i}(G) = \{ q \in \mathbb{Q} ; \exists p \in G \ i(p) \leq_{\mathbb{Q}} q \}$ for any subset G .

Then we have the followings:

1. For any \mathbb{P} -generic G , $\tilde{i}(G)$ is \mathbb{Q} -generic and $G = i^{-1}\tilde{i}(G)$, moreover we have $M[G] = M[\tilde{i}(G)]$,
2. For any \mathbb{Q} -generic H , $i^{-1}(H)$ is \mathbb{P} -generic and $H = \tilde{i}i^{-1}(H)$, moreover we have $M[H] = M[i^{-1}(H)]$,
3. For any formula $\varphi(x_1, \dots, x_{n-1})$ of $\mathcal{L} = \{\in\}$, $p \in \mathbb{P}$ and $\tau_1, \dots, \tau_{n-1} \in M^{\mathbb{P}}$, we have

$$p \Vdash_{\mathbb{P}} \varphi(\tau_1, \dots, \tau_{n-1}) \text{ iff } i(p) \Vdash_{\mathbb{Q}} \varphi(i_*(\tau_1), \dots, i_*(\tau_{n-1})).$$

Definition II.2.3 ([Jec03]). A forcing poset \mathbb{P} is *separative* provided that for any conditions $p, q \in \mathbb{P}$ if $p \not\leq q$ then there is $r \leq p$ such that $r \perp q$.

Theorem II.2.4 ([Jec03]). Let \mathbb{P} be a forcing poset.

1. If \mathbb{P} is separative, $p \leq q$ iff $p \Vdash q \in \dot{G}$ for any $p, q \in \mathbb{P}$.
2. There is a separative forcing poset \mathbb{Q} and dense embedding $i: \mathbb{P} \rightarrow \mathbb{Q}$.

Proof. Let us show the first assertion.

Let us show the second assertion. Define an equivalent relation $\sim \subset \mathbb{P} \times \mathbb{P}$ such that for any $p, q \in \mathbb{P}$:

$$p \sim q \Leftrightarrow \forall z \in \mathbb{P} \ (z \not\leq p \leftrightarrow z \not\leq q)$$

Then we have the following property:

$$x_0 \sim x_1, y_0 \sim y_1 \text{ and } \forall z \leq_{\mathbb{P}} x_0 \ (z \not\leq y_0) \text{ implies } \forall z \leq_{\mathbb{P}} x_1 \ (z \not\leq y_1).$$

Define a forcing poset $\mathbb{Q} = (\mathbb{P}/\sim, \leq_{\mathbb{Q}})$ where $[q] \leq_{\mathbb{Q}} [p]$ if and only if r is compatible with p for any extension $r \leq_{\mathbb{P}} q$. Then it is easy to see that \mathbb{Q} is separative and to see that i is dense embedding since i is surjective, it suffices to see the followings:

- (a) $p \leq_{\mathbb{P}} q$ implies $i(p) \leq_{\mathbb{Q}} i(q)$ for any $p, q \in \mathbb{P}$,
- (b) $p \not\leq_{\mathbb{P}} q$ iff $i(p) \not\leq_{\mathbb{Q}} i(q)$ for any $p, q \in \mathbb{P}$.

It is obvious that condition (a) and sufficient condition in (b). To see the necessary condition, let $r \in \mathbb{P}$ such that $[r] \leq_{\mathbb{Q}} [p], [q]$. $[r] \leq_{\mathbb{Q}} [p]$ asserts that there is a common extension $r' \leq_{\mathbb{P}} r, p$, and moreover $[r] \leq_{\mathbb{Q}} [q]$ asserts that there is a common extension $r'' \leq_{\mathbb{P}} r', q$. Therefore, r'' witnesses that $p \not\leq q$. \square

Definition II.2.5. Let \mathbb{P} be a forcing notion, θ be a cardinal and M be a ctm:

- 1. \mathbb{P} has θ -cc provided that the size of antichain in \mathbb{P} is less than θ ,
- 2. \mathbb{P} *preseves cofinalities* $\leq \theta$ provided that $M[G] \models \text{cf}(\gamma) = \text{cf}^M(\gamma)$ holds for every limit $\gamma \in o(M)$ with $\text{cf}^M(\gamma) \geq \theta$,
- 3. \mathbb{P} *preseves cardinals* $\leq \theta$ provided that $M \models \text{“} \beta \text{ is a cardinal. ”}$ iff $M[G] \models \text{“} \beta \text{ is a cardinal. ”}$ for $\theta \leq \beta \in o(M)$.

Lemma II.2.6. Let $\mathbb{P} \in M$ be a forcing notion with $M \models \text{“} \theta \text{ is a regular cardinal. ”}$ then we have:

- 1. \mathbb{P} *preseves cofinalities* $\geq \theta$ iff for arbitrary limit β with $\theta \leq \beta \in o(M)$ we have that $M \models \text{“} \beta \text{ is regular. ”}$ implies $M[G] \models \text{“} \beta \text{ is regular. ”}$,
- 2. If \mathbb{P} *preseves cofinalities* $\leq \theta$, so does *cardinals* $\leq \theta$.

Theorem II.2.7. Let $\mathbb{P} \in M$ and assume that $M \models \text{“} \theta \text{ is a regular cardinal and } \mathbb{P} \text{ is } \theta\text{-cc. ”}$. Then \mathbb{P} *preseves cofinalities* $\geq \theta$.

Proof. By the previous lemma, it suffice to show that suppose that there is $\theta \leq \beta \in o(M)$ such that $M \models \text{“} \beta \text{ is regular. ”}$ and $M[G] \models \text{“} \beta \text{ is singular. ”}$, a contradiction.

Let β be such an ordinal. Let X be a set in $M[G]$ and a function $f: \alpha \rightarrow X$ in $M[G]$ with $\alpha \in \beta$ and $p \Vdash_{\mathbb{P}} \text{“} \check{f}: (\alpha, \in) \approx (X, \in) \text{ and } \cup X = \beta. \text{”}$ for some condition p . Define $F: \alpha \rightarrow \mathcal{P}(\beta)$ which assign the ξ the $\{\eta \in \beta; \exists q \leq p (q \Vdash_{\mathbb{P}} \check{f}(\xi) = \eta)\}$ then $F \in M$. Now we assrt that $f(\xi) \in F(\xi)$ and $|F(\xi)| < \theta$ for arbitrary $\xi \in \alpha$. The former is immediate. For the later, for $\eta \in F(\xi)$ choose $p_\eta \leq p$ such that $p_\eta \Vdash_{\mathbb{P}} \check{f}(\xi) = \eta$. Then note that $p_\eta, \eta \in F(\xi)$, is pairwise incompatible, thus we obtain that $|F(\xi)| < \theta$. To conclude the proof, let $Y = \cup\{F(\xi); \xi \in \alpha\}$, then we obtain that $Y \subset \beta$ and $\cup Y = \beta$. However since $|F(\xi)| < \beta$, $\alpha < \theta$ and β is regular, we have $|Y| < \beta$. This contrary to that $\beta = \text{type}(Y)$. \square

Definition II.2.8 ([Pal13]). A *dominating real* in a generic extension is a real $x \in {}^\omega\omega$ which eventually dominates all funcions $f \in {}^\omega\omega$ in V . Similarly a *unbounded real* in a generic extension is a real $x \in {}^\omega\omega$ which does not eventually dominated by any function $f \in {}^\omega\omega$ in V .

II.3 Iterated Forcing

Definition II.3.1 ([Kun11]). Let \mathbb{P} be a forcing notion. The \mathbb{P} -name for a forcing notion is a triple pair of \mathbb{P} -names, $(\dot{\mathbb{Q}}, \dot{\leq}_{\mathbb{Q}}, \dot{1}_{\mathbb{Q}})$, such that $\dot{1}_{\mathbb{Q}} \in \text{dom}(\dot{\mathbb{Q}})$ and for any conditions $p \in \mathbb{P}$ force:

- $\dot{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \dot{1}_{\mathbb{Q}} \in \dot{\mathbb{Q}}$, and
- $\dot{\leq}_{\mathbb{Q}}$ is a pre-order of $\dot{\mathbb{Q}}$ with the largest element $\dot{1}_{\mathbb{Q}}$.

We write $\dot{\mathbb{Q}}$ for $(\dot{\mathbb{Q}}, \dot{\leq}_{\mathbb{Q}}, \dot{1}_{\mathbb{Q}})$, $\dot{\leq}$ for $\dot{\leq}_{\mathbb{Q}}$ and $\dot{1}$ for $\dot{1}_{\mathbb{Q}}$.

Definition II.3.2 ([Kun11]). Let \mathbb{P} be a forcing notion and $(\dot{\mathbb{Q}}, \dot{\leq}_{\mathbb{Q}}, \dot{1}_{\mathbb{Q}})$ be a \mathbb{P} -name for a forcing notion, define the *product* $\mathbb{P} * \dot{\mathbb{Q}}$ is the triple pair $(\mathbb{R}, \leq, \mathbb{1})$ such that;

1. $\mathbb{R} = \{(p, \dot{q}) \in \mathbb{P} \times \text{dom } \dot{\mathbb{Q}}; p \Vdash \dot{q} \in \dot{\mathbb{Q}}\}$,
2. $\leq = \{ \langle (p_0, \dot{q}_0), (p_1, \dot{q}_1) \rangle; p_0 \leq_{\mathbb{P}} p_1 \wedge p_0 \Vdash (\dot{q}_1 \leq_{\dot{\mathbb{Q}}} \dot{q}_2) \}$,
3. $\mathbb{1} = (\dot{1}_{\mathbb{P}}, \dot{1}_{\dot{\mathbb{Q}}})$,

We simply write $(p, \dot{q}) \in \mathbb{P} * \dot{\mathbb{Q}}$ for $(p, \dot{q}) \in \mathbb{R}$ and define $i: \mathbb{P} \rightarrow \mathbb{P} * \dot{\mathbb{Q}}$ which assign the p the $(p, \dot{1}_{\dot{\mathbb{Q}}})$.

Theorem II.3.3. Using the notion of **Definition II.3.2**, with $p_0, p_1 \in \mathbb{P}$ and $\dot{q}_0, \dot{q}_1 \in \dot{\mathbb{Q}}$. Then we have the following facts:

1. $\mathbb{P} * \dot{\mathbb{Q}}$ is a forcing notion,
2. $p_0 \leq_{\mathbb{P}} p_1$ iff $i(p_0) \leq_{\mathbb{P} * \dot{\mathbb{Q}}} i(p_1)$,
3. $i(\dot{1}_{\mathbb{P}}) = \mathbb{1}_{\mathbb{P} * \dot{\mathbb{Q}}}$,
4. $p_0 \perp_{\mathbb{P}} p_1$ implies $(p_0, \dot{q}_0) \perp_{\mathbb{P} * \dot{\mathbb{Q}}} (p_1, \dot{q}_1)$ if $(p_0, \dot{q}_0), (p_1, \dot{q}_1) \in \mathbb{P} * \dot{\mathbb{Q}}$,
5. $p_0 \perp_{\mathbb{P}} p_1$ iff $(p_0, \dot{1}_{\dot{\mathbb{Q}}}) \perp_{\mathbb{P} * \dot{\mathbb{Q}}} (p_1, \dot{q}_1)$ whenever $(p_1, \dot{q}_1) \in \mathbb{P} * \dot{\mathbb{Q}}$,
6. $p_0 \perp_{\mathbb{P}} p_1$ iff $i(p_0) \perp_{\mathbb{P} * \dot{\mathbb{Q}}} i(p_1)$,
7. i is a complete embedding.

Definition II.3.4. Let G be \mathbb{P} -generic over M and H be subset of $\dot{\mathbb{Q}}_G$. Then $G * H = \{(p, \dot{q}) \in \mathbb{P} * \dot{\mathbb{Q}}; p \in G \wedge \dot{q}_G \in H\}$.

Theorem II.3.5. Let K be a $\mathbb{P} * \dot{\mathbb{Q}}$ -generic over M . Define $G = i^{-1}(K)$ and let $H = \{\dot{q}_G; \exists p \in \mathbb{P} (p, \dot{q}) \in K\}$. Then we have the following:

1. G is \mathbb{P} -generic over M ,
2. H is $\dot{\mathbb{Q}}_G$ -generic over $M[G]$,
3. $K = G * H$,

4. $M[K] = (M[G])[H]$.

Proof. For (1). To see that G is a filter. Let $p, q \in G$. Let $(r, \dot{r}') \in K$ such that $(r, \dot{r}') \leq_{\mathbb{P} * \dot{\mathbb{Q}}} i^{-1}(p), i^{-1}(q)$. Thus we have $r \leq_{\mathbb{P}} p, q$ and $r \in G$. To see that meets every dense subset. Let D be a dense subset of \mathbb{P} in M . By letting $D' = \{(p, \dot{q}); p \in D \wedge \dot{q} \in \dot{\mathbb{Q}}_G\}$, D' is dense over $\mathbb{P} * \dot{\mathbb{Q}}$. Therefore there is $(p, \dot{q}) \in D' \cap K$, furthermore, $p \in D \cap G$.

For (2). To see that H is a filter. Let $(\dot{q}_i)_G \in H$ with $p_i \in G$ such that $(p_i, \dot{q}_i) \in K$ for $i \in 2$. Then there is a common extension $(r, \dot{r}) \in K$. Then we have $r \Vdash_{\mathbb{P}} \dot{r} \leq_{\dot{\mathbb{Q}}} \dot{q}_0$ and $\dot{r} \leq_{\dot{\mathbb{Q}}} \dot{q}_1$. Moreover since $r \in G$, $M[G] \models \dot{r}_G \leq_{\dot{\mathbb{Q}}_G} (\dot{q}_0)_G$ and $\dot{r}_G \leq_{\dot{\mathbb{Q}}_G} (\dot{q}_1)_G$. To see that H meets every dense subset. Let D be dense subset of $\dot{\mathbb{Q}}_G$ in $M[G]$. By letting $D' = \{(p, \dot{q}); p \in G \wedge \dot{q}_G \in H\}$, then since D' is dense over $\mathbb{P} * \dot{\mathbb{Q}}$, there is $(p, \dot{q}) \in D' \cap K$. Thus we obtain that $\dot{q}_G \in D \cap H$.

For (3). To see that (\subset) . Let $(p, \dot{q}) \in K$. By the definition we have $p \in G$ and $\dot{q}_G \in H$ thus $(p, \dot{q}_G) \in G * H$. To see that (\supset) . Let $p \in G$ and $\dot{q}_G \in H$ with $(p', \dot{q}) \in K$ for some $p' \in \mathbb{P}$. Since $(p, \dot{1}_{\dot{\mathbb{Q}}}) \in K$, there is a common extension $(p'', \dot{q}'') \leq (p, \dot{1}_{\dot{\mathbb{Q}}}), (p', \dot{q})$. Then we obtain that $p'' \leq_{\mathbb{P}} p$ and $p'' \Vdash_{\mathbb{P}} \dot{q}'' \leq_{\dot{\mathbb{Q}}} (p, \dot{q})$. This shows that $(p, \dot{q}) \in K$.

For (4). To see (\subset) , by the minimality of extension, it suffices to show that $G * H \in (M[G])[H]$. By letting $\Gamma_{\mathbb{P} * \dot{\mathbb{Q}}} = \{(\langle p, \dot{q} \rangle, \dot{q}_G); \dot{q}_G \in H\}$, we have $G * H = (\Gamma_{\mathbb{P} * \dot{\mathbb{Q}}})_K \in (M[G])[H]$. To see (\supset) , it suffices to show that $G \in M[K]$ and $H \in M[K]$. For the former, by letting $\Gamma_G = \{(\langle p, (p, \dot{1}_{\dot{\mathbb{Q}}}) \rangle); p \in G\}$, we have $G = (\Gamma_G)_K \in M[K]$. For the later, by letting $\Gamma_H = \{(\langle \dot{q}_G, \langle \dot{1}_{\mathbb{P}}, \dot{q}_G \rangle \rangle); \dot{q}_G \in H\}$, we have $H = (\Gamma_H)_K \in M[K]$. \square

Definition II.3.6. For ordinal α , an α -stage iterated forcing is a pair $((\mathbb{P}_\xi, \leq_\xi, \dot{1}_\xi); \xi \leq \alpha), (\langle \dot{\mathbb{Q}}_\xi, \leq_{\dot{\mathbb{Q}}_\xi}, \dot{1}_{\dot{\mathbb{Q}}_\xi} \rangle; \xi < \alpha)$ with following properties:

- I-1 Every $(\mathbb{P}_\xi, \leq_\xi, \dot{1}_\xi)$ is a forcing notion,
- I-2 Every $(\dot{\mathbb{Q}}_\xi, \leq_{\dot{\mathbb{Q}}_\xi}, \dot{1}_{\dot{\mathbb{Q}}_\xi})$ is a $(\mathbb{P}_\xi, \leq_\xi, \dot{1}_\xi)$ -name for a forcing notion,
- I-3 Every $p \in \mathbb{P}_\xi$ is a sequence of the form $\langle \dot{q}_\mu; \mu < \xi \rangle$ where each $\dot{q}_\xi \in \text{dom}(\dot{\mathbb{Q}}_\mu)$,
- I-4 $\xi < \eta$ and $p \in \mathbb{P}_\eta$ implies $p|_\xi \in \mathbb{P}_\xi$,
- I-5 Let $\xi < \eta$, $p \in \mathbb{P}_\xi$ and p' be an η -sequence such that $p'|_\xi = p$ and $(p')|_\mu = \dot{1}_{\dot{\mathbb{Q}}}$, then $\xi \leq \mu < \eta$ implies $p' \in \mathbb{P}_\eta$ and we write $i_\xi^\eta(p)$ for p' ,
- I-6 $\dot{1}_\xi$ is the sequence $\langle \dot{q}_\mu; \mu < \xi \rangle$, where each $\dot{q}(\mu) = \dot{1}_{\dot{\mathbb{Q}}_\mu}$,
- I-7 For $p = \langle \dot{q}_\mu; \mu < \xi \rangle \in \mathbb{P}_\xi$ and $p' = \langle \dot{q}'_\mu; \mu < \xi \rangle \in \mathbb{P}_\xi$. $p \leq_\xi p'$ iff $p|_\mu \Vdash_{\mathbb{P}_\mu} \dot{q}(\mu) \leq_{\dot{\mathbb{Q}}_\mu} \dot{q}'(\mu)$ for all $\mu < \xi$,
- I-8 $\mathbb{P}_{\xi+1} = \{p \frown \langle \dot{q} \rangle; p \in \mathbb{P}_\xi \wedge \dot{q} \in \text{dom}(\dot{\mathbb{Q}}_\xi) \wedge p \Vdash_{\mathbb{P}_\xi} \dot{q} \in \dot{\mathbb{Q}}_\xi\}$ for every $\xi < \alpha$.

Definition II.3.7 ($<\kappa$ -support iteration). For a sequence p with length ξ , the *support* of p is

$$\text{supt}(p) = \{\mu < \xi; (p)_\mu \neq \mathbb{1}_{\mathbb{Q}_\mu}\}.$$

For an infinite cardinal κ , the iteration is $<\kappa$ -support provided that for all limit $\eta(\leq \alpha)$,

$$\mathbb{P}_\eta = \{p; \text{"}p \text{ is a sequence of length } \eta \text{"} \wedge |\text{supt}(p)| < \kappa \wedge \forall \xi < \eta (p|_\xi \in \mathbb{P}_\xi)\}.$$

A finite support iteration is $<\aleph_0$ -support and a countable support iteration is $<\aleph_1$ -support iteration.

Definition II.3.8. For a limit ordinal α . \mathbb{P}_α is an *inverse limit* of $\langle \mathbb{P}_\beta; \beta \in \alpha \rangle$ provided that

$$\forall p(p \in \mathbb{P}_\alpha \leftrightarrow \forall \beta \in \alpha p|_\beta \in \mathbb{P}_\beta).$$

\mathbb{P}_α is a *direct limit* provided that

$$\forall p(p \in \mathbb{P}_\alpha \leftrightarrow \exists \beta \in \alpha (p|_\beta \in \mathbb{P}_\beta \wedge \forall \xi \in [\beta, \alpha) p(\xi) = \mathbb{1}_{\mathbb{Q}_\xi}).$$

Theorem II.3.9 ([Jec03]). Let \mathbb{P}_α be an iterated forcing and $\beta \in \alpha$ a limit ordinal. Then we have:

1. \mathbb{P}_β is a set of finite support iff
 - (1-a) \mathbb{P}_β is a direct limit.
2. \mathbb{P}_β is a set of countable support iff
 - (2-a) If $\text{cf}(\beta) = \omega$, \mathbb{P}_β is an inverse limit,
 - (2-b) If $\text{cf}(\beta) \geq \omega$, \mathbb{P}_β is a direct limit.

Proof. At the beginning of the proof, We write \mathbb{P}'_γ for an inverse limit or a direct limit.

To see the statement 1, note that $\mathbb{P}_\gamma \subset \mathbb{P}'_\gamma$ and $\mathbb{P}'_\omega \subset \mathbb{P}_\omega$ hold. To see that $\mathbb{P}'_\gamma \subset \mathbb{P}_\gamma$ for $\gamma \geq \omega$, let $p \in \mathbb{P}'_\gamma$ and let $\beta \in \gamma$ such that $p(\xi) = \mathbb{1}_{\mathbb{Q}_\xi}$ for each $\xi \in [\beta, \gamma)$. Let δ be a limit ordinal and n a finite ordinal such that $\beta = \delta + n$. Then we obtain that $\text{supt}(p) \subset \text{supt}(p|_\delta) \cup [\delta, \delta + n)$ and this is a finite set by induction hypothesis.

To see the statement 2, we shall verify by the induction on γ and note that the case of $\gamma = \omega$ is obvious.

Case 2-a. $\omega \in \gamma$ with $\text{cf}(\gamma) = \omega$. To see the $(\mathbb{P}_\gamma \subset \mathbb{P}'_\gamma)$. Let $p \in \mathbb{P}_\gamma$. Since for arbitrary $\beta \in \gamma$ we have $\text{supt}(p|_\beta) \subset \text{supt}(p)$ is countable. Thus by induction hypothesis, we have that $p|_\beta \in \mathbb{P}'_\beta = \mathbb{P}_\beta$. To see the $(\mathbb{P}_\gamma \supset \mathbb{P}'_\gamma)$. Let $p \in \mathbb{P}'_\gamma$ and $\langle \xi_j; j \in \omega \rangle$ with $\xi_j \nearrow \gamma$. Note that $i(p|_{\xi_j}) \rightarrow p$ where $i(p|_{\xi_j}) \in \mathbb{P}_\gamma$ which eventually assigns $\mathbb{1}$. Then we obtain that $\text{supt}(p) = \cup \{\text{supt}(p|_{\xi_j}); j \in \omega\}$ and by induction hypothesis this is a countable set. Thus we have $p \in \mathbb{P}_\gamma$.

Case 2-b. γ with $\text{cf}(\gamma) \geq \omega$. To see $(\mathbb{P}_\gamma \subset \mathbb{P}'_\gamma)$. Let $\langle \xi_\alpha; \alpha \in |\text{cf}(\gamma)| \rangle$ with $\xi_\alpha \nearrow \gamma$. For $p \in \mathbb{P}_\gamma$ there is $\alpha \in |\text{cf}(\gamma)|$ such that $\cup \text{supt}(p) \in \xi_\alpha$ and this

α witnesses that $p \in \mathbb{P}'_\gamma$. To see $(\mathbb{P}_\gamma \supset \mathbb{P}'_\gamma)$. Let $p \in \mathbb{P}'_\gamma$ and $\beta \in \gamma$ such that $p|_\beta \in \mathbb{P}_\beta$ and $p(\xi) = \dot{1}_{\dot{\mathbb{Q}}_\xi}$ for $\xi \in [\beta, \gamma)$. Then by induction hypothesis, we obtain that $\text{supt}(p)$ is countable. \square

Definition II.3.10 (Intermediate stage, [Bau83]). Let $p \in \mathbb{P}_\alpha$. We define the followings:

1. $p^\beta = p|_{\{\gamma \in \alpha; \beta \leq \gamma\}}$ for $\beta \in \alpha$,
2. For $\beta \in \alpha$, $\mathbb{P}_{\beta\alpha} = \{p^\beta; p \in \mathbb{P}_\alpha\}$,
3. For \mathbb{P}_β -generic G_β , $f \leq g$ over $\mathbb{P}_{\beta\alpha}$ provided that there is $p \in G_\beta$ such that $p \cap f \leq_{\mathbb{P}_\alpha} p \cup g$.

$\mathbb{P}_{\beta\alpha}$ is a forcing notion with order in 3.

III Type of Forcings

III.1 ccc Forcing

Definition III.1.1. For a cardinal κ , a forcing notion is $< \kappa$ -cc forcing provided that there are no antichains of size κ . A forcing notion is ccc provided that it is a \aleph_1 -cc.

Example III.1.2.

1. Cohen forcing is a ccc forcing. (see [Theorem IV.1.2](#)),
2. σ -centerd forcing is ccc forcing.

Theorem III.1.3 ([Kun11]). ccc forcing preseves cofinalities.

Proof. [IV.3.4](#) \square

Theorem III.1.4 ([Jec03]). Let κ be an uncountable regular cardinal and \mathbb{P}_α be a finite support iteration. If \Vdash_β “ $\dot{\mathbb{Q}}_\beta$ is $< \kappa$ -cc.”, for each $\beta \in \alpha$, so does \mathbb{P}_α .

Proof. We shall show by induction on α .

Case I. $\alpha = \gamma + 1$. Let $A = \{p_\xi \widehat{\langle \dot{q}_\xi \rangle}; \xi \in \kappa\}$ be a subset of \mathbb{P}_α of size κ such that $p_\xi \in \mathbb{P}_\gamma$ and $\dot{q}_\xi \in \dot{\mathbb{Q}}_\gamma$, respectively for each $\xi \in \kappa$. Since κ is a regular and $\mathbb{P}_\gamma \models$ is $< \kappa$ -cc, there is $B \in [A]^\kappa$ such that for any $\xi, \eta \in B$ $p_\xi = p_\eta$. Thus we obtain a κ size subset $\{\dot{q}_\xi; \xi \in B\}$ of $\dot{\mathbb{Q}}_\gamma$. Moreover, since \Vdash_γ “ $\dot{\mathbb{Q}}_\gamma$ is $< \kappa$ -cc.”, there are distinct ξ and η which is compatible. Let $\dot{r} \in \dot{\mathbb{Q}}_\gamma$ be a common extension. Then we obtain a common $p_\xi \widehat{\langle \dot{r} \rangle}$ for $p_\xi \widehat{\langle \dot{q}_\xi \rangle}$ and $p_\eta \widehat{\langle \dot{q}_\eta \rangle}$. This shows that A is not an antichain, i.e., \mathbb{P}_α is $< \kappa$ -cc.

Case II-i. α is limit and $\text{cf}(\alpha) \geq \kappa$. Let $A = \{p_\xi \in \mathbb{P}_\alpha; \xi \in \kappa\}$ be a subset of size κ . Since \mathbb{P}_α is a finite support iteration, there is a $\beta \in \alpha$ such that

$$\forall \xi \in \kappa \forall \eta \in [\beta, \alpha) \ p_\xi(\eta) = \dot{1}_{\dot{\mathbb{Q}}_\eta}$$

moreover, we obtain that

$$\forall \xi \in \kappa \forall \eta \in [\beta, \alpha) \ p_\xi(\eta) = \dot{1}_{\mathbb{Q}_\eta}$$

for a limit ordinal $\beta \in \alpha$. Now by induction hypothesis, there are distinct $\xi, \xi' \in \kappa$ and $p \in \mathbb{P}_\beta$ such that $p \leq_\beta p_\xi|_\beta, p_{\xi'}|_\beta$. Furthermore, $p \wedge \langle \dot{1} \rangle \wedge \dots \wedge \langle \dot{1} \rangle \leq_\alpha p_\xi, p_{\xi'}$. Therefore, A is not an antichain, i.e., \mathbb{P}_α is $< \kappa$ -cc.

Case II–ii. α is limit and $\text{cf}(\alpha) = \kappa$. Let $A = \{p_\xi; \xi \in \kappa\}$ be a subset of size κ and $\langle \alpha_\xi; \xi \in \kappa \rangle$ be a sequence with following properties:

- $\cup\{\text{supt}(p_\eta); \eta \in \xi\} \subset \alpha_\xi$ for all $\xi \in \kappa$,
- $\cup\{\alpha_\xi; \xi \in \kappa\} = \alpha$.

Then for each limit $\xi \in \kappa$, since $\text{supt}(p_\xi)$ is finite and $\cup\{\text{supt}(p_\eta); \eta \in \xi\} \subset \alpha_\xi$, there is $\gamma_\xi \in \xi$ such that $\text{supt}(p_\xi) \cap \alpha_\xi \subset \alpha_{\gamma_\xi}$.

Since the set $\kappa \cap \text{Lim}$ is a stationary subset of κ , applying [Theorem I.1.4](#), there is a stationary subset $S \subset C$ of size κ and $\gamma \in \kappa$ such that $\text{supt}(p_\xi) \cap \alpha_\xi \subset \alpha_\gamma$ for any $\xi \in S$ and thus we obtain a subset $\{p_\xi|_\gamma; \xi \in S\} \subset \mathbb{P}_\gamma$. Now induction hypothesis and κ is regular assert that there are ξ and η such that $\gamma \in \xi \in \eta$ and $p_\xi|_\gamma$ and $p_\eta|_\gamma$ possesses a common extension, named $p \in \mathbb{P}_\gamma$.

Define an r by:

$$r(\beta) = \begin{cases} p(\beta) & \beta \in \alpha_\gamma \\ p_\xi(\beta) & \beta \in [\alpha_\gamma, \alpha_\eta) \\ p_\eta(\beta) & \beta \in [\alpha_\eta, \alpha) \end{cases}$$

then obviously we have $r \in \mathbb{P}_\alpha$. To see that r is an extension for both p_ξ and p_η , the former is derived by the fact $\eta \in \xi$ and the later is derived by the fact $\eta \in S$. Therefore, A possesses a compatible pair, i.e., \mathbb{P}_α is $< \kappa$ -cc. \square

III.2 σ -centerd Forcing

Definition III.2.1. Let \mathbb{P} be a forcing notion and C a subset. C is *centered* provided that for any finite subset of C there is a common extension $p \in \mathbb{P}$. A forcing notion \mathbb{P} is *σ -centered* provided that there are countably many centered subsets $C_i \subset \mathbb{P}$, $i \in \omega$, whose union is \mathbb{P} .

Example III.2.2.

1. Hechler forcing is σ -centered forcing. (see [Theorem IV.2.4](#))

III.3 σ -closed Forcing

Definition III.3.1. A forcing notion \mathbb{P} is a *σ -closed forcing* (or ω_1 -closed, *countably closed*) provided that for every decreasing sequence $\langle p_n \in \mathbb{P}; n \in \omega \rangle$ with $p_{n+1} \leq p_n$ there is a condition $p \in \mathbb{P}$ such that $p \leq p_n$ for each $n \in \omega$.

III.4 Axiom A Forcing

Definition III.4.1 ([Bau83]). For a forcing notion \mathbb{P} , \mathbb{P} satisfies Axiom A provided that there is an order sequence $\{\leq_n ; n \in \omega\}$ with following properties:

- A-1** $\leq_{\mathbb{P}} = \leq_0$,
- A-2** For every $n \in \omega$ and $p, q \in \mathbb{P}$, $q \leq_{n+1} p \implies p \leq_n q$,
- A-3** For every sequence $\langle p_n \in \mathbb{P} ; n \in \omega \rangle$ with $p_{n+1} \leq_n p_n$ for each $n \in \omega$, there is $q \in \mathbb{P}$ such that $p \leq_n p_n$ for every $n \in \omega$. We call that the sequence is a *fusion sequence* and q is a *fusion*,
- A-4** Let I be an antichain over \mathbb{P} , $p \in \mathbb{P}$ and $n \in \omega$, there is $q \in \mathbb{P}$ such that $q \leq_n p$ and $\{r \in I ; r \not\leq q\}$ is countable.

Example III.4.2.

1. A ccc forcing satisfies Axiom A.
2. A σ -closed forcing satisfies Axiom A.
3. A Sacks forcing satisfies Axiom A. (see [Theorem IV.4.3](#))
4. A Mathias forcing satisfies Axiom A. (see [Theorem IV.3.7](#))
5. A shooting a club by finite conditions forcing does not satisfy Axiom A. (see [Theorem IV.5.4](#))

Theorem III.4.3. The following is equivalent to **A-4**:

- A-4'** Let \dot{a} be a \mathbb{P} -name, $p \in \mathbb{P}$ and $n \in \omega$. If $p \Vdash_{\mathbb{P}} \dot{a} \in V$, equivalently $\exists b \in V (p \Vdash_{\mathbb{P}} \dot{a} \in \check{b})$, then there is $X \in V$ and $q \in \mathbb{P}$ such that X is countable, $q \leq_n p$ and $q \Vdash_{\mathbb{P}} \dot{a} \in \check{X}$.

Proof. For **A-4** \implies **A-4'**. Assume $p \Vdash_{\mathbb{P}} \dot{a} \in V$ and let $n \in \omega$. Since $\{p \in \mathbb{P} ; \exists b \in V (p \Vdash_{\mathbb{P}} \dot{a} \in \check{b})\}$ is not empty let I be a maximal antichain of it and for $r \in I$ let $a_r \in V$ be the unique element which satisfies $r \Vdash_{\mathbb{P}} \dot{a} = \check{a}_r$. Applying **A-4**, there is $q \leq_n p$ such that $\{r \in I ; r \not\leq q\}$ is countable. By letting $X = \{a_r ; r \in I \wedge r \not\leq q\}$, X is countable. Finally we check $q \Vdash_{\mathbb{P}} \dot{a} \in \check{X}$. For arbitrary $q_1 \leq_{\mathbb{P}} q$ there is $r \in I$ such that $r \not\leq_{\mathbb{P}} q_1$, moreover there is $q_2 \in \mathbb{P}$ such that $q_2 \leq_{\mathbb{P}} q_1, r$. Then we obtain $q_2 \Vdash_{\mathbb{P}} \dot{a} = \check{a}_r$.

For **A-4'** \implies **A-4**. Let I be an antichain, $p \in \mathbb{P}$ and $n \in \omega$ and let I' be a maximal antichain finer than I . By letting $\dot{a} = \{(\check{r}, r) ; r \in I'\}$, for any generic G we have $(\dot{a})_G \in I'$, so we obtain $p \Vdash_{\mathbb{P}} \dot{a} \in V$. By **A-4'** there is a countable $X \in V$ and $q \in \mathbb{P}$ such that $q \leq_n p$ and $q \Vdash_{\mathbb{P}} \dot{a} \in \check{X}$. Now we show that $\{r \in I' ; r \not\leq q\} \subset X$. Let $r \in I'$ with $r \perp_{\mathbb{P}} q$. By $q \Vdash_{\mathbb{P}} \dot{a} \in \check{X}$ there is q' and $x \in X$ such that $q' \Vdash_{\mathbb{P}} \dot{a} = \check{x}$. Then for any generic G with $q' \in G$, since $r \in G$, we obtain that $r = (\dot{a})_G = (\check{x})_G = x$. Thus, we have the subset relation and we complete the proof. \square

Theorem III.4.4. Let \mathbb{P}_α is an α -stage iteration over M and $\beta \in \alpha$. Let $i: \mathbb{P}_\alpha \rightarrow \mathbb{P}_\beta$ which assign the p the $p \frown \langle \mathbb{1} \rangle \frown \dots \frown \langle \mathbb{1} \rangle \in \mathbb{P}_\alpha$. For \mathbb{P}_α -generic K , define $G = i^{-1}(K)$ and $H = \{p^\beta; p \in K\}$. Then we have the followings:

1. G is \mathbb{P}_β -generic over M ,
2. H is $\mathbb{P}_{\beta\alpha}$ -generic over $M[G]$,
3. $K = G \frown H = \{p \frown q; p \in G \wedge q \in H\}$,
4. $M[K] = (M[G])[H]$.

Theorem III.4.5. Let \mathbb{P}_α is an α -stage iteration with countable support which satisfies $\mathbb{1} \Vdash_{\mathbb{P}_\beta} \text{“}\dot{\mathbb{Q}}_\beta \text{ satisfies Axiom A.”}$ for all $\beta \in \alpha$. Then \mathbb{P}_α does not collapse ω_1 , i.e., preserves ω_1 . And moreover if we have $\alpha < \omega_2$, $V \models \text{ZFC} + \text{CH}$ and $\Vdash_{\mathbb{P}_\beta} |\dot{\mathbb{Q}}_\beta| \leq \aleph_1$ for all $\beta \in \alpha$, then \mathbb{P}_α has an \aleph_2 -chain condition.

Before we show the theorem, we make some definition and lemmata. (See the proof.)

Definition III.4.6. For $\gamma \leq \alpha$, $F \in [\alpha]^{<\omega}$ and $n \in \omega$, we define the followings:

- (i) For $p, q \in \mathbb{P}$, $p \leq_{F,n} q$ provided that $\forall \beta \in F (p|_\beta \Vdash_{\mathbb{P}_\beta} p(\beta) \leq_n^{\circ} q(\beta))$.
- (ii) $D \subset \mathbb{P}_\gamma$ is (F, n) -dense provided that $\forall p \in \mathbb{P}_\gamma \exists q \in D (q \leq_{F,n} p)$.
- (iii) A sequence $\langle (p_n, F_n); n \in \omega \rangle$ is an (F, n) -fusion sequence of \mathbb{P}_α provided that
 - (iii-a) $p_n \in \mathbb{P}_\alpha$ and $F_n \in [\alpha]^{<\omega}$ for each $n \in \omega$,
 - (iii-b) $\forall n \in \omega (p_{n+1} \leq_{F_n, n} p_n)$,
 - (iii-c) $\forall n \in \omega (F_n \subset F_{n+1})$,
 - (iii-d) $\cup\{F_n; n \in \omega\} = \cup\{\text{supt}(p_n); n \in \omega\}$.

Lemma III.4.7. Let $\langle (p_n, F_n); n \in \omega \rangle$ be an (F, n) -fusion sequence of \mathbb{P}_α . There is $p \in \mathbb{P}_\alpha$ such that $\forall n \in \omega (p \leq_{F_n, n} p_n)$. Moreover we may assume that $\text{supt}(p) = \cup\{F_n; n \in \omega\}$.

Proof. First we see the “moreover” part. Suppose there is the desired $q \in \mathbb{P}_\alpha$. By letting

$$q' = \{(\eta, q(\eta)); \eta \notin \cup\{\text{supt}(p_n); n \in \omega\}\} \cup \{(\eta, \dot{\mathbb{1}}_{\dot{\mathbb{Q}}_\eta}); \eta \in \cup\{\text{supt}(p_n); n \in \omega\}\}$$

we obtain that $\text{supt}(q) = \cup\{\text{supt}(p_n); n \in \omega\}$.

To see the first statement, we show the following by the induction on β .

$$\forall \beta \leq \alpha \forall \langle (p_n, F_n); n \in \omega \rangle \exists p \in \mathbb{P}_\beta \forall n \in \omega (p \leq_{F_n, n} p_n)$$

Where $\langle (p_n, F_n); n \in \omega \rangle$ is an (F, n) -fusion sequence of \mathbb{P}_β .

(i) $\beta = 0$. For an (F, n) -fusion sequence $\langle (p_n, F_n); n \in \omega \rangle$ of \mathbb{P}_0 , by letting $p = \emptyset$, we have $p \in \mathbb{P}_0$ and for arbitrary $n \in \omega$, $p \leq_{0,n} p_n$ is trivial.

(ii) $\beta = \gamma + 1$. Let $\langle (p_n, F_n); n \in \omega \rangle$ be an (F, n) -fusion sequence of $\mathbb{P}_{\gamma+1}$. By induction hypothesis, there is $p' \in \mathbb{P}_\gamma$ such that $\forall n \in \omega$ ($p' \leq_{F_n, n} p_n$) holds. We distinguish two cases, according to whether $\gamma \in \cup \{\text{supt}(p_n); n \in \omega\}$. If $\gamma \notin \cup \{\text{supt}(p_n); n \in \omega\}$, by letting $p = p' \cup \{(\gamma, \dot{1}_{\dot{\mathbb{Q}}_\gamma})\}$, we have $p \in \mathbb{P}_{\gamma+1}$ and $\forall n \in \omega$ ($p \leq_{F_n, n} p_n$) holds. If $\gamma \in \cup \{\text{supt}(p_n); n \in \omega\}$. Let n_0 be an \in -minimal such that $\gamma \in F_{n_0}$. For $m \in \omega$, define \dot{q}_m for $p_{n_0}(\gamma)$ if $m \in n_0$ and $p_m(\gamma)$ if $m \notin n_0$. Then we note that $\dot{1} \Vdash_{\mathbb{P}_\gamma} \langle \dot{q}_m; m \in \omega \rangle$ is a sequence of $\dot{\mathbb{Q}}_\gamma$." and we assert that $p' \Vdash_{\mathbb{P}_\gamma} \forall n \in \omega$ ($\dot{q}_{m+1} \leq_m^\gamma \dot{q}_m$). To see the later, for $m \in \omega$, we distinguish two cases, according to whether $m \in n_0$. If $m \notin n_0$. Since we have $\gamma \in F_{n_0} \subset F_m$, $\dot{q}_{m+1}(\gamma) = p_m(\gamma)$ and $\dot{q}_{m+1}(\gamma) = p_{m+1}(\gamma)$, by the condition (a) in an (F, n) -fusion sequence, we obtain that $p_{m+1} \upharpoonright_\gamma \Vdash_{\mathbb{P}_\gamma} p_{m+1}(\gamma) \leq_m^\gamma p_m(\gamma)$. Since $p' \leq_{F_{m+1}, m+1} p_{m+1} \upharpoonright_\gamma$ and $\gamma \in F_{n_0} \subset F_{m+1}$ hold, we have $\forall \delta < \gamma$ ($p' \Vdash_{\mathbb{P}_\delta} p'(\delta) \leq^\delta p_{m+1}(\delta)$), moreover, $p' \leq_{\mathbb{P}_\gamma} p_{m+1} \upharpoonright_\gamma$. Therefore we obtain that $p' \Vdash_{\mathbb{P}_\gamma} p_{m+1}(\gamma) \leq_m^\gamma p_m(\gamma)$. If $m \in n_0$. Since $\dot{q}_m(\gamma) = p_{n_0}(\gamma) = \dot{q}_{m+1}(\gamma)$, we have $p' \Vdash_{\mathbb{P}_\gamma} p_{m+1}(\gamma) \leq_m^\gamma p_m(\gamma)$. Now since $p' \Vdash_{\mathbb{P}_\gamma} \dot{\mathbb{Q}}_\gamma$ satisfies Axiom A.", by A-4', there is a $\dot{q} \in V^{\mathbb{P}_\gamma}$ such that $p' \Vdash_{\mathbb{P}_\gamma} \dot{q} \in \dot{\mathbb{Q}}_\gamma$ and $\forall m \in \omega$ ($\dot{q} \leq_m^\gamma \dot{q}_m$) holds. Thus by letting $p = p' \dot{\wedge} \dot{q}$, we obtain that $p \in \mathbb{P}_\beta$ and $\forall n \in \omega$ ($p \leq_{F_n, n} p_n$).

(iii) β is limit. Let $\langle (p_n, F_n); n \in \omega \rangle$ be an (F, n) -fusion sequence of \mathbb{P}_β . By induction hypothesis for every $\xi \in \beta$, choose $p'_\xi \in \mathbb{P}_\xi$ such that $\forall n \in \omega$ ($p'_\xi \leq_{F_n, n} p_n$). We distinguish two cases, according to whether $\omega \in \text{cf}(\beta)$. If $\omega \in \text{cf}(\beta)$. Since every F_n is finite there is $\eta \in \beta$ such that $\cup \{F_n; n \in \omega\} \subset \eta$. Since we have $\forall n \in \omega$ ($p_\eta \leq_{F_n, n} p_n$), we obtain that $\forall n \in \omega$ ($p_\eta \leq_{F_n, n} p_n$). If $\omega = \text{cf}(\beta)$. Let $\xi_n \in \beta$ such that $\xi_n \nearrow \beta$ and $\cup \{F_m; m \leq n\} \subset \xi_n$. First, for an (F, n) -fusion sequence $\langle (p_n|_{\xi_0}, \cap \{F_n, \xi_0\}); n \in \omega \rangle$ of \mathbb{P}_{ξ_0} , there is $q_0 \in \mathbb{P}_{\xi_0}$ such that $q_0 \leq_{\cap \{F_n, \xi_0\}, n} p_n|_{\xi_0}$ for every $n \in \omega$. Secondly, for $n \in \omega$, define $p_n^1 = q_0 \cup f|_{[\xi_0, \xi_1)}$ then we obtain an (F, n) -fusion sequence $\langle (p_n^1, \cap \{F_n, \xi_1\}); i \in \omega \rangle$, there is $q_1 \in \mathbb{P}_{\xi_1}$ such that $q_1 \leq_{\cap \{F_n, \xi_n\}, n} p_n^1$ for every $n \in \omega$. And by letting $q'_1 = q|_{\xi_0} \cup q_1|_{[\xi_0, \xi_1)}$, we have $q_0 \subset q'_1$. Continuing this process, we obtain $q_m \in \mathbb{P}_{\xi_m}$, $m \in \omega$, with following properties:

1. $q_0 \leq_{\cap \{F_n, \xi_0\}, n} p_n|_{\xi_0}$ holds for each $n \ni 1$,
2. $q_m \leq_{\cap \{F_n, \xi_n\}, i} p_n^m$ for every $n \in \omega$ holds for each $m \ni 0$, where $p_n^{m+1} = q_m \cup p_n|_{[\xi_m, \xi_{m+1})}$.

Let $q = \cup \{q_m; m \in \omega\}$. To conclude the proof we remain to show that $q \in \mathbb{P}_\beta$ and $q \leq_{F_n, n} p_n$ for every $n \in \omega$. The former is immediate by the fact \mathbb{P}_α is countable support and $q_m \subset q_{m+1}$ for every $m \in \omega$. To see the later statement. Let $n \in \omega$ and $\gamma \in F_n$. If $\gamma \in \xi_0$, we obtain $q \leq_{F_n, n} p_n$ immediately by $q_0 \subset q$. Thus we may assume that $\xi_0 \leq \gamma \in \beta$ and choose $m \in \omega$ such that $\gamma \in [\xi_{m-1}, \xi_m)$. Thus we obtain that $q_m \upharpoonright_\gamma \Vdash_{\mathbb{P}_\gamma} q_m(\gamma) \leq_n^\gamma p_n^m(\gamma)$, furthermore, by $q \supset q_m$ and $p_n^m(\gamma) = p_n(\gamma)$, we have that $q \upharpoonright_\gamma \Vdash_{\mathbb{P}_\gamma} q(\gamma) \leq_n^\gamma p_n(\gamma)$. Now we complete the proof of the limit case.

□

Lemma III.4.8. For $\beta \leq \alpha$ we have the followings:

- (a) If $\Vdash_{\mathbb{P}_\beta} \dot{a} \in V$, then the following set is (F, n) -dense for all finite $F \subset \beta$ and $n \in \omega$;

$$\{q \in \mathbb{P}_\beta; \exists X (\text{“ } X \text{ is countable in } V. \text{”} \wedge q \Vdash_{\mathbb{P}_\beta} \dot{a} \in \check{X})\},$$

- (b) If $\Vdash_{\mathbb{P}_\beta}$ “ \dot{X} is a countable subset of V . ”, then the following set is (F, n) -dense for all finite $F \subset \beta$ and $n \in \omega$;

$$\{q \in \mathbb{P}_\beta; \exists X (\text{“ } X \text{ is countable in } V. \text{”} \wedge q \Vdash_{\mathbb{P}_\beta} \dot{X} \subset \check{X})\},$$

- (c) If $\beta \in \gamma \leq \alpha$ and $\Vdash_{\mathbb{P}_\beta} \dot{f} \in \mathbb{P}_{\beta\gamma}$, then the following set is (F, n) -dense for all finite $F \subset \beta$ and $n \in \omega$;

$$\{q \in \mathbb{P}_\beta; \exists f \in \mathbb{P}_{\beta\gamma} (q \Vdash_{\mathbb{P}_\beta} \dot{f} = f)\}.$$

Proof. At the beginning of the proof, if $\beta \leq \alpha$ satisfies (a) (resp. (b), (c)), we say that (a) holds at a β -stage. To conclude the proof, thanks to induction, it suffices to show the followings:

1. (a) holds at a β -stage then so does (b),
2. (b) holds at a β -stage then so does (c),
3. (a) holds at the leading-stage, $\beta = 0$,
4. (b) holds at a β -stage then (a) holds at the $\beta + 1$ -stage,
5. For any limit $\delta \leq \alpha$, if (b) and (c) hold at a $\beta \in \delta$ then (a) holds at a δ -stage.

We shall show 1. Assume that $\Vdash_{\mathbb{P}_\beta}$ “ \dot{X} is countable in V . ” and $F \in [\beta]^{<\omega}$, $n \in \omega$ and $p \in \mathbb{P}_\beta$. Let $\dot{a}_n, n \in \omega$, be \mathbb{P}_β -names such that $\Vdash_{\mathbb{P}_\beta} \dot{X} = \{\dot{a}_n; n \in \omega\}$. We define an (F, n) -fusion sequence $\langle (p_k, F_k); k \in \omega \rangle$ of \mathbb{P}_β , recursively. Let $p_0 = p \in \mathbb{P}_\beta$ and $F_0 = F \cap \text{supt}(p_0)$. Since \mathbb{P}_α is countable support, let $\dot{\alpha}_i^0, i \in \omega$, such that $\{\dot{\alpha}_i^0; i \in \omega\} = \text{supt}(p_0)$. Assume that $p_k \in \mathbb{P}_\beta$ and F_k are defined. Applying (a) at a β -stage, there are $p_{k+1} \in \mathbb{P}_\beta$ and countable X_k in V such that $p_{k+1} \leq_{F_k, n+k} p_k$ and $p_{k+1} \Vdash_{\mathbb{P}_\beta} \dot{a}_k \in \check{X}_k$ and let $\dot{\alpha}_i^k, i \in \omega$, be \mathbb{P}_β -names such that $\{\dot{\alpha}_i^k; i \in \omega\} = \text{supt}(p_k)$. Then by letting $F_{k+1} = \cup\{F_k, \{\dot{\alpha}_i^k; i \in \omega\}\}$. Therefore we obtain an (F, n) -fusion sequence $\langle (p_k, F_k); k \in \omega \rangle$. Now by applying [Lemma III.4.7](#), there is $q \in \mathbb{P}_\beta$ such that $q \leq_{F_k, n+k} p_k$ for every $k \in \omega$. Moreover, we have $q \leq_{F_0, n} p$. Thus we obtain that $q \leq_{F, n} p$ and we obtain that $q \Vdash_{\mathbb{P}_\beta} \dot{X} \subset \cup\{\check{X}_k; k \in \omega\}$.

We shall show 2. Let γ be $\beta \in \gamma \leq \alpha$, $\Vdash_{\mathbb{P}_\beta} \dot{f} \in \mathbb{P}_{\beta\gamma}$, $F \in [\beta]^{<\omega}$, $n \in \omega$ and $p \in \mathbb{P}_\beta$. Since $\Vdash_{\mathbb{P}_\beta}$ “ $\text{supt}(p)$ is countable in V . ”, by applying (b) at a β -stage, there is $q \in \mathbb{P}_\beta$ and countable X in V such that $q \leq_{F, n} p$ and $q \Vdash_{\mathbb{P}_\beta} \text{supt}(p) \subset \check{X}$. To conclude the proof, we shall construct a function $f \in \mathbb{P}_{\beta\gamma}$ as follows:

- $f(\xi) = \mathbb{1}_{\dot{Q}_\xi}$ for all $\xi \notin X$

- $f(\xi)$ be a P_ξ -name such that $q \cup f|_\xi \Vdash_{\mathbb{P}_\xi} f(\xi) = \dot{f}(\xi)$ for all $\xi \in X$

Then since $\text{supt}(f)$ is countable we obtain $f \in \mathbb{P}_{\beta\gamma}$. We remain to show that $q \Vdash_\beta \dot{f} = f$ and it suffices to verify that $q \Vdash_\beta \forall \xi (\beta \leq \xi \wedge \xi \in \gamma \implies \dot{f}(\xi) = f(\xi))$. (i) If $\xi = \beta$. Since $q \cup f|_\xi = q$, we obtain that $q \Vdash_{\mathbb{P}_\beta} f(\xi) = \dot{f}(\xi)$. (ii) If ξ is successor or limit. We distinguish two cases, according to whether $\xi \in X$. If $\xi \notin X$. By the construction, we obtain that $q \Vdash_{\mathbb{P}_\beta} f(\xi) = \dot{f}(\xi)$. If $\xi \in X$. By induction hypothesis, for arbitrary η with $\beta \leq \eta \in \xi$, we obtain $q \Vdash_{\mathbb{P}_\beta} (f|_\eta \Vdash_{\mathbb{P}_\eta} f(\eta) = \dot{f}(\eta))$ and $q \Vdash_{\mathbb{P}_\beta} (\dot{f}|_\eta \Vdash_{\mathbb{P}_\eta} f(\eta) = \dot{f}(\eta))$. Thus we obtain that $q \Vdash_{\mathbb{P}_\beta} f(\xi) \leq_{\mathbb{P}_\xi} \dot{f}(\xi)$ and $q \Vdash_{\mathbb{P}_\beta} f(\xi) \geq_{\mathbb{P}_\xi} \dot{f}(\xi)$, furthermore, $q \Vdash_{\mathbb{P}_\beta} f(\xi) = \dot{f}(\xi)$.

We shall show 3. First we note that $\mathbb{P}_\beta = \{\emptyset\}$ and for any condition there are no extensions without it self. By letting $X = \{\dot{a}\} \subset V$ we obtain that $\emptyset \Vdash_{\mathbb{P}_0} \dot{a} \in \check{X}$. This shows (a) holds at the 0-stage.

We shall show 4. Assume $\Vdash_{\mathbb{P}_{\beta+1}} \dot{a} \in V$ and let $F \in [\beta+1]^{<\omega}$, $n \in \omega$ and $p \in \mathbb{P}_{\beta+1}$. Since we have $\mathbb{P}_{\beta+1} \text{ embbends } \mathbb{P}_\beta * \dot{Q}_{\beta+1}$ and $\Vdash_{\mathbb{P}_\beta} \dot{a} \in V$, we obtain that $\mathbb{1}_{\dot{Q}_{\beta+1}} \Vdash_{\mathbb{P}_{\beta+1}} (\mathbb{1}_{\dot{Q}_\beta} \Vdash_{\mathbb{P}_\beta} \dot{a} \in V)$ furthermore, $\mathbb{1}_{\dot{Q}_{\beta+1}} \Vdash_{\mathbb{P}_{\beta+1}} (p(\beta)_{\dot{Q}_\beta} \Vdash_{\mathbb{P}_\beta} \dot{a} \in V)$. Since, $\Vdash_{\mathbb{P}_\beta}$ “ \dot{Q} satisfies Axiom A.”, $\mathbb{1}_{\dot{Q}_{\beta+1}}$ forces “ $\exists q \leq_n^\beta p(\beta) \exists X$ (“ X is countable.” $\wedge q \Vdash_{\mathbb{P}_\beta} \dot{a} \in \check{X}$).” Moreover, by letting $X' = X \cap V$, since $\Vdash_{\mathbb{P}_{\beta+1}} \dot{a} \in V$, we can forces “ $\exists q \leq_n^\beta p(\beta) \exists X'$ (“ X' is countable in V .” $\wedge q \Vdash_{\mathbb{P}_\beta} \dot{a} \in \check{X}'$).” . Now by applying (b), there is countable Y in V and $q' \leq_{F,n} p|_\beta$ such that $q'' \Vdash_{\mathbb{P}_\beta} (\exists q \leq_n^\beta p(\beta) \wedge q \Vdash_{\dot{Q}_\beta} \dot{a} \in \check{Y})$. Then we obtain that $q'' \wedge \langle q \rangle \leq_{F,n} p$ and $q'' \wedge \langle q \rangle \Vdash_{\mathbb{P}_{\beta+1}} \dot{a} \in \check{Y}$.

We shall show 5. Let $\delta \leq \alpha$ be a limit. Assume that $\Vdash_{\mathbb{P}_\gamma} \dot{a} \in V$ and let $F \in [\delta]^{<\omega}$, $n \in \omega$ and $p \in \mathbb{P}_\delta$ and let γ be an ordinal with $F \subset \gamma \in \delta$. Since $\mathbb{P}_\delta \text{ embbends } \mathbb{P}_\gamma * \dot{\mathbb{P}}_{\gamma\delta}$ and $\Vdash_{\mathbb{P}_\gamma} \dot{a} \in V$, we have $\mathbb{1}_{\dot{Q}_\gamma} \Vdash_{\mathbb{P}_\gamma} (\mathbb{1}_{\mathbb{P}_{\gamma\delta}} \Vdash_{\mathbb{P}_{\gamma\delta}} \dot{a} \in V)$, moreover, $\mathbb{1}_{\dot{Q}_\gamma} \Vdash_{\mathbb{P}_\gamma} (p^\gamma \Vdash_{\mathbb{P}_{\gamma\delta}} \dot{a} \in V)$. Then we obtain that $\mathbb{1}_{\dot{Q}_\gamma}$ forces:

$$\text{there are } \mathbb{P}_{\gamma\delta}\text{-names } \dot{f} \leq_n^\gamma p^\gamma \text{ and } b \in V \text{ such that } \dot{f} \Vdash_{\mathbb{P}_{\gamma\delta}} \dot{a} = \check{b} \quad (*)$$

For (*), applying (c) at the γ -stage, there are $q \in \mathbb{P}_\gamma$ and $f \in \mathbb{P}_{\gamma\delta}$ such that $q \leq_{F,n} p|_\gamma$ and $q \Vdash_{\mathbb{P}_\gamma} \dot{f} = f \cdots (*\text{-i})$. And for (*), applying (a) at the γ -stage there are $q' \in \mathbb{P}_\gamma$ and countable X in V such that $q' \leq_{F,n} q$ and $q' \Vdash_{\mathbb{P}_\gamma} b \in \check{X} \cdots (*\text{-ii})$. Therefore from (*), (*-i) and (*-ii), for $f \in \mathbb{P}_{\gamma\delta}$, $q' \in \mathbb{P}_\gamma$ and countable X , $q' \Vdash_{\mathbb{P}_\gamma} (f \leq_{F,n} p^\gamma \wedge f \Vdash_{\mathbb{P}_{\gamma\delta}} \dot{a} \in \check{X})$. Recall that we have $q \leq_{F,n} p|_\gamma$, $f \leq_{\mathbb{P}_{\gamma\delta}} p^\gamma$ and $F \subset \gamma$, we obtain that $q \cup f \leq_{F,n} p|_\gamma \cup p^\gamma = p$. \square

Since we have already seen that the equivalent for A-4 and A-4', we have the following result:

Lemma III.4.9. The following is equivalent to (a) in Lemma III.4.8.

(a') If I is an antichain in \mathbb{P}_β , then the following set is (F, n) -dense for all finite $F \subset \beta$ and $n \in \omega$;

- $\{q \in \mathbb{P}_\beta; \text{ “ } \{q \in I; p \not\leq q\} \text{ is countable in } V. \text{” } \}$,

Lemma III.4.10. Assume that CH and let $\alpha \in \omega_2$ with $\|\mathbb{P}_\beta\| \leq \aleph_1$. Then there is $D_\alpha \subset \mathbb{P}_\alpha$ such that $|D_\alpha| \leq \aleph_1$ and D_α is (F, n) -dense for arbitrary $F \in [\alpha]^{<\omega}$ and $n \in \omega$.

Proof. We show by induction on α .

For $\alpha = 0$. By letting $D_\alpha = \{\emptyset\}$, we are done.

The case of $\alpha = \beta + 1$. Let $D_\beta \subset \mathbb{P}_\beta$ with $\|\mathbb{P}_\beta\| \leq \aleph_1$ and D_β is (F, n) -dense for all $F \in [\beta]^{<\omega}$ and $n \in \omega$. Let $\{\dot{d}_\xi; \xi \in \omega_1\}$ be a set such that $\Vdash_\beta \dot{Q}_\beta = \{\dot{d}_\xi; \xi \in \omega_1\}$. Fix a \mathbb{P}_β -generic G_β . For antichain $I \subset D_\beta$ which is countable in V and a function $f \in {}^I \omega_1$ in V , define a \mathbb{P}_β -name $\dot{q}(f)$ is; $\dot{d}_{f(p)}$ if $I \cap G_\beta = \{p\}$ and $\dot{1}_{\dot{Q}_\beta}$ if $I \cap G_\beta = \emptyset$. Since we have $V \models \text{CH}$, the number of such a function f is at most \aleph_1 , so the cardinality of $T = \{\dot{q}(f); f\}$ is $\leq \aleph_1$. Then by letting $D_\alpha = \{p \in \mathbb{P}_\alpha; p|_\beta \in D_\beta \wedge p(\beta) \in T\}$ we obtain that $|D_\alpha| \leq \aleph_1$. To see that D_α is (F, n) -dense for every $F \in [\alpha]^{<\omega}$ and $n \in \omega$. Let I be a maximal antichain of $\{d \in D_\beta; d \leq_{\mathbb{P}_\beta} p|_\beta \wedge \exists \xi \in \omega_1 (d \Vdash_\beta p(\beta) = \dot{d}_\xi)\}$. By Lemma III.4.9 and induction hypothesis, there is $d \in D_\beta$ such that $d \leq_{F, n} p$ and $J = \{q \in I; d \not\leq q\}$ is countable in V . Note that J is a dense antichain. So by letting $f: J \rightarrow \omega_1$ which assign the $q \in J$ the ξ if $q \Vdash_\beta p(\beta) = \dot{d}_\xi$, the $\dot{1}_{\dot{Q}}$ if otherwise. Note that since I is maximal, there is $d' \in J$ with $d \not\leq d'$ and $d' \Vdash_\beta p(\beta) = \dot{d}'_\xi$ for some $\xi \in \omega_1$. Thus for a common extension d'' for d and d' , we obtain that $d'' \Vdash_\beta p(\beta) = \dot{q}(f)$. Then by letting $r = d'' \wedge \langle \dot{q}(f) \rangle \in D_\alpha$, since we have $d'' \leq_{F \cap \beta, n} p|_\beta$, we obtain that $r \leq_{F, n} p$.

The case of α is limit. For $\beta \in \alpha$ and $q \in D_\beta$, define $\bar{q} = q \wedge \langle \dot{1} \rangle \wedge \dots \wedge \langle \dot{1} \rangle \in \mathbb{P}_\alpha$. Define $\bar{D}_\alpha = \{\bar{q}; \exists \beta \in \alpha q \in D_\beta\}$ and we shall find $D_\alpha \subset \mathbb{P}_\alpha$ with following conditions:

- $|D_\alpha| \leq \aleph_1$,
- $\bar{D}_\alpha \subset D_\alpha$,
- For an (F, n) -fusion sequence $\{p_n, F_n; n \in \omega\}$ in \mathbb{P}_α with $p_n \in D_\alpha$, there is a fusion p in D_α .

Fix $d \in D_\alpha$ and a fusion for (F, n) -fusion sequences in \mathbb{P}_α . For $D \subset \mathbb{P}_\alpha$ let $f: {}^{<\omega_1} D \rightarrow D$ be a fusion which assigns to x the:

$$\begin{cases} p & \text{if there is } F: \omega \rightarrow [\omega]^{<\omega} \text{ such that } \{\langle x(n), F(n) \rangle; n \in \omega\} \text{ is an} \\ & (F, n)\text{-fusion sequence and } p \text{ is the fusion for } \{\langle x(n), F(n) \rangle; n \in \omega\} \\ d & \text{if there is } n \in \omega \text{ } x(n) \text{ is not a singleton or otherwise.} \end{cases}$$

Then there is a closure D_α such that $\bar{D}_\alpha \subset D_\alpha$ and D_α is closed under f and note that $|D_\alpha| \leq \omega_1^{<\omega} = \omega_1$ and triviality hold the conditions.

To see that D_α is (F, n) -dense, let $p \in \mathbb{P}_\alpha$ $F \in [\alpha]^{<\omega}$ and $n \in \omega$ and we distinguish two cases, according to whether $\text{cf}(\alpha) = \omega$. If $\omega \in \text{cf}(\alpha)$. Let $\beta \in \alpha$ with $\text{supt}(p) \subset \beta$ and $F \subset \beta$. By induction hypothesis there is $q \in D_\beta$ such that $q \leq_{F,n} p|_\beta$ moreover, we obtain that $\bar{q} \leq_{F,n} p$. If $\text{cf}(\alpha) = \omega$. Let $\xi_m \in \alpha$, $m \in \omega$, with $\xi_m \nearrow \alpha$ and $F \subset \xi_0$. Define $q_m \in D_{\xi_m}$, $m \geq n$, recursively, by induction hypothesis there is $q_n \in D_{\xi_0}$ such that $q_n \leq_{F \cap \beta, n} p|_{\xi_0}$ and for $q_m \in D_{\xi_m}$ there is $q_{m+1} \in D_{\xi_{m+1}}$ such that $q_{m+1} \leq_{F,n} q_m \cup p|_{[\xi_m, \xi_{m+1})}$ for $m \geq n$. Then we obtain an (F, n) -fusion sequence $\langle (p_m, F_m); m \in \omega \rangle$ of \mathbb{P}_α such that $p_m = \bar{q}_m$ and $F_m = F$. Then by [Lemma III.4.7](#), there is $q \in D_\alpha$ such that $q \leq_{F,m} p_m$ for all $m \in \omega$. Therefore we obtain that $q \leq_{F,n} p$. \square

Proof of [Theorem III.4.5](#). To see that $\mathbb{1} \Vdash \omega_1 = \omega_1^V$, we assume that if not, i.e., there is $p \in \mathbb{P}$ such that $p \Vdash (\exists f: \omega_1^V \rightarrow \omega \text{ “ } f \text{ is an injection.”})$. Then we obtain that $p \Vdash \{f(n); n \in \omega_1^V\} \subset \omega$. Now by applying (b) in [Lemma III.4.8](#), we can show the general case in the similar way, there is an extension such that $q \in \mathbb{P}_\alpha$ and countable X in V such that $q \Vdash \{f(n); n \in \omega_1^V\} \subset \check{X}$. This shows that $\omega_1 \subset X$ in V , a contradiction.

For the moreover part. We distinguish two cases, according whether $\alpha = \omega_2$. If $\alpha \in \omega_2$. Assume that there is an antichain $I \subset \mathbb{P}_\alpha$ of size $> \aleph_2$. We may assume that I is maximal. [Lemma III.4.9](#) asserts that there is D_α such that for every $x \in I$, $F \in [\alpha]^{<\omega}$ and $n \in \omega$, there is $x_D \in D_\alpha$ so that $x_D \leq_{F,n} x$. Since $|D_\alpha| \leq \aleph_2$ and $|I| > \aleph_2$, Pigeonhole principle asserts that there is $\{x, y\} \in [I]^2$ such that $x_D = y_D$. Then for every $\eta \in \alpha$, for a finite set $\{\eta\}$ and $0 \in \omega$, we obtain that $x_D|_\eta \Vdash x_D(\eta) \leq_{\mathbb{P}_\eta} x(\eta), y(\eta)$. Thus we obtain that $x_D \leq_{\mathbb{P}_\alpha} x, y$. This contraly that $x \perp y$. If $\alpha = \omega_2$. Assume that there is an antichain I of size \aleph_2 . Let $\mathcal{A} = \{\text{supt}(r); r \in I\}$. We assert that there is an ordinal $\beta \in \alpha$ and $\mathcal{I} \in [I]^{\aleph_2}$ such that $\forall \{p, q\} \in [\mathcal{I}]^2$ $\text{supt}(p) \cap \text{supt}(q) \in \beta$. If this holds, since $p|_\beta \not\leq q|_\beta$ iff $p \not\leq q$ for $p, q \in \mathcal{I}$, a contradiction. To see there is an ordinal and a family, we distinguish two cases, according to whether $|\cup \mathcal{A}| = \aleph_2$. If $|\cup \mathcal{A}| \in \aleph_2$, Pigeonhole principle shows that there is $\mathcal{I} \in [I]^{\aleph_2}$ which elements are the same support and there is an ordinal β greater than the support them all. If $|\cup \mathcal{A}| = \aleph_2$. Since we have CH, we obtain that $\forall \theta \in \aleph_1$ ($\theta^{<\aleph_1} \in \aleph_2$) holds. Therefore, by [Delta System Lemma](#), there is $\mathcal{I} \in [I]^{\aleph_2}$ and a root R such that $\forall \{p, q\} \in [\mathcal{I}]^2$ $\text{supt}(p) \cap \text{supt}(q) = R \in \beta$ for some ordinal $\beta \in \alpha$. \square

III.5 Proper Forcing

In this section we assume that a forcing notion is separative.

Definition III.5.1 ([\[Kun11\]](#)). For a regular cardinal κ , the collection of sets of *cardinality hereditarily* $< \kappa$ is $H_\kappa = \{x; |\text{trcl}(x)| < \kappa\}$. Note that $H_\kappa \models \text{ZFC} - P$ if κ is an uncountable regular cardinal.

Definition III.5.2 ([\[Abr10\]](#)). Let \mathbb{P} be a forcing notion, λ be an uncountable regular cardinal and $M \preceq H_\lambda$ with $\mathbb{P} \in M$. $q \in \mathbb{P}$ is (M, \mathbb{P}) -generic provided that for any dense subset, D , of \mathbb{P} if $D \in M$ $D \cap M$ is predense below q .

Lemma III.5.3 ([Abr10]). For \mathbb{P} and M as per in previous definition, $q \in \mathbb{P}$ is (M, \mathbb{P}) -generic iff for every dense subset D of \mathbb{P} in M and (V, \mathbb{P}) -generic G with $p \in G$ there is $\dot{p} \in V^{\mathbb{P}}$ such that $q \Vdash \dot{p} \in M \cap D \cap \dot{G}$.

Proof. It suffices to show that for $q \in \mathbb{P}$ and $E \subset \mathbb{P}$, E is predense below q iff $q \Vdash E \cap \dot{G} \neq \emptyset$ for arbitrary (V, \mathbb{P}) -generic G with $p \in G$.

(\Rightarrow) Let G be (V, \mathbb{P}) -generic with $q \in G$. Since $F = \{p \in \mathbb{P}; p \perp q \vee \exists r \in E \ p \leq r\}$ is dense subset of \mathbb{P} , there is $p \in G \cap F$. Since $p \perp q$, we have $r \in G$. This shows that $q \Vdash E \cap \dot{G} \neq \emptyset$.

(\Leftarrow) Let $p \in q$ and G a (V, \mathbb{P}) -generic with $p \in G$. Since $q \in G$ there is $r \in E \cap G$, which witnesses E is predense below q . \square

Definition III.5.4 ([Abr10]). A forcing notion \mathbb{P} is *proper* provided that for an arbitrary uncountable regular cardinal $\lambda > 2^{|\mathbb{P}|}$, or simply says sufficiently large regular cardinal λ , and a countable elementary submodel $M \preceq H_\lambda$ with $\mathbb{P} \in M$, we have:

$$\forall p \in \mathbb{P} \cap M \exists q \leq p \text{ “} q \text{ is } (M, \mathbb{P})\text{-generic.”}$$

Example III.5.5 ([Abr10]).

1. A ccc forcing is proper.
2. A σ -closed forcing is proper.
3. An Axiom A forcing is proper.
4. A shooting a club by finite conditions forcing is proper. (see [Theorem IV.5.2](#))

Proof. 1. Let λ be an uncountable regular cardinal with $\lambda > 2^{|\mathbb{P}|}$, $M \preceq H_\lambda$ with $\mathbb{P} \in M$ and $p \in \mathbb{P} \cap M$. We shall verify that p is (M, \mathbb{P}) -generic. For a dense subset $D \subset \mathbb{P}$ with $D \in M$, let $A \subset D$ be a maximal antichain and by elementarity we may choose as $A \in M$. Since \mathbb{P} is a ccc forcing, there is a surjection $f: \omega \rightarrow A$ in V . Furthermore, since $f \in H_\lambda$, we may assume that $f \in M$ and therefore, $A = f(\omega) \subset M$. Then A is predense below p and since $A \subset D \cap M$, so does D .

2. Let λ be an uncountable regular cardinal with $\lambda > 2^{|\mathbb{P}|}$, $M \preceq H_\lambda$ with $\mathbb{P} \in M$ and $p \in \mathbb{P} \cap M$. Let $\mathcal{D} = \{D \in M; “D \subset \mathbb{P} \text{ is dense.”}\} \subset M$ and enumerate $\mathcal{D} = \{D_i; i \in \omega\}$ in V . Note that, by elementarity, for each $q \in \mathbb{P} \cap M$ there is an extension $r \in D \cap M$ for any $D \in \mathcal{D}$. Thus we obtain a decreasing sequence $\langle p_n \in \mathbb{P} \cap M; n \in \omega \rangle$ such that:

- $p_0 = p$,
- $p_{n+1} \leq p_n$ such that $p_{n+1} \in D_n \cap M$.

and moreover since \mathbb{P} is σ -closed, there is $q \in \mathbb{P}$ such that $q \leq p_n$ for all $n \in \omega$. The statement that q is (M, \mathbb{P}) -generic is immediate by the construction, thus we conclude the proof.

3. \square

Theorem III.5.6 ([Jec03]). A forcing notion $(\mathbb{P}, <)$ is proper iff for every uncountable cardinal λ every stationary subset of $[\lambda]^\omega$ remains stationary in the generic extension.

Before, we prove the theorem, we shall make some lemmata.

Lemma III.5.7. Let \mathbb{P} be a ccc forcing. Every club subset $C \subset [\lambda]^\omega$ in $V[G]$ has a subset $D \in V$ which is club $D \in V$. Moreover, every stationary set $S \subset [\lambda]^\omega$ remains stationary in $V[G]$.

Lemma III.5.8. Let \mathbb{P} be a σ -closed forcing. Every stationary set $S \subset [\lambda]^\omega$ remains stationary in $V[G]$.

Lemma III.5.9.

Definition III.5.10 ([Jec03]). Let \mathbb{P} be a forcing notion and $p \in \mathbb{P}$. The *proper game* for \mathbb{P} , below p , is played as follows:

- I plays P -names $\dot{\alpha}_n$ for ordinal numbers, and
- II plays ordinal numbers β_n .

Player II wins provided that there exists an extension $q \leq p$ which forces $\forall n \in \omega \exists k \in \omega \dot{\alpha}_n = \beta_k$.

Theorem III.5.11 ([Jec03]). A forcing notion \mathbb{P} is proper iff II has a winning strategy for the proper game for any $p \in \mathbb{P}$.

Theorem III.5.12 ([Abr10]). A proper forcing preserves ω_1 .

Proof. Let \dot{f} be a \mathbb{P} -name and p a condition such that $p \Vdash \dot{f}: \omega \rightarrow \omega_1^V$. We shall show that $p \Vdash \dot{f}$ is not a surjection. Fix an uncountable regular cardinal $\lambda > 2^{|\mathbb{P}|}$ such that $\dot{f}, \mathbb{P}, \omega_1^V \in H_\lambda$ and let M be a countable elementary submodel of H_λ such that $\dot{f}, \mathbb{P}, p, \omega_1^V \in M$. Since \mathbb{P} is proper, there is an extension $q \leq p$ which is (M, \mathbb{P}) -generic. Fix an $n \in \omega$. Define $D_n = \{r \in \mathbb{P}; (r \perp p) \vee (\exists \alpha \in \omega_1^V r \Vdash \dot{f}(n) = \alpha)\}$ in H_λ . By elementary we obtain that $D_n \in M$, thus for an extension $q_1 \leq q$ there is $r \in D_n \cap M$ such that $q_1 \not\leq r$ and moreover since $q_1 \leq p$, there is $\alpha \in \omega_1$ such that $r \Vdash \dot{f}(n) = \alpha$, furthermore, by elementary we may assume that $\alpha \in \omega_1^V \cap M$.

This shows that $q \Vdash \forall n \in \omega \dot{f}(n) \in M$, i.e., $p \Vdash \dot{f}$ is not a surjection. Therefore, $\Vdash \omega_1 = \omega_1^V$. \square

Hereafter, let λ be a sufficiently large regular cardinal such that $\lambda > 2^{|\mathbb{P}|}$.

Theorem III.5.13 ([Abr10]). Let \mathbb{P} be a forcing notion and $M \prec H_\lambda$ be countable with $\mathbb{P} \in M$ and suppose that $p \in \mathbb{P}$ is a (M, \mathbb{P}) -generic condition. For $x, y \in M \cap V^{\mathbb{P}}$, if we have $p \Vdash x = y$, then there is an extension $r \leq p$ and $(z, q) \in y \cap M$ such that $z \in \mathbb{P}$, $q \in V^{\mathbb{P}}$, $r \leq q$ and $r \Vdash x = z$.

Proof. Define $E = \{r \in \mathbb{P}; \exists(z, q) \in x (r \leq q \wedge r \Vdash x = z)\}$ and $E' = \{r \in \mathbb{P}; r \in E \vee r \perp E\}$. Then E' is a dense set in \mathbb{P} with $E' \in M$. Then there is $r \in M \cap E'$ which is compatible with p and there is $(z, q) \in x$ such that $r \leq q$ and $r \Vdash x = z$. Furthermore, since $M \preceq H_\lambda$, we may assume that $(z, q) \in M$, we are done. \square

Lemma III.5.14 ([Abr10]). Let \mathbb{P} be a forcing notion and $\Vdash_{\mathbb{P}}$ “ \dot{Q} is a forcing notion.”, $M \preceq H_\lambda$ countable with $\mathbb{P}, \dot{Q}_G \in M$ and $p \in \mathbb{P} \cap M$ and $\dot{q} \in \dot{Q} \cap M$ where G is (V, \mathbb{P}) -generic. Then

(p, \dot{q}) is $(M, \mathbb{P} * \dot{Q})$ -generic iff

p is (M, \mathbb{P}) -generic and $p \Vdash$ “ \dot{q} is $(M[\dot{G}], \dot{Q})$ -generic.”.

Proof. (\Rightarrow) . To see that p is (M, \mathbb{P}) -generic. Let $D \subset \mathbb{P}$ be a dense subset with $D \in M$. Then $D' = \{(d, \dot{q}); d \in D\}$ be a dense subset of $\mathbb{P} * \dot{Q}$ with $D' \in M$. Therefore there is $(d, \dot{q}) \in D' \cap M$ such that $(d, \dot{q}) \leq_{\mathbb{P} * \dot{Q}} (p, \dot{q})$. Moreover, by the elementary there is $d \in D \cap M$ such that $d \leq_{\mathbb{P}} p$.

To see that $p \Vdash_{\mathbb{P}}$ “ \dot{q} is $(M[\dot{G}], \dot{Q})$ -generic.”, it suffices show that for any dense open subset $\dot{D} \in M[\dot{G}]$ in \dot{Q} and $\dot{q}' \leq_{\dot{Q}} \dot{q}$, p forces:

$$\exists \dot{q}'' \in \dot{D} \cap M[\dot{G}] \dot{q}'' \not\leq_{\dot{Q}} \dot{q}.$$

Let $p' \leq_{\mathbb{P}} p$. Define $E = \{(s, \dot{t}) \in \mathbb{P} * \dot{Q}; s \Vdash \dot{t} \in \dot{D}\}$ and $F = \{(s, \dot{t}) \in \mathbb{P} * \dot{Q}; (s, \dot{t}) \in E \vee (s, \dot{t}) \perp_{\mathbb{P} * \dot{Q}} E\}$. Then $F \in M$ and F is dense in $\mathbb{P} * \dot{Q}$. Then there is $(s, \dot{t}) \in F \cap M$ which compatible with (p', \dot{q}) and let (p_2, \dot{q}_2) be a common extension and $\dot{q}'' \in \dot{D}$ with $\dot{q}'' \leq_{\dot{Q}} \dot{q}_2$. Then we obtain that $p_2 \Vdash \dot{q}'' \leq_{\dot{Q}} \dot{t}$. So, $(s, \dot{t}) \in E$, moreover, $p_2 \Vdash \dot{t} \in \dot{D}$ and note that $\dot{t} \in M$.

This argument shows that there is $(p_2, \dot{t}) \in \mathbb{P} * \dot{Q}$ such that $p_2 \leq p'$, $p_2 \Vdash \dot{q}' \not\leq_{\dot{Q}} \dot{t} \wedge \dot{t} \in \dot{D} \wedge \dot{t} \in M[\dot{G}]$. Furthermore, since there is densely many extension, we have:

$$p \Vdash \exists \dot{q}'' \in \dot{D} \cap M[\dot{G}] \dot{q} \not\leq_{\dot{Q}} \dot{q}''.$$

(\Leftarrow) . Let $D \subset \mathbb{P} * \dot{Q}$ be a dense open set with $D \in M$ and let (p_1, q_1) be an extension for (p, q) . We shall find a $(p', q') \in D \cap M$ such that $(p_1, q_1) \not\leq_{\mathbb{P} * \dot{Q}} (p', q')$. Let $\dot{E} = \{\dot{q} \in \dot{Q}; \exists p' \in \dot{G} (p', \dot{q}) \in D\}$. To see that $\Vdash_{\mathbb{P}}$ “ \dot{E} is dense.”, it suffices to verify that:

$$\forall (s, \dot{t}) \in \mathbb{P} * \dot{Q} \exists s' \leq_{\mathbb{P}} s \ s' \Vdash_{\mathbb{P}} (\dot{q} \in \dot{E} \wedge \dot{q} \leq_{\dot{Q}} \dot{t}).$$

For (s, \dot{t}) , since D is dense, there is an extension (s', \dot{t}') and $s' \Vdash_{\mathbb{P}} (\dot{t}' \leq_{\dot{Q}} \dot{t} \wedge s \in \dot{G})$ holds. Thus, $s \Vdash_{\mathbb{P}} \dot{t} \in \dot{E}$, and we obtain that $\Vdash_{\mathbb{P}}$ “ \dot{E} is dense.” and note that $\Vdash_{\mathbb{P}} \dot{E} \in M[\dot{G}]$, so by assumption we have $p_1 \Vdash_{\mathbb{P}} \exists \dot{r} \in \dot{E} \cap M[\dot{G}] \dot{r} \not\leq_{\dot{Q}} \dot{q}_1$, furthermore there is $\dot{r} \in V^{\mathbb{P}} \cap M$ and $\dot{t}' \in \dot{Q}$ such that $p_1 \Vdash_{\mathbb{P}} \dot{r} \in \dot{E} \wedge \dot{t}' \leq_{\dot{Q}} \dot{r}, \dot{q}_1$.

Define $F = \{s \in \mathbb{P}; (s, \dot{r}) \in D\}$ and $F' = \{s \in \mathbb{P}; s \in F \vee s \perp_{\mathbb{P}} F\}$ and note that F' is predense in \mathbb{P} and $F' \in M$. Then there is $s \in F' \cap M$ such that $s \not\leq_{\mathbb{P}} p_1$ and by letting s' be a common extension, we have $s' \Vdash_{\mathbb{P}} \dot{r} \in \dot{E}$. Hence,

there is a t such that $s' \Vdash_{\mathbb{P}} (t \in \dot{G} \wedge (t, \dot{r}) \in D)$ and note that we obtain $s' \leq_{\mathbb{P}} t$ and $t \in F$ witnesses $s \in F$.

Therefore, $(s, \dot{r}) \in D \cap M$ and (s', \dot{t}) witnesses $(s, \dot{r}) \not\leq_{\mathbb{P} * \dot{\mathbb{Q}}} (p_1, \dot{q}_1)$ \square

Theorem III.5.15 ([Abr10]). Let \mathbb{P}_0 be a proper forcing notion, $\dot{\mathbb{P}}_1 \in V^{\mathbb{P}_0}$ with $\Vdash_{\mathbb{P}_0}$ “ $\dot{\mathbb{P}}_1$ is proper.”, $\mathbb{Q} = \mathbb{P}_0 * \dot{\mathbb{P}}_1$ and $\pi: \mathbb{Q} \rightarrow \mathbb{P}_0; (p, \dot{q}) \mapsto p$. Then for a countable model $M \preceq H_\lambda$ with $\mathbb{Q} \in M$ and (M, \mathbb{P}_0) -generic $p_0 \in \mathbb{P}_0$ we have the following.

For any $\dot{r} \in V^{\mathbb{P}_0} \cap M$ if $p_0 \Vdash_{\mathbb{P}_0} (\dot{r} \in \mathbb{Q} \wedge \pi(\dot{r}) \in \dot{G}_0)$, then there is $\dot{p}_1 \in V^{\mathbb{P}_0}$ such that $(p_0, \dot{p}_1) \in \mathbb{Q}$ is (M, \mathbb{Q}) -generic and $(p_0, \dot{p}_1) \Vdash_{\mathbb{Q}} \dot{r} \in \dot{G}$. Where G_0 and G is \mathbb{P}_0 and \mathbb{Q} generic, respectively.

Moreover, \mathbb{Q} is proper.

Proof. Let $p_0 \in \mathbb{P}_0$ be (M, \mathbb{P}_0) -generic and $(r_0, \dot{r}_1) \in R \cap M$ with $p_0 \Vdash_{\mathbb{P}_0} r_0 \in \dot{G}_0$. Since $p_0 \Vdash_{\mathbb{P}_0}$ “ $\dot{\mathbb{P}}_1$ is proper.” and $M[\dot{G}] \preceq H_\lambda[\dot{G}]$, there is $\dot{p}_1 \in V^{\mathbb{P}_0}$ such that $p_0 \Vdash_{\mathbb{P}_0} (\dot{p}_1 \Vdash_{\dot{\mathbb{P}}_1} \dot{r}_1) \wedge$ “ \dot{p}_1 is $(M[\dot{G}], \dot{\mathbb{P}}_1)$ -generic.” and since $p_0 \leq_{\mathbb{P}_0} r_0$, we obtain that $(p_0, \dot{p}_1) \leq_{\mathbb{Q}} (r_0, \dot{r}_1)$ and $p_0 \Vdash_{\mathbb{P}_0}$ “ \dot{p}_1 is $(M[\dot{G}], \dot{\mathbb{P}}_1)$ -generic.”. Therefore, by the previous lemma, we obtain that (p_0, \dot{p}_1) is (M, \mathbb{Q}) -generic and $(p_0, \dot{p}_1) \Vdash_{\mathbb{Q}} (r_0, \dot{r}_1) \in \dot{G}$.

The moreover part is immediate by the first assertion. \square

Lemma III.5.16 (The Properness Extension Lemma, [Abr10]). Let γ be a limit ordinal, $\mathbb{P}_\gamma = \langle \mathbb{P}_\alpha; \alpha \in \gamma \rangle$ a countable support iteration of proper forcing and $M \preceq H_\lambda$ a countable submodel with sufficiently large cardinal λ with $\mathbb{P}_\gamma, \gamma \in M$.

For any $\gamma_0 \in \gamma \cap M$, $q_0 \in \mathbb{P}_{\gamma_0}$ and name $\dot{p}_0 \in V^{\mathbb{P}_{\gamma_0}}$.

If $q_0 \in \mathbb{P}_{\gamma_0}$ forces at \mathbb{P}_{γ_0} stage:

$$\dot{p}_0 \in \mathbb{P}_\gamma \cap M \text{ and } \dot{p}_0|_{\gamma_0} \in \dot{G}_0$$

then there is an (M, \mathbb{P}_γ) -generic condition q such that $q|_{\gamma_0} = q_0$ and $q \Vdash_{\mathbb{P}_\gamma} i_*(\dot{p}_0) \in \dot{G}_\gamma$

Proof. For $\gamma_0 \in \gamma \cap M$, $\dot{p}_{\gamma_0} \in V^{\mathbb{P}_{\gamma_0}}$ and $(M, \mathbb{P}_{\gamma_0})$ -generic q_{γ_0} we assume that

$$q_{\gamma_0} \Vdash_{\mathbb{P}_{\gamma_0}} (\dot{p}_{\gamma_0} \in \mathbb{P}_\gamma \cap M \wedge \dot{p}_{\gamma_0}|_{\gamma_0} \in \dot{G}_{\gamma_0})$$

We shall found the desired (M, \mathbb{P}_γ) -generic condition by induction on γ .

(i) $\gamma = 0$. It is immediate that there are no any elements in $\gamma \cap M$.

(ii) $\gamma = \gamma' + 1$. We have seen the case of $\gamma' = \gamma_0$ in **Theorem III.5.15**, so we may assume that $\gamma_0 \in \gamma'$.

Note that we have $\gamma', \mathbb{P}_{\gamma'} \in M$ and $\lambda > 2^{|\mathbb{P}_{\gamma'}|}$ and since q_{γ_0} forces

$$\dot{p}_{\gamma_0}|_{\gamma'} \in \mathbb{P}_{\gamma'} \cap M \text{ and } (\dot{p}_{\gamma_0}|_{\gamma'})|_{\gamma_0} \in \dot{G}_{\gamma_0},$$

by induction hypothesis, there is an $(M, \mathbb{P}_{\gamma'})$ -generic $q_{\gamma'}$ such that

$$q_{\gamma'}|_{\gamma_0} = q_{\gamma_0} \text{ and } q_{\gamma'} \Vdash_{\mathbb{P}_{\gamma'}} i_*(\dot{p}_{\gamma_0})|_{\gamma'} = \dot{p}_{\gamma_0}|_{\gamma'} \in \dot{G}_{\gamma'}.$$

Theorem III.5.15 asserts that there is (M, \mathbb{P}_γ) -generic $q \in \mathbb{P}_\gamma$ such that

$$q|_{\gamma'} = q_{\gamma'} \text{ and } q \Vdash_{\mathbb{P}_\gamma} i_*(p_{\gamma_0}^\circ) \in \mathring{G}_\gamma$$

and moreover, we have $q|_{\gamma_0} = q_{\gamma_0}$.

(ii) γ is limit. Since $\gamma_0 \in \gamma \cap M$ and $\text{cf}(\gamma \cap M) = \omega$, there is $\langle \gamma_n; n \in \omega \rangle \nearrow \gamma \cap M$ and enumerate the dense sets in $\mathbb{P}_\gamma \cap M$, $\{D_n; n \in \omega\}$. Now we define $q_n \in \mathbb{P}_{\gamma_n}$ and $p_n \in V^{\mathbb{P}_{\gamma_n}}$ recursively such as:

1. $q_0 = q_{\gamma_0}$,
2. q_n is $(M, \mathbb{P}_{\gamma_n})$ -generic such as $q_{n+1}|_{\gamma_n} = q_n$,
3. $p_0 = p_{\gamma_0}^\circ$,
4. q_{n+1} forces, over $\mathbb{P}_{\gamma_{n+1}}$, and
 - (4-a.) $p_{n+1} \in \mathbb{P}_\gamma \cap M$,
 - (4-b.) $p_{n+1}^\circ|_{\gamma_{n+1}} \in G_{\gamma_{n+1}}^\circ$,
 - (4-c.) $p_{n+1}^\circ \leq_{\mathbb{P}_{\gamma_n}} i_*(p_n)$, and
 - (4-d.) $p_{n+1}^\circ \in D_n$.

Suppose we have p_n and q_n , we shall find p_{n+1} and q_{n+1} .

Fix a $q' \leq_{\mathbb{P}_{\gamma_n}} q_n$. There is $q'' \leq_{\mathbb{P}_{\gamma_n}} q'$ and $p \in \mathbb{P}_\gamma \cap M$ such that $q' \Vdash p_n^\circ = p$. By letting $E = \{r|_{\gamma_n} \in \mathbb{P}_{\gamma_n}; r \in D_n \wedge r \leq_{\mathbb{P}_{\gamma_n}} p\}$, we have $E \in M$ and is dense below $p|_{\gamma_n}$. Since $q'' \Vdash_{\mathbb{P}_{\gamma_n}} p|_{\gamma_n} \in G_{\gamma_n}^\circ$ and q'' is $(M, \mathbb{P}_{\gamma_n})$ -generic, by **Lemma III.5.3** there is $\dot{r} \in V^{\mathbb{P}_{\gamma_n}}$ such that $q'' \Vdash_{\mathbb{P}_{\gamma_n}} \dot{r} \in E \cap M \cap G_{\gamma_n}^\circ$ and moreover, there is $\dot{p} \in V^{\mathbb{P}_{\gamma_{n+1}}}$ such that q'' forces on \mathbb{P}_{γ_n}

$$\dot{p} \in D_n \cap M, \dot{p}|_{\gamma_n} \in G_{\gamma_n}^\circ \text{ and } \dot{p} \leq_{\mathbb{P}_{\gamma_n}} p_n^\circ.$$

Therefore we obtain that q_n forces on \mathbb{P}_{γ_n} :

$$\dot{p}|_{\gamma_{n+1}} \in \mathbb{P}_{\gamma_{n+1}} \cap M \text{ and } (p|_{\gamma_{n+1}})|_{\gamma_n} \in G_{\gamma_n}^\circ$$

and hence applying induction hypothesis there is $(M, \mathbb{P}_{\gamma_{n+1}})$ -generic $q_{n+1} \in \mathbb{P}_{\gamma_{n+1}}$ such that

$$q_{n+1}|_{\gamma_n} = q_n \text{ and } q_{\gamma_{n+1}} \Vdash_{\mathbb{P}_{\gamma_{n+1}}} i_*(p|_{\gamma_{n+1}}) \in G_{\gamma_{n+1}}^\circ.$$

By letting $p_{n+1}^\circ = i_*\dot{p}$, we have that $q_{n+1} \leq_{\mathbb{P}_{\gamma_{n+1}}} i(q_n)$ and $i(q_n)$ forces on $\mathbb{P}_{\gamma_{n+1}}$

$$p_{n+1} \in \mathbb{P}_\gamma \cap M, p_{n+1}^\circ \leq_{\mathbb{P}_{\gamma_{n+1}}} i_*(p_n^\circ), p_{n+1}^\circ|_{\gamma_{n+1}} \in G_{\gamma_{n+1}}^\circ \text{ and } p_{n+1}^\circ \in D_n.$$

Now we complete the construction of p_n and q_n for $n \in \omega$.

Before we conclude the proof, we shall show that for $q = \cup\{q_n \in \mathbb{P}_{\gamma_n}; n \in \omega\} \in \mathbb{P}_\gamma$, we have

$$\forall n \in \omega \ q \Vdash_{\mathbb{P}_\gamma} i_*(p_n) \in \mathring{G}_{\gamma_n} \quad (*)$$

Fix an $n \in \omega$. For $m \ni n$, since we have $q_m \Vdash_{\mathbb{P}_{\gamma_m}} p_m^\circ \leq_{\mathbb{P}_\gamma} i_*(p_n^\circ)$ and $q_m \Vdash_{\mathbb{P}_{\gamma_m}} p_m^\circ|_{\gamma_m} \in \mathring{G}_{\gamma_m}$, we have

$$q \Vdash_{\mathbb{P}_\gamma} i_*(p_n^\circ) = i_*(i_*(p_n^\circ)|_{\gamma_m}) \in i_*(\mathring{G}_{\gamma_m})$$

For any extension $q' \leq_{\mathbb{P}_\gamma} q$ there is an extension $q'' \leq_{\mathbb{P}_\gamma} q'$ and $p \in \mathbb{P}_\gamma \cap M$ such that $q'' \Vdash_{\mathbb{P}_\gamma} i_*(p_n^\circ) = p$ and thus we have

$$q'' \Vdash_{\mathbb{P}_\gamma} p|_{\gamma_n} \in i_*(\mathring{G}_{\gamma_m}), \text{ moreover } q''|_{\gamma_m} \leq_{\mathbb{P}_{\gamma_m}} p|_{\gamma_m}.$$

Therefore we obtain $q'' \leq_{\mathbb{P}_\gamma} p$, i.e., $q'' \Vdash_{\mathbb{P}_\gamma} p \in \mathring{G}_\gamma$. Hence, we have $q'' \Vdash_{\mathbb{P}_\gamma} i_*(p_n) \in \mathring{G}_\gamma$ and $(*)$ hold.

To see that q is (M, \mathbb{P}_γ) -generic thanks to [Lemma III.5.3](#), we shall verify that

$$\forall n \in \omega \ q \Vdash_{\mathbb{P}_\gamma} i_*(p_{n+1}^\circ) \in D_n \cap M \cap \mathring{G}_\gamma.$$

For an $n \in \omega$, we have that $i(q_{n+1}) \Vdash_{\mathbb{P}_\gamma} i_*(p_{n+1}) \in D_n \cap M$ and $q \leq_{\mathbb{P}_\gamma} i_*(q_{n+1})$, we obtain that

$$q \Vdash_{\mathbb{P}_\gamma} i_*(p_{n+1}^\circ) \in D_n \cap M \in \mathring{G}_\gamma.$$

Thus, q is (M, \mathbb{P}_γ) -generic and since we have $q|_{\gamma_0} = q_{\gamma_0}$, $(*)$ asserts that we have

$$q \Vdash_{\mathbb{P}_\gamma} i_*(p_{\gamma_0}^\circ) \in \mathring{G}_\gamma.$$

□

Theorem III.5.17 ([\[Abr10\]](#)). For a limit ordinal δ and a countable support iteration of proper forcings $\langle \mathbb{P}_\alpha; \alpha \in \delta \rangle$, \mathbb{P}_δ is proper.

Proof. Let $\lambda > 2^{|\mathbb{P}_\delta|}$ be a regular uncountable cardinal and $M \preceq H_\lambda$ be an elementary substructure with $\mathbb{P}_\delta \in M$ and we may assume that $\mathbb{P}_\delta \cap M \neq \emptyset$.

Let $p \in \mathbb{P}_\delta \cap M$. Since \mathbb{P}_0 is proper, there is an (M, \mathbb{P}_0) -generic $q_0 \in \mathbb{P}_0$ with $q_0 \leq_{\mathbb{P}_0} p|_0$. Thus by applying [The Properness Extension Lemma](#), there is an (M, \mathbb{P}_δ) -generic $q \in \mathbb{P}_\delta$ such that $q|_1 = q_0$ and $q \Vdash_{\mathbb{P}_\delta} p \in \mathring{G}_\delta$. Thus we obtain that $q \leq_{\mathbb{P}_\delta} p$.

□

Theorem III.5.18 ([\[Abr10\]](#)). Assume CH and δ be a limit ordinal. For a countable support iteration $\mathbb{P}_\delta = \langle \mathbb{P}_i; i \in \delta \rangle$ with:

$$\Vdash_\gamma (\text{“ } \mathring{\mathbb{Q}}_\gamma \text{ is proper.” } \wedge |\mathring{\mathbb{Q}}_\gamma| = \aleph_1) \text{ for any } \gamma \in \delta.$$

Then, \mathbb{P}_δ satisfies an \aleph_2 -cc.

Proof. Fix $\{r_\xi; \xi \in \aleph_2\}$, regular $\lambda \geq \max\{2^{|\mathbb{P}_\delta|}, \aleph_3\}$ and well-order \triangleleft over H_λ . For each $\xi \in \aleph_2$ let $M_\xi = \langle M_\xi, \in|_{M_\xi}, \triangleleft|_{M_\xi}, \mathbb{P}_\delta, r_\xi \rangle$ be a countable submodel of H_λ .

First of all, we shall found an $I \in [\omega_2]^{\omega_2}$ and cuntable $C \in [\omega_2]^\omega$ such that:

1. $\forall \{\xi_0, \xi_1\} \in [I]^2 \exists f: (M_{\xi_0}, \in) \rightarrow (M_{\xi_1}, \in)$
 f is order-isomorphism and $f(r_{\xi_0}) = r_{\xi_1}$,
2. $\forall \{\xi_0, \xi_1\} \in [I]^2 \ M_{\xi_0} \cap M_{\xi_1} \cap \omega_2 = C$,
3. $\forall \xi_0, \xi_1 \in I \ \xi_0 \in \xi_1 \implies (\cup C \in \mu_{\xi_0} \wedge \cup(M_{\xi_0} \cap \omega_2) \in \mu_{\xi_1})$,
where $\mu_\xi = \cap((M_\xi \cap \omega_2) \setminus C)$.
4. C is an initial segment of $M_\xi \cap \omega_1$ for any $\xi \in I$, that is:

$$\forall y \in C \forall x \in M_\xi \cap \omega_1 \ x \in y \implies x \in C.$$

To see the condition 4. Fix a distinct $\eta \in I$. Let $y \in C$ and $x \in y \cap M_3 \cap \omega_1$. We shall show that $x \in M_\eta$. $y \in M_\xi \cap \omega_2$ asserts that there is a function $f_\xi \in M_\xi$ such that $M_\xi \models (\text{“} f_\xi: \omega_1 \rightarrow y \text{ is surjective.”} \wedge \text{“} f_\xi \text{ is } \triangleleft\text{-minimal.”})$. So does over H_λ . Similarly we found a funciton $f_\eta \in M_\eta$. Since the minimality assert that $f_\xi = f_\eta$ in H_λ and an $\alpha \in M_\xi \cap \omega_1$ such that $f_\xi(\alpha) = x$ witnesses $f_\eta(\alpha) = x \in M_\eta$.

Let $\xi, \eta \in I$ such that $\mu := \mu(\xi) \leq \mu(\eta)$ and define $r_1 = r_\xi|_{\cup C}$ and $r_2 = r_\eta|_{\cup C}$. Note that $r_\xi = i(r_\xi|_{\cup C})$ for any $\xi \in I$. Since $r_1|_\mu \in M_\xi \cap \mathbb{P}_\mu$, there is an (M_ξ, \mathbb{P}_μ) -generic extension $p \in \mathbb{P}_\mu$ and thanks to the result of the following lemma, we have:

If $p \leq_{\mathbb{P}_\mu} r_1|_\mu$ is (M_ξ, \mathbb{P}_μ) -generic, then $i(p) \leq_{\mathbb{P}_{\mu_\eta}} r_2|_{\mu_\eta}$.

Therefore, we obtain a common extension $r' \in \mathbb{P}_\mu$ such that $r' \leq_{\mathbb{P}_{\cup C}} r_1|_{\cup C}$ and $r' \leq_{\mathbb{P}_{\cup C}} r_2|_{\cup C}$. Define a fucntion r which assigns:

$$\alpha \mapsto \begin{cases} r'(\alpha) & \alpha \in \cup C \\ r_\xi(\alpha) & \alpha \in \text{supt}(r_\xi) \setminus \cup C \\ r_\eta(\alpha) & \alpha \in \text{supt}(r_\eta) \setminus \cup C \\ \mathbb{1}_{\dot{Q}_\alpha} & \text{otherwise} \end{cases}$$

We remain to see that $r \in \mathbb{P}_\delta$ and r is a common extension for r_ξ and r_η in \mathbb{P}_δ . \square

Lemma III.5.19. Let M_1, M_2 be two isomorphic countable elementary submodels of H_λ . Let $h: M_1 \rightarrow M_2$ be an isomorphism and $\mu \in M_1 \cap \omega_2$ be such that $\mu \leq h(\mu)$ and identity on $\mu \cap M_1$. Then if $p \in \mathbb{P}_\mu$ is any (M_1, \mathbb{P}_μ) -generic condition then for any condition $r \in \mathbb{P}_\mu \cap M_1$, $p \leq_{\mathbb{P}_\mu} r$ implies $p \leq_{\mathbb{P}_{h(\mu)}} h(r)$.

IV List of Forcing Notions

IV.1 Cohen Forcing

Definition IV.1.1. Let $\text{Fn}(I, J) = \cup\{^AJ; A \in [I]^{<\omega}\}$ be a forcing notion with order $\leq_{\mathbb{P}} = \supset$.

Theorem IV.1.2. $\text{Fn}(I, J)$ has ccc iff I is empty or J is countable.

Proof. (\Rightarrow). We shall verify that if I is not empty, J is countable. Fix an $n \in I$ then since $\{\langle n, j \rangle; j \in J\}$ is an antichain, we obtain that J is countable.

(\Leftarrow). Since the case of I is an empty set is clear, we may assume that $I \neq \emptyset$ and J is countable. Suppose that there is an uncountable antichain A , the Delta system lemma asserts that there is $B \in [A]^{\aleph_1}$ such that $\forall \{f, g\} \in [B]^2$ $f \cap g = h$ for some $h \in {}^IJ$. This shows that there are distinct $f, g \in B$ such that $f \not\leq g$, a contradiction. \square

IV.2 Hechler Forcing

Definition IV.2.1 ([Bla10]). Let \mathbb{P} be a forcing notion such that:

1. $(s, f) \in \mathbb{P}$ provided that $s \in {}^{<\omega}\omega$ and $f \in {}^\omega\omega$,
2. $(s', f') \leq (s, f)$ provided that
 - (a) $s' \supset s$,
 - (b) $f' \geq f$,
 - (c) $\forall n \in \text{dom}(s') \setminus \text{dom}(s)$ $s'(n) \ni f(n)$.

Theorem IV.2.2 ([Pal13]). Let G be a generic in \mathbb{P} , as per **Definition IV.2.1**, a Hechler real x is a function $x = \cup\{s; \exists f (s, f) \in G\}: \omega \rightarrow \omega$. Then, G can be recovered from x .

Proof. First, we show that x is a function with domain ω . Since G is a filter, manifestly, x is a function and since $D_n = \{(s, f) \in \mathbb{P}; n \in |s| \wedge f \in {}^\omega\omega\}$ is dense for every $n \in \omega$, $\text{dom}(x) = \omega$.

Secondly, let $G' = \{(s, f) \in \mathbb{P}; s \subset x \wedge \forall n \in \omega \setminus \text{dom}(s) f(n) \leq x(n)\}$ and we assert that $G = G'$ holds. To see $G \subset G'$, let $(s, f) \in G$ and we show that $f(n) \leq x(n)$ for all $n \in \omega \setminus \text{dom}(s)$. Since $\{(t, g) \in \mathbb{P}; n \in \text{dom}(t)\}$ is dense, there is $(t, g) \in G$ with $n \in \text{dom}(t)$ and let (u, h) be a common extension. Then we obtain that $f(n) \leq t(n) = x(n)$. To see $G \supset G'$, let $(s, f) \in G'$. Since $\{(t, g); t \supset s \wedge g \geq f\}$ is dense, there is $(t, g) \in G \cap D$. Then we have $(t, g) \leq (s, f)$, i.e., $(s, f) \in G$. \square

Theorem IV.2.3 ([Bla10]). A Hechler real x is a dominating real.

Proof. Let $f \in {}^\omega\omega \cap V$. Since $D = \{(s, g) \in \mathbb{P}; f \leq g\}$ is dense, there is $(s, g) \in G$ with $f \leq g$. Thanks to the previous theorem we have that $M[G] \models G = G'$, furthermore, $M[G] \models g \leq x$. Thus we obtain that $M[G] \models \forall f \in {}^\omega\omega \cap V f \leq x$. \square

Theorem IV.2.4 ([Bla10]). A Hechler forcing notion is σ -centered, in particular ccc poset.

Proof. Manifestly, by letting $\mathbb{P} = \cup\{(s, f) \in \mathbb{P} ; s \in {}^{<\omega}\omega\}$. \square

Definition IV.2.5 ([Jec03]). Fix a non-empty family $\mathcal{G} \subset {}^\omega\omega$ in the ground model. Let \mathbb{P} be a forcing notion such that:

1. $(s, E) \in \mathbb{P}$ provided that $s \in {}^{<\omega}\omega$ and $E \in [\mathcal{G}]^{<\omega}$,
2. $(s', E') \leq (s, E)$ provided that
 - (a) $s' \supset s$,
 - (b) $E' \supset E$,
 - (c) $\forall n \in \text{dom}(s') \setminus \text{dom}(s) \forall h \in E' h(n) \in s'(n)$.

Theorem IV.2.6. Let G be a generic in \mathbb{P} , as per **Definition IV.2.5**, a *Hechler real* x is a function $x = \cup\{s ; \exists E (s, E) \in G\} : \omega \rightarrow \omega$. Then, V can be recovered from x .

Proof. At the beginning of the proof, we note that since G is a filter and $\{(s, E) \in \mathbb{P} ; \exists n \ni m \ n \in \text{dom}(s)\}$ is dense for every n , x is a function with domain ω .

Let $G' = \{(s, E) \in \mathbb{P} ; s \subset x \wedge \forall f \in E \forall n \in \omega \setminus \text{dom}(s) (f(n) \leq x(n))\}$ and we verify that $G = G'$. To see $G \subset G'$. Let $(s, E) \in G$. We shall show that $h(n) \leq x(n)$ for any $h \in E$ and $n \in \omega \setminus \text{dom}(s)$. Let $n \in \omega \setminus \text{dom}(s)$. Since $D_n = \{(t, F) \in \mathbb{P} ; t \supset s \wedge n \in \text{dom}(t) \wedge F \supset E\}$ is dense, there is $(t, F) \in D \cap G$ and let (u, H) be a common extension in G . Then $(u, H) \leq (s, E)$ shows that $h(n) \leq u(n) = x(n)$ for any $h \in E$. To see $G' \subset G$. Let $(s, E) \in G'$. Since $\{(u, H) \in \mathbb{P} \supset u \supset t \wedge H \supset F\}$ is dense, there is $(u, H) \in G \cap D$. Then since for $n \in \text{dom}(u) \setminus \text{dom}(t)$ we have $u(n) = x(n) \geq h(n)$, we obtain that $(u, H) \leq (t, F)$, i.e., $(t, F) \in G$. \square

Similarly to **Theorem IV.2.3**, we have the following theorem:

Theorem IV.2.7 ([Jec03]). A Hechler real x is a dominating real.

Theorem IV.2.8. A Hechler forcing notion is σ -centered, in particular ccc poset.

Proof. Manifestly, by letting $\mathbb{P} = \cup\{(s, f) \in \mathbb{P} ; s \in {}^{<\omega}\omega\}$. \square

IV.3 Mathias Forcing

Definition IV.3.1. Let \mathbb{P} be a forcing notion such that $(s, A) \in \mathbb{P}$ provided that $s \in [\omega]^{<\omega}$, $A \in [\omega]^\omega$ and $\cup s \in \cap A$ and the order $\leq_{\mathbb{P}}$ is given by $(s, A) \leq_{\mathbb{P}} (t, B)$ provided that $s \supset t$, $A \subset B$ and $s \setminus t \subset B$.

Theorem IV.3.2. For a generic G the *Mathias real* x is an infinite subset of ω defined by $\cup\{s \in [\omega]^{<\omega} ; \exists A \in [\omega]^\omega (s, A) \in G\}$. G can be recovered from x .

Proof. At the beginning of the proof, since $D_n = \{(s, A) \in \mathbb{P}; \exists m \ni n \ m \subset s\}$ is dense for every $n \in \omega$, x is an infinite set.

We shall show that $G = G'$ for $G' = \{(s, A) \in \mathbb{P}; s \subset x \wedge x \subset s \cup A\}$. To see $G \subset G'$. Let $(s, A) \in G$. It suffices to verify that $x \subset s \cup A$. For any $a \in x$, there is $(t, B) \in G$ such that $a \in t$ and there is a common extension (u, C) . Since $a \in u$, we must have $a \in s$ or $a \in A$. Therefore, we have the subset relation. To see $G' \subset G$. Let $(s, A) \in G'$. Since a set

$$D_A = \{(t, B) \in \mathbb{P}; |A \cap B| < \aleph_0 \vee B \subset A\}$$

is dense, there is $(t, B) \in G \cap D$. Since $(t, B) \in G'$, elements in x are eventually contained in both A and B . This shows $B \subset A$. On the other hand, since a set

$$D_s = \{(t, B) \in \mathbb{P}; t \supset \vee s \not\subset \cup \{t, B\}\}$$

is dense, there is $(t', B') \in D_s \cap G$. Since we have $(t', B') \in G'$ we have $t' \supset s$.

Let (u, C) be a common extension for (t, B) and (t', B') in G . To see that (u, C) is an extension of (s, A) , we remain to verify that $u \setminus s \subset A$ but this is immediate from since $(u, c) \in G'$ and $(s, A) \in G'$ implies $u \subset x \subset s \cup A$. \square

Theorem IV.3.3. A Mathias real x is a dominating real, i.e., a function $x: n \mapsto x_n$ where $\text{rank}_{\in, x}(x_n) = n$ is a dominating real.

Proof. Let $f \in {}^\omega \omega$ in V . We may assume that f is a strictly increasing function, $f(0) = 0$ and identify f with an infinite set $\{f(i); i \in \omega\}$. Since a set, D_f ,

$$\{(t, B) \in \mathbb{P}; \text{“every interval in } \Pi_B \text{ possesses more than two elements in } f\text{.”}\}$$

is a dense set, where Π_f is an interval partition endowed with f , let $(t, B) \in G \cap D_f$. Thanks to the previous theorem, since we have $M[G] \models G = G'$, we obtain that $M[G] \models x \subset s \cup f$. This shows that, in $M[G]$, by construction there is some $n \in \omega$ such that the n -th least element in x is greater than in f , i.e., $M[G] \models \forall^\infty n \in \omega f(n) \leq x(n)$. \square

Theorem IV.3.4. Let \mathbb{P}_{ω_2} a ω_2 -countable support iteration with

$$\Vdash_{\mathbb{P}_\alpha} \text{“}\dot{\mathbb{Q}}_\alpha \text{ is a Mathias ordering.”}$$

Then we have $\Vdash_{\mathbb{P}_{\omega_2}} \mathfrak{c} = \aleph_2$.

Proof. To see (\geq) . Since the iteration adjoining ω_2 many Mathias reals, dominating reals, we have $\Vdash_{\omega_2} \mathfrak{c} \geq \aleph_2$. To see (\leq) . Note that by Theorem III.4.5 \mathbb{P}_{ω_2} preserves all cardinals. \square

Theorem IV.3.5 ([Bla10]). For the Mathias real X and a dense open family \mathcal{D} , there is $D \in \mathcal{D}$ such that $\Vdash X \subset D$.

Proof. Let G be generic, X the Mathias real and \mathcal{D} a dense open family. Let (s, A) be any condition. Since \mathcal{D} is dense, there is an extension $(s, A') \leq (s, A)$ with $A' \in \mathcal{D}$. We assert that $(s, A') \Vdash X \setminus s \subset A'$. Let $n \in \omega \setminus s$. If $n \in \dot{X}$. There is $(\dot{t}, \dot{B}) \in \dot{G}$ such that $n \in \dot{t}$ and $(\dot{t}, \dot{B}) \leq (s, A')$, then we have $n \in \dot{t} \setminus s \subset A'$. This shows that $\{(t, B) \in \mathbb{P}; (t, B) \Vdash (n \in X \implies n \in A')\}$ is dense below (s, A') . Therefore we obtain that $(s, A') \Vdash (n \in \dot{X} \implies n \in A')$ for each $n \in \omega \setminus s$. Moreover, $(s, A') \Vdash \dot{X} \subset \cup\{A', s\}$ and we have $\cup\{A', s\} \in \mathcal{D}$ since \mathcal{D} is dense open. Therefore, there is $D \in \mathcal{D}$ such that $\Vdash X \subset \dot{D}$. \square

Theorem IV.3.6 ([Bla10]). Let \mathbb{P}_{ω_2} be a forcing notion as per [Theorem IV.3.4](#). We have $\mathfrak{h} = \mathfrak{N}_2 = \mathfrak{c}$. Moreover, $\mathfrak{b} = \mathfrak{g} = \mathfrak{s} = \mathfrak{r} = \mathfrak{d} = \mathfrak{a} = \mathfrak{u} = \mathfrak{i} = \mathbf{non}(\mathcal{B}) = \mathbf{cof}(\mathcal{B}) = \mathbf{non}(\mathcal{L}) = \mathbf{cof}(\mathcal{L}) = \mathfrak{c}$.

Proof. For the first statement, it suffices to show that $\Vdash_{\omega_2} \mathfrak{N}_1 < \mathfrak{h}$. Let $\{\mathcal{D}_\xi; \xi \in \omega_1\}$ be a family of dense open sets. Let $\alpha \in \omega_2$ be an ordinal such that D_ξ in $V[G_\alpha]$ and let X be a Mathias real adjoined in $\alpha + 1$ step. Previous theorem asserts that there is $D_\xi \in \mathcal{D}_\xi$ such that $X \subset D$ for each $\xi \in \omega_1$. Then we obtain that $\Vdash_{\alpha+1} \forall \xi \in \omega_1 \exists D \in \mathcal{D}_\xi X \subset D$, this shows that $\Vdash_{\alpha+1} \forall \xi \in \omega_1 X \in \mathcal{D}_\xi$. Therefore, $\Vdash_{\omega_2} \mathfrak{N}_1 < \mathfrak{h}$ holds.

For the moreover part is immediate by the general consequence $(\mathfrak{h} \leq \mathfrak{s}, \mathfrak{b}, \mathfrak{g})$, $(\mathfrak{s} \leq \mathbf{non}(L))$, $(\mathfrak{b} \leq \mathfrak{a})$ and $(\mathfrak{b} \leq \mathfrak{r} \leq \mathfrak{u})$. (see 3.8, 5.19, 6.9, 6.27, 8.4 and 9.7 in [Bla10].) \square

Theorem IV.3.7. \mathbb{P} satisfies Axiom A.

Proof. Define \leq_n as

1. $\leq_0 = \leq_{\mathbb{P}}$
2. $(s, A) \leq_n (t, B)$ provided that
 - (2-a) $(s, A) \leq_{\mathbb{P}} (t, B)$,
 - (2-b) $s = t$,
 - (2-c) $\{x \in B; \text{rank}_{B, \in}(x) \in n\} \subset A$ for each $n \ni 0$.

For the conditions [A-1](#), [A-2](#) are clear. We remain to check the conditions [A-3](#) and [A-4'](#).

For [A-3](#). Let $\langle (s_i, A_i); i \in \omega \rangle$ be a fusion sequence. Define $s = s_0$ and $A = \cap\{A_i; i \in \omega\}$. To show $(s, A) \in \mathbb{P}$ and $\forall i \in \omega ((s, A) \leq_n (s_n, A_n))$, it remain to check that $A \in [\omega]^\omega$. For $n \in \omega$ there is $m \in A_{n+1}$ such that $\text{rank}_{A_{n+1}, \in}(m) = n$. Then we have $n \in m$ and $\forall i \in \omega (m \in A_i)$. Therefore we obtain that $\exists^\infty n \in \omega (n \in A)$, i.e., $A \in [\omega]^\omega$.

For [A-4'](#). Let $(s, A) \in \mathbb{P}$ with $(s, A) \Vdash \dot{a} \in V$ and let $n \in \omega$. Define $t = \{a \in A; \text{rank}_{A, \in}(a) \in n\}$ and enumerate $\mathcal{P}(t)$ such that $\{t_i; i \in k\} = \mathcal{P}(t)$. By the following lemma, there are B_i , $i \in k$ and X_{i+1} , $i \in k - 1$, recursively, satisfying:

1. $A \supset B_i \supset B_{i+1}$ for $i \in k - 1$,

$$2. (s \cup t_i, B_{i+1}) \Vdash \dot{a} \in \check{X}_{i+1}.$$

Define $B = t \cup B_k$ and $X = \cup\{X_{i+1}; i \in k-1\}$ and we verify that $(s, B) \leq_n (s, A)$ and $(s, B) \Vdash \dot{a} \in \check{X}$. The former is clear by the definition of t . To see that $(s, B) \Vdash \dot{a} \in \check{X}$. Let $(u, C) \leq_{\mathbb{P}} (s, B)$ since $u \setminus s \subset B$ there are $t_i \subset t$ and $u' \subset B_k \subset B$ such that $u \setminus s = t_i \cup u'$. We shall show that $(t, C \setminus t) \in \mathbb{P}$, $(u, C \setminus t) \leq (t, C)$ and $(u, C \setminus t) \Vdash \dot{a} \in X$. The first two properties are immediate. To see the last property, we shall show $(u, C \setminus t) \leq (s \cup t_i, B_{i+1})$ and this is immediate by the properties $B = t \cup B_k$ and $B_k \subset B_{i+1}$. \square

Lemma IV.3.8. For $(s, A) \in \mathbb{P}$, suppose that $(s, A) \Vdash \dot{a} \in V$. Then there are $B \subset A$ and countable set X in V such that $(s, B) \Vdash \dot{a} \in \check{X}$. Moreover, B may be chosen such that if $(t, C) \leq_{\mathbb{P}} (s, B)$, $a \in V$ and $(t, C) \Vdash \dot{a} = a$, then $(t, B \setminus (\cup t + 1)) \Vdash \dot{a} = a$.

Proof. First we construct $b_j \in A$ and $B_j \subset A$, recursively, which satisfies:

1. $B_0 \supset B_1 \supset \dots$,
2. $\forall b \in B_{j+1} (b_j \in b)$.

For the leading stage, let $B_0 = A$. Suppose that b_j and B_j , $j \in n$, are defined. Enumerate the subsets of $\{b_j; j \in n\}$, $\{s_{n,0}, \dots, s_{n,k-1}\} = \mathcal{P}(\{b_j; j \in n\})$. We construct a sequence $B_0^n \supset B_1^n \supset \dots \supset B_k^n$ as $B_0^n = B_{n-1}$ and

$$B_{i+1}^n = \begin{cases} C \subset B_i^n & \text{such that } (s \cup s_{n,i}, C) \Vdash \dot{a} \in V, \\ B_i^n & \text{"if such a } C \text{ does not exist."} \end{cases}$$

Let $b_n = \cap B_k^n$, $B_{n+1} = B_k^n \setminus \{b_n\}$. Then by letting, $B = \{b_j; j \in \omega\}$ and $X = \{a \in V; \exists t \subset B (\cup\{s, t\}, B \setminus (\cup t + 1)) \Vdash \dot{a} = a\}$, we assert that X is countable and $(s, B) \Vdash \dot{a} \in X$.

We assert that $(s \cup s_{n,i}, B_{i+1}^n) \Vdash \dot{a} \in V$ for every $n \in \omega$ and $i \in k_n$. Let $(t, C) \leq (s, B)$. Since $t \setminus s \subset B \subset \{b_i; i \in \omega\}$, there is $n \in \omega$ and $i \in k_n$ such that $t \setminus s = s_{n,i}$. Then we have $t = s \cup s_{n,i}$. Note that $(t, C) \leq (t, B_{i+1}^n)$ and $(t, B \setminus (\cup t + 1)) \leq (t, B_{i+1}^n)$ (we check at the end of the proof). The first inequality asserts that there is an $a \in V$ such that $(t, C) \Vdash \dot{a} = a$ and second inequality asserts that the a in X .

For the moreover part, by letting t as in above, let $a' \in V$ such that $(t, B_{i+1}^n) \Vdash \dot{a} = a'$. The first inequality asserts that $a = a'$ and the second inequality asserts that $a' \in X$.

Now we show the inequalities. To see $(t, C) \leq (t, B_{i+1}^n)$. Immediate by $t \subset \{b_i; i \in n\}$, $B^{n+1} \subset B_{i+1}^n$ and $B^{n+1} \subset B \setminus \{b_i; i \in n\}$. To see $(t, B \setminus (\cup t + 1)) \leq (t, B_{i+1}^n)$. Immediate by $B \setminus (\cup t + 1) \subset B^{n+1}$. \square

Theorem IV.3.9 (Pre-decision property, [Bau84]). Let φ be a sentence of the language of forcing. For any $(s, A) \in P$ there is $B \subset A$ such that either $(s, B) \Vdash \varphi$ or $(s, B) \Vdash \neg \varphi$.

Proof. Let A_0 and A_1 be maximal antichains in $\{p \in \mathbb{P}; p \Vdash \varphi\}$ and $\{p \in \mathbb{P}; p \Vdash \neg\varphi\}$, respectively and let \dot{a} be a nice name induced by antichain $\cup\{A_i; i \in 2\}$, then note that we have $(s, A) \Vdash \dot{a} \in V$.

Lemma IV.3.8 asserts that there is countable $B' \subset A$ with follows:

- (a) For $(t, C) \leq (s, B')$, $(t, C) \Vdash \varphi$ implies $(t, B' \setminus (\cup t + 1)) \Vdash \varphi$, and
- (b) for $(t, C) \leq (s, B')$, $(t, C) \Vdash \neg\varphi$ implies $(t, B' \setminus (\cup t + 1)) \Vdash \neg\varphi$.

Now, we define a sequence $\langle b_n \in B'; n \in \omega \rangle$ and $\langle B_n \in [B']^\omega; n \in \omega \rangle$, recursively, with following properties:

- $b_n \in b_{n+1}$ for each $n \in \omega$,
- $B_n \supset B_{n+1}$ for each $n \in \omega$.

Let $B_0 = B'$ and assume that $B_n \subset B'$ and b_n are defined for $n \in \omega$. We shall find an infinite B'_{n+1} such that for arbitrary $s' \subset \{b_i; i \in n\}$ we have the one of the following properties:

- (i) $\forall b \in B'_{n+1} (s \cup s' \cup \{b\}, B' \setminus (\cup \cup \{s', \{b\}\} + 1)) \Vdash \varphi$,
- (ii) $\forall b \in B'_{n+1} (s \cup s' \cup \{b\}, B' \setminus (\cup \cup \{s', \{b\}\} + 1)) \Vdash \neg\varphi$,
- (iii) both of (i) and (ii) does not hold.

Let $\{s_{n,i}; i \in k_n\} = \mathcal{P}(\{b_i; i \in n\})$ and define a decreasing sequence of infinite sets $B^n = B = B_0^n \supset B_1^n \supset \dots \supset B_{k_n}^n$ such that B_{i+1}^n and $s_{n,i}$ satisfies one of the condition in (i), (ii), (iii) by following processes:

Assume B_i^n is defined, for $i \in k_n$, since the one of the following sets are infinite, let B_i^{n+1} be such an infinite set:

- $T = \{b \in B_i^n; (\cup\{s, s_{n,i}, \{b\}\}, B' \setminus (\cup \cup \{s_{n,i}, \{b\}\} + 1)) \Vdash \varphi\}$,
- $F = \{b \in B_i^n; (\cup\{s, s_{n,i}, \{b\}\}, B' \setminus (\cup \cup \{s_{n,i}, \{b\}\} + 1)) \Vdash \neg\varphi\}$,
- $B_i^n \setminus \cup\{T, F\}$.

And let $B'_{n+1} = B_{k_n}^n$ and $b_n = \cap B'_{n+1}$ and $B_n = B'_{n+1} \setminus \{b_n\}$. Define $B = \{b_n; n \in \omega\}$.

To conclude the proof, we shall show that (s, B) forces φ or its negation. Assume that it does not force $\neg\varphi$. Then there is an extension $(t, C) \leq (s, B)$ which forces φ and we may assume that t has the least cardinality among them. We distinguish two case, according to whether $|s| = |t|$. If $|s| = |t|$. Since we have $s = t$, (s, C) forces φ , so does (s, B) , since $(t, B' \setminus (\cup t + 1))$ is an extension of (t, B) and by (a). If $|s| < |t|$. Since $t \setminus s \subset B$ there is an $n \in \omega$ such that $\cup(t \setminus s) = b_n$. Let $s' = t \setminus (s \cup \{b_n\})$. If we have $(s \cup s', B'_{n+1})$ forces φ , Since $(s \cup s', B'_{n+1}) \leq (s, B)$, the minimality for the cardinality asserts that $|t| \leq |s \cup s'| = |t| - 1$, a contradiction and we complete the proof.

Thus, we remain to show that $(s \cup s', B'_{n+1})$ forces φ . Since (t, C) forces φ , by (a), so does $(t, B' \setminus (\cup t + 1))$. Furthermore, $s' = t \setminus (s \cup \{b_n\})$ and $b_n \in B'_{n+1}$ witnesses that B'_{n+1} holds the condition (i). To see that $(s \cup s', B'_{n+1})$ forces φ , let (u, D) be an extension. We distinguish two cases, according to whether $u \setminus (s \cup s') = \emptyset$. Case I. $u \setminus \cup\{s, s'\} = \emptyset$. Let $b = \cap D$. Then we have $(u \cup \{b\}, D \setminus \{b\})$ is an extension for (u, D) and $(s \cup s' \cup \{d\}, B'_{n+1} \setminus (\cup(s' \cup \{d\} + 1)))$. And the later condition forces φ , we obtain that $(\cup\{s, s'\}, B'_{n+1}) \Vdash \varphi$. Case II. $u \setminus (s \cup s') \neq \emptyset$. Let $b = u \setminus \cup\{s, s'\}$ and $d = u \setminus (s \cup s' \cup \{b\})$. Since $(s \cup s' \cup \{b\}, B' \setminus \cup(s \cup \{b\} + 1))$ forces φ , we shall show that this condition possesses an extension (u, D) . To see that $D \subset B' \setminus (\cup(s' \cup \{b\} + 1))$, we note that $D \subset B'_{n+1} \subset B'$ and for arbitrary $c \in D$, $\cup u \in x$, $\cup s' \leq \cup u \in x$ and $b \leq \cup u \in x$ imply $x \notin \cup(s' \cup \{b\} + 1)$. To see that $u \setminus (s \cup s' \cup \{b\}) \subset B' \setminus (\cup(s' \cup \{b\} + 1))$. It suffices to see that $d \subset B' \setminus (\cup(s' \cup \{b\} + 1))$. For $x \in d$, $x \in u \setminus \{s, s'\}$ shows that $x \in B'_{n+1} \subset B'$ and $x \notin s' \cup \{b\}$ shows that $x \notin \cup(s' \cup \{b\} + 1)$. \square

Corollary IV.3.10. Let X be a finite set in V , $(s, A) \Vdash \dot{a} \in X$ then there is $B \subset A$ and $a \in X$ such that $(s, B) \Vdash \dot{a} = a$. Moreover, for $n \in \omega$ there is $(t, B) \leq_n (s, A)$ and $Y \subset X$ of size $\leq 2^n$ such that $(t, B) \Vdash \dot{a} \in Y$.

Proof. Let $X = \{a_i; i \in k\}$ in V and we define the squence $A \supset B_0 \supset B_1 \supset \dots \supset B_k$, recursively, such that $(s, B_i) \Vdash \dot{a} \in X \setminus \{a_j; j \in i\}$ or $\exists j \leq i$ $(s, B_i) \Vdash \dot{a} = a_j$. For the leading stage. **Theorem IV.3.9** asserts that there is finite $B_0 \subset A$ such that either

- (i) $(s, B_0) \Vdash \dot{a} = a_0$, or
- (ii) $(s, B_0) \Vdash \dot{a} \neq a_0$.

Note that $(s, B_0) \Vdash \dot{a} \neq a_0$ implies $(s, B_0) \Vdash \dot{a} \in X \setminus \{a_0\}$. For the successor stages. If B_i have the condition (i), let $B_{i+1} = B_i$. If B_i have condition (ii), by **Theorem IV.3.9** there is $B_{i+1} \subset B_i$ such that either $(s, B_{i+1}) \Vdash \dot{a} = a_{i+1}$ or $(s, B_{i+1}) \Vdash \dot{a} \in \{a_j; j \in n+1\}$. Since we have $(s, B_n) \not\Vdash \dot{a} \in \emptyset$. There is $j \leq k$ such that $(s, B_n) \Vdash \dot{a} = a_j$. To see the moreover part, fix an n . For the leading stage, since $(s, A) \Vdash \dot{a} \in X$ there is an extension (t, B) and $x \in X$ such that $(t, B) \Vdash \dot{a} = x$. So by letting $Y = \{x\}$ we have the $|Y| \leq 2^0 = 1$ and $(s, B) \Vdash \dot{a} \in Y$. For the successor stages. Let $U_n = \{a \in A; \text{rank}_{\in, A}(a) \in n\}$ and enumerate $\{s_i^n; i \in 2^n\} = \mathcal{P}(U_n)$. Define $B_0^n, B_{i+1}^n \subset A$ and $a_i \in X$ for $i \in 2^n$, recursively:

- $B_0^n = A$,
- B_{i+1}^n and a_i be sets such that $(\cup\{s, s_i^n\}, B_{i+1}^n) \Vdash \dot{a} = a_i$.

note that for the successor stage, since $(\cup\{s, s_i^n\}, B_i^n \setminus (\cup s_i^n + 1))$ is an extension for (s, A) , the first statement asserts that there are desired sets B_{i+1}^n and a_i . Let $B = \cup\{B_{2^n}^n, U_n\}$ and $Y = \{a_i; i \in 2^n\}$. To conclude the proof, we shall verify that $|Y| \leq 2^n$, $(s, B) \leq_n (s, A)$ and $(s, B) \Vdash \dot{a} \in Y$. The first two statements are obvious, we see the last condition. Let (t, C) be an extension for (s, B) .

By letting $W = (t \setminus s) \cap B_{2^n}^n$ and $s_j^n = (t \setminus s) \cap U_n$ for some $j \in 2^n$, we have $t \setminus s = \cup\{W, s_j^n\}$. To conclude the proof, we shall show that $(t, C \cap B_{j+1}^n)$ is an extension for (t, C) and $(\cup\{s, s_j^n\}, B_{j+1}^n)$. The former is immediate and the later is immediate by the fact that $W \subset B_{2^n}^n \subset B_{j+1}^n$. Then, since the later condition forces $\dot{a} \in Y$, we obtain that $(s, B) \Vdash \dot{a} \in Y$. \square

IV.4 Sacks Forcing (Perfect Set Forcing)

Definition IV.4.1. Let \mathbb{S} be a poset with order $\leq_{\mathbb{S}} = \subset$ such that $p \in \mathbb{S}$ provided that

- (a) $\emptyset \neq p \subset \cup\{^n 2; n \in \omega\}$,
- (b) $\forall s \in p \forall n \in \omega (s|_n \in p)$,
- (c) $\forall q \in p \exists s, t \in p (q \subset s \wedge q \subset t \wedge \exists n \in \omega s(n) \neq t(n))$.

Note that the condition (c) asserts that every $p \in \mathbb{S}$ is a perfect set in $2^{<\omega}$.

For $p \in \mathbb{S}$ and $s \in p$ define *the degree of s in p* , $\deg(s, p)$, by the cardinality of $\{n \in \omega; \exists t \in p t(n) \neq s(n)\}$. $s \in p$ is a *splitting node of p* provided that $s \smallfrown \langle 0 \rangle, s \smallfrown \langle 1 \rangle \in p$ in addition if we have $\deg(s, p) = n$ then s is an *n -splitting node of p* . $p|_s$ denote the codition $\{t \in p; t \subset s \vee s \subset t\}$.

Theorem IV.4.2 ([Jec03]). For generic G the Sacks real, or generic branch, is a function $f: \omega \rightarrow 2$ given by $f = \cap\{\cup p; p \in G\}$. Then G can be recovered from x .

Proof. First, we assert that f is a function with $\omega \rightarrow 2$. Since G is a filter, for any $n \in \omega$ and for distinct $p, q \in G$ we obtain that $p \cap q \cap ^n 2 \neq \emptyset$. Moreover since $^n 2$ is finite, we obtain that $S_n = (\cap G) \cap ^n 2 \neq \emptyset$. Furthermore, since $D_n = \{p \in \mathbb{S}; \text{“} \cup p \text{ does not possess either } \langle n-1, 0 \rangle \text{ or } \langle n-1, 1 \rangle \text{”}\}$ is dense in \mathbb{S} for $n \geq 0$, S_n must be a singleton. Therefore, $f: \omega \rightarrow 2$ is a function.

Second, we assert that $M[f] = M[G]$. Since $f \in M[G]$ manifestly holds, it suffices to verify that $G = G'$ for $G' = \{p \in \mathbb{S}; f \subset \cup p\}$. The relation (\subset) is obvious. To see the (\supset) . Since $D = \{q \in G; q \subset p \vee \neg(\exists f \in (\cup p) \cap (\cup q) f \in ^\omega 2)\}$ is dense, there is $q \in G \cap D$. Since $q \in G'$, we have $f \in (\cup p) \cap (\cup q)$. Thus we obtain that $q \subset p$, i.e., $p \in G$. \square

Theorem IV.4.3.

- (i) For $p \in \mathbb{S}$, p possesses a binary subtree,
- (ii) Define \leq_n as $p \leq_n q$ provided that $p \leq_{\mathbb{S}} q$ and $\forall s \in q (\deg(s, q) \leq n \implies s \in p)$. Note that $\leq_0 = \leq_{\mathbb{S}}$ holds. Then,
 - \mathbb{S} satisfies Axiom A. Moreover we can say that $|\{r \in I; r \not\leq q\}| \leq 2^n$ in the condition A-4.

Proof. For (i). Let $p \in \mathbb{S}$. Define a binary subtree $T_p = \cup\{T_p^n = {}^{(n)} 2; n \in \omega\}$, recursively on a height $n \in \omega$,

1. For a leading stage. $T_p^0 = \{\emptyset\}$.
2. For a successor stages. For $s^\frown \langle i \rangle \in {}^{(n+1)}2$, let $N = \min\{m \in \omega; \exists t, u \in p (s \subset t_0, t_1 \wedge \text{dom}(t_0) = \text{dom}(t_1) = m + 1 \wedge t_0(m) = 0 \wedge t_1(m) = 1)\}$ and $t_0, t_1 \in p$ which endow N . Define $T_p^{n+1}(s^\frown \langle i \rangle) = t_i$.

Then we obtain a subtree T_p . Note that the heigh of s in T_p and the degree of s in p are the same.

For (ii). The conditions [A-1](#) and [A-2](#) are manifestly. We check the conditions [A-3](#) and [A-4](#).

For condition [A-3](#). Let $\langle p_n \in \mathbb{S}; n \in \omega \rangle$ with $p_{n+1} \leq_n p_n$ for each $n \in \omega$. Let $q = \cap \{p_n; n \in \omega\}$. First we check that $q \in \mathbb{S}$. Since the condition (a) and (b) are obvious, we remain to see the condition (c). Let $s \in q$ and $n = \deg(s, p_0)$. Since we have $\deg(s, p_{n+1}) \leq n$, $p_{n+1} \in \mathbb{S}$ and by the result of (i), there are $t, u \in p_{n+1}$ such that $\deg(t, p_{n+1}) = \deg(u, p_{n+1}) = n + 1$. To see that $t, u \in q$, let $m \in \omega$ and we shall distinguish two cases, according to whether $m \leq n + 1$. If $m \leq n + 1$. Immediate by the fact $p_{n+1} \subset p_m$. If $n + 1 \in m$. Note that for $s, t \in p_{n+1}$ with $\deg(s, p_{n+1}) \leq n + 1$, $p_{n+2} \leq_{n+1} p_{n+1}$ implies $s, t \in p_{n+2}$. Thus, by inductively, we obtain that $s, t \in p_m$. Note that the conditions $q \subset s, t$, $s \not\subset t$ and $t \not\subset s$ are clear, thus $q \in \mathbb{S}$. To conclude the proof for condition [A-3](#), we show that $q \subset p_n$, for every $n \in \omega$. This can prove by the mimik the proof of $t, u \in q$.

For condition [A-4](#). Let I be an antichain over \mathbb{S} , $p \in \mathbb{S}$ and $n \in \omega$. Let $\Gamma(p, n) = \{s \in p; \deg(s, p) = n \wedge s^\frown \langle 0 \rangle \in p \wedge s^\frown \langle 1 \rangle \in p\}$ and $p|_s = \{t \in p; t \subset s \vee s \subset t\} \in \mathbb{S}$. For $s \in \Gamma(p, n)$ since I is an antichain there is $q_s \leq_{\mathbb{S}} p|_s$ such that either $\exists r_s \in I (q_s \leq_{\mathbb{S}} r_s)$ or $\forall r \in I (q_s \perp_{\mathbb{S}} r)$. We choose q_s and r_s (if exists and note that uniquely exists) for each $s \in \Gamma(p, n)$ and define $q = \cup \{q_s; s \in \Gamma(p, n)\}$ and we check the q is the desired one. For $q \leq_n p$. Let $t \in p$ with $\deg(t, p) \leq n$, by the result of (i), there is $s \in \Gamma(p, n)$ such that $t \subset s$. Since we have $t \in p|_s$, $\deg(t, p|_s) \leq \deg(s, p|_s) = 0$ and $q_s \leq_0 p|_s$, we obtain $t \in q_s \subset q$. Thus $q \leq_n p$. For $|\{r \in I; r \not\leq_{\mathbb{S}} q\}| \leq 2^n$. First we note that $|\Gamma(p, n)| \leq 2^n$ since the number of nodes in a binary tree with height n is 2^n . If we obtain the equation $\{r \in I; r \not\leq_{\mathbb{S}} q\} = \{r \in I; \exists s \in \Gamma(p, n) (q_s \not\leq_{\mathbb{S}} r)\}$, we have $|\{r \in I; r \not\leq_{\mathbb{S}} q\}| \leq 2^n$.

Therefore we remain to show the equality. The relation, (\supset) , is immediate by $q_s \subset q$. Before we show the (\subset) , we assert that the following trivial facts.

1. For $\{s_0, s_1\} \in [\Gamma(p, n)]^2$, $q_{s_0} \perp_{\mathbb{S}} q_{s_1}$.
2. For $\mathcal{D}, \mathcal{F} \in [\Gamma(p, n)]^{<\omega}$, if $\mathcal{D} \cap \mathcal{F} = \emptyset$ then $\cup \{q_s; s \in \mathcal{D}\} \perp_{\mathbb{S}} \cup \{q_s; s \in \mathcal{F}\}$.

To see (\subset) . Let $r \in I$ and let $q' \leq_{\mathbb{S}} p$ with $q' \leq_{\mathbb{S}} q, r$. Suppose that we have $\forall s \in \Gamma(p, n) (q_s \perp_{\mathbb{S}} q')$, inductively and by the above facts, we obtain that $\cup \{q_s; s \in \Gamma(p, n)\} \perp_{\mathbb{S}} q'$, i.e., $q \perp_{\mathbb{S}} q'$, a contradiction. Thus for some $s \in \Gamma(p, n)$ we have $q_s \not\leq_{\mathbb{S}} q'$ moreover, $q_s \not\leq_{\mathbb{S}} r$. \square

Definition IV.4.4 ([[Jec03](#)]). A generic filter G is *minimal over the ground model* M provided that every set of ordinals in $M[G]$, either $X \in M$ or $G \in M[X]$. Here $M[X]$ is the least extension with $M \subset M[X]$ and $X \in M[X]$.

Theorem IV.4.5 ([Jec03]). A generic filter G of a Sacks forcing notion is minimal over the ground model.

Proof. Let $X \in V[G]$ be a set of ordinals and assume that $\nVdash \dot{X} \in V$ and let p be a condition which forces $\dot{X} \notin V$.

Define a fusion sequence $\langle p_n; n \in \omega \rangle$, and ordinals γ_s in $V[G]$, recursively. Let $p_0 = p$. Let S_n be the set of an n -splitting nodes of p_{n-1} . For each $s \in S_n$ let γ_s be an ordinal which does not decide $\gamma_s \in \dot{X}$ by $p_{n-1}|_s$. Then for $i \in 2$, since $p_{n-1}|_{s \smallfrown \langle i \rangle}$ is an extension of $p_{n-1}|_s$, there is an extension q_s^i such that $q_s^0 \Vdash \gamma_s \notin \dot{X}$ and $q_s^1 \Vdash \gamma_s \in \dot{X}$. Then by letting p be the amalgamation $\{q_s^i; s \in S_n \wedge i \in 2\}$ into p_{n-1} :

$$p_n = \cup \{ \{ t \in p_{n-1}; (t \subset s) \vee (s \subset t \wedge t \in q_s^i) \}; s \in S_n \wedge i \in 2 \} \in \mathbb{P}$$

we have $p_n \leq_n p_{n-1}$ and $q_s^i \leq p_n$ for each $s \in S_n$ and $i \in 2$. And moreover, there is a fusion q such that $q \leq_n p_n$ for each $n \in \omega$. \square

Theorem IV.4.6 (Sacks property, [Bla10]). Let $s \in \mathbb{S}$ and $Y \in V$ satisfying $s \Vdash \dot{f}: \omega \rightarrow \check{Y}$ there is $g: \omega \rightarrow X$ in V with $X \in V$ such that $f(n) \in g(n)$ and $|g(n)| \leq 2^n$ for each $n \in \omega$.

Proof. For every $n \in \omega$ we have $s \Vdash (\dot{f})(n) \in \check{Y}$, so by applying **A-4'**, there is q_n and $X_n \in V$ with $|X_n| \leq 2^n$ such that $q_n \leq_n s$ and $q_n \Vdash (\dot{f})(n) \in \check{X}_n$. By letting the function g with assign the n the X_n , we obtain the desired one. \square

IV.5 Shooting A Club by Finite Conditions

Definition IV.5.1 ([Bau84]). Let \mathbb{P} be a forcing notion with an order $\leq_{\mathbb{P}}$ such that:

1. $p \in \mathbb{P}$ provided that p is a function $p: 2n \rightarrow \omega_1$ for $n \in \omega \setminus 1$ such that $p(2i+1) \in p(2i+2)$ for any $i \in n-1$, we simply write $p = \{ \langle \xi_{2i}, \xi_{2i+1} \rangle; i \in n \}$,
2. For $p, q \in \mathbb{P}$, $q \leq_{\mathbb{P}} p$ provided that $q \supset p$.

Theorem IV.5.2. \mathbb{P} is proper.

Proof. Let $\lambda > 2^{|\mathbb{P}|}$ be regular uncountable, $M \preceq H_\lambda$ with $\mathbb{P} \in M$ and $p \in M \cap \mathbb{P}$. We shall find an extension $q \leq p$ which is (M, \mathbb{P}) -generic. Fix a $\beta_0 \in \omega_1$ such that $\omega_1 \cap M \in \beta_0$ and let $p' = p \cup \{ \langle \omega_1 \cap M, \beta_0 \rangle \}$. We assert that p' is a witness. Let $D \subset \mathbb{P}$ be a dense set with $D \in M$ and let $I \subset D$ be a maximal antichain. Let $q \leq p'$ be an extension. Since there is $r \in I$ such that $r \not\leq q$. Now we note that the following properties (we shall proved at the end of the proof):

- (a.) $s \cap M \in M$ for any condition $s \in \mathbb{P}$,
- (b.) $s \in M$ implies $\cup s \subset M$ for any condition $s \in \mathbb{P}$.

Thus, there is an $r' \in I \cap M$ such that $r \cap M = r' \cap M$ and $r' \subset M$. Then $r' \cup q = (r \cap M) \cup q \in \mathbb{P}$, i.e., $r' \not\perp q$.

To see that (a). For $s \in \mathbb{P}$. Since $s \cap M$ is finite, there is a bijection $f: |s \cap M| \rightarrow s \cap M$ and thus $s \cap M \in M$. To see that (b). For $s \in M \cap \mathbb{P}$, there is a surjection $f: \omega \rightarrow s$ in M . Thus we obtain that $\cup s \subset M$. \square

Theorem IV.5.3. For any generic G , $\mathbb{1} \Vdash_{\mathbb{P}} \text{“} \dot{C} = \{\alpha \in \omega_1; \exists \beta \in \omega_1 \langle \alpha, \beta \rangle \in \cup \dot{G}\} \text{ is club in } \omega_1\text{”}$.

Proof. To see \dot{C} is unbounded. Let $\gamma \in \omega_1$. Since $D_\gamma = \{p \in \mathbb{P}; \exists \langle \alpha, \beta \rangle \in \cup p \gamma \in [\alpha, \beta]\}$ is dense in \mathbb{P} , there is $p \in \dot{G}$ and $\langle \alpha, \beta \rangle \in p$ such that $\gamma \in [\alpha, \beta]$. And, similarly, for $\beta + 1$ there is $q \in \dot{G}$ such that $\beta + 1 \in [\alpha', \beta']$ for some $\langle \alpha', \beta' \rangle \in q$. Then since $\beta + 1 \leq \alpha'$, $\beta + 1$ witnesses that \dot{C} possesses an element $\geq \gamma$. Thus $\models \cup \dot{C} = \omega_1$.

To see that \dot{C} is closed set, it suffices show that $\omega_1 \setminus \dot{C} = \cup \{[\alpha + 1, \beta]; \langle \alpha, \beta \rangle \in \cup \dot{G}\}$ since $[\alpha, \beta]$ is an open interval $(\alpha, \beta + 1)$. For (\subset) . Let $\gamma \in \omega_1 \setminus \dot{C}$. By genericity there is $\langle \alpha, \beta \rangle \in \cup \dot{G}$ such that $\gamma \in [\alpha, \beta]$, moreover since $\gamma \notin \dot{C}$, $\gamma \in [\alpha + 1, \beta]$. For (\supset) . Let $\gamma \in [\alpha + 1, \beta]$ and $\langle \alpha, \beta \rangle \in p \in G$. If $\gamma \in \dot{C}$, there is $\delta \in \omega_1$ such that $\langle \gamma, \delta \rangle \in q$ for some $q \in G$. However, this contradicts that $\alpha = \gamma$ since p and q are compatible. \square

Theorem IV.5.4. There are no sequence of order relations $\langle \leq_n; n \in \omega \rangle$ which satisfies Axiom A.

Proof. Assume that there is a sequence of order relations $\langle \leq_n; n \in \omega \rangle$ which satisfies Axiom A. At the beginning of the proof, we assert that for $p \in \mathbb{P}$ there is an uncountable antichain I_p such that every $q \in I_p$ is compatible with p . Since p is a finite set there is $\alpha \in \omega_1$ such that $\cup \cup p \in \alpha$. Then by letting $I_p = \{\cup \{p, \{\langle \alpha, \beta \rangle\}\}; \alpha \in \beta \in \omega_1\}$ we obtain the desired uncountable antichain.

To have a contradiction, we define a fusion sequence $p_n, n \in \omega$ with $|p_n| \geq n$, recursively,

1. For a leading stage. Fix arbitrary $p_0 \in \mathbb{P}$,
2. For a successor stages. For p_n and I_{p_n} , by applying the condition A-4, we obtain a $q \in \mathbb{P}$ such that $q \leq_{n+1} p_n$ and $\{r \in I_{p_n}; r \perp_{\mathbb{P}} q\}$ is countable. If $|q| \leq n$, by $q \leq_{n+1} p_n$, $|p_n| \geq n$ and the condition A-2 shows that $q = p_n$. However this contrary to that $\{r \in I_{p_n}; r \perp_{\mathbb{P}} p_n\}$ is countable. Thus we have $q \in \mathbb{P}$ such that $q \leq_{n+1} p_n$ and $|q| \geq n + 1$.

Therefore, we obtain a specific fusion sequence. However the condition A-3 asserts that there is a fusion $p \in \mathbb{P}$ such that $q \leq_n p_n$ for every n , moreover $|q| \geq n$ for every n , a contradiction. \square

V The Concrete Forcing Examples

V.1 Axiom of Choice

V.2 Continuum Hypothesis

V.3 Martin Axiom

V.4 Borel Conjecture

Definition V.4.1. A subset of reals A is *strong measure zero* provided that for each sequence of positive reals $\langle \varepsilon_n; n \in \omega \rangle$ there is a sequence of intervals $\langle I_n \subset \mathbb{R}; n \in \omega \rangle$ such that $|I_n| \leq \varepsilon_n$ for each $n \in \omega$ and $A \subset \bigcup \{I_n; n \in \omega\}$.

Theorem V.4.2. X is strongly measure zero over ${}^\omega 2$ iff $\forall f \in {}^\omega \omega \exists \langle g_n: f(n) \rightarrow 2; n \in \omega \rangle (\forall h \in X \exists n \in \omega \ g_n \subset h)$.

Lemma V.4.3 ([JSW90]). $\mathfrak{b} = \aleph_1$ implies there is an uncountable strongly measure zero set.

Proof. At the beginning of the proof, we shall define a set $X \subset \mathbb{R}$ is *concentrated on* \mathbb{Q} provided that for arbitrary open set \mathcal{U} with $\mathcal{U} \supset \mathbb{Q}$ we have $X \setminus \mathcal{U}$ is countable and we assert that X is concentrated on \mathbb{Q} implies that X is a strongly measure zero set.

Enumerate $\mathbb{Q} = \{q_i; i \in \omega\}$ and let $\langle \varepsilon_i; i \in \omega \rangle$ be a sequence of positive reals. Since $\mathcal{U} = \bigcup \{(q_i - \varepsilon_i, q_i + \varepsilon_i) \subset \mathbb{R}; i \in \omega\}$ is an open set with $\mathcal{U} \supset \mathbb{Q}$, $X \setminus \mathcal{U}$ is countable. Moreover since X have the form of the union of \mathcal{U} and countable $X \setminus \mathcal{U}$, so X can covered by countable intervals.

To conclude the proof, let \mathcal{F} be an unbounded family of size ω_1 and assume that $f_i <^* f_j$ holds for $i \in j \in \omega$. Identify ${}^\omega \omega$ with $P = [0, 1] \setminus \mathbb{Q}$. To see that \mathcal{F} is concentrated on \mathbb{Q} , let \mathcal{U} be an open set with $\mathcal{U} \supset \mathbb{Q}$. Since $K = [0, 1] \setminus \mathcal{U}$ is compact and K is covered by $\bigcup \{f \in {}^\omega \omega; f < g\}; g \in {}^\omega \omega\}$, there is $g \in {}^\omega \omega$ such that K is covered by $\{f \in {}^\omega \omega; f <^* g\}$ for some $g \in {}^\omega \omega$. Furthermore, we have $\mathcal{F} \setminus \mathcal{U} \subset \{f \in \mathcal{F}; f <^* g\}$. Therefore, since \mathcal{F} is unbounded family, $\mathcal{F} \setminus \mathcal{U}$ is countable of size \aleph_1 , \mathcal{F} is an uncountable strongly measure zero set. \square

Hereafter, let \mathbb{P}_α be an α -stage iteration with countable support such that $\beta \in \alpha$:

$$1 \Vdash_{\mathbb{P}_\beta} \text{“}\dot{\mathbb{Q}}_\beta \text{ is the Mathias ordering.”}$$

Lemma V.4.4. Let X be finite in V and $p \Vdash_\alpha \dot{a} \in X$. For any finite $F \subset \alpha$ and $n \in \omega$ there is $q \leq_{F,n} p$ and $Y \subset X$ such that $|Y| \leq 2^{2 \cdot |F|}$ and $q \Vdash_\alpha \dot{a} \in Y$.

Proof. We prove by induction on α . The leading stage is immediate by **Theorem IV.3.10**. For the successor stages, $\alpha = \beta + 1$. We distinguish two cases, according to whether $\beta \notin F$. If $\beta \notin F$, note that we have $F \subset \beta$. Since we have $p \Vdash_\alpha \dot{a} \in X$, there is $\dot{f}, \dot{b} \in V^{\mathbb{P}_\beta}$ such that $p \restriction_\beta$ forces, at the β -stage,

$$\dot{f} \leq_{\dot{\mathbb{Q}}_\beta} p(\beta), \dot{b} \in X \text{ and } \dot{f} \Vdash_{\dot{\mathbb{Q}}_\beta} \dot{a} = \dot{b}.$$

Then, by induction hypothesis, there is an extension $q' \leq_{F,n} p|_\beta$ and $Y \subset X$ such that $|Y| \leq 2^{2 \cdot |F|}$ and q' forces, at the β -stage,

$$\dot{f} \leq_{\dot{Q}_\beta} p(\beta) \text{ and } \exists \dot{b} \in Y \ \dot{f} \Vdash_{\dot{Q}_\beta} \dot{a} = \dot{b}.$$

letting $q = \cup\{q', \langle \beta, \dot{f} \rangle\} \in \mathbb{P}_\alpha$, we have $q \leq_{F,n} p$ and $q \Vdash \dot{a} \in Y$. If $\beta \in F$. Applying [Theorem IV.3.10](#) to $p|_\beta \Vdash_{\mathbb{P}_\beta} (p(\beta) \Vdash_{\dot{Q}_\beta} \dot{a} \in X)$, there is $\dot{q}, \dot{f}, \dot{Y} \in V^{\mathbb{P}_\beta}$ such that $p|_\beta$ forces, at β -stage,

$$\dot{q} \leq_n^\beta p(\beta), \ \dot{Y} \subset X, \ \exists \dot{f}: \dot{Y} \rightarrow 2^n \text{ ("}\dot{f} \text{ is injective" and } \dot{q} \Vdash_{\dot{Q}_\beta} \dot{a} \in \dot{Y}).$$

Since for each $i \in k_n$ we have $p|_\beta \Vdash_{\mathbb{P}_\beta} \dot{f}(i) \in X$ by induction hypothesis we obtain the sequence q_i and Y_i with

- $p|_\beta = q_0 \geq_{F \cap \beta, n} q_1 \geq_{F \cap \beta, n} q_2 \geq_{F \cap \beta, n} \dots \geq_{F \cap \beta, n} q_{k_n}$,
- $|Y_i| \leq 2^{n \cdot (|F|-1)}$,
- $q_{i+1} \Vdash_{\mathbb{P}_\beta} \dot{f}(\beta) \in Y_{i+1}$.

Then we have $q_{k_n} \geq_{F \cap \beta, n} p|_\beta$ and $q_{k_n} \Vdash_{\mathbb{P}_\beta} \dot{f}(i) \in Y_{i+1}$ for each $i \in k_n$. Then by letting $Y = \cup\{Y_{i+1}; i \in 2^n\}$ and $q = \cup\{q_{k_n}, \langle \beta, \dot{q} \rangle\}$, we obtain that $q \in \mathbb{P}_\alpha$, $q \geq_{F \cap \beta, n} p$ and $|Y| \leq 2^{n \cdot (|F|-1)} \cdot 2^n = 2^{n \cdot |F|}$. For the limit stages. Since there is β such that $F \subset \beta \subset \alpha$. [We can prove by mimik the case of \$\beta \notin F\$ in the successor stages.](#) \square

Lemma V.4.5. Let $\langle X_n; n \in \omega \rangle$ be a sequence of finite sets and assume that $p \Vdash_\alpha \forall n \in \omega \ \dot{f}(n) \in X_n$. Then there is $q \leq p$ and a sequence $\langle Y_n; n \in \omega \rangle$ in V such that $Y_n \subset X_n$ and $|Y_n| \leq 2^{(n^2)}$ for each $n \in \omega$ and $q \Vdash_\alpha \forall n \in \omega \ \dot{f}(n) \in Y_n$.

Proof. We shall define a sequence of sets $\langle Y_n; n \in \omega \rangle$ and an (F, n) -fusion sequence, $\langle (p_n, F_n); n \in \omega \rangle$ such that $|F_n| \leq n$ and $p_{n+1} \Vdash_{\mathbb{P}_\alpha} \dot{f}(n) \in Y_n$.

Let $p_0 = p$ and $F_0 = \emptyset$. Suppose we have p_n and F_n , we shall find a p_{n+1} and F_{n+1} . Since $p_n \Vdash \dot{f}(n) \in X_n$ by [Lemma V.4.4](#) there is $p_{n+1} \leq_{F_n, n} p_n$ and $Y_n \subset X_n$ such that $|Y_n| \leq 2^{n \cdot |F_n|}$ and $p_{n+1} \Vdash \dot{f}(n) \in Y_n$. Fix functors h_1 and h_2 which assigns to $\Sigma\{j; j \in n\} + i$ to n and to j , respectively, for $i \in n$ and enumerate $\text{supt}(p_m) = \{\xi_i^m; i \in \omega\}$. $Y_{n+1} = \cup\{F_n, \{\xi_{h_2(n)}^{h_1(n)}\}\}$. Then we obtain that $\cup\{F_n; n \in \omega\} = \cup\{\text{supt}(p_n); n \in \omega\}$. Thus we obtain the (F, n) -fusion sequence $\langle (p_n, F_n); n \in \omega \rangle$.

[Lemma III.4.7](#) asserts that there is a fusion $q \in \mathbb{P}_\alpha$. Then we have $q \Vdash_{\mathbb{P}_\alpha} \forall n \in \omega \ \dot{f}(n) \in Y_n$ and moreover we have $q \leq_{\mathbb{P}_\alpha} p$, $Y_n \subset X_n$ and $|Y_n| \leq 2^{(n^2)}$ for any $n \in \omega$. \square

Theorem V.4.6. Assume that CH. \Vdash_{ω_2} “Borel conjecture.” + $\mathfrak{c} = \aleph_2$.

Proof. We have seen that $\Vdash_{\omega_2} \mathfrak{c} = \aleph_2$ in [Theorem IV.3.4](#).

At the beginning of the proof, we show that there is a $\beta \in \omega_2$ such that $X \in [G_\beta]$. To see this statement, enumerate $X = \{x_\alpha; \alpha \in \omega\}$ in $V[G_{\omega_2}]$. For arbitrary α and $n \in \omega$, since $\{p \in \mathbb{P}_{\omega_2}; \exists i \in 2 \ p \Vdash_{\omega_2} x_\alpha(n) = i\}$ is dense in \mathbb{P}_{ω_2} , let A_n^α be a maximal antichain in it dense set and $f_n^\alpha: A_n^\alpha \rightarrow 2$ a function in V which assign to condition p the witness $i \in 2$. Then we obtain that $p \Vdash_{\omega_2} x_\alpha(n) = f_n^\alpha(p)$. Note that since \mathbb{P}_{ω_1} has an \aleph_2 -cc, there is a $\beta \in \omega_2$ such that

$$\bigcup_{\alpha \in \omega_1} \bigcup_{n \in \omega} \bigcup_{p \in A_n^\alpha} \text{supt}(p) \in \beta.$$

Therefore, for a name $\mathring{Y} = \{\langle n, \check{f}_\alpha^n(p) \rangle; p \in A_n \wedge n \in \omega\}; \alpha \in \omega\}$, we obtain that $Y \in V[G_\beta]$ and $\Vdash_{\omega_2} \mathring{Y} = X$. \square

Index

Symbols

\nsubseteq	3
\perp	3

A

amalgamation	32
antichain	3

C

cardinality hereditarily	18
ccc	9
centered	11
closed	1
club set	1
compatible	3
complete embedding	4
concentrated	34
countably closed	11

D

degree	30
Delta System Lemma	3
dense embedding	5
dense open	3
dense set	3
diagonal intersection	2
direct limit	8
dominating real	6

E

Extension Lemma (proper forcing)	21
--	----

F

(F, n) -dense	12
(F, n) -fusion sequence	12
forcing notion	3
fusion	11
fusion sequence	11

H

Hechler real	23, 24
Hodor Theorem	2

I

incompatible	3
--------------------	---

inverse limit	8
iterated forcing	8

K

$< \kappa$ -support	8
$< \kappa$ -cc forcing	9

M

(M, \mathbb{P}) -generic	18
Mathias real	25
maximal antichain	3
minimal (extension)	32

O

ω_1 -closed	11
open	3

P

\mathbb{P} -name	4
Pre-decision property	28
predense	3
preserve cardinal	5
preserve cofinality	5
product	6
proper	18
proper game	19

R

regressive	2
------------------	---

S

Sacks property	32
Sacks real	30
separative	5
σ -centered	11
σ -closed	11
splitting node	30
stationary set	1
strong measure zero set	34
support	8

T

θ -cc	5
--------------------	---

U	
unbounded	1
unbounded real	6
V	
(V, \mathbb{P}) -generic filter	3

References

- [Abr10] Uri Abraham. Proper forcing. In *Handbook of set theory. Vols. 1, 2, 3*, pages 333–394. Springer, Dordrecht, 2010.
- [Bau83] James E. Baumgartner. Iterated forcing. In *Surveys in set theory*, volume 87 of *London Math. Soc. Lecture Note Ser.*, pages 1–59. Cambridge Univ. Press, Cambridge, 1983.
- [Bau84] James E. Baumgartner. Applications of the proper forcing axiom. In *Handbook of set-theoretic topology*, pages 913–959. North-Holland, Amsterdam, 1984.
- [Bla10] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In *Handbook of set theory. Vols. 1, 2, 3*, pages 395–489. Springer, Dordrecht, 2010.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, millennium edition, 2003.
- [JSW90] Haim Judah, Saharon Shelah, and W. H. Woodin. The Borel conjecture. *Ann. Pure Appl. Logic*, 50(3):255–269, 1990.
- [Kun11] Kenneth Kunen. *Set theory*, volume 34 of *Studies in Logic (London)*. College Publications, London, 2011.
- [Pal13] Justin Thomas Palumbo. *Hechler forcing and its relatives*. ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)—University of California, Los Angeles.