

Chapter 2. Probability

1. Sample Space

In the study of statistics we are concerned with the presentation and interpretation of **chance outcomes** that occur in a planned study or scientific investigation.

1. Sample Space

In the study of statistics we are concerned with the presentation and interpretation of **chance outcomes** that occur in a planned study or scientific investigation.

Examples:

- record the number of accidents that occur monthly at the intersection of Driftwood Lane and Royal Oak Drive, hoping to justify the installation of a traffic light;

1. Sample Space

In the study of statistics we are concerned with the presentation and interpretation of **chance outcomes** that occur in a planned study or scientific investigation.

Examples:

- record the number of accidents that occur monthly at the intersection of Driftwood Lane and Royal Oak Drive, hoping to justify the installation of a traffic light;
- classify items coming off an assembly line as "defective" or "nondefective";

1. Sample Space

In the study of statistics we are concerned with the presentation and interpretation of **chance outcomes** that occur in a planned study or scientific investigation.

Examples:

- record the number of accidents that occur monthly at the intersection of Driftwood Lane and Royal Oak Drive, hoping to justify the installation of a traffic light;
- classify items coming off an assembly line as "defective" or "nondefective";
- the volume of gas released in a chemical reaction when the concentration of an acid is varied.

The statistician is often dealing with either **experimental data**, representing **counts** or **measurements**, or perhaps with **categorical** data that can be classified according to some criterion.

The statistician is often dealing with either **experimental data**, representing **counts** or **measurements**, or perhaps with **categorical** data that can be classified according to some criterion.

We shall refer to any recording of information, whether it be numerical or categorical, as an **observation**.

- The number 2, 0, 1 and 2, representing the number of accidents that occurred for each month from January through April during the past year at the intersection of Driftwood Lane and Royal Oak Drive, constitute a set of **observations**.

- The number 2, 0, 1 and 2, representing the number of accidents that occurred for each month from January through April during the past year at the intersection of Driftwood Lane and Royal Oak Drive, constitute a set of **observations**.
- The categorical data N, D, N, D and D , representing the items found to be defective or nondefective when five items are inspected, are recorded as **observations**.

experiment

experiment: any process that generates a set of data.

experiment

experiment: any process that generates a set of data.

Examples:

- The tossing of a coin is a statistical experiment. There are only two possible outcomes, heads or tails.

experiment

experiment: any process that generates a set of data.

Examples:

- The tossing of a coin is a statistical experiment. There are only two possible outcomes, heads or tails.
- the launching of a missile and observing its velocity at specified times.

experiment

experiment: any process that generates a set of data.

Examples:

- The tossing of a coin is a statistical experiment. There are only two possible outcomes, heads or tails.
- the launching of a missile and observing its velocity at specified times.
- The opinions of voters concerning a new sales tax can also be considered as observations of an experiment.

experiment

Statisticians are particularly interested in the observations obtained by **repeating the experiment** several times.

experiment

Statisticians are particularly interested in the observations obtained by **repeating the experiment** several times.

Example: a coin is tossed repeatedly

experiment

Statisticians are particularly interested in the observations obtained by **repeating the experiment** several times.

Example: a coin is tossed repeatedly

- cannot predict the result of a given toss
(i.e. the outcome depend on chance)

experiment

Statisticians are particularly interested in the observations obtained by **repeating the experiment** several times.

Example: a coin is tossed repeatedly

- cannot predict the result of a given toss (i.e. the outcome depend on chance)
- know the entire set of possibilities for each toss.

sample space

Definition 2.1 The set of all possible outcomes of a statistical experiment is called the **sample space** and is represented by S .

sample space

Definition 2.1 The set of all possible outcomes of a statistical experiment is called the **sample space** and is represented by S .

Each outcome in a sample space is called an **element** or a **sample point**.

sample space

What's the possible outcomes when a coin is tossed?

sample space

What's the possible outcomes when a coin is tossed?

The sample space may be written

$$S = \{H, T\},$$

where H and T corresponds to 'heads' and 'tails', respectively.

sample space

Example 2.1 Consider the experiment of tossing a die.

interested in the number that shows on the top face

$$S_1 = \{1, 2, 3, 4, 5, 6\}$$

sample space

Example 2.1 Consider the experiment of tossing a die.

interested in the number that shows on the top face

$$S_1 = \{1, 2, 3, 4, 5, 6\}$$

interested only in whether the number is even or odd

$$S_2 = \{\text{even}, \text{odd}\}$$

sample space

Example 2.1 Consider the experiment of tossing a die.

interested in the number that shows on the top face

$$S_1 = \{1, 2, 3, 4, 5, 6\}$$

interested only in whether the number is even or odd

$$S_2 = \{\text{even}, \text{odd}\}$$

S_1 provides **more information** than S_2 .

sample space

Remarks:

- more than one sample space can be used to describe the outcomes of an experiment.

sample space

Remarks:

- more than one sample space can be used to describe the outcomes of an experiment.
- It is desirable to use a sample space that gives the most information.

sample space

Example 2.3 Suppose that three items are selected at random from a manufacturing process. Each item is inspected and classified defective, D , or nondefective, N . List the elements of the sample space providing the most information.

sample space

Example 2.3 Suppose that three items are selected at random from a manufacturing process. Each item is inspected and classified defective, D , or nondefective, N . List the elements of the sample space providing the most information.

The sample space is

$$S = \{DDD, DDN, DND, DNN, NDD, NDN, NND, NNN\}.$$

sample space

sample space with a large or infinite number of sample points

sample space

sample space with a large or infinite number of sample points

How to describe?

sample space

sample space with a large or infinite number of sample points

How to describe?

by a statement or rule

sample space

sample space with a large or infinite number of sample points

How to describe?

by a statement or rule

Examples:

- $S = \{x | x \text{ is a city with a population over 1 million}\}$

sample space

sample space with a large or infinite number of sample points

How to describe?

by a statement or rule

Examples:

- $S = \{x | x \text{ is a city with a population over 1 million}\}$
- all points (x, y) on the boundary or the interior of a circle of radius 2 with center at the origin

sample space

sample space with a large or infinite number of sample points

How to describe?

by a statement or rule

Examples:

- $S = \{x | x \text{ is a city with a population over 1 million}\}$
- all points (x, y) on the boundary or the interior of a circle of radius 2 with center at the origin

$$S = \{(x, y) | x^2 + y^2 \leq 4\}$$

2. Events

We may be interested in the occurrence of certain **events** rather than in the outcome of a specific element in the sample space.

2. Events

We may be interested in the occurrence of certain **events** rather than in the outcome of a specific element in the sample space.

Example: a die is tossed
event A : the outcome is divisible by 3.

2. Events

We may be interested in the occurrence of certain **events** rather than in the outcome of a specific element in the sample space.

Example: a die is tossed

event A : the outcome is divisible by 3.

Event A will occur if the outcome is an element of the set $A = \{3, 6\}$.

2. Events

We may be interested in the occurrence of certain **events** rather than in the outcome of a specific element in the sample space.

Example: a die is tossed

event A : the outcome is divisible by 3.

Event A will occur if the outcome is an element of the set $A = \{3, 6\}$.

$A = \{3, 6\}$ is the subset of the sample space

$S_1 = \{1, 2, 3, 4, 5, 6\}$ in Example 2.1.

event

Example: three items are selected at random from a manufacturing process

event

Example: three items are selected at random from a manufacturing process

event B : the number of defectives is greater than 1

event

Example: three items are selected at random from a manufacturing process

event B : the number of defectives is greater than 1

Event B will occur if the outcome is an element of the subset

$$B = \{DDN, DND, NDD, DDD\}.$$

event

Example: three items are selected at random from a manufacturing process

event B : the number of defectives is greater than 1

Event B will occur if the outcome is an element of the subset

$$B = \{DDN, DND, NDD, DDD\}.$$

Definition 2.2 An **event** is a subset of a sample space.

event

Example 2.4. Given the sample space $S = \{t | t \geq 0\}$, where t is the life in years of a certain electronic component. Event A is that the component fails before the end of the fifth year.

event

Example 2.4. Given the sample space $S = \{t | t \geq 0\}$, where t is the life in years of a certain electronic component. Event A is that the component fails before the end of the fifth year.

$$\text{subset } A = \{t | 0 \leq t < 5\}$$

event

Two special subset

event

Two special subset

a subset that includes the entire sample space S

event

Two special subset

- a subset that includes the entire sample space S

- a subset contains no elements at all, denoted by \emptyset , called **null set**.

event

Two special subset

- a subset that includes the entire sample space S

- a subset contains no elements at all, denoted by \emptyset , called **null set**.

Example

$$B = \{x | x \text{ is an even factor of } 7\},$$

event

Two special subset

- a subset that includes the entire sample space S

- a subset contains no elements at all, denoted by \emptyset , called **null set**.

Example

$B = \{x | x \text{ is an even factor of } 7\}$,

then $B = \emptyset$, since the only possible factors of 7 are odd numbers 1 and 7.

complement

We now consider certain operations with events that will result in the formation of new events.

complement

We now consider certain operations with events that will result in the formation of new events.

Definition 2.3 The **complement** of an event A with respect to S is the subset of all elements of S that are not in A .

complement

We now consider certain operations with events that will result in the formation of new events.

Definition 2.3 The **complement** of an event A with respect to S is the subset of all elements of S that are not in A .

the complement of A : A'

complement

Example 2.5 Let R be the event that a red card is selected from an ordinary deck of 52 playing cards and let S be the entire deck. What is R' ?

complement

Example 2.5 Let R be the event that a red card is selected from an ordinary deck of 52 playing cards and let S be the entire deck. What is R' ?

R' is the event that the card selected from the deck is not red but a black card.

intersection

Definition 2.4 The **intersection** of two events A and B , denoted by the symbol $A \cap B$, is the event containing all elements that are common to A and B .

intersection

Definition 2.4 The **intersection** of two events A and B , denoted by the symbol $A \cap B$, is the event containing all elements that are common to A and B .

Example 2.7

Let P be the event that a person selected at random while dining at a popular cafeteria is a taxpayer, and let Q be the event that the person is over 65 years of age. Then the event $P \cap Q$ is the set of all taxpayers in the cafeteria who are over 65 years of age.

intersection

Example 2.8 Let $M = \{a, e, i, o, u\}$ and $N = \{r, s, t\}$; then it follows that $M \cap N = \emptyset$

intersection

Example 2.8 Let $M = \{a, e, i, o, u\}$ and $N = \{r, s, t\}$; then it follows that $M \cap N = \emptyset$

That is, M and N have no elements in common and, therefore, cannot both occur simultaneously.

intersection

Definition 2.5 Two events A and B are **mutually exclusive**, or **disjoint** if $A \cap B = \emptyset$, that is, if A and B have no elements in common.

union

Definition 2.6 The **union** of the two events A and B , denote by symbol $A \cup B$, is the event containing all the elements that belong to A or B or both.

union

Definition 2.6 The **union** of the two events A and B , denote by symbol $A \cup B$, is the event containing all the elements that belong to A or B or both.

Example 2.10 Let $A = \{a, b, c\}$ and $B = \{b, c, d, e\}$; then

$$A \cup B = \{a, b, c, d, e\}.$$

union

Example 2.11 Let P be the event that an employee selected at random from an oil drilling company smokes cigarettes. Let Q be the event that the employee selected drinks alcoholic beverages. Then the event $P \cup Q$ is the set of all employees who either drink or smoke, or do both.

union

Example 2.11 Let P be the event that an employee selected at random from an oil drilling company smokes cigarettes. Let Q be the event that the employee selected drinks alcoholic beverages. Then the event $P \cup Q$ is the set of all employees who either drink or smoke, or do both.

Example 2.12 If $M = \{x|3 < x < 9\}$, $N = \{x|5 < x < 12\}$,
then $M \cup N = \{x|3 < x < 12\}$.

The relationship between events and the corresponding sample space can be illustrated graphically by means of **Venn diagrams**.

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

$$A \cap A' = \emptyset \qquad A \cup A' = S$$

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

$$A \cap A' = \emptyset \qquad A \cup A' = S$$

$$S' = \emptyset \qquad \emptyset' = S$$

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

$$A \cap A' = \emptyset \qquad A \cup A' = S$$

$$S' = \emptyset \qquad \emptyset' = S$$

$$(A')' = A$$

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

$$A \cap A' = \emptyset \qquad A \cup A' = S$$

$$S' = \emptyset \qquad \emptyset' = S$$

$$(A')' = A$$

$$(A \cap B)' = A' \cup B'$$

$$(A \cup B)' = A' \cap B'$$

several results

The following results may easily be verified by means of **Venn diagrams**

$$A \cap \emptyset = \emptyset \qquad A \cup \emptyset = A$$

$$A \cap A' = \emptyset \qquad A \cup A' = S$$

$$S' = \emptyset \qquad \emptyset' = S$$

$$(A')' = A$$

$$(A \cap B)' = A' \cup B'$$

$$(A \cup B)' = A' \cap B'$$