

## 4. Chebyshev's Theorem

The Russian mathematician P.L. Chebyshev (1821-1894) discovered that the fraction of the area between any two values symmetric about the mean is related to the standard deviation.

## 4. Chebyshev's Theorem

The Russian mathematician P.L. Chebyshev (1821-1894) discovered that the fraction of the area between any two values symmetric about the mean is related to the standard deviation.

**Theorem 4.11 (Chebyshev's Theorem)** The probability that any random variable  $X$  will assume a value within  $k$  standard deviations of the mean  $\mu$  is at least  $1 - 1/k^2$ . That is,

$$P(\mu - k\sigma < X < \mu + k\sigma) \geq 1 - \frac{1}{k^2}.$$

# Chebyshev's Theorem

The theorem gives a conservative estimate of the probability that a random variable assumes a value within  $k$  standard deviations of its mean for any real number  $k$ .

# Chebyshev's Theorem

**PROOF** (only for continuous case)

By our previous definition of the variance of  $X$  we can write

$$\begin{aligned}
 \sigma^2 &= E[(X - \mu)^2] \\
 &= \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx \\
 &= \int_{-\infty}^{\mu - k\sigma} (x - \mu)^2 f(x) dx + \int_{\mu - k\sigma}^{\mu + k\sigma} (x - \mu)^2 f(x) dx \\
 &\quad + \int_{\mu + k\sigma}^{\infty} (x - \mu)^2 f(x) dx \\
 &\geq \int_{-\infty}^{\mu - k\sigma} (x - \mu)^2 f(x) dx + \int_{\mu + k\sigma}^{\infty} (x - \mu)^2 f(x) dx
 \end{aligned}$$

since the second of the three integrals is nonnegative. Now, since  $|x - \mu| \geq k\sigma$  wherever  $x \geq \mu + k\sigma$  or  $x \leq \mu - k\sigma$ , we have  $(x - \mu)^2 \geq k^2\sigma^2$  in both remaining integral. It follows that

# Chebyshev's Theorem

$$\sigma^2 \geq \int_{-\infty}^{\mu-k\sigma} k^2 \sigma^2 f(x) dx + \int_{\mu+k\sigma}^{\infty} k^2 \sigma^2 f(x) dx$$

and that

$$\frac{1}{k^2} \geq \int_{-\infty}^{\mu-k\sigma} f(x) dx + \int_{\mu+k\sigma}^{\infty} f(x) dx$$

Hence

$$P(\mu - k\sigma < X < \mu + k\sigma) \geq 1 - \frac{1}{k^2}.$$

# Chebyshev's Theorem

For  $k = 2$  the theorem states that  $3/4$  or more of the observations of any distribution lie in the interval  $\mu \pm 2\sigma$ .

# Chebyshev's Theorem

For  $k = 2$  the theorem states that  $3/4$  or more of the observations of any distribution lie in the interval  $\mu \pm 2\sigma$ .

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 1 - \frac{1}{2^2}.$$

# Chebyshev's Theorem

For  $k = 2$  the theorem states that  $3/4$  or more of the observations of any distribution lie in the interval  $\mu \pm 2\sigma$ .

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 1 - \frac{1}{2^2}.$$

Similarly, at least eight-ninths of the observations of any distribution fall in the interval  $\mu \pm 3\sigma$ .



# Chebyshev's Theorem

For  $k = 2$  the theorem states that  $3/4$  or more of the observations of any distribution lie in the interval  $\mu \pm 2\sigma$ .

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 1 - \frac{1}{2^2}.$$

Similarly, at least eight-ninths of the observations of any distribution fall in the interval  $\mu \pm 3\sigma$ .

$$P(\mu - 3\sigma < X < \mu + 3\sigma) \geq 1 - \frac{1}{3^2}.$$

# Chebyshev's Theorem

**Example 4.22** A random variable  $X$  has a mean  $\mu = 8$ , a variance  $\sigma^2 = 9$ , and an unknown probability distribution. Find  
(a)  $P(-4 < X < 20)$ ,      (b)  $P(|X - 8| \geq 6)$

# Chebyshev's Theorem

**Example 4.22** A random variable  $X$  has a mean  $\mu = 8$ , a variance  $\sigma^2 = 9$ , and an unknown probability distribution. Find  
(a)  $P(-4 < X < 20)$ ,      (b)  $P(|X - 8| \geq 6)$

**Solution.....**

$$P(-4 < X < 20) \geq \frac{15}{16}, \quad P(|X - 8| \geq 6) \leq \frac{1}{4}.$$

# Chebyshev's Theorem

**The Chebyshev's Theorem is a distribution-free result.**

Therefore, the results are usually weak.

# Chebyshev's Theorem

**The Chebyshev's Theorem is a distribution-free result.**

Therefore, the results are usually weak.

The value given by the theorem is a **lower bound only**.

# Chebyshev's Theorem

**The Chebyshev's Theorem is a distribution-free result.**

Therefore, the results are usually weak.

The value given by the theorem is **a lower bound only**.

We know that the probability of a random variable falling within two standard deviations of the mean can be no less than  $3/4$ , but we never know how much more it might actually be.

# Chebyshev's Theorem

**The Chebyshev's Theorem is a distribution-free result.**

Therefore, the results are usually weak.

The value given by the theorem is **a lower bound only**.

We know that the probability of a random variable falling within two standard deviations of the mean can be no less than  $3/4$ , but we never know how much more it might actually be.

Only when the probability distribution is known can we determine exact probabilities.