

3.Areas Under the Normal Curve

$$P(x_1 < X < x_2) = \int_{x_1}^{x_2} f(x; \mu, \sigma) dx = \frac{1}{\sqrt{2\pi}\sigma} \int_{x_1}^{x_2} e^{-(x-\mu)^2/(2\sigma^2)} dx$$

is represented by the area of the shaded region for the normal curve in Figure 6.6.




Figure 6.6: $P(x_1 < X < x_2)$ = area of the shaded region.

Areas Under the Normal Curve

The difficulty encountered in solving integrals of normal density functions necessitates the tabulation of normal curve areas for quick reference.

Areas Under the Normal Curve

The difficulty encountered in solving integrals of normal density functions necessitates the tabulation of normal curve areas for quick reference.

However, it would be a hopeless task to attempt to set up separate tables for every conceivable value of μ and σ .

Areas Under the Normal Curve

Transform all the observations of any normal random variable X to a new set of observations of a normal random variable Z with mean 0 and variance 1.

Areas Under the Normal Curve

Transform all the observations of any normal random variable X to a new set of observations of a normal random variable Z with mean 0 and variance 1.

This can be done by the transformation

$$Z = \frac{X - \mu}{\sigma}.$$

Areas Under the Normal Curve

Therefore, if X falls between the values $x = x_1$ and $x = x_2$, the random variable Z will fall between the corresponding values $z_1 = (x_1 - \mu)/\sigma$ and $z_2 = (x_2 - \mu)/\sigma$.

Areas Under the Normal Curve

Therefore, if X falls between the values $x = x_1$ and $x = x_2$, the random variable Z will fall between the corresponding values $z_1 = (x_1 - \mu)/\sigma$ and $z_2 = (x_2 - \mu)/\sigma$. Consequently, we may write

$$\begin{aligned} P(x_1 < X < x_2) &= P\left(\frac{x_1 - \mu}{\sigma} < \frac{X - \mu}{\sigma} < \frac{x_2 - \mu}{\sigma}\right) \\ &= P(z_1 < Z < z_2) \end{aligned}$$

Areas Under the Normal Curve

Definition 6.1

The distribution of a normal random variable with mean zero and variance 1 is called a **standard normal distribution**.

Areas Under the Normal Curve

Definition 6.1

The distribution of a normal random variable with mean zero and variance 1 is called a **standard normal distribution**.

We have now reduced the required number of tables of normal curve areas to one, that of the standard normal distribution.

Areas Under the Normal Curve

Definition 6.1

The distribution of a normal random variable with mean zero and variance 1 is called a **standard normal distribution**.

We have now reduced the required number of tables of normal curve areas to one, that of the standard normal distribution.

Table A.3 indicates the area under the standard normal curve corresponding to $P(Z < z)$.

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74.

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74. First, we locate a value of z equal to 1.7 in the left column, then move across the row to the column under 0.04 , where we read 0.9591.

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74. First, we locate a value of z equal to 1.7 in the left column, then move across the row to the column under 0.04 , where we read 0.9591. Therefore, $P(Z < 1.74) = 0.9591$.

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74. First, we locate a value of z equal to 1.7 in the left column, then move across the row to the column under 0.04, where we read 0.9591. Therefore, $P(Z < 1.74) = 0.9591$.

To find a z -value corresponding to a given probability, the process is reversed.

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74. First, we locate a value of z equal to 1.7 in the left column, then move across the row to the column under 0.04, where we read 0.9591. Therefore, $P(Z < 1.74) = 0.9591$.

To find a z -value corresponding to a given probability, the process is reversed. For example, $P(Z < z) = 0.2148$,

Areas Under the Normal Curve

Let us find the probability that Z is less than 1.74. First, we locate a value of z equal to 1.7 in the left column, then move across the row to the column under 0.04, where we read 0.9591. Therefore, $P(Z < 1.74) = 0.9591$.

To find a z -value corresponding to a given probability, the process is reversed. For example, $P(Z < z) = 0.2148$, is seen to be -0.79 .

Areas Under the Normal Curve

Example 6.2 Given a standard normal distribution, find the area under the curve that lies

(a) to the right of $z = 1.84$,

(b) between $z = -1.97$ and $z = 0.86$.





Figure 6.9: Areas for Example 6.2.

Areas Under the Normal Curve

Example 6.3 Given a standard normal distribution, find the value of k such that

(a) $P(Z > k) = 0.3015$,

(b) $P(k < Z < -0.18) = 0.4197$.



Areas Under the Normal Curve

Example 6.4 Given a random variable X having a normal distribution with $\mu = 50$ and $\sigma = 10$, find the probability that X assumes a value between 45 and 62.

Areas Under the Normal Curve

Example 6.4 Given a random variable X having a normal distribution with $\mu = 50$ and $\sigma = 10$, find the probability that X assumes a value between 45 and 62.

Solution

$$P(45 < X < 62) = P\left(\frac{45 - 50}{10} < \frac{X - 50}{10} < \frac{62 - 50}{10}\right)$$

Areas Under the Normal Curve

Example 6.4 Given a random variable X having a normal distribution with $\mu = 50$ and $\sigma = 10$, find the probability that X assumes a value between 45 and 62.

Solution

$$\begin{aligned} P(45 < X < 62) &= P\left(\frac{45 - 50}{10} < \frac{X - 50}{10} < \frac{62 - 50}{10}\right) \\ &= P(-0.5 < Z < 1.2) \end{aligned}$$

Areas Under the Normal Curve

Example 6.4 Given a random variable X having a normal distribution with $\mu = 50$ and $\sigma = 10$, find the probability that X assumes a value between 45 and 62.

Solution

$$P(45 < X < 62) = P\left(\frac{45 - 50}{10} < \frac{X - 50}{10} < \frac{62 - 50}{10}\right)$$

$$= P(-0.5 < Z < 1.2)$$

$$= 0.8849 - 0.3085 = 0.5764$$

Areas Under the Normal Curve

Example 6.5 Given that X has a normal distribution with $\mu = 300$ and $\sigma = 50$, find the probability that X assumes a value greater than 362 .

Areas Under the Normal Curve

Example 6.5 Given that X has a normal distribution with $\mu = 300$ and $\sigma = 50$, find the probability that X assumes a value greater than 362 .

Solution

$$P(X > 362) = P\left(\frac{X - 300}{50} > \frac{362 - 300}{50}\right)$$

Areas Under the Normal Curve

Example 6.5 Given that X has a normal distribution with $\mu = 300$ and $\sigma = 50$, find the probability that X assumes a value greater than 362 .

Solution

$$\begin{aligned} P(X > 362) &= P\left(\frac{X - 300}{50} > \frac{362 - 300}{50}\right) \\ &= P(Z > 1.24) = 1 - P(Z < 1.24) \end{aligned}$$

Areas Under the Normal Curve

Example 6.5 Given that X has a normal distribution with $\mu = 300$ and $\sigma = 50$, find the probability that X assumes a value greater than 362 .

Solution

$$P(X > 362) = P\left(\frac{X - 300}{50} > \frac{362 - 300}{50}\right)$$

$$= P(Z > 1.24) = 1 - P(Z < 1.24)$$

$$= 1 - 0.8925 = 0.1075$$

Areas Under the Normal Curve

According to Chebyshev's theorem, the probability that a random variable assumes a value within 2 standard deviation of the mean is at least $3/4$. That is

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 3/4.$$

Areas Under the Normal Curve

According to Chebyshev's theorem, the probability that a random variable assumes a value within 2 standard deviation of the mean is at least $3/4$. That is

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 3/4.$$

If the random variable has a **normal distribution**,

$$\begin{aligned} & P(\mu - 2\sigma < X < \mu + 2\sigma) \\ &= P\left(\frac{(\mu - 2\sigma) - \mu}{\sigma} < \frac{X - \mu}{\sigma} < \frac{(\mu + 2\sigma) - \mu}{\sigma}\right) \\ &= P(-2 < Z < 2) = 0.9772 - 0.0228 = 0.9544 \end{aligned}$$

Areas Under the Normal Curve

According to Chebyshev's theorem, the probability that a random variable assumes a value within 2 standard deviation of the mean is at least $3/4$. That is

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \geq 3/4.$$

If the random variable has a **normal distribution**,

$$\begin{aligned} & P(\mu - 2\sigma < X < \mu + 2\sigma) \\ &= P\left(\frac{(\mu - 2\sigma) - \mu}{\sigma} < \frac{X - \mu}{\sigma} < \frac{(\mu + 2\sigma) - \mu}{\sigma}\right) \\ &= P(-2 < Z < 2) = 0.9772 - 0.0228 = 0.9544 \end{aligned}$$

This is a much stronger statement than that stated by Chebyshev's theorem.

Areas Under the Normal Curve

Example 6.6

Given a normal distribution with $\mu = 40$ and $\sigma = 6$, find the value of x that has

- (a) 45% of the area to the left,
- (b) 14% of the area to the right.