

Solving the quantum master equation of coupled harmonic oscillators with Lie algebra methods

November 4, 2021

Introduction

Hilbert space \mathcal{H} and Liouville space \mathfrak{L}

Hilbert space \mathcal{H}

- Density operator $\hat{\rho}$
- Hamiltonian $\hat{\mathcal{H}}$
- von Neumann equation :

$$\frac{\partial}{\partial t} \hat{\rho} = -i[\hat{\mathcal{H}}, \hat{\rho}] \quad (1)$$

\implies Eq. (1) as the problem in linear algebra.

$$\boldsymbol{x} = A\boldsymbol{b}$$

Liouville space \mathfrak{L}

- Superket $|\rho\rangle\rangle$
- Liouvillian $\hat{\mathcal{L}}$
- von Neumann equation :

$$\frac{\partial}{\partial t} |\rho\rangle\rangle = \hat{\mathcal{L}} |\rho\rangle\rangle \quad (2)$$

Quantum master equation

Von Neumann equation : $\partial_t |\rho\rangle\rangle = -i[\mathcal{H}, \hat{\rho}]$

- + Total Hamiltonian : $\hat{H}_T = \hat{H} + \hat{H}_E + \hat{H}_I$.
- + Markovian interaction.
- + Initial state is separable : $\hat{\rho}_T(0) \approx \hat{\rho}(0) \otimes \hat{\rho}_E(0)$.
- + Born approximation : $\hat{\rho}_T(t) \approx \hat{\rho}(t) \otimes \hat{\rho}_E(0)$.
- + Weak coupled regime : $\hat{H}_I \ll \hat{H}, \hat{H}_E$.
- + Rotating wave approximation.

Quantum master equation¹ : ($\hat{\mathcal{L}}_i$ is Lindblad jump operator)

$$\frac{\partial}{\partial t} \hat{\rho}(t) = -i[\mathcal{H}, \hat{\rho}] + \frac{1}{2} \sum_i \left(2\hat{\mathcal{L}}_i \hat{\rho}(t) \hat{\mathcal{L}}_i^\dagger - \{ \hat{\mathcal{L}}_i^\dagger \hat{\mathcal{L}}_i, \hat{\rho}(t) \} \right) \quad (3)$$

¹Manzano2020.

Lie algebra \mathfrak{g} and Lie bracket $[\cdot, \cdot]$

Lie algebra \mathfrak{g} is vector space with Lie bracket $[\cdot, \cdot]$.

- Alternativity : $[X, Y] = -[Y, X]$.
- Linearity : $[aX + bY, Z] = a[X, Z] + b[Y, Z]$.
- Jacobi identity : $[[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0$.

ex) Commutator $[X, Y] := XY - YX$.

Lie algebra is not commutative.

\implies Left translation and right transition :

$$\hat{L}[\hat{O}]|A\rangle\rangle := \hat{O}\hat{A}, \quad \hat{R}[\hat{O}]|A\rangle\rangle := \hat{A}\hat{O} \quad (4)$$

A linear chain of N coupled harmonic oscillators

Hamiltonian :

$$\hat{\mathcal{H}} = \sum_{k=1}^N \varepsilon_k \hat{a}_k^\dagger \hat{a}_k + \sum_{k=1}^{N-1} \kappa_k (\hat{a}_k^\dagger \hat{a}_{k+1} + \hat{a}_k \hat{a}_{k+1}^\dagger) \quad (5)$$

Quantum master equation :

$$\frac{\partial}{\partial t} \hat{\rho}(t) = -i[\hat{\mathcal{H}}, \hat{\rho}] + \sum_{k=1}^N \gamma_k (2\hat{a}_k \hat{\rho} \hat{a}_k^\dagger - \{\hat{a}_k^\dagger \hat{a}_k, \hat{\rho}\}) \quad (6)$$

where $\hat{a}_k, \hat{a}_k^\dagger$ is bosonic mode operators,

ε_k : energy, κ_k : coupling constant, γ_k : loss rate

Induced Lie algebra

Effective non-Hermite Hamiltonian $\hat{\mathcal{H}}_{\text{eff}}$

Effective non-Hermite Hamiltonian $\hat{\mathcal{H}}_{\text{eff}}$:

$$-i(\hat{\mathcal{H}}_{\text{eff}}\hat{\rho} - \hat{\rho}\hat{\mathcal{H}}_{\text{eff}}^{\dagger}) := -i[\hat{\mathcal{H}}, \hat{\rho}] - \sum_{k=1}^N \gamma_k \{\hat{a}_k^{\dagger} \hat{a}_k, \hat{\rho}\}_+ \quad (7)$$

Eigenvalue (energy) :

$$\varepsilon_k \longrightarrow \varepsilon_k - i\gamma_k \quad (8)$$

Bosonic mode superoperators

Bosonic mode superoperators :

$$\hat{\hat{L}}_k^- |A\rangle\rangle = \hat{a}_k \hat{A}, \quad \hat{\hat{L}}_k^+ |A\rangle\rangle = \hat{a}_k^\dagger \hat{A}, \quad (9)$$

$$\hat{\hat{R}}_k^- |A\rangle\rangle = \hat{A} \hat{a}_k, \quad \hat{\hat{R}}_k^+ |A\rangle\rangle = \hat{A} \hat{a}_k^\dagger. \quad (10)$$

As well as $[\hat{a}_j, \hat{a}_k^\dagger] = \delta_{jk}$,

$$[\hat{\hat{L}}_i^-, \hat{\hat{L}}_j^+] = \delta_{ij}, \quad [\hat{\hat{R}}_i^-, \hat{\hat{R}}_j^+] = \delta_{ij}. \quad (11)$$

The set

$$\{\hat{\mathbb{1}}_{\mathcal{L}}, \hat{\hat{L}}_i^\pm, \hat{\hat{R}}_i^\pm, \hat{\hat{L}}_j^+ \hat{\hat{L}}_k^-, \hat{\hat{R}}_j^- \hat{\hat{R}}_k^+, \hat{\hat{L}}_j^- \hat{\hat{R}}_k^-\} \quad (12)$$

spans Lie algebra \mathfrak{g} .

Bosonic mode superoperators

Bosonic mode superoperators :

$$\hat{\hat{L}}_k^- |A\rangle\rangle = \hat{a}_k \hat{A}, \quad \hat{\hat{L}}_k^+ |A\rangle\rangle = \hat{a}_k^\dagger \hat{A}, \quad (9)$$

$$\hat{\hat{R}}_k^- |A\rangle\rangle = \hat{A} \hat{a}_k, \quad \hat{\hat{R}}_k^+ |A\rangle\rangle = \hat{A} \hat{a}_k^\dagger. \quad (10)$$

As well as $[\hat{a}_j, \hat{a}_k^\dagger] = \delta_{jk}$,

$$[\hat{\hat{L}}_i^-, \hat{\hat{L}}_j^+] = \delta_{ij}, \quad [\hat{\hat{R}}_i^-, \hat{\hat{R}}_j^+] = \delta_{ij}. \quad (11)$$

The set

$$\{\hat{\mathbb{1}}_{\mathcal{L}}, \hat{\hat{L}}_i^\pm, \hat{\hat{R}}_i^\pm, \hat{\mathcal{L}}\} \quad (12)$$

spans Lie algebra \mathfrak{g} .

Liouvillian $\hat{\mathcal{L}}$ for harmonic oscillators

Liouvillian (right action) :

$$\begin{aligned}\hat{\mathcal{L}} = \sum_{k=1}^N & \left[(i\varepsilon_k - \gamma_k) \hat{R}_k^+ \hat{R}_k^- - (i\varepsilon_k + \gamma_k) \hat{L}_k^+ \hat{L}_k^- + 2\gamma_k \hat{L}_k^- \hat{R}_k^- \right] \\ & + \sum_{k=1}^N i\kappa_k \left(\hat{L}_k^+ \hat{L}_{k+1}^- + \hat{L}_{k+1}^+ \hat{L}_k^- - \hat{R}_{k+1}^+ \hat{R}_k^- - \hat{R}_k^+ \hat{R}_{k+1}^- \right)\end{aligned}\quad (13)$$

Time evolution superoperator :

$$\frac{\partial}{\partial t} \hat{\mathcal{U}}(t) = \hat{\mathcal{L}} \hat{\mathcal{U}}(t) \quad (14)$$

where $\hat{\mathcal{U}}(0) = \hat{\mathbb{1}}_{\mathcal{L}}$.

Approaches to solve master equation

Two approaches to solve master equation.

1. Eigendecomposition (time-independent)
2. Wei-Norman expansion (time-dependent)

Eigendecomposition

Regular representation in Lie algebra

Regular representation $\hat{\mathcal{R}}(Z)$:

$$[\hat{Z}, \hat{X}_i] = \hat{\mathcal{R}}_{ij}(\hat{Z})\hat{X}_j. \quad (15)$$

Interested in $\hat{\mathcal{R}}(\hat{\mathcal{L}})$,

$$\{\hat{\mathbb{1}}_{\mathcal{L}}, \hat{L}_i^{\pm}, \hat{R}_i^{\pm}, \hat{\mathcal{L}}\} \longrightarrow \{\hat{L}_i^{\pm}, \hat{R}_i^{\pm}\} \quad (16)$$

Eigenvalues of $\hat{\mathcal{R}}(\hat{\mathcal{L}})$:

$$\{\pm\lambda_1, \dots, \pm\lambda_N, \pm\lambda_1^*, \dots, \pm\lambda_N^*\} \quad (17)$$

where $\lambda = (\lambda_1, \dots, \lambda_N)$ are the eigenvalues of $\hat{\mathcal{H}}_{\text{eff}}$.

Collective creation and annihilation operators

Eigenvectors of $\hat{\mathcal{R}}(\hat{\mathcal{L}})$:

$$\hat{P}_i^+ = \sum_k c_{ik}(\hat{L}_k^+ - \hat{R}_k^-), \quad \hat{P}_i^- = \sum_k c_{ik}\hat{L}_k^-, \quad (18)$$

$$\hat{Q}_i^+ = \sum_k c_{ik}(\hat{R}_k^+ - \hat{L}_k^-), \quad \hat{Q}_i^- = \sum_k c_{ik}^*\hat{R}_k^-. \quad (19)$$

\Rightarrow Collective creation and annihilation operators :

$$[\hat{P}_i^-, \hat{P}_j^+] = \delta_{ij}, \quad [\hat{Q}_i^-, \hat{Q}_j^+] = \delta_{ij} \quad (20)$$

$$\begin{aligned} [\hat{\mathcal{L}}, \hat{P}_i^+] &= \lambda_i \hat{P}_i^+ & [\hat{\mathcal{L}}, \hat{P}_i^-] &= -\lambda_i \hat{P}_i^- \\ [\hat{\mathcal{L}}, \hat{Q}_i^+] &= \lambda_i^* \hat{Q}_i^+ & [\hat{\mathcal{L}}, \hat{Q}_i^-] &= -\lambda_i^* \hat{Q}_i^- \end{aligned} \quad (21)$$

Ground state and ladder state

Ground state in \mathcal{L} :

$$\hat{P}_i^- |0\rangle\rangle = \hat{Q}_i^- |0\rangle\rangle = 0_{\mathcal{H}} \quad (22)$$

$$\hat{P}_i^+ \langle\langle 0| = \hat{Q}_i^+ \langle\langle 0| = 0_{\mathcal{H}} \quad (23)$$

Higher rungs of the ladder :

$$|\alpha, \beta\rangle\rangle = \frac{1}{\sqrt{\alpha! \beta!}} \hat{\mathbf{P}}^{+\alpha} \hat{\mathbf{Q}}^{+\beta} |0\rangle\rangle \quad (24)$$

$$\langle\langle \alpha, \beta| = \frac{1}{\sqrt{\alpha! \beta!}} \langle\langle 0| \hat{\mathbf{P}}^{+\alpha} \hat{\mathbf{Q}}^{+\beta} \quad (25)$$

where $\alpha = (\alpha_1, \dots, \alpha_N), \beta = (\beta_1, \dots, \beta_N)$.

Collective creation and annihilation operators

Baker – Campbell – Hausdorff formula :

$$e^A B e^{-A} = \sum_{n=0}^{\infty} \frac{1}{n!} \underbrace{[A, [A, \dots, [A, B] \dots]]}_n \quad (26)$$

Ladder state :

$$e^{\mathcal{L}t} |\alpha, \beta\rangle\rangle = e^{(\alpha \cdot \lambda + \beta \cdot \lambda^*)t} |\alpha, \beta\rangle\rangle \quad (27)$$

Density superket $|\rho\rangle\rangle$ evolved as

$$|\rho(t)\rangle\rangle = \sum_{\alpha, \beta} e^{(\alpha \cdot \lambda + \beta \cdot \lambda^*)t} |\alpha, \beta\rangle\rangle \langle\langle \alpha, \beta | \rho_0 \rangle\rangle . \quad (28)$$

Wei-Norman expansion

Collective creation and annihilation operators

Expansion of $\hat{\mathcal{L}}$ by basis :

$$\hat{\mathcal{L}}(t) = \sum_{k=1}^m c_k(t) \hat{X}_k. \quad (29)$$

Time evolution superoperator :

$$\hat{\mathcal{U}}(t) = \prod_k^n \hat{\mathcal{U}}_k(t) = \prod_k^n \exp g_k(t) \hat{X}_k \quad (30)$$

where

$$\frac{\partial}{\partial t} \hat{\mathcal{U}}_k(t) = g_k(t) \hat{X}_k \hat{\mathcal{U}}_k(t), \quad \hat{\mathcal{U}}_k(0) = \hat{\mathbb{1}}_{\mathcal{L}}. \quad (31)$$

Collective creation and annihilation operators

Liouvillian is split as :

$$\hat{\mathcal{L}} = \underbrace{\hat{\mathcal{L}}_S}_{\text{solvable}} + \underbrace{\hat{\mathcal{L}}_R}_{\text{remaining}} \quad (32)$$

Time evolution superoperator :

$$\frac{\partial}{\partial t} \hat{\mathcal{U}}_S(t) = \hat{\mathcal{L}}_S \hat{\mathcal{U}}_S(t), \quad \frac{\partial}{\partial t} \hat{\mathcal{U}}_R(t) = (\hat{\mathcal{U}}_S^{-1} \hat{\mathcal{L}}_R \hat{\mathcal{U}}_S) \hat{\mathcal{U}}_R(t) \quad (33)$$

However, $\hat{\mathcal{U}}_R(t)$ is not always solvable.