



Nanotechnological applications in medicine

Shelton D Caruthers^{1,2}, Samuel A Wickline¹ and Gregory M Lanza¹

Nanotechnology-based tools and techniques are rapidly emerging in the fields of medical imaging and targeted drug delivery. Employing constructs such as dendrimers, liposomes, nanoshells, nanotubes, emulsions and quantum dots, these advances lead toward the concept of personalized medicine and the potential for very early, even presymptomatic, diagnoses coupled with highly-effective targeted therapy. Highlighting clinically available and preclinical applications, this review explores the opportunities and issues surrounding nanomedicine.

Addresses

¹ Consortium for Translational Research in Advanced Imaging and Nanomedicine, Washington University School of Medicine, 4444 Forest Park Avenue, Campus Box 8215, St Louis, MO 63108-2212, USA ² Philips Medical Systems, 3000 Minuteman Road, Andover, MA 01810, USA

Corresponding author: Caruthers, Shelton D (scaruthers@cmrl.wustl.edu)

Current Opinion in Biotechnology 2007, 18:26-30

This review comes from a themed issue on Analytical biotechnology Edited by Joseph C Wu and Jagat Narula

Available online 24th January 2007

0958-1669/\$ - see front matter
© 2007 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.copbio.2007.01.006

Introduction

'Molecular imaging' is a phrase that has come into heavy use within the past decade. It is a broad term, difficult to define. Although the initial use implied imaging novel contrast agents to probe molecular information [1], the current use has evolved to include a wide scope of imaging techniques that use molecular agents or rely on molecular signatures. These include virtually all positron emission tomography imaging (indeed all nuclear medicine techniques requiring radioactive or radiolabeled molecules), magnetic resonance spectroscopy, and other parametric magnetic resonance imaging techniques (like diffusion-weighted imaging). Similarly, the term 'nanomedicine' is difficult to define precisely, as much of what transpires in human biology happens at the nanometer scale. Thus, one might successfully argue that all medicine is 'nanomedicine'. Herein, we discuss nanomedicine as the application of technologies on the scale of approximately 1 to 500 nm toward the goal of diagnosing and treating disease.

Within the past decade, there has been a plethora of new research, development and patent activities around nanoscaled technologies in the health sciences field [2°]. One indicator of this is the flurry of activity around the National Nanotechnology Initiative [3] of the US government and, in particular, those initiatives of the National Institutes of Health (NIH). Within the National Cancer Institute (NCI) alone, the Alliance for Nanotechnology in Cancer program (http://nano.cancer.gov/) has launched eight Centers of Cancer Nanotechnology Excellence (http://nano.cancer. gov/programs/ccne.asp) and twelve Cancer Nanotechnol-Platform Partnerships (http://nano.cancer.gov/ programs/nanotech_platforms.asp) [4]. The Program for Excellence in Nanotechnology (http://www.nhlbi-pen. info/) of the National Heart, Lung and Blood Institute is another instance of the profuse programs fueling this growing research arena. To help facilitate the transfer of nanotechnological research into clinical practice, the NCI, in conjunction with the National Institute of Standards and Technology and the US Food and Drug Administration (FDA), has also established the National Characterization Laboratory (http://ncl.cancer.gov/).

The overall goal of nanomedicine is the same as it always has been in medicine: to diagnose as accurately and early as possible, to treat as effectively as possible without side effects, and to evaluate the efficacy of treatment noninvasively. The promise that nanotechnology brings is multifaceted, offering not only improvements to current techniques but also providing entirely new tools and capabilities. By manipulating drugs and other materials at the nanometer scale, the fundamental properties and bioactivity of materials can be altered. These tools can permit control over characteristics of drugs or agents such as solubility, blood pool retention times, controlled release over short or long durations, environmentally triggered controlled release or highly specific site-targeted delivery. Furthermore, by using nanometer-sized particles, the increased functional surface area per unit volume can be exploited in various ways.

This review presents some of the more recent successes of applying various new nanotechnological techniques and tools in diagnostic imaging and therapeutics, including both current products and late-stage preclinical research. It will not delve into screening or *in vitro* diagnostics, although nanotechnological applications certainly abound in these areas (e.g. improvements to 'laboratory on a chip' technologies like microarrays and microfluidics [5,6]). Nor will this review focus on the many advances in biomedical laboratory research tools and techniques resulting from nanotechnology.

Diagnostic imaging in nanomedicine

Combining advances in related fields such as genomics, proteomics, drug delivery and molecular imaging, nanomedicine offers the potential to move from a 'one-sizefits-all' approach to one more individually tailored for higher efficacy [1,7]. For diagnosis, this translates to recognition and characterization of very early (even pre-symptomatic) disease providing assessment, preferably non-invasively, akin to that of immunohistochemistry. Of course doing so is complex, requiring simultaneous detection and quantification of multiple markers. Starting with single biomarkers as sine quo non for disease, initial successes in nanotechnology-enabled molecular imaging have been made in all imaging modalities including optical [8–10], nuclear [9,11], ultrasound [12,13], computed tomography (CT) [14,15°], and magnetic resonance imaging (MRI) [16-18]. Optical and nuclear techniques are exquisitely sensitive to their respective agents, but are complicated by high background noise. The other major modalities (i.e. MRI, CT and ultrasound) are hampered by relative insensitivity to low concentrations of imaging agent, as would be the case for agents bound to molecular markers. The increased surface area per volume of nanoparticles helps overcome this limitation by coupling each targeted particle with very large payloads of the imaging agent [17].

One of the earliest applications of nanotechnology in MRI was the use of paramagnetic iron oxide particles; when taken up by healthy hepatocytes, these particles could help to distinguish between normal and cancerous liver cells [19]. Other applications of these nanoparticulate iron oxides have been used to highlight lymph node metastases in prostate cancer [20] and atherosclerotic plaques in hyperlipidemic rabbit models [21]. More recently, nanometer-sized constructs such as dendrimers [18,22,23], liposomes [24,25], nanoshells [26], nanotubes [27], nanoemulsions [28], quantum dots [29,30], and even viruses [31] have been developed as imaging agents intended as noninvasive probes or to target biomarkers of disease. Many have moved from in vitro to in vivo applications, targeting biomarkers such as the overexpressed folate receptor of cancer cells [22], human growth factor receptors (HER2) on tumors [32°], or integrins crucial to angiogenesis around tumors [33,34] and atherosclerotic vessels [35].

Expanding on targeting single biomarkers, recent research has addressed the need for quantification and simultaneous identification of multiple markers. Quantum dots offer a full spectrum of colors to achieve this, but are limited to *in vitro*, small animal or surface applications where optical techniques excel. In a method analogous to multicolored quantum dots, researchers exploiting the unique magnetic resonance spectra of various liquid-core perfluorocarbon nanoparticle emulsions have shown the potential of MRI for quantifying biomarkers (i.e. fibrin deposits, a biomarker for ruptured atherosclerotic plaques) [36] and simultaneously distinguishing and uniquely imaging multiple targeted agents bound in the same location [37°].

Nanotherapeutics

Controlled drug delivery

Unfortunately, early diagnosis is futile if not coupled with effective therapy. Owing in part to the national nanotechnology initiatives, there has been much activity in applying nanotechnology to the rapeutics [38–41,42°]. Currently, there are limited numbers of nanomedical products on the market [2°], with the majority being pharmaceuticals that are formulated (or re-formulated) into nanosized structures to manipulate the pharmacodynamics, biodistribution and overall effectiveness. Cosmetological and dermatological applications are well represented, especially with liposomal versions of exisiting drugs [25]. Liposomal packaging of chemotherapeutic agents (e.g. doxorubicin) are also on the market [43]. The surface addition of polyethylene glycol (PEG) of varying dimensions is a common technique used to alter chemical formulations (liposome or otherwise) into 'stealthy' nanoparticles with increased circulating half lives. This technique is employed for many current products with indications including leukemia, neutropenia, hepatitis, and macular degeneration [2°]. Abraxane, a nanoparticle version of paclitaxel bound to albumin, achieves water solubility without harsh excipients, has been shown to be more effective than previous formulations of paclitaxel. and is the first such chemotherapeutic nanoparticle approved by the FDA [44**]. In addition to these drug delivery techniques, advances in nanotechnology and polymer sciences have contributed to techniques such as implantable nanoporous membranes [45,46], controlledrelease biochips [47], and transdermal microneedle systems [48].

Site-targeted nanotherapeutics

Beyond clever pharmacokinetic manipulations, a more complex goal of nanotherapeutics is to devise agents that travel within the body undetected to deliver specific therapy — whether chemical, radioactive, genetic or other — to uniquely identified sites having minimal untoward effect elsewhere. Additionally, agents that can, upon their success, 'report back' through non-invasive signaling (i.e. imaging) to provide confirmation and predicted efficacy would greatly alter the current paradigm for treating and following patients. Many novel agents, currently under development and perhaps soon to enter clinical trials, share this ultimate goal [22,32[•],38,40,42[•],49–54].

An in vitro model of atherosclerosis illustrates the potential targeted nanotherapeutic paradigm from diagnosis, through therapy, to serial non-invasive follow-up. In hyperlipidemic rabbits, Winter et al. [55°] used integrin-targeted paramagnetic nanoparticles that had previously been shown [35] to detect angiogenesis in very early atherosclerotic disease with MRI. This novel agent provides quantification of the extent of disease and, at the same time, can convey therapeutic doses of an anti-angiogenic drug, fumagillin, which could slow plaque progression. Moreover, following treatment, MRI scans were performed using the diagnostic version of the targeted paramagnetic nanoparticles to monitor the effect longitudinally. The targeted therapeutic nanoparticles not only had a significant anti-angiogenic effect that could be monitored via non-invasive imaging, but the magnitude of response correlated linearly with the degree of image enhancement measured at the time of initial treatment; that is, the images at time of treatment confirmed drug delivery and were predictive of therapeutic efficacy. Furthermore, as the targeted nanoparticle drug delivery concentrated the drug at the desired site, the results were achieved with a total drug dose thousands of times lower than similar studies using conventional drug delivery schemes. Thus, targeted nanoparticles can drastically improve the safety profile of the drug by lowering overall dosage and concentrating levels in disease sites.

Limitations and considerations in nanomedicine

Obviously, not all attempts to apply new nanotechnology approaches in medicine have met with the same success as those cited herein. The new tools are not necessarily intuitive and bring with them new challenges and hurdles. Nanometer-sized structures do not behave in the same predictive ways that single, small-molecule interactions occur. Nanoconstructs, especially multifunctional ones, are complex three-dimensional objects with critical dependence on position, size, shape and charge of interrelated components. Currently, what we know about nanomedicine is dwarfed by what we do not know; most of the challenges have not been defined.

The potential that nanotechnology offers for improved circulating half-life, greater functional surface area, and other benefits is superb, but offset by new, and yetunknown, constraints. Coupled with changing circulation times, for example, are changes with clearance from the body. If the clearance mechanism for a radiotherapeutic agent is shifted from being cleared in minutes via the kidneys to hours via the liver, this has potential ramifications. Other nanoparticles might be retained not only for days, but potentially for years. In that case, the safety profile for all components becomes of paramount significance. Although the long-term safety of the components of some nanotechnology devices like perfluorocarbons or semiconductors is well described, others, like carbon nanotubes, are not well appreciated. Furthermore, as physical properties of materials are altered at the nanometer level, current safety information cannot be assumed to be accurate for nanoversions.

Safety and drug effects are not limited solely to the patient population receiving the final nanomedicine product. The entire manufacture and disposal process also needs to be considered. For example, as nanoparticulates can be very easily aerosolized, standard filtering or protection equipment might not be appropriate. Waste water and other environmental impacts might be different than with customary techniques. The promise of nanotechnology applications in medicine is the breadth of entirely novel techniques; the responsibility is to redefine the safety, function, and environmental effects of the new tools and techniques.

Conclusions

Nanotechnology, in general, is experiencing a rapid growth period with major advances arriving quickly. Accordingly, these advances are applied in the biomedical field in numerous diverse ways. Already, a few medical products are providing a glimmer of the overwhelming benefits nanomedicine will surely provide. Current preclinical research promises new ways to diagnose disease, to deliver specific therapy, and to monitor the effects acutely and non-invasively. This rapid onset and drastic change in methods will create new challenges in the regulatory process, but will also provide a fruitful ground from which many exciting, and yet unimagined, applications of nanotechnology in medicine will arise.

Disclosure statement

SDC is an employee of Philips Medical Systems. SAW and GML receive research grant funding from Philips Medical Systems and are equity shareholders in Kereos.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest
- Weissleder R: Molecular imaging: exploring the next frontier. Radiology 1999, 212:609-614.
- 2. Wagner V, Dullaart A, Bock AK, Zweck A: The emerging
- nanomedicine landscape. Nat Biotechnol 2006, 24:1211-1217. Good review, including commercial activities in nanomedicine.
- The National Nanotechnology Initiative: Research and development leading to a revolution in technology and industry - supplement to the President's 2007 budget. Edited by National Science and Technology Council Subcommittee on Nanoscale Science, Engineering and Technology; 2006: http://www.nano.gov/NNI_07Budget.pdf.
- Brower V: Is nanotechnology ready for primetime? J Natl Cancer Inst 2006. 98:9-11.
- Jain KK: Nanodiagnostics: application of nanotechnology in molecular diagnostics. Expert Rev Mol Diagn 2003,
- Craighead H: Future lab-on-a-chip technologies for interrogating individual molecules. Nature 2006,
- Pettigrew RI, Fee CA, Li KC: Changes in the world of biomedical research are moving the field of 'personalized medicine' from concept to reality. J Nucl Med 2004, 45:1427.

- Bremer C, Ntziachristos V, Weissleder R: Optical-based molecular imaging: contrast agents and potential medical applications. Eur Radiol 2003, 13:231-243.
- Herschman HR: Molecular imaging: looking at problems, seeing solutions. Science 2003, 302:605-608.
- Tsien RY: Imagining imaging's future. Nat Rev Mol Cell Biol (Suppl):2003:SS16-S
- 11. Britz-Cunningham SH, Adelstein SJ: Molecular targeting with radionuclides: state of the science. J Nucl Med 2003, 44:1945-1961.
- 12. Lanza GM, Wickline SA: Targeted ultrasonic contrast agents for molecular imaging and therapy. Curr Probl Cardiol 2003, 28:625-
- Sakamoto JH, Smith BR, Xie B, Rokhlin SI, Lee SC, Ferrari M: The molecular analysis of breast cancer utilizing targeted nanoparticle based ultrasound contrast agents. Technol Cancer Res Treat 2005, 4:627-636.
- 14. Bergman A: Hepatocyte-specific contrast media for CT. An experimental investigation. Acta Radiol Suppl 1997, 411:1-27.
- Winter PM, Shukla HP, Caruthers SD, Scott MJ, Fuhrhop RW, Robertson JD, Gaffney PJ, Wickline SA, Lanza GM: Molecular imaging of human thrombus with computed tomography. Acad Radiol 2005, 12:9-13.

One of the first CT agents specifically targeted to atherosclerotic plaques.

- Caruthers SD, Winter PM, Wickline SA, Lanza GM: Targeted magnetic resonance imaging contrast agents. Methods Mol Med 2006, 124:387-400.
- 17. Wickline SA, Lanza GM: Nanotechnology for molecular imaging and targeted therapy. Circulation 2003, 107:1092-1095.
- Kobayashi H, Brechbiel MW: Nano-sized MRI contrast agents with dendrimer cores. Adv Drug Deliv Rev 2005, 57:2271-2286.
- Saini S, Edelman RR, Sharma P, Li W, Mayo-Smith W, Slater GJ, Eisenberg PJ, Hahn PF: Blood-pool MR contrast material for detection and characterization of focal hepatic lesions: initial clinical experience with ultrasmall superparamagnetic iron oxide (AMI-227). AJR Am J Roentgenol 1995, 164:1147-1152.
- 20. Weissleder R, Elizondo G, Wittenberg J, Lee AS, Josephson L, Brady TJ: Ultrasmall superparamagnetic iron oxide: an intravenous contrast agent for assessing lymph nodes with MR imaging. Radiology 1990, 175:494-498.
- 21. Ruehm SG, Corot C, Vogt P, Cristina H, Debatin JF: Ultrasmall superparamagnetic iron oxide-enhanced MR imaging of atherosclerotic plaque in hyperlipidemic rabbits. Acad Radiol 2002, 9(Suppl 1):S143-S144.
- 22. Quintana A, Raczka E, Piehler L, Lee I, Myc A, Majoros I, Patri AK, Thomas T, Mule J, Baker JR Jr: Design and function of a dendrimer-based therapeutic nanodevice targeted to tumor cells through the folate receptor. Pharm Res 2002, 19:1310-1316.
- 23. Svenson S, Tomalia DA: Dendrimers in biomedical applications-reflections on the field. Adv Drug Deliv Rev 2005, **57**:2106-2129
- 24. Mulder WJ, Strijkers GJ, Habets JW, Bleeker EJ, van der Schaft DW, Storm G, Koning GA, Griffioen AW, Nicolay K: MR molecular imaging and fluorescence microscopy for identification of activated tumor endothelium using a bimodal lipidic nanoparticle. FASEB J 2005. 19:2008-2010.
- 25. Banerjee R: Liposomes: applications in medicine. J Biomater Appl 2001, 16:3-21.
- 26. Hirsch LR, Gobin AM, Lowery AR, Tam F, Drezek RA, Halas NJ, West JL: Metal nanoshells. Ann Biomed Eng 2006, 34:15-22.
- 27. Toth E, Bolskar RD, Borel A, Gonzalez G, Helm L, Merbach AE, Sitharaman B, Wilson LJ: **Water-soluble gadofullerenes: toward** high-relaxivity, pH-responsive MRI contrast agents. $J\ Am$ Chem Soc 2005, 127:799-805.
- Wickline SA, Neubauer AM, Winter P, Caruthers S, Lanza G: Applications of nanotechnology to atherosclerosis,

- thrombosis, and vascular biology. Arterioscler Thromb Vasc Biol
- 29. Gao X, Yang L, Petros JA, Marshall FF, Simons JW, Nie S: In vivo molecular and cellular imaging with quantum dots. Curr Opin Biotechnol 2005. 16:63-72.
- 30. Michalet X, Pinaud FF, Bentolila LA, Tsay JM, Doose S, Li JJ, Sundaresan G, Wu AM, Gambhir SS, Weiss S: **Quantum dots for** live cells, in vivo imaging, and diagnostics. Science 2005, 307:538-544.
- 31. Manchester M, Singh P: Virus-based nanoparticles (VNPs): platform technologies for diagnostic imaging. Adv Drug Deliv Rev 2006. 58:1505-1522.
- Shukla R, Thomas TP, Peters JL, Desai AM, Kukowska-Latallo J, Patri AK, Kotlyar A, Baker JR Jr: **HER2 specific tumor targeting** with dendrimer conjugated anti-HER2 mAb. Bioconjug Chem 2006. 17:1109-1115.

Demonstration of in vivo targeting of HER2-positive tumors with PAMAM dendrimers to deliver therapeutic or imaging agents.

- Schmieder AH, Winter PM, Caruthers SD, Harris TD, Williams TA Allen JS, Lacy EK, Zhang H, Scott MJ, Hu G et al.: Molecular MR imaging of melanoma angiogenesis with alphanubeta3targeted paramagnetic nanoparticles. Magn Reson Med 2005, **53**:621-627.
- 34. Winter PM, Caruthers SD, Kassner A, Harris TD, Chinen LK, Allen JS, Lacy EK, Zhang H, Robertson JD, Wickline SA et al.: Molecular imaging of angiogenesis in nascent Vx-2 rabbit tumors using a novel alpha(nu)beta3-targeted nanoparticle and 1.5 tesla magnetic resonance imaging. Cancer Res 2003, 63:5838-5843.
- Winter PM, Morawski AM, Caruthers SD, Fuhrhop RW, Zhang H, Williams TA, Allen JS, Lacy EK, Robertson JD, Lanza GM et al.: Molecular imaging of angiogenesis in early-stage atherosclerosis with alpha(v)beta3-integrin-targeted nanoparticles. Circulation 2003, 108:2270-2274.
- 36. Morawski AM, Winter PM, Yu X, Fuhrhop RW, Scott MJ, Hockett F, Robertson JD, Gaffney PJ, Lanza GM, Wickline SA: Quantitative 'magnetic resonance immunohistochemistry' with ligand-targeted ¹⁹F nanoparticles. *Magn Reson Med* 2004, **52**:1255-1262.
- 37. Caruthers SD, Neubauer AM, Hockett FD, Lamerichs R, Winter PM, Scott MJ, Gaffney PJ, Wickline SA, Lanza GM: *In vitro* demonstration using ¹⁹F magnetic resonance to augment molecular imaging with paramagnetic perfluorocarbon nanoparticles at 1.5 Tesla. Invest Radiol 2006, 41:305-312. Demonstration of multispectral MRI for simultaneous detection and imaging of multiple-targeted nanoparticle agents.
- Cyrus T, Winter PM, Caruthers SD, Wickline SA, Lanza GM: Magnetic resonance nanoparticles for cardiovascular molecular imaging and therapy. Expert Rev Cardiovasc Ther 2005. **3**:705-715.
- 39. Farokhzad OC, Langer R: Nanomedicine: developing smarter therapeutic and diagnostic modalities. Adv Drug Deliv Rev 2006, 58:1456-1459.
- 40. Lanza G, Winter P, Cyrus T, Caruthers S, Marsh J, Hughes M, Wickline S: Nanomedicine opportunities in cardiology. Ann NY Acad Sci 2006, 1080:451-465.
- 41. Leary SP, Liu CY, Apuzzo ML: Toward the emergence of nanoneurosurgery: part III-nanomedicine: targeted nanotherapy, nanosurgery, and progress toward the realization of nanoneurosurgery. Neurosurgery 2006, 58:1009-1026 discussion 1009-1026.
- 42. Pison U, Welte T, Giersig M, Groneberg DA: Nanomedicine for respiratory diseases. Eur J Pharmacol 2006, 533:341-350. Good review with basics of nanomedicine.
- 43. Cattel L, Ceruti M, Dosio F: From conventional to stealth liposomes: a new Frontier in cancer chemotherapy. J Chemother 2004, 16(Suppl 4):94-97.
- 44. Gradishar WJ: Albumin-bound paclitaxel: a next-generation taxane. Expert Opin Pharmacother 2006, 7:1041-1053. First nanoparticle-based chemotherapeutic of its kind to receive FDA approval.

- 45. Tao SL, Desai TA: Microfabricated drug delivery systems: from particles to pores. Adv Drug Deliv Rev 2003,
- 46. Lesniak MS, Langer R, Brem H: Drug delivery to tumors of the central nervous system. Curr Neurol Neurosci Rep 2001, 1.210-216
- 47. Santini JT Jr, Cima MJ, Langer R: A controlled-release microchip. Nature 1999, 397:335-338.
- 48. Prausnitz MR: Microneedles for transdermal drug delivery. Adv Drug Deliv Rev 2004, 56:581-587.
- 49. Majoros IJ, Myc A, Thomas T, Mehta CB, Baker JR Jr: PAMAM dendrimer-based multifunctional conjugate for cancer therapy: synthesis, characterization, and functionality. Biomacromolecules 2006, 7:572-579.
- 50. Roy I, Ohulchanskyy TY, Pudavar HE, Bergey EJ, Oseroff AR, Morgan J, Dougherty TJ, Prasad PN: **Ceramic-based** nanoparticles entrapping water-insoluble photosensitizing anticancer drugs: a novel drug-carrier system for photodynamic therapy. J Am Chem Soc 2003, 125:7860-7865.
- 51. Reddy GR, Bhojani MS, McConville P, Moody J, Moffat BA, Hall DE, Kim G, Koo YE, Woolliscroft MJ, Sugai JV et al.: Vascular

- targeted nanoparticles for imaging and treatment of brain tumors. Clin Cancer Res 2006, 12:6677-6686.
- 52. Kannan R, Rahing V, Cutler C, Pandrapragada R, Katti KK, Kattumuri V, Robertson JD, Casteel SJ, Jurisson S, Smith C et al.: Nanocompatible chemistry toward fabrication of targetspecific gold nanoparticles. J Am Chem Soc 2006, **128**:11342-11343.
- 53. Veiseh O, Sun C, Gunn J, Kohler N, Gabikian P, Lee D, Bhattarai N, Ellenbogen R, Sze R, Hallahan A *et al.*: Optical and MRI multifunctional nanoprobe for targeting gliomas. Nano Lett 2005, 5:1003-1008.
- 54. Rossin R, Pan D, Qi K, Turner JL, Sun X, Wooley KL, Welch MJ: ⁶⁴Culabeled folate-conjugated shell cross-linked nanoparticles for tumor imaging and radiotherapy: synthesis, radiolabeling, and biologic evaluation. J Nucl Med 2005, 46:1210-1218.
- 55. Winter PM, Neubauer AM, Caruthers SD, Harris TD, Robertson JD, Williams TA, Schmieder AH, Hu G, Allen JS, Lacy EK et al.: Endothelial alpha(v)beta3 integrin-targeted fumagillin nanoparticles inhibit angiogenesis in atherosclerosis. Arterioscler Thromb Vasc Biol 2006, 26:2103-2109.

One of the first examples of site-targeted drug delivery with longitudinal image-based monitoring using multifunction nanoparticles.