
## Specific Studies on Varying Factors



## The influence of musical training experience

☐ Music major students showed higher emotional experience than non-music major students, specifically, stronger activation of the bilateral inferior parietal lobules. This region is an important structure involved in emotional information processing.

				Valence		Arousal		
Tempo	Musician	Non-musician	Musician	Non-musician	Mean scores of musicians and non-musicians	Musician	Non-musician	Mean scores of musicians and non-musicians
Fast	$5.79 \pm 0.56$	$5.34 \pm 0.60$	$6.08 \pm 0.84$	$5.57 \pm 0.72$	$5.77 \pm 0.80$	$5.51 \pm 0.81$	$5.12 \pm 0.76$	$5.27 \pm 0.80$
Medium	$5.05 \pm 0.64$	$5.21 \pm 0.58$	$4.47 \pm 0.79$	$4.56 \pm 0.61$	$4.52 \pm 0.68$	$5.63 \pm 1.05$	$5.86 \pm 0.87$	$5.77 \pm 0.94$
Slow	$5.05 \pm 1.20$	$5.23 \pm 1.13$	$5.10 \pm 1.28$	$5.16 \pm 1.24$	$5.13 \pm 1.24$	$5.01 \pm 1.20$	$5.31 \pm 1.11$	$5.19 \pm 1.14$

Average scores of three tempi in two emotional dimensions between musicians and non-musicians https://www.frontiersin.org/articles/10.3389/fpsyg.2018.02118/full