
Magnitude and Phase of a Linear System

Tamas Kis | kis@stanford.edu | <https://github.com/tamaskis>

Contents

1	Magnitude and Phase of a Linear System	2
1.1	Complex Numbers	2
1.2	Continuous-Time Transfer Functions	2
1.3	Discrete-Time Transfer Functions	2
1.4	State-Space Representation	2

Copyright © 2021 Tamas Kis

Permission is hereby granted, free of charge, to any person obtaining a copy of this software and associated documentation files (the "Software"), to deal in the Software without restriction, including without limitation the rights to use, copy, modify, merge, publish, distribute, sublicense, and/or sell copies of the Software, and to permit persons to whom the Software is furnished to do so, subject to the following conditions:

The above copyright notice and this permission notice shall be included in all copies or substantial portions of the Software.

THE SOFTWARE IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE SOFTWARE OR THE USE OR OTHER DEALINGS IN THE SOFTWARE.



1 MAGNITUDE AND PHASE OF A LINEAR SYSTEM

1.1 Complex Numbers

A complex number, z , can be written as

$$z = x + jy = \text{Re}(z) + j \text{Im}(z)$$

In the complex plane, this can be represented as a vector $z = (x, y)^T$, where this vector has magnitude $|z|$ and phase¹ (i.e. direction) $\angle z$.

$$|z| = \sqrt{x^2 + y^2}$$

$$\angle z = \text{arctan2}(y, x)$$

1.2 Continuous-Time Transfer Functions

A continuous-time transfer function, $H(s)$, evaluates to a complex number.

$$H(s) = \text{Re}[H(s)] + j \text{Im}[H(s)]$$

Thus,

$$|H(s)| = \sqrt{\text{Re}[H(s)]^2 + \text{Im}[H(s)]^2}$$

$$\angle H(s) = \text{arctan2}(\text{Im}[H(s)], \text{Re}[H(s)])$$

1.3 Discrete-Time Transfer Functions

Finding the magnitude and phase of a discrete-time transfer function can be done in an identical matter as continuous-time transfer functions:

$$|G(z)| = \sqrt{\text{Re}[G(z)]^2 + \text{Im}[G(z)]^2}$$

$$\angle G(z) = \text{arctan2}(\text{Im}[G(z)], \text{Re}[G(z)])$$

1.4 State-Space Representation

Consider the case where we have a linear system expressed in the state-space form

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{B}u$$

$$y = \mathbf{C}\mathbf{x} + Du$$

where $u, y, D \in \mathbb{R}$. This can readily be converted to a continuous-time transfer function $G(s)$ in MATLAB using the `tf2ss` function. However, we can simply define the linear system using

$$\text{sys} = \text{ss}(\mathbf{A}, \mathbf{B}, \mathbf{C}, D)$$

and then use the `evalfr` function to evaluate the system at a specific point s in the frequency domain.

¹ Note that the phase is often defined as

$$\angle z = \arctan\left(\frac{y}{x}\right)$$

This will not always return the correct phase since the more general definition is

$$\tan(\angle z) = \frac{y}{x}$$