

# Newton's Method

## *MATLAB Implementation*

---

Tamas Kis | [kis@stanford.edu](mailto:kis@stanford.edu)

TAMAS KIS  
<https://github.com/tamaskis>

Copyright © 2021 Tamas Kis

*Permission is hereby granted, free of charge, to any person obtaining a copy of this software and associated documentation files (the "Software"), to deal in the Software without restriction, including without limitation the rights to use, copy, modify, merge, publish, distribute, sublicense, and/or sell copies of the Software, and to permit persons to whom the Software is furnished to do so, subject to the following conditions:*

*The above copyright notice and this permission notice shall be included in all copies or substantial portions of the Software.*

*THE SOFTWARE IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE SOFTWARE OR THE USE OR OTHER DEALINGS IN THE SOFTWARE.*



# Contents

<b>1</b>	<b>Download and Installation</b>	<b>4</b>
1.1	Download from MATLAB Central's File Exchange . . . . .	4
1.2	Download from GitHub . . . . .	4
1.3	Files Included With Download . . . . .	4
1.4	Accessing the <code>newtons_method</code> Function in a MATLAB Script . . . . .	4
<b>2</b>	<b><code>newtons_method</code></b>	<b>5</b>
<b>3</b>	<b>Newton's Method</b>	<b>7</b>
	<b>References</b>	<b>9</b>

# 1 Download and Installation

---

## 1.1 Download from MATLAB Central's File Exchange

The `newtons_method` function is available for download on MATLAB® Central's File Exchange at [https://www.mathworks.com/matlabcentral/fileexchange/85735-newton-s-method-newtons\\_method](https://www.mathworks.com/matlabcentral/fileexchange/85735-newton-s-method-newtons_method).

## 1.2 Download from GitHub

The `newtons_method` function is available for download on GitHub® at [https://github.com/tamaskis/newtons\\_method-MATLAB](https://github.com/tamaskis/newtons_method-MATLAB).

## 1.3 Files Included With Download

There are **five** files included in the downloaded zip file:

1. `EXAMPLE.M` – *example for using the `newtons_method` function*
2. `LICENSE` – *license for the `newtons_method` function*
3. `Newton's Method - MATLAB Implementation.pdf` – *this PDF*
4. `newtons_method.m` – *MATLAB function implementing Newton's method*
5. `README.md` – *markdown file for GitHub documentation*

## 1.4 Accessing the `newtons_method` Function in a MATLAB Script

There are **four** options for accessing the `newtons_method` function in a MATLAB script:

1. Copy the `newtons_method` function to the *end* of your MATLAB script.
2. Place the `newtons_method.m` file in the same folder as the MATLAB script.
3. Place the `newtons_method.m` file into whatever folder you want, and then use the `addpath(folderName)` command<sup>1</sup> where the `folderName` parameter is a string that stores the filepath of the folder that `newtons_method.m` is in *relative to* the folder that your script is in.
4. Make a toolbox by first opening `newtons_method.m`, then going to the HOME tab in MATLAB, and finally selecting **Package Toolbox** in the drop-down menu under **Add-Ons**. Once you package the `newtons_method` function as a toolbox, you can use it in any script.

---

<sup>1</sup> <https://www.mathworks.com/help/matlab/ref/addpath.html>

## 2 newtons\_method

Calculates the root of a differentiable, univariate function using Newton's method.

### Syntax

```
root = newtons_method(f,df,x0)
root = newtons_method(f,df,x0,TOL)
root = newtons_method(f,df,x0,[],imax)
root = newtons_method(f,df,x0,TOL,imax)
```

### Description

`root = newtons_method(f,df,x0)` returns the root of a differentiable function  $f(x)$  specified by the function handle `f`, where `df` is the derivative of  $f(x)$  (i.e.  $f'(x)$ ) and `x0` is an initial guess of the root. The default tolerance and maximum number of iterations are `TOL = 1e-12` and `imax = 1e6`, respectively.

`root = newtons_method(f,df,x0)` returns the root of a differentiable function  $f(x)$  specified by the function handle `f`, where `df` is the derivative of  $f(x)$  (i.e.  $f'(x)$ ), `x0` is an initial guess of the root, and `TOL` is the tolerance. The default maximum number of iterations is `imax = 1e6`.

`root = newtons_method(f,df,x0,[],imax)` returns the root of a differential function  $f(x)$  specified by the function handle `f`, where `df` is the derivative of  $f(x)$  (i.e.  $f'(x)$ ), `x0` is an initial guess of the root, and `imax` is the maximum number of iterations. The default tolerance is `TOL = 1e-12`.

`root = newtons_method(f,df,x0,TOL,imax)` returns the root of a differentiable function  $f(x)$  specified by the function handle `f`, where `df` is the derivative of  $f(x)$  (i.e.  $f'(x)$ ), `x0` is an initial guess of the root, `TOL` is the tolerance, and `imax` is the maximum number of iterations.

### Example

#### Example 2.1

Find the root(s) of  $f(x) = x^2 - 1$ .

#### ■ SOLUTION

To apply Newton's method to find the root(s) of  $f(x)$ , we first need to find  $f'(x)$ .

$$f'(x) = \frac{d}{dx} (x^2 - 1) = 2x$$

Defining  $f(x)$  and  $f'(x)$  in MATLAB,

```
% f(x) and its derivative
f = @(x) x^2-1;
df = @(x) 2*x;
```

We know  $f(x) = x^2 - 1$  has roots at  $x = \pm 1$ , but let's pretend we don't know this, and solve this problem using a more general approach. Since  $f(x)$  is a quadratic function, we know that it will have either 0 roots (in the case where  $f(x)$  does not cross the  $x$ -axis) or 2 roots. Let's assume the latter case (otherwise it would be pointless to try and find roots of  $f(x)$ ). Therefore, we use Newton's method twice, with two different guesses. Let's pick  $-10$  and  $10$  as our initial guesses.

```
% finds first root of f(x)=x^2-1 using Newton's method
root1 = newtons_method(f,-10)

% finds second root of f(x)=x^2-1 using Newton's method
root2 = newtons_method(f,10)
```

This yields the result

```
root1 =
    -1

root2 =
     1
```

---

### 3 Newton's Method

**Newton's method** is a technique used to find the root (based on an initial guess<sup>2</sup>  $x_0$ ) of a *differentiable*, univariate function  $f(x)$ . The equation of the tangent line to the curve  $y = f(x)$  at  $x = x_0$  is

$$y = f'(x_0)(x - x_0) + f(x_0)$$

where  $f'(x_0)$  is the derivative of  $f(x)$  evaluated at  $x_0$ . The  $x$ -intercept of this tangent line,  $x = x_1$ , can be solved by setting  $y = 0$ .

$$0 = f'(x_0)(x_1 - x_0) + f(x_0)$$

$$\therefore x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$x_1$  is an updated estimate of the root of  $f(x)$ . To keep refining our estimate, we can keep iterating through this procedure using Eq. (1).

$$\boxed{x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}} \quad (1)$$

So how do we actually use Eq. (1)? Given an initial guess  $x_0$ , we can keep coming up with new estimates of the root. But how do we know when to stop? To resolve this issue, we define the **error**<sup>3</sup> as

$$\boxed{\varepsilon = |x_{i+1} - x_i|} \quad (2)$$

Once  $\varepsilon$  is small enough, we say that the estimate of the root has **converged** to the true root, within some **tolerance** (which we denote as TOL). Therefore, if we predetermine that, at most, we can *tolerate* an error of TOL, then we will keep iterating Eq. (1) until  $\varepsilon < \text{TOL}$ . In some cases, the error may never decrease below TOL, or take too long to decrease to below TOL. Therefore, we also define the **maximum number of iterations** ( $i_{\max}$ ) so that the algorithm does not keep iterating forever, or for too long of a time.

The algorithm itself is shown below in Algorithm 1 [1, 2].

<sup>2</sup> Often, a function  $f(x)$  will have multiple roots. Therefore, Newton's method typically finds the root closest to the initial guess  $x_0$ . However, this is not always the case; the algorithm depends heavily on the derivative of  $f(x)$ , which, depending on its form, may cause it to converge on a root further from  $x_0$ .

<sup>3</sup> Note that  $\varepsilon$  is an *approximate* error. The motivation behind using this definition of  $\varepsilon$  is that as  $i$  gets large (i.e.  $i \rightarrow \infty$ ),  $x_{i+1} - x_i$  approaches  $x_{i+1} - x^*$  (assuming this sequence is convergent), where  $x^*$  is the true root (and therefore  $x_{i+1} - x^*$  represents the *exact* error).

---

**Algorithm 1:** Newton's method.

---

**1** Given:  $f(x)$ ,  $f'(x)$ ,  $x_0$ , TOL,  $i_{\max}$ *// initializes the error so the loop will be entered***2**  $\text{err} = (2)(\text{TOL})$ **3** Preallocate an  $i_{\max} \times 1$  vector  $\mathbf{x}$ , where  $\mathbf{x}$  is the vector storing the root estimate at each iteration.*// inputs initial guess for root into  $\mathbf{x}$  vector***4**  $x_1 = x_0$ *// Newton's method***5**  $i = 1$ **6 while**  $\varepsilon > \text{TOL}$  **and**  $i < i_{\max}$  **do***// updates estimate of root***7**  $x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$ *// calculates error***8**  $\varepsilon = |x_{i+1} - x_i|$ *// increments loop index***9**  $i = i + 1$ **10 end***// returns root***11**  $\text{root} = x_i$ **12 return** root

---



## References

---

- [1] James Hateley. *Nonlinear Equations*. MATH 3620 Course Reader (Vanderbilt University). 2019.
- [2] *Newton's method*. [https://en.wikipedia.org/wiki/Newton%27s\\_method](https://en.wikipedia.org/wiki/Newton%27s_method). (accessed: June 10, 2020).