

Pharmacovigilance & Adverse Drug Reaction (ADR) Analysis

Case Study: Tirzepatide

Intern	P. Tamilselvan
Task	Task 3 – Pharmacovigilance & ADR Analysis
Drug	Tirzepatide
Primary sources	FDA Prescribing Information + DrugBank
Prepared	15 August 2026

Prepared as part of the CodeAlpha Pharmaceutical Research Internship. The report covers mechanism of action, reported ADRs and measures to reduce risks as required for Task 3.

Abstract

Tirzepatide is a dual glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1 (GLP-1) receptor agonist used in the management of type 2 diabetes mellitus and, under the Zepbound brand, chronic weight management in eligible adults. Its pharmacological activity produces glucose-dependent enhancement of insulin secretion and reduction of glucagon secretion. Because pharmacovigilance evaluates the safety of medicines throughout their lifecycle, tirzepatide provides a useful case study of common adverse reactions, clinically important warnings, and risk-minimization measures. The most frequently reported adverse reactions in clinical studies are gastrointestinal, particularly nausea, diarrhea and decreased appetite; vomiting, constipation, dyspepsia and abdominal discomfort are also reported. Clinically important risks include severe gastrointestinal reactions, acute pancreatitis, gallbladder disease, hypoglycemia when used with insulin or insulin secretagogues, hypersensitivity reactions, acute kidney injury related to volume depletion, and complications in patients with diabetic retinopathy. Postmarketing reports also include serious gastrointestinal, hypersensitivity and renal events. This report summarizes the mechanism of action, adverse drug reactions and practical measures used to reduce risk, based primarily on current FDA prescribing information and DrugBank pharmacovigilance resources.

1. Drug Overview

Tirzepatide is a modified peptide administered by subcutaneous injection. It is a dual agonist at the GIP and GLP-1 receptors and is marketed as Mounjaro for improving glycemic control in adults with type 2 diabetes mellitus. The same active ingredient is marketed as Zepbound for chronic weight management in eligible adults. The molecule contains a C20 fatty diacid moiety that promotes albumin binding and contributes to its prolonged half-life, supporting once-weekly dosing. For pharmacovigilance purposes, tirzepatide is particularly relevant because its therapeutic effects are accompanied by predictable gastrointestinal tolerability issues and a set of less frequent but clinically important safety concerns. DrugBank provides structured clinical safety information, including adverse effects, black-box warnings, contraindications and drug interactions, while FDA labeling provides the regulatory safety framework and recommended risk-management measures.

2. Mechanism of Action

Tirzepatide selectively binds to and activates both the GIP and GLP-1 receptors. These receptors are involved in incretin signaling and metabolic regulation. According to FDA prescribing information, tirzepatide enhances first- and second-phase insulin secretion and reduces glucagon levels in a glucose-dependent manner. The drug's pharmacological effects are therefore linked to improved glucose regulation. Tirzepatide also acts on pathways involved in appetite and energy balance, contributing to reductions in food intake and body weight. Because gastrointestinal effects are among the most common adverse reactions, the pharmacological actions that produce therapeutic benefit must be considered alongside tolerability during dose escalation. The once-weekly regimen is supported by the molecule's modified structure and prolonged exposure. Dose escalation is used clinically to reduce the risk of gastrointestinal adverse reactions. This is an important pharmacovigilance principle: the way a medicine is initiated and titrated can influence the occurrence and severity of adverse reactions.

3. Reported Adverse Drug Reactions (ADRs)

The adverse-effect profile of tirzepatide is dominated by gastrointestinal reactions. FDA clinical-trial labeling identifies nausea, diarrhea and decreased appetite among the most common reactions, with vomiting, constipation, dyspepsia and abdominal pain or discomfort also reported. These events are generally more prominent during dose escalation and may decrease over time for many patients. Clinically important adverse reactions and warnings include:

- Severe gastrointestinal adverse reactions: Tirzepatide can cause severe gastrointestinal symptoms. The current FDA label warns about severe gastrointestinal adverse reactions and advises against use in patients with severe gastroparesis.
- Acute pancreatitis: Acute pancreatitis has been reported. Patients should be monitored for persistent or severe abdominal pain, particularly when associated with vomiting, and the medicine should be discontinued if pancreatitis is suspected.
- Gallbladder disease: Acute gallbladder disease, including cholelithiasis and cholecystitis, has been reported. Rapid weight loss can also be associated with gallbladder events, so clinical assessment is important when compatible symptoms occur.
- Hypoglycemia: The risk of hypoglycemia increases

when tirzepatide is used with insulin or an insulin secretagogue such as a sulfonylurea. The FDA recommends considering reduction of the dose of the concomitant insulin secretagogue or insulin when initiating tirzepatide. • Hypersensitivity: Serious hypersensitivity reactions, including anaphylaxis and angioedema, have been reported. Patients with a history of serious hypersensitivity to tirzepatide or its excipients should not receive the medicine. • Acute kidney injury: Kidney injury can occur, particularly when gastrointestinal adverse reactions cause dehydration or volume depletion. Monitoring renal function is important in patients at increased risk, especially when severe vomiting or diarrhea occurs. • Diabetic retinopathy complications: Rapid improvement in glycemic control can be associated with a temporary worsening of diabetic retinopathy in susceptible patients. Patients with a history of diabetic retinopathy require appropriate monitoring. • Thyroid C-cell tumor warning: FDA labeling contains a boxed warning because tirzepatide caused thyroid C-cell tumors in rats. It is unknown whether this effect occurs in humans. Tirzepatide is contraindicated in patients with a personal or family history of medullary thyroid carcinoma or in patients with Multiple Endocrine Neoplasia syndrome type 2. • Postmarketing events: FDA postmarketing information includes acute pancreatitis, including severe forms; ileus or intestinal obstruction; severe constipation including fecal impaction; anaphylaxis and angioedema; pulmonary aspiration during anesthesia or deep sedation; and acute renal failure or worsening of chronic renal failure. Postmarketing reports are voluntary, so their frequency and causal relationship cannot always be reliably estimated.

4. Risk-Minimization and ADR Management

Risk minimization should combine appropriate patient selection, gradual dose escalation, patient education, monitoring and prompt clinical action when warning symptoms occur. 1. Appropriate patient selection: Avoid tirzepatide in patients with a personal or family history of medullary thyroid carcinoma or MEN 2 and in patients with serious hypersensitivity to the drug or its components. Use caution and clinical judgment in patients with severe gastrointestinal disease or relevant comorbidities. 2. Gradual dose escalation: Starting at the recommended low dose and increasing gradually can reduce gastrointestinal adverse reactions. Patients should follow the prescribed escalation schedule rather than increasing the dose independently. 3. Gastrointestinal monitoring: Patients should be counseled about nausea, vomiting, diarrhea, constipation and abdominal discomfort. Persistent or severe symptoms require clinical evaluation because dehydration can lead to renal complications. 4. Pancreatitis vigilance: Patients should be instructed to seek medical assessment for persistent severe abdominal pain, especially if pain radiates to the back or is accompanied by vomiting. Suspected pancreatitis requires prompt evaluation and discontinuation according to the prescribing information. 5. Hypoglycemia prevention: When tirzepatide is used with insulin or insulin secretagogues, clinicians should consider lowering the dose of the concomitant agent to reduce hypoglycemia risk. Patients should be educated about symptoms and appropriate monitoring. 6. Renal monitoring: In patients who experience severe gastrointestinal reactions, particularly those with pre-existing renal impairment or other risk factors, renal function should be monitored because volume depletion can precipitate acute kidney injury. 7. Retinopathy monitoring: Patients with a history of diabetic retinopathy should receive appropriate ophthalmic monitoring, particularly when glycemic control improves rapidly. 8. Patient education and pharmacovigilance reporting: Patients should understand the purpose of treatment, common adverse effects, warning symptoms and when to contact a healthcare professional. Suspected serious or unexpected adverse drug reactions should be documented and reported through the applicable pharmacovigilance system.

5. Pharmacovigilance Perspective

Pharmacovigilance does not end when a drug receives marketing authorization. Clinical trials identify common and important risks under controlled conditions, whereas postmarketing surveillance can detect rare, delayed or population-specific safety signals. Tirzepatide illustrates this lifecycle approach: FDA labeling contains both clinical-trial adverse reactions and postmarketing reports. DrugBank's clinical safety resources include adverse effects, black-box warnings, contraindications and drug interactions, demonstrating how structured drug-safety data can support adverse-event identification and risk assessment. However, DrugBank and FDA labeling should be treated as complementary sources: DrugBank provides a structured knowledge resource, while the FDA prescribing information is the regulatory source for U.S. approved labeling and official warnings. A useful pharmacovigilance workflow for tirzepatide therefore includes recognizing the event, assessing its seriousness and temporal relationship to exposure, considering alternative causes and concomitant medicines, documenting the case,

managing the patient appropriately, and reporting eligible suspected ADRs through the relevant national reporting system. Signal detection from accumulated reports can then contribute to updates in labeling, clinical guidance and risk-minimization strategies.

6. Conclusion

Tirzepatide demonstrates how pharmacovigilance integrates pharmacology, clinical monitoring and regulatory risk management. Its dual GIP/GLP-1 receptor agonist mechanism provides therapeutic metabolic effects, while gastrointestinal reactions are the most prominent tolerability issue. More serious or clinically important risks include pancreatitis, gallbladder disease, hypoglycemia with certain concomitant medicines, hypersensitivity, renal injury related to volume depletion, diabetic retinopathy complications and the boxed warning concerning thyroid C-cell tumors. The most effective risk-reduction approach is proactive: appropriate patient selection, gradual dose escalation, patient counseling, monitoring of vulnerable patients and prompt response to warning symptoms. Continued postmarketing surveillance remains essential because rare events may become more apparent only after wider clinical use. This case study highlights the central role of pharmacovigilance in maintaining a favorable benefit-risk balance throughout a medicine's lifecycle.

References

1. U.S. Food and Drug Administration. MOUNJARO (tirzepatide) Prescribing Information. FDA. Revised 2025.
2. U.S. Food and Drug Administration. ZEPBOUND (tirzepatide) Prescribing Information. FDA. Revised January 2026.
3. DrugBank Online. Tirzepatide (DB15171) and DrugBank Clinical Insights / Adverse Effects resources. DrugBank.
4. Coskun T, Sloop KW, Loghin C, et al. LY3298176, a novel dual GIP and GLP-1 receptor agonist for the treatment of type 2 diabetes mellitus: From discovery to clinical proof of concept. *Molecular Metabolism*. 2018;18:3-14. doi:10.1016/j.molmet.2018.09.009.
5. Rosenstock J, Wysham C, Frías JP, et al. Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1). *The Lancet*. 2021;398:143-155. doi:10.1016/S0140-6736(21)01324-6.