

A Predictive electrode attribution
Proximal
Distal
F1
F2
F3
F4
F5
F6
F7
F8
F9
F10
F11
F12
F13
F14
F15
F16
1
2
3
4
5
6
7
8
Circumferential position
Magnitude (%)
8
6
B Which gestures lose performance?
G7: Index-middle Extension
G11: Wrist Extension
G3: Thumb-little Opposition
G5: Thumb-index Extension
G15: Hand Open
G4: Thumb-index Opposition
G13: Forearm Supination
G8: Little Finger Extension
G12: Wrist Flexion
G6: Thumb-little Extension
G1: Lateral Prehension
G16: Hand Close
G10: Thumb Finger Extension
G2: Thumb Adduction
G9: Index Finger Extension
G14: Forearm Pronation
−0.06
−0.05
−0.04
−0.03
−0.02
−0.01
0.00
0.01
SHAP12 - full16 gesture F1
C Deletion with retraining
Mean Macro-F1 (lighter model)
0.50
0.45
0.40
0.35
0.30
remove top
remove random
remove bottom
2
6
10
14
Electrodes remaining after deletion
D Participant-level penalty
Penalized - unpenalized F1
0.006
0.004
0.002
0.000
lambda=0.25
lambda=0.5
lambda=1
4
6
8
12
Retained electrodes