

Copyright Warning

COMMONWEALTH OF AUSTRALIA
Copyright Regulation 1969

WARNING

This material has been copied and communicated to you by or on behalf
of **Curtin University of Technology** pursuant to Part VB of the
Copyright Act 1968 (the Act)

The material in this communication may be subject to copyright under the
Act. Any further copying or communication of this material by you
may be the subject of copyright protection under the Act.

Do not remove this notice

Theoretical Foundations of Computer Science

Lecture 3b

The Pumping Lemma Context-Free Grammars

Topics

- Non-regular expressions
 - Pumping lemma

Unit Learning Outcome

- Synthesize FA, PDA, CFG, and TMs with specific properties, and to relate and convert from one form to another.

Assessment Criteria

- Use the pumping lemma to prove a language to be a non-regular language.

Non-regular languages

- Limitations of finite automata
 - certain languages cannot be recognized by any finite automaton
- Example: Language $B = \{0^n 1^n \mid n \geq 0\}$
 - Claim:
 - a machine recognizing B need to remember how many 0s have been seen so far as it reads input
 - an unlimited number of states needed for this
 - Thus non-regular

Failure of Intuition

- Another non-regular language:
 - $C = \{w \mid w \text{ has an equal number of 0s and 1s}\}$
- However, the following language is regular:
 - $D = \{w \mid w \text{ has an equal number of occurrences of 01 and 10 as sub-strings}\}$
- Thus claims that a language is regular (or not) must be proven.

Pumping lemma

- All regular languages have a special property
 - If we can show that a language does not have this property, then it is guaranteed to be non-regular
- The property is that all strings in the language can be “pumped” if they are longer than a certain special value, called the *pumping length*.
 - Each such string contains a section that can be repeated any number of times and the resulting strings will remain in the given language.

Pumping lemma

- If A is a regular language, there is a pumping length p such that if s is any string in A of at least length p , then s may be divided into three pieces, $s = xyz$, satisfying the following conditions:
 1. for each $i \geq 0$, $xy^iz \in A$,
 2. $|y| > 0$, and
 3. $|xy| \leq p$.
- Note on notation:
 - y^i means i copies of y concatenated together ($y^0 = \varepsilon$)
 - either x or z may be ε , but y cannot be ε

Using the Lemma

- To show that a language C is not regular,
 - first assume it is regular
 - That is: there is a pumping length p for C
 - WE DO NOT KNOW A VALUE FOR p
 - Select a pattern typically at start of string
 - EXCLUDE something that is required
 - Concatenate start string p times.
 - Now complete the string so it is part of language

Using the Lemma

- To show that a language C is not regular,
 - We have a string of length greater than p
 - Such that strings of length p or greater can be pumped
 - Aim to show our suitably chosen string s in C in fact cannot be pumped, thus contradicting the assumption that C is regular.
 - Pumping means we can add (subtract) portion of start.
 - Our tail is unaltered – it contains essential element
 - Some rule connecting start with tail should break

Example

- Example: Language $B = \{0^n 1^n \mid n \geq 0\}$
 - Assume regular \Rightarrow exist p
 - Start consists of 0, but 1's are important
 - Let 0 be base starting pattern
 - Use p by raising starting pattern to power p – 0^p
 - Complete an element of the language: $0^p 1^p$
 - Now have trial string
 - Pumping
 - y of the lemma must be sequence of 0
 - Pumping means adding 0
 - But no 1's added
 - So new string will not have equal 0's and 1's

Proof

Let $s = 0^p 1^p$

By 2) xy must be within the p letters of s
But first p letters of s are all 0's.

By PL:

$$1) s = xyz$$

$$2) |xy| \leq p$$

$$3) |y| > 0$$

$$4) xy^kz \in B$$

So from 1), there is i, j such that

$$x = 0^i \quad 0 \leq i < p \text{ by 2) 3)}$$

$$y = 0^j \quad 1 \leq j \leq p-1 \text{ by 2) 3)}$$

$$z = 0^{(p-i-j)} 1^p$$

$$xy^2z = 0^i 0^{2j} 0^{(p-i-j)} 1^p = 0^{(p+j)} 1^p$$

As (4) this is in \mathbf{B} , so $p+j = p$ that is $j=0$

BUT (3) says $j \geq 1$, CONTRADICTION

We can choose k :

Usually 0 or 2.

Say 2.

Hence \mathbf{B} is not regular.

- Non-regular expressions
 - Pumping lemma
 - Know the theorem
 - <ULO> Use the theorem