

Assignment 4

Tanay Yadav - AI20BTECH11026

Download the python codes from:

https://github.com/tanayyadav28/EE3900-Assignments/blob/main/Assignment_4/code/Assignment_4.py

Download the latex-tikz codes from:

https://github.com/tanayyadav28/EE3900-Assignments/blob/main/Assignment_4/Assignment_4.tex

The lines L_1 and L_2 are intersecting if,

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} + \lambda_2 \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \quad (2.0.5)$$

$$\lambda_1 \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} - \lambda_2 \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad (2.0.6)$$

$$\begin{pmatrix} 1 & 2 \\ -3 & 3 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ -\lambda_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \quad (2.0.7)$$

1 PROBLEM

[Linear Forms 2.23] Find the shortest distance between the lines:

$$L_1 : \mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda_1 \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} \quad (1.0.1)$$

$$L_2 : \mathbf{x} = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} + \lambda_2 \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \quad (1.0.2)$$

2 SOLUTION

Given,

$$\mathbf{A}_1 = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad (2.0.1)$$

$$\mathbf{m}_1 = \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{A}_2 = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} \quad (2.0.3)$$

$$\mathbf{m}_2 = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \quad (2.0.4)$$

Row reducing the augmented matrix,

$$\begin{pmatrix} 1 & 2 & 3 \\ -3 & 3 & 3 \\ 2 & 1 & 3 \end{pmatrix} \xleftrightarrow{R_2=R_2+3R_1} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 9 & 12 \\ 1 & 2 & 3 \end{pmatrix} \quad (2.0.8)$$

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 9 & 12 \\ 2 & 1 & 3 \end{pmatrix} \xleftrightarrow{R_3=R_3-2R_1} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 9 & 12 \\ 0 & -3 & -3 \end{pmatrix} \quad (2.0.9)$$

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 9 & 12 \\ 0 & -3 & -3 \end{pmatrix} \xleftrightarrow{R_3=R_3+\frac{R_2}{3}} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 9 & 12 \\ 0 & 0 & 1 \end{pmatrix} \quad (2.0.10)$$

The above matrix has $rank = 3$. Hence, the given lines are skew-lines.

The normal to both the lines is:

$$\mathbf{n} = \mathbf{m}_1 \times \mathbf{m}_2 \quad (2.0.11)$$

The equation of the plane is then obtained as:

$$\mathbf{n}^T \mathbf{x} = \mathbf{n}^T \mathbf{A}_2 \quad (2.0.12)$$

The distance of the above line from \mathbf{A}_2 is obtained as:

$$\frac{\mathbf{n}^T (\mathbf{A}_2 - \mathbf{A}_1)}{\|\mathbf{n}\|} = \frac{(\mathbf{A}_2 - \mathbf{A}_1)^T (\mathbf{m}_1 \times \mathbf{m}_2)}{\|\mathbf{m}_1 \times \mathbf{m}_2\|} \quad (2.0.13)$$

$$= \frac{\begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}^T \begin{pmatrix} -9 \\ 3 \\ 9 \end{pmatrix}}{\sqrt{\begin{pmatrix} -9 \\ 3 \\ 9 \end{pmatrix}^T \begin{pmatrix} -9 \\ 3 \\ 9 \end{pmatrix}}} \quad (2.0.14)$$

$$= \frac{9}{\sqrt{171}} \quad (2.0.15)$$

$$= \frac{3}{\sqrt{19}} \quad (2.0.16)$$

Hence, the shortest distance between the given lines is $\frac{3}{\sqrt{19}}$.




Fig. 0: Plot from Python Code.